



UNIVERSITÀ  
DEGLI STUDI  
FIRENZE

## FLORE

# Repository istituzionale dell'Università degli Studi di Firenze

### **Liver resection with thrombectomy as a treatment of hepatocellular carcinoma with major vascular invasion: results from a retrospective**

Questa è la versione Preprint (Submitted version) della seguente pubblicazione:

*Original Citation:*

Liver resection with thrombectomy as a treatment of hepatocellular carcinoma with major vascular invasion: results from a retrospective multicentric study / Pesi, B; Ferrero, A; Grazi, GI; Cescon, M; Russolillo, N; Leo, F; Boni, L; Pinna, Ad; Capussotti, L; Batignani, G. - In: THE AMERICAN JOURNAL OF SURGERY. - ISSN 0002-9610. - ELETTRONICO. - (2015), pp. 35-44.

*Availability:*

The webpage <https://hdl.handle.net/2158/1014519> of the repository was last updated on 2015-12-21T10:01:41Z

*Terms of use:*

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

*Publisher copyright claim:*

Conformità alle politiche dell'editore / Compliance to publisher's policies

Questa versione della pubblicazione è conforme a quanto richiesto dalle politiche dell'editore in materia di copyright.

This version of the publication conforms to the publisher's copyright policies.

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

## **Torquetenovirus (TTV) load [is associated with mortality](#) in Italian elderly subjects**

Robertina Giacconi<sup>1\*</sup>°, Fabrizio Maggi<sup>2°</sup>, Lisa Macera<sup>2</sup>, Mauro Pistello<sup>2</sup>, Mauro Provinciali<sup>1</sup>, Simone Giannecchini<sup>3</sup>, Francesco Martelli<sup>3</sup>, Pietro Giorgio Spezia<sup>2</sup>, Erminia Mariani<sup>4,5</sup>, [Roberta Galeazzi<sup>6</sup>](#), [Laura Costarelli<sup>6</sup>](#), Lorenzo Iovino<sup>7</sup>, Sara Galimberti<sup>7</sup>, Lorenzo Nisi<sup>1</sup>, Francesco Piacenza<sup>1</sup>, Marco Malavolta<sup>1</sup>

<sup>1</sup> Advanced Technology Center for Aging Research, Scientific and Technological Pole, Italian National Institute of Health and Science on Aging (INRCA), Ancona, Italy.

<sup>2</sup> Retrovirus Center and Virology Section, Department of Translational Research, University of Pisa, Pisa, Italy.

<sup>3</sup> Department of Experimental and Clinical Medicine, University of Florence, Florence, Italy.

<sup>4</sup> Laboratory of Immunorheumatology and Tissue Regeneration, Rizzoli Orthopaedic Institute, Bologna, Italy.

<sup>5</sup> Department of Medical and Surgical Sciences, University of Bologna, Bologna, Italy.

<sup>6</sup> [Clinical Laboratory & Molecular Diagnostics, INRCA-IRCCS, Ancona, Italy](#)

<sup>7</sup> Section of Hematology, Department of Clinical and Experimental Medicine, University of Pisa, Pisa, Italy.

° Both authors contributed equally to this manuscript.

\*Address correspondence to this author at the Advanced Technology Center for Aging Research, IRCCS-INRCA, Via Birarelli 8, 60121, Ancona, Italy; Tel: +39-071-8004213; Fax +39-071-206791; E-mail [r.giacconi@inrca.it](mailto:r.giacconi@inrca.it)

## Abstract

An age-related dysregulation of immune response, known as immunosenescence, contributes to increased susceptibility to infections, frailty and high risk of mortality in the elderly. Torquetenovirus (TTV), a circular, single-stranded DNA virus, is highly prevalent in the general population and it may persist in the organism, [also in association with other viruses such as Cytomegalovirus \(CMV\)](#), causing chronic viremia. The relationship that TTV establishes with the immune system of infected hosts is not clear. It is known that TTV encodes microRNAs (miRNAs) that might contribute to immune evasion and that the highest viral loads are found in peripheral blood cells. Moreover, it is suspected that TTV infection lead to increased production of inflammatory mediators, thus playing a role in immunosenescence. We investigated the association of TTV load and miRNAs expression with inflammatory and immune markers and the influence of TTV load on mortality within a cohort of 379 elderly subjects who were followed up for 3 years. TTV DNA load in polymorphonuclear leukocytes was [slightly](#) positively correlated with age and negatively associated with serum albumin levels and NK cell activity. A [marginal](#) positive correlation between TTV DNA load, monocytes and IL-8 plasma levels was found in females and males respectively. [TTV DNA copies  \$\geq 4.0\$  log](#) represented a strong predictor of mortality (Hazard ratio = 4.78, 95% CI: 1.70-13.44, after adjusting for age, sex and the main predictors of mortality rate) [and this association remained significant even after the CMV IgG antibody titer was included in the model \(HR = 9.83; 95% CI: 2.48-38.97; N=343 subjects\)](#). Moreover, multiple linear regression model showed that TTV miRNA-t3b of genogroup 3 was inversely associated with triglycerides, monocytes and C-reactive protein, and directly associated with IL6. Overall these findings suggest a role of TTV in immunesenescence and in the prediction of all-cause mortality risk in Italian elderly subjects. Further studies are needed to fully understand the pathogenic mechanisms of TTV infection during aging.

Keywords: TTV, aging, immunesenescence, mortality, NK cell activity

## 1. Introduction

Aging is associated with a variety of physiological changes, including a decline in immune function known as “immunosenescence.” This condition is characterized by age-related dysfunction of the immune system, affecting both innate and adaptive immunity, causing a reduced response to pathogens and to vaccines with greater risk of frailty and mortality in elderly subjects (Pawełec et al., 2014). NK cells are involved in the elimination of viral-infected cells as well as in the innate and adaptive immunological regulation through the production of cytokines and chemokines (Camous et al., 2012). During aging, NK cell dysfunction may contribute to the imbalance between inflammatory and anti-inflammatory signals contributing to a state defined as “inflammaging” (Ventura et al., 2017). The measure of Cu/Zn ratio may represent a sensitive biomarker of systemic inflammation and “inflammaging” status, because it is strongly related to circulating inflammatory biomarkers in old population (Malavolta et al., 2010; Giacconi et al., 2017).

Several herpesviruses, such as cytomegalovirus (CMV), cause chronic antigen stimulation, evade NK cell recognition (Charpak-Amikam et al., 2017) and have been associated with immunosenescence and telomere length attrition (Nikolich-Zugich and van Lier, 2017; Stowe et al., 2012). Some evidence shows that CMV seropositivity, associated with higher C-reactive protein, increased the risk for all-cause mortality and cardiovascular diseases (CVD)-related mortality (Simanek et al., 2011), while the higher anti-CMV titres increased the risk of death in elderly over 80 years (Matheï et al., 2015). Currently there are no simple biomarkers for the overall level of immune function decline conferred on a patient with the age. Thus, new biomarkers are needed to accurately assess the degree of immunosenescence in elderly people, and to help physicians in the management of aged patients. Recently, Torquetenovirus (TTV) viremia has been proposed as a potentially simple and general measure of immune system function of infected host. TTV, originally isolated from a patient suffering from posttransfusion hepatitis (Nishizawa et al., 1997), is the prototype of a vast group of viruses with a small, circular single-stranded DNA virus belonging to Anelloviridae family and it is the most representative and abundant virus of the human virome (Focosi et al., 2016).

Since TTV was first detected in blood samples, it was referred to as transfusion-transmitted (TT) virus (Nishizawa et al., 1997), but other ways of TTV transmission including parenteral, oro-fecal, respiratory, sexual and vertical transmission have been reported (Haloschan et al., 2014). Due to its high prevalence (70-90 %) in the general population, a causative role

of TTV in human diseases remains uncertain. However, a higher prevalence of TTV was found to be associated with several pathologies such as viral hepatitis, respiratory diseases, sepsis, enteritis, and cancer (Rosa et al., 2017; Feyzioğlu et al., 2014; Walton et al., 2014; Brassard et al., 2016; Szládek et al., 2005). In addition, certain TTV species/genotypes were suggested to be more pathogenic than others in patients with liver disease (Tokita et al., 2001). More recently, it has been shown that TTV encodes microRNAs (miRNAs) (Kincaid et al., 2013; Vignolini et al., 2016) and that some of these miRNAs may interfere with cytokine signalling promoting virus persistence and immune evasion (Kincaid et al., 2013).

However, the possible involvement of TTV and its miRNAs in immunesenescence or increased risk of mortality in elderly people is currently unknown. The purpose of this study was to investigate the association of TTV load and miRNAs with age, inflammatory and immune markers as well as the risk for all-cause mortality in elderly population after three-year follow-up.

## **2. Materials and methods**

### **2.1. Subject population**

The study included n. 379 noninstitutionalised elderly subjects (age range 60–105 years) born and living in east central Italy (the Marche Region) recruited in the years 2007-2008.

The exclusion criteria included autoimmune diseases, neurodegenerative diseases, infections, cancer, Crohn's disease and acrodermatitis enteropathica, kidney disease, liver disease, sickle cell anaemia, chronic skin ulcerations and endocrine disorders. Subjects also had to be free of anticonvulsants, anti-depressives drug and antibiotics. A standardised questionnaire was administered to collect data regarding the medical history, anti-inflammatory medication (acetylsalicylic acid, corticosteroids and non-steroidal anti-inflammatory drugs (NSAIDs), statins, ACEinhibitors and fibrates) and personal habits. To ensure validity, the family doctor was requested to check the completed questionnaire.

Subjects with CVD (angina, arrhythmia, compensated heart failure) were included in the study if these conditions did not compromise the individual's independence. The healthy status was evaluated by a specific questionnaire on health and morbidity planned for the study and filled in by the general practitioner after medical examination and analysis of the clinical record. Hypertension was defined as seated systolic blood pressure  $\geq 160$  mmHg, diastolic pressure  $\geq 95$  mmHg or both, or self-reported hypertension and if the subject was already taking anti-hypertensive medication. The cohort was followed up for 3 years using

mortality records held by the General Register Office in the subject's place of residence. Death certificates were examined by a medical doctor to determine the main cause of death. All subjects signed an informed consent, which was approved by the Ethical Committee of INRCA. Biological samples were collected in the INRCA biobank and made anonymous.

## **2.2. Laboratory determinations**

Venous peripheral blood samples, collected after an overnight fast, underwent basal biochemical laboratory determinations. Serum, plasma, and polymorphonuclear leukocytes, were separated, aliquoted and stored frozen at  $-80^{\circ}\text{C}$  in the Biological Bank of INRCA until analysis. The blood haematocrit and haemoglobin concentration, were determined by automated blood analysers. Serum total cholesterol, HDL-cholesterol, triglycerides, were measured using an automatic analyser under conventional enzymatic methodology. Glucose, creatinine and albumin levels were determined using standard laboratory methods. C-reactive protein (CRP) levels were detected by CardioPhase hsCRP assay (Dade Behring Inc Deerfield, IL). [Immunoglobulin G levels of CMV-specific antibodies were determined in plasma samples by the Liaison assay \(LIAISON® CMV IgG II; Diasorin\) according to the manufacturer's specifications and using the Liaison analyzer \(DiaSorin, Italy\).](#)

## **2.3. Polymorphonuclear leukocytes (PMNLs) sample preparation**

PMNLs were separated by sedimentation of 3–7 mL of heparinized blood sample in a 6% dextran solution (5 vol. blood/1 vol. dextran solution). Leukocytes were centrifuged at 2500 rev/min for 10 min at room temperature, and the resulting pellet was washed in 0.8%  $\text{NH}_4\text{Cl}$  for 1–2 min to eliminate red blood cell contamination. PMNLs were then centrifuged and resuspended in 1 mL of phosphate-buffered saline (PBS), pH 7.4. PMNLs aliquots were stored at  $-80^{\circ}\text{C}$  until DNA extraction.

## **2.4. TTV DNA detection and quantification**

[TTV load was assessed in PMNLs where it has been reported that the viral load is highest \(Takahashi et al., 2002\).](#) Viral DNA was extracted from blood PMNLs using The Wizard® Genomic DNA Purification Kit (Promega Italia s.r.l., Milan, Italy), as specified by the manufacturer. Presence and load of TTV DNA were determined in a single step TaqMan PCR assay as described elsewhere (Maggi et al., 2005). This assay uses primers designed on a highly conserved segment of the untranslated region of the viral genome and has therefore the capacity to detect all the species in which TTV is actually classified. TTV loads were expressed as the number of viral DNA copies per  $\mu\text{g}$  of genomic DNA extracted by PMNLs. The lower limit of detection was [10](#) copies of TTV DNA per  $\mu\text{g}$  genomic DNA. The

procedures used to quantitate the copy numbers and assess specificity, sensitivity, intra- and inter-assay precision, and reproducibility have been previously described (Maggi et al., 2005). In a limited group of patients, viral DNA was also extracted from 200 µl plasma samples using QIAamp DNA minikit® (Qiagen, Chatsworth, CA, USA), and amplified as described above. The lower limit of sensitivity was 10 TTV genomes per ml of plasma.

## **2.5. TTV miRNAs detection and quantification**

Total RNA was isolated from plasma exosomes obtained from 250 µl plasma processed with Plasma/Serum Circulating and Exosomal RNA Purification kit (Norgen Biotek Corp., ON, Canada) and following the manufacturer's protocol. miRNA expression was analysed and quantified with a commercial quantitative RT-PCR miRNA assay (Life Technologies, Foster City, CA) and using primers targeting the miRNA mature region previously described (Vignolini et al., 2016). In particular, TTV miRNAs t1a of genogroup 1, t3b of genogroup 3, and tth8 of genogroup 5 were selected according to predictive computational analysis (Kincaid et al., 2013) and because found to be the most prevalent in plasma samples of patients previously studied (Vignolini et al. 2016). Each reaction was carried out in triplicate with 10 ng RNA and included negative controls (no template) and  $10^1$ – $10^6$  copies of positive control (synthetic oligonucleotide template). The lower limit of detection was 10 copies of TTV miRNA per ng RNA. As assessed by several reactions performed in preliminary experiments and various conditions, the assay proved specific and reproducible. Preliminarily each TTV miRNA oligonucleotide standard was tested without amplification of unrelated target and with <0.5Ct value interassay variation.

## **2.6. Zinc and copper plasma level determinations**

The plasma zinc (Zn) and copper (Cu), were determined by a Thermo XII Series ICP-MS (Thermo Electron Corporation, Waltham, MA, USA) by adapting methods used for the measurement of trace elements in human plasma with slight modifications (Malavolta et al., 2010). Plasma samples were diluted 1:10, with a diluent containing 0.1% triton and 0.15% HNO<sub>3</sub>, to ensure that trace elements are maintained in solution and to favour the washout of these elements between samples. External calibration solutions containing Zn, and Cu (blank to 2000 ppb) were prepared by serial dilution of a parent multi-element solution (1000 ppm for Zn and Cu) (VHG Labs, Manchester, USA), using the same diluent used for the samples. Rhodium (Rh), at 200 ng/ml was used as internal standard. Data were acquired for <sup>66</sup>Zn, <sup>65</sup>Cu.

Quality of the analysis was assured by assessment of “quality standard samples” (SERONORM™ TRACE ELEMENT SERUM, Sero AS, Billingstad, Norway). Zn and Cu

levels of the quality standard samples were within 10% of the certified levels, as previously reported (Malavolta et al., 2010). The instrument was operated with a Peltier cooled impact bead spray chamber, single piece quartz torch (1.5 mm i.d. injector) together with Xi interface cones and a Cetac-ASX 100 autosampler (CETAC Technologies, Omaha, NE). A Burgener Trace nebulizer was used as this device does not block during aspiration of clinical samples. The instrument was operated in standard mode (non-CCT), using 1400 WRF power, 1.10 L min<sup>-1</sup> nebulizer gas flow, 0.70 L min<sup>-1</sup> auxiliary gas flow, 13.0 L min<sup>-1</sup> cool gas flow, 70 ms dwell time, 30 s sample uptake and 35 s wash time (2 repeats per sample).

## **2.7. NK cell lytic activity**

K562 tumor target cells ( $1 \times 10^6$ ) were labelled with 15  $\mu$ M calcein-AM (Molecular Probe, Eugene, OR) for 30 min at 37 °C and washed 2 times with cold complete medium. Labelled target cells ( $5 \times 10^3/50 \mu$ l) and varying numbers of cryopreserved and thawed effector cells (from  $5 \times 10^6$  to  $5 \times 10^3$  PBMC/100  $\mu$ l) were incubated in triplicate in V-bottom 96-well plates (Effector/Target E/T ratios from 50/1 to 1/1). After incubation at 37 °C in 5% CO<sub>2</sub> for 4 h, 75  $\mu$ l of each supernatant were harvested, transferred into new plates and measured by a dual-scanning microplate spectrofluorimeter (Spectramax Gemini – Molecular Devices, Sunnyvale, CA) (Neri et al., 2001).

Data were expressed as arbitrary fluorescent units (AFU). Percent lytic activity was calculated according to the formula  $[(\text{test AFU} - \text{spontaneous AFU})/(\text{maximum AFU} - \text{spontaneous AFU})] \times 100$ . Spontaneous AFU represents Calcein AM released from target cells in medium alone and maximum AFU is Calcein AM released from target cells lysed in medium plus 2% Triton X-100, each measured in at least six replicate wells.

## **2.8. Multiple immunoassay for chemokine and cytokine plasma assessment**

Fourfold diluted plasma samples were assayed in duplicate using commercially available multiplex bead-based immunoassay kits. IL-6 and IL-8 concentrations were simultaneously evaluated using multiplex reagent kits (Bio-Rad Laboratories, USA). Values presenting a coefficient of variation beyond 10% were discarded before the final data analysis. Data were analysed using the Bio-Plex Manager software version 6.0 (Bio-Rad Laboratories, USA). Values were expressed as pg/ml.

## **2.9. Telomere length analysis**

Telomere length was determined [in peripheral blood mononuclear cells \(PBMC\)](#) by flow-FISH methodology by using the Telomere PNA kit/FITC from Dako (Glostrup, Denmark) [according to the manufacturer's instructions](#). Samples were then analyzed by flow cytometry

(Coulter Epics XL, Coulter, USA) and relative telomere length was determined by comparing isolated test cells with a control cell line (Human T-cell leukaemia 1301 cell line).

### 3. Statistical analysis

Pearson Correlation between TTV DNA loads (log-transformed data) and age was investigated stratifying by gender. A linear regression model for determining the associations between TTV DNA loads and albumin, NK cell activity, monocytes percentage and IL-8 plasma levels was performed. Log transformation of the variables was carried out if they were not normally distributed as assessed by the Kolmogorov-Smirnov test. The relationship between TTV DNA loads and all-cause and CV mortality was investigated by Kaplan–Meier analysis and by Cox regression analyses. The predictive model included age, sex, Cu/Zn, CRP, triglycerides, total cholesterol, albumin and glucose serum levels as potential covariates.

The automatic linear modelling procedures implemented in SPSS were carried out to explore the main predictors of TTV DNA loads or miRNAs following a procedure previously described (Giacconi et al., 2017). The variables inserted were: age, gender, TTV DNA loads or miRNAs, C-reactive protein, albumin, Cu/Zn ratio, erythrocytes count, hemoglobin, leucocyte count, neutrophils, monocytes, lymphocytes percentages, BMI, lipid profile, NK cell activity, IL-6 and IL8. Main predictors of TTV DNA loads or miRNA were also confirmed using linear regression models. All statistical analyses were performed with SPSS Statistics version 22.0.0 (IBM, IL, USA).

## 4. RESULTS

### 4.1. TTV DNA detection and clinical and biochemical parameters of elderly subjects

Of 379 patients, 273 (72%) showed TTV DNA positivity in PMNLs. Table 1 outlines the baseline characteristics of the study population, grouped by TTV detection. **TTV-positive subjects had a very slightly higher mean age than the TTV-negative group.** No significant differences were observed for gender distribution, BMI, white blood cell count, erythrocytes number, haemoglobin, CRP, creatinine, albumin, fasting glucose and lipid serum levels. Conversely, the percentage of lymphocytes was increased, while neutrophils decreased in TTV positive subjects as compared to the negative ones (Table1,  $p < 0.01$ ). The prevalence of hypertension, cardiovascular diseases and type 2 diabetes was similar between the two classes of subjects, **while an association between CMV serostatus and TTV DNA positivity was found and 76.4% of subjects were positive for both viruses (Table1,  $p < 0.001$ ).**

#### 4.2. TTV DNA load association with age, albumin, leukocytes, NK cell activity, inflammatory markers and CMV IgG antibody titer

TTV levels were measured in PMNLs samples from the 273 subjects found positive for viral DNA presence. Overall, mean TTV load was 3.0 log copies per  $\mu\text{g}$  of total DNA extracted but, among individual subjects, it varied widely, ranging from 1.1 to 5.4 log DNA copies (confidence interval: 2.9 – 3.1 log TTV DNA copies per  $\mu\text{g}$  of total DNA extracted). In 65/273 subjects (24%), values  $>4$  log DNA copies/ $\mu\text{g}$  of total DNA were measured, which represents a very high virus load, considering that the mean TTV levels measured in healthy blood donors range from 1 to 3 log TTV copies per ml of plasma or serum (Maggi et al., unpublished results).

TTV DNA loads were slightly, but significantly positively associated with age ( $R=0.267$ ,  $p<0.001$ ), independent of the sex of the subject (Fig. 1, females:  $R=0.283$ ,  $p<0.001$ ; males:  $R=0.264$ ,  $p<0.001$ ). Linear regression correcting for age evidenced a slight negative association among TTV DNA loads, albumin and NK cell activity (Fig. 2A:  $R = -0.244$ ,  $p<0.0001$ ; Fig. 2B:  $R = -0.269$ ,  $p<0.05$ ). A slight positive correlation was found among TTV DNA loads and monocytes in females (Fig. 3A:  $R = 0.206$ ,  $p<0.01$ ) and IL8 in males (Fig. 3B:  $R = 0.178$ ,  $p<0.05$ ). No significant association among TTV DNA loads and CMV IgG antibody titer or other inflammatory markers (CRP, Cu/Zn ratio and IL-6) (Fig. 2S, Fig. 3S, Fig. 4S, Fig. 5S, Supplementary material) or between TTV DNA load and other leukocytes populations (neutrophils and lymphocytes) was observed (Fig. 6S, Supplementary materials). Moreover, we observed a significant positive correlation between TTV DNA loads in PMNL and in plasma samples (log-transformed data) in a subgroup of 50 elderly subjects ( $N=50$ ; 36 females and 14 males mean age:  $80.4 \pm 9.7$ ) ( $R=0.571$ ,  $P<0.0001$ ). Of these fifty subjects, six (12%) had TTV negative plasma samples (Fig. 7S, Supplementary materials).

#### 4.3. Survival analysis

Based on the Kaplan-Meier method, survival at three years differed significantly by TTV load (log-rank test:  $p < 0.0001$ ; Fig. 4). The proportion of patients that died after three years of follow-up was estimated as 21.9% for patients with TTV DNA copies  $\geq 4.0$  log and 5.4% for patients with TTV copies  $< 4.0$  log.

The Cox proportional hazards model revealed that, after controlling for age, sex, Cu/Zn, CRP, triglycerides, total cholesterol, albumin and glucose serum levels, the three year

mortality risk of elderly subjects increased significantly when TTV DNA copies were  $\geq 4.0$  log. (HR = 4.78; 95% CI: 1.70-13.44). Other factors independently associated with three year mortality were: age (HR = 1.07; 95% CI: 1.01-1.15), Cu/Zn (HR = 4.94; 95% CI: 1.62-15.07), CRP (HR = 0.69; 95% CI: 0.56-0.87) and serum albumin (HR = 0.09; 95% CI: 0.02-0.41) (Table 2). Interestingly, the Cox proportional hazards analysis showed a stronger association between TTV DNA copies  $\geq 4.0$  log and three year mortality risk when CMV IgG antibody titer was included in the model (HR = 9.83; 95% CI: 2.48-38.97; N=343 subjects) (Table 1S, Supplementary material). Moreover, a slight association with three year mortality risk was also confirmed using the continuous variable TTV DNA levels (log-transformed data) in the Cox proportional hazards regression, correcting for age, sex, Cu/Zn, CRP, triglycerides, total cholesterol, albumin and glucose serum levels (HR = 1.62; 95% CI: 1.05-2.51) (Table 2S, Supplementary material).

#### 4.4. TTV miRNA detection and quantification.

TTV miRNAs were measured in plasma exosomes, microvesicles released by the cells to transfer signalling competent proteins and functional microRNAs to other cells playing a role in cell-cell communication and in the modulation of immune response (Schorey et al., 2015) also during virus infection, as even suggested for miRNA-tth8 (Vignolini et al., 2016; Kincaid et al., 2013). Plasma exosome samples of 103 elderly subjects were examined for presence and loads of miRNA t1a, t3b, and tth8 belonging to TTV genogroups 1, 3, and 5, respectively. Overall, blood samples of 30 subjects (29%) contained at least one TTV miRNA, while only 3 samples revealed the simultaneous presence of two miRNAs (2%). miRNA-t3b was detected in blood of 28% of subjects at mean level of 566 copies per ng of RNA ( $\pm$  standard error (SE): 156 copies per ng of RNA), followed by miRNA-tth8 in 5% (mean level  $\pm$  SE: 720  $\pm$  279 copies per ng of RNA). No elderly subject was found to be positive for the presence of TTV miRNA-t1a (Table 3S Supplementary material).

#### 4.5. Automatic linear modelling for variables independently associated with TTV DNA loads and miRNA-t3b

According to the automatic linear modelling, we have categorized the top 5 important predictors of miRNA-t3b variations in the plasma of elderly subjects as follow: triglycerides, monocytes, CRP, IL6 and lymphocyte percentage (Fig. 5A). To confirm the independent contributions of the variables to miRNA-t3b and TTV DNA load, a multiple linear regression model was performed (Table 4S Supplementary materials). The variability in the miRNA-t3b was best explained by triglycerides, monocytes, CRP, and IL6, while TTV DNA load

variability was dependent on age, triglycerides and albumin levels (Fig. 5B, automatic linear modelling), although linear regression did not confirm triglycerides serum concentrations as predictor for TTV viremia (Table 5S Supplementary materials).

## 5. Discussion

Susceptibility to infections increases significantly with age and persistent virus infection is responsible for immune perturbations leading to changes in the T cell pool that contributes to immunosenescence (Hassouneh et al., 2016; Di Benedetto et al., 2015).

Indeed, subclinical infections with viruses of the herpesviridae family, particularly CMV, are very common during ageing and have been associated with accelerated immunosenescence, telomere length shortening, elevated pro-inflammatory response, increased morbidity and mortality (Pawelec, 2014; Dowd et al., 2017; Westman et al., 2014; Simanek et al., 2011). Moreover, persistent CMV infection has been also linked to increased TTV DNA loads (Haloschan et al., 2014). TTV is an opportunistic pathogen that is often associated with co-infections, idiopathic pulmonary fibrosis, sepsis, cancer (Rosa et al., 2017; Walton et al., 2014; Szládek et al., 2005), as well as with COPD severity (Feyzioğlu et al., 2014). It has been suggested that TTV plasma DNA level reflects the balance between a high daily virion generation rate and a host's clearance rate of over 90 % per day (Maggi et al. 2001) and, thus, indirectly reflects the antiviral immune response against TTV.

According to previous evidence, we observed a positive correlation between age and TTV viremia, suggesting a relationship between degree of immunosenescence and virus replication (Haloschan et al., 2014). These findings indicate a decreased antiviral immune response against chronic TTV infection with advancing age, supporting previous studies, showing an impaired immune response in the elderly (Pawelec et al., 2014). In fact, TTV genome replication seems to be controlled by the immune system and higher viral titers were found in individuals with immunodeficiencies, such as AIDS (Shibayama et al., 2001) or immunosuppression condition (Focosi et al., 2015). Although the significance of TTV load on increased risk of morbidity and mortality is still unclear, in the present study we show, for the first time, that TTV load in PMNLs is independently associated with all-cause mortality, after controlling for the main confounders related to death (Malavolta et al., 2010; Barger et al., 2016) including CMV IgG antibody titer (Table 2S, Supplementary Material) that was not associated with the increased risk mortality in contrast to a previous study (Simanek et al., 2011). The reason for this lack of association may be attributed to intrinsic

or environmental features of the Italian population involved in this study. Interestingly, most Italian centenarians were previously found to be positive for CMV (94.7%) and their offspring seem to be less susceptible to CMV-driven changes to the immune system (Sansonetti et al. 2014).

Serum albumin levels represent a strong predictor of mortality in the elderly (Liu et al., 2015), as well as during some viral infections (Saroj et al., 2017; Lang et al., 2013) and this is consistent with the negative association among TTV viremia and serum albumin concentrations observed in our cohort. Another result, supporting the implication of TTV load in the increased risk of mortality in our elderly population, may be represented by the positive correlation between TTV DNA and IL8 (although only in females), because higher IL-8 plasma levels have been associated with an increased risk of long-term all-cause mortality in patients with coronary-heart-disease (Cavusoglu et al., 2015). However, no significant association between TTV load and markers of systemic inflammation (i. e. Cu/Zn ratio, IL6, CRP; Supplementary Materials), previously demonstrated as strong predictors of mortality (Malavolta et al., 2010; Puzianowska-Kuźnicka et al., 2016), was observed in our cohort.

Moreover, a possible exhaustion of the CD4/CD8 clonal repertoire, during persistent TTV infection, could explain the decreased survival in elderly population as observed in octo-nagenarian subjects with CMV chronic infection (Hadrup et al., 2006). In fact, similarly to CMV infection, it is likely that a constant immune surveillance, necessary to keep the TTV under control, might lead to clonal expansions of TTV-specific T cells, at the expense of a naïve/memory highly diversified repertoire (Di Benedetto et al., 2015). In our study, we have no data on TTV-specific T cells subsets, but we found a negative correlation between TTV reactivation and NK cell activity. NK cells exert a critical role in the control of viral infections and the age-related NK cell impairment predisposes to virus reactivation and persistence (Brandstadter and Yang 2011), therefore it is not surprising that elderly individuals with impaired NK cell activity presented higher TTV viral loads. Furthermore, TTV encodes for miRNA targeting N-myc interactor (NMI) (Kincaid et al., 2013), a transcription cofactor regulating the activity of STAT members in response to interferon- $\gamma$  (IFN- $\gamma$ ) and cytokine signaling (Zhu et al., 1999). Consequently, TTV miRNA could mediate immune evasion leading to a reduced IFN- $\gamma$  production by NK cells with less efficient DC maturation and T cell polarization (Hazeldine and Lord 2013). Interestingly, a recent investigation shows a similar mechanism of viral immune escape from IFN- $\gamma$  responses in CMV infection (Feng et al., 2018).

Considering that persistent antigen stimulation leads to T-cell exhaustion and seems to favor telomere attrition during chronic viral infections (Bellon et al., 2017; Dowd et al., 2017), and because a close correlation has been demonstrated between granulocyte and lymphocyte telomere length (TL) (Kimura et al., 2010), we evaluated the association between TL in PBMCs and TTV viral load to demonstrate the role of TTV chronic infection in immunosenescence. However, we did not find an association between TL in PBMCs and TTV loads in a subgroup of 87 subjects (Fig 1S Supplementary materials), that might also depend on the small sample size.

We have also investigated TTV miRNA that might be involved in viral immune evasion. Conversely to previous evidence (Vignolini et al., 2016), we did not find any blood sample positive for miRNA-t1a of TTV group 1, besides miRNA-t3b and miRNA-tth8 showed a frequency in elderly people lower than the one found in healthy donors previously examined (Vignolini et al. 2016; Table 5S, Supplementary materials). To date, role and significance of TTV miRNAs expression is yet essentially unknown, but our results demonstrate that significant differences could appear in production of some TTV miRNAs when different groups of subjects are compared to each other. What these findings mean is still unexplored, but the different expression of these miRNAs in relation to the immune status of differently aged subjects encourages further studies in this direction.

In addition, no correlation between TTV DNA loads and miRNA-t3b was found. This finding might be related to the different samples used (PMNLs for TTV DNA load and plasma for miRNA-t3b), but also depend from the circumstance that miRNA-t3b may be mainly produced by TTV localized in body districts different from PMNLs.

We have also demonstrated by an automatic linear modelling, that the levels of miRNA-t3b are inversely related to triglycerides, monocytes and CRP, as well as positively correlated to IL6.

Therefore, lipid levels, CRP and monocytes could play a role in suppressing virus infection and miRNA production, as observed during HCV or HIV infection (Atta et al., 2012; Dai et al., 2008; Kuwata et al., 2007), whereas IL6 might be upregulated by viral miRNA as observed in other viral diseases (Qi et al., 2013). However, the regulatory role of TTV miRNAs in inflammatory pathways is unknown and deserves specific functional investigations.

This study presents a limitation consisting in the small sample size, therefore the association between TTV DNA load and mortality needs to be confirmed in larger cohorts of elderly individuals.

In conclusion, this study supports a role of TTV in immunesenescence and increased risk for all-cause mortality in a cohort of Italian elderly people. TTV may represent an additional virus that establishes latency after primary infection and reactivates in aging when the immune system is compromised. Further research examining if TTV may be causally related to immunosenescence and age-related diseases is warranted.

**Table 1 Laboratory parameters in elderly subjects**

	<b>TTV negative (n.106)</b>	<b>TTV positive (n.273)</b>	<b>P value</b>
Age (yrs)	75.0±6.7	77.5±9.0	0.012
Females (%)	58.5%	57.1%	NS
BMI	25.1	24.6	NS
WBC (10 <sup>3</sup> /μl)	7.1±4.9	6.7±3.8	NS
Lymphocytes (%)	27.7±7.8	30.1±8.6	0.007
Neutrophils (%)	61.1±8.4	58.6±9.4	0.009
Erythrocytes (10 <sup>6</sup> / μl)	4.6±0.5	4.5±0.5	NS
Hemoglobin (g/dl)	13.8±1.5	13.5±1.6	NS
CRP (pg/ml)	0.74 ±1.85	0.77±1.94	NS
Creatinine (mg/dl)	0.94±0.42	1.00±0.27	NS
Albumin (g/dl)	3.75±0.43	3.62±0.38	NS
Glycemia (mg/dl)	103±29	100±29	NS

TG (mg/dl)	116±48	122±77	NS
TC (mg/dl)	201±36	205±43	NS
HDL-C (mg/dl)	55±14	53±17	NS
Hypertension	59.1%	61.7%	NS
CVD diseases	34.4%	36.2%	NS
Diabetes mellitus	17.6	13.2	NS
CMV seropositivity*	23.6%	76.4%	0.001

---

Data are mean ± standard deviation

The laboratory parameter analysis were adjusted for age and gender

WBC, white blood cells; CRP, C-reactive protein; TG, triglycerides; TC, total cholesterol; HDL-C, high-density lipoprotein cholesterol; CVD cardiovascular diseases

\* analyzed in 343 subjects

**Table 2. Multivariate cox regression analysis of TTV DNA loads in PMNLs and all-cause mortality in elderly subjects**

	<b>HR</b>	<b>CI 95%</b>	<b>P value</b>
<b>TTV <math>\geq</math> 4.0 DNA log copies/??g</b>	4.78	1.70-13.44	0.003
<b>Age</b>	1.07	1.01-1.15	0.026
<b>gender</b>	0.86	0.28-2.64	0.796
<b>Cu/Zn</b>	4.94	1.62-15.07	0.005
<b>CRP</b>	0.69	0.56-0.87	0.001
<b>TG</b>	1.00	0.99-1.01	0.540
<b>TC</b>	0.99	0.98-1.01	0.714
<b>Albumin</b>	0.09	0.02-0.41	0.002
<b>Glycemia</b>	0.98	0.95-1.02	0.350

Cox regression model for all-cause mortality (N = 379) adjusted for age, gender, Cu/Zn (Copper Zinc ratio), CRP (C-reactive protein), TG, triglycerides, TC, total cholesterol, albumin and fasting glucose levels

**Fig 1 Correlation between TTV DNA loads in PMNL and age in elderly subjects**

Pearson Correlation between TTV DNA loads (log-transformed data) and age was analyzed both in males and females). Viral loads were expressed as the number of viral genomic copies per  $\mu\text{g}$  of genomic DNA extracted by PMNL (Copies/  $\mu\text{g}$ ). R coefficient and P value are shown.

**Fig 2 Linear regression between TTV DNA loads in PMNL serum albumin levels and NK cell activity in elderly subjects**

A negative linear regression between TTV DNA loads (log-transformed data) and serum albumin levels (Ln-transformed data) after adjusting for age (Fig. 2A). A negative linear regression between TTV DNA loads (log-transformed data) and NK cell activity (Ln-transformed data) after adjusting for age in a subgroup of 71 elderly subjects (27 males: mean age  $76.8 \pm 7.9$ ; 44 females, mean age:  $76.1 \pm 8.4$ ) (Fig. 2B) . Viral loads were expressed as the number of viral genomic copies per  $\mu\text{g}$  of genomic DNA extracted by PMNLs (copies/  $\mu\text{g}$ ). Standardized Beta coefficient and P value are shown.

**Fig 3 Linear regression between TTV DNA loads in PMNL , monocytes and IL8 in elderly subjects**

Significant positive linear regression between TTV DNA loads (log-transformed data) and monocyte percentage (Ln-transformed data) after adjusting for age in females (Fig. 3A). Significant positive linear regression between TTV DNA loads (log-transformed data) and IL8 (Ln-transformed data) after adjusting for age in males (Fig. 3B).

Viral loads were expressed as the number of viral genomic copies per million of cells (copies/M). Standardized Beta coefficient and P value are shown.

**Fig. 4 Kaplan-Meier survival estimates according to TTV DNA loads in elderly patients**

TTV cat 1 = TTV log copies/ $\mu$ g <4.0; TTV cat 2= TTV log copies/  $\mu$ g  $\geq$  4.0. This figure shows Kaplan-Meier plots comparing all-cause-mortality between TTV log copies/ $\mu$ g <4.0 (blue line) and TTV log copies/ $\mu$ g  $\geq$ 4.0 (green line)

**Fig. 5 Coefficients (SPSS automatic Linear modeling) for miRNA-t3b and TTV DNA loads.**

These charts display the intercept first and then sorts effects from top to bottom in decreasing predictor importance. Within effect containing factors, coefficients are sorted by ascending order of data values. The width of connecting lines in the diagram reflects the coefficient significance, with greater line width corresponding to more significant coefficients (smaller p values).

**Acknowledgments**

This study was supported by INRCA, CARIVERONA Foundation (Call for Scientific Research 2008), and Progetto di Ricerca di Ateneo 2017-2018 of University of Pisa, project no. PRA\_2017\_38, DR n. 83/2017.

**Conflict of Interest:** On behalf of all authors, the corresponding author states that there is no conflict of interest.

## References

- Atta M, Cabral M, Santos G, Paraná R, Atta A. Inflammation biomarkers in chronic hepatitis C: association with liver histopathology, HCV genotype and cryoglobulinemia. *Inflamm Res*. 2012; 61:1101-1106. doi: 10.1007/s00011-012-0502-2.
- Barger SD, Cribbet MR, Muldoon MF. Participant-Reported Health Status Predicts Cardiovascular and All-Cause Mortality Independent of Established and Nontraditional Biomarkers: Evidence From a Representative US Sample. *J Am Heart Assoc*. 2016; 5(9). pii: e003741. doi: 10.1161/JAHA.116.003741.
- Bellon M, Nicot C. Telomere Dynamics in Immune Senescence and Exhaustion Triggered by Chronic Viral Infection. *Viruses*. 2017; 9(10). pii: E289. doi: 10.3390/v9100289.
- Brandstadter JD, Yang Y. Natural killer cell responses to viral infection. *J Innate Immun*. 2011; 3:274-279. doi: 10.1159/000324176.
- Brassard J, Gagné MJ, Leblanc D, Poitras É, Houde A, Boras VF, Inglis GD. Association of age and gender with Torque teno virus detection in stools from diarrheic and non-diarrheic people. *J Clin Virol*. 2015;72:55-59. doi: 10.1016/j.jcv.2015.08.020
- Camous X, Pera A, Solana R, Larbi A. NK cells in healthy aging and age-associated diseases. *J Biomed Biotechnol*. 2012;2012:195956. doi: 10.1155/2012/195956.
- Cavusoglu E, Marmur JD, Yanamadala S, Chopra V, Hegde S, Nazli A, Singh KP, Zhang M, Eng C. Elevated baseline plasma IL-8 levels are an independent predictor of long-term all-cause mortality in patients with acute coronary syndrome. *Atherosclerosis*. 2015;242:589-594. doi: 10.1016/j.atherosclerosis.2015.08.022.
- Charpak-Amikam Y, Kubsch T, Seidel E, Oiknine-Djian E, Cavaletto N, Yamin R, Schmiedel D, Wolf D, Gribaudo G, Messerle M, Cicin-Sain L, Mandelboim O. Human cytomegalovirus escapes immune recognition by NK cells through the downregulation of B7-H6 by the viral genes US18 and US20. *Sci Rep*. 2017 Aug 17;7(1):8661. doi: 10.1038/s41598-017-08866-2.
- Dai CY, Chuang WL, Ho CK, Hsieh MY, Huang JF, Lee LP, Hou NJ, Lin ZY, Chen SC, Hsieh MY, Wang LY, Tsai JF, Chang WY, Yu ML. Associations between hepatitis C viremia and low serum triglyceride and cholesterol levels: a community-based study. *J Hepatol*. 2008;49:9-16. doi: 10.1016/j.jhep.2008.03.016.


- Di Benedetto S, Derhovanessian E, Steinhagen-Thiessen E, Goldeck D, Müller L, Pawelec G. Impact of age, sex and CMV-infection on peripheral T cell phenotypes: results from the Berlin BASE-II. *Biogerontology*. 2015; 16:631-643. doi: 10.1007/s10522-015-9563-2.
- Dowd JB, Bosch JA, Steptoe A, Jayabalasingham B, Lin J, Yolken R, Aiello AE. Persistent Herpesvirus Infections and Telomere Attrition Over 3 Years in the Whitehall II Cohort. *J Infect Dis*. 2017 Sep 1;216(5):565-572. doi: 10.1093/infdis/jix255.
- Feng L, Sheng J, Vu GP, Liu Y, Foo C, Wu S, Trang P, Paliza-Carre M, Ran Y, Yang X, Sun X, Deng Z, Zhou T, Lu S, Li H, Liu F. Human cytomegalovirus UL23 inhibits transcription of interferon- $\gamma$  stimulated genes and blocks antiviral interferon- $\gamma$  responses by interacting with human N-myc interactor protein. *PLoS Pathog*. 2018; 14:e1006867. doi: 10.1371/journal.ppat.1006867.
- Feyzioğlu B, Teke T, Ozdemir M, Karaibrahimoğlu A, Doğan M, Yavşan M. The presence of Torque teno virus in chronic obstructive pulmonary disease. *Int J Clin Exp Med*. 2014 ; 7:3461-3466.
- Focosi D, Antonelli G, Pistello M, Maggi F. Torquetenovirus: the human virome from bench to bedside. *Clin Microbiol Infect* 2016 ;22:589-593. doi: 10.1016/j.cmi.2016.04.007.
- Focosi D, Macera L, Boggi U, Nelli LC, Maggi F. Short-term kinetics of torque teno virus viraemia after induction immunosuppression confirm T lymphocytes as the main replication-competent cells. *J Gen Virol*. 2015; 96:115-117. doi: 10.1099/vir.0.070094-0.
- Giacconi R, Costarelli L, Piacenza F, Basso A, Rink L, Mariani E, Fulop T, Dedoussis G, Herbein G, Provinciali M, Jajte J, Lengyel I, Mocchegiani E, Malavolta M. Main biomarkers associated with age-related plasma zinc decrease and copper/zinc ratio in healthy elderly from ZincAge study. *Eur J Nutr*. 2017; 56:2457-2466. doi: 10.1007/s00394-016-1281-2.
- Hadrup SR, Strindhall J, Køllgaard T, Seremet T, Johansson B, Pawelec G, Thor Straten P, Wikby A. Longitudinal studies of clonally expanded CD8 T cells reveal a repertoire shrinkage predicting mortality and an increased number of dysfunctional cytomegalovirus-specific T cells in the very elderly. *J Immunol*. 2006; 176:2645-2653.


- Haloschan M, Bettesch R, Görzer I, Weseslindtner L, Kundi M, Puchhammer-Stöckl E. TTV DNA plasma load and its association with age, gender, and HCMV IgG serostatus in healthy adults. *Age (Dordr)*. 2014; 36:9716. doi: 10.1007/s11357-014-9716-2.
- Hassounah F, Campos C, López-Sejas N, Alonso C, Tarazona R, Solana R, Pera A. Effect of age and latent CMV infection on CD8+ CD56+ T cells (NKT-like) frequency and functionality *Mech Ageing Dev*. 2016;158:38-45. doi: 10.1016/j.mad.2015.12.003.
- Hazeldine J, Lord JM. The impact of ageing on natural killer cell function and potential consequences for health in older adults. *Ageing Res Rev*. 2013; 12:1069-1078. doi:
- Kimura M, Gazitt Y, Cao X, Zhao X, Lansdorp PM, Aviv A. Synchrony of telomere length among hematopoietic cells. *Exp Hematol*. 2010;38: 854-859. doi: 10.1016/j.exphem.2010.06.010. 10.1016/j.arr.2013.04.003.
- Kincaid RP, Burke JM, Cox JC, de Villiers EM, Sullivan CS. A human torque teno virus encodes a microRNA that inhibits interferon signaling. *PLoS Pathog*. 2013; 9:e1003818. doi: 10.1371/journal.ppat.1003818.
- Kuwata T, Kodama M, Sato A, Suzuki H, Miyazaki Y, Miura T, Hayami M. Contribution of monocytes to viral replication in macaques during acute infection with simian immunodeficiency virus. *AIDS Res Hum Retroviruses*. 2007;23:372-380.
- Lang J, Scherzer R, Weekley CC, Tien PC, Grunfeld C, Shlipak MG. Serum albumin and short-term risk for mortality and cardiovascular disease among HIV-infected veterans. *AIDS*. 2013;27:1339-1343. doi: 10.1097/QAD.0b013e32835f1dd6.
- Liu Z, Zhong G, Li S, Deng W, Zhang Y, Qian D, Jin L, Wang X. Use of serum albumin and activities of daily living to predict mortality in long-lived individuals over 95 years of age: a population-based study. *Age (Dordr)*. 2015; 37:9809. doi: 10.1007/s11357-015-9809-6.
- Maggi F, Pistello M, Vatteroni M, Presciuttini S, Marchi S, Isola P, Fornai C, Fagnani S, Andreoli E, Antonelli G, Bendinelli M. Dynamics of persistent TT virus infection, as determined in patients treated with alpha interferon for concomitant hepatitis C virus infection. *J Virol*. 2001;75:11999-12004.
- Maggi F, Andreoli E, Lanini L, Fornai C, Vatteroni M, Pistello M, Presciuttini S, Bendinelli M. Relationships between total plasma load of torquetenovirus (TTV) and TTV genogroups carried. *J Clin Microbiol*. 2005; 43:4807-4810.

- Malavolta M, Giacconi R, Piacenza F, Santarelli L, Cipriano C, Costarelli L, Tesei S, Pierpaoli S, Basso A, Galeazzi R, Lattanzio F, Mocchegiani E. Plasma copper/zinc ratio: an inflammatory/nutritional biomarker as predictor of all-cause mortality in elderly population. *Biogerontology*. 2010; 11:309-319. doi: 10.1007/s10522-009-9251-1.
- Matheï C, Adriaensen W, Vaes B, Van Pottelbergh G, Wallemacq P, Degryse J. No relation between CMV infection and mortality in the oldest old: results from the Belfrail study. *Age Ageing*. 2015; 44:130-135. doi: 10.1093/ageing/afu094.
- Neri S, Mariani E, Meneghetti A, Cattini L, Facchini A. Calcein-acetyoxymethyl cytotoxicity assay: standardization of a method allowing additional analyses on recovered effector cells and supernatants. *Clin Diagn Lab Immunol*. 2001; 8:1131-1135.
- Nikolich-Žugich J, van Lier RAW. Cytomegalovirus (CMV) research in immune senescence comes of age: overview of the 6th International Workshop on CMV and Immunosenescence. *Geroscience*. 2017; 39:245-249. doi: 10.1007/s11357-017-9984-8.
- Nishizawa T, Okamoto H, Konishi K, Yoshizawa H, Miyakawa Y, Mayumi M. A novel DNA virus (TTV) associated with elevated transaminase levels in posttransfusion hepatitis of unknown etiology. *Biochem Biophys Res Commun*. 1997; 241:92–97. doi: 10.1006/bbrc.1997.7765
- Pawelec G. Immunosenescence: role of Cytomegalovirus. *Exp. Gerontol*. 2014; 54: 1–5.
- Puzianowska-Kuźnicka M, Owczarz M, Wieczorowska-Tobis K, Nadrowski P, Chudek J, Slusarczyk P, Skalska A, Jonas M, Franek E, Mossakowska M. Interleukin-6 and C-reactive protein, successful aging, and mortality: the PolSenior study. *Immun Ageing*. 2016; 13:21. doi: 10.1186/s12979-016-0076-x. eCollection 2016.
- Qi Y, Li Y, Zhang L, Huang J. microRNA expression profiling and bioinformatic analysis of dengue virus-infected peripheral blood mononuclear cells. *Mol Med Rep*. 2013; 7:791-798. doi: 10.3892/mmr.2013.1288.
- Rosa AS, Araujo OC, Savassi-Ribas F, Fernandes CA, Coelho HS, Niel C, Villela-Nogueira CA, Araujo NM. Prevalence of occult hepatitis B virus infection and Torque teno virus infection and their association with hepatocellular carcinoma in chronic hepatitis C patients. *Virus Res*. 2017;242:166-172. doi: 10.1016/j.virusres.2017.09.022.


- Sansoni P, Vescovini R, Fagnoni FF, Akbar A, Arens R, Chiu YL, Čičin-Šain L, Dechanet-Merville J, Derhovanessian E, Ferrando-Martinez S, Franceschi C, Frasca D, Fulöp T, Furman D, Gkrania-Klotsas E, Goodrum F, Grubeck-Loebenstien B, Hurme M, Kern F, Lilleri D, López-Botet M, Maier AB, Marandu T, Marchant A, Matheï C, Moss P, Muntasell A, Remmerswaal EB, Riddell NE, Rothe K, Sauce D, Shin EC, Simanek AM, Smithey MJ, Söderberg-Nauclér C, Solana R, Thomas PG, van Lier R, Pawelec G, Nikolic-Zugich J. New advances in CMV and immunosenescence. *Exp Gerontol*. 2014;55:54-62. doi: 10.1016/j.exger.2014.03.020.
- Schorey JS, Cheng Y, Singh PP, Smith VL. Exosomes and other extracellular vesicles in host-pathogen interactions. *EMBO Rep*. 2015;16 :24-43. doi: 10.15252/embr.201439363.
- Saroach A, Arya V, Sinha N, Taneja RS, Sahai P, Mahajan RK. Clinical and laboratory factors associated with mortality in dengue. *Trop Doct*. 2017; 47:141-145. doi: 10.1177/0049475517693993.
- Shibayama T, Masuda G, Ajisawa A, Takahashi M, Nishizawa T, Tsuda F, Okamoto H. Inverse relationship between the titre of TT virus DNA and the CD4 cell count in patients infected with HIV. *AIDS*. 2001;15:563-70.
- Simanek AM, Dowd JB, Pawelec G, Melzer D, Dutta A, Aiello AE. Seropositivity to cytomegalovirus, inflammation, all-cause and cardiovascular disease-related mortality in the United States. *PLoS One*. 2011; 6:e16103. doi: 10.1371/journal.pone.0016103.
- Stowe RP, Peek MK, Cutchin MP, Goodwin JS. Reactivation of herpes simplex virus type 1 is associated with cytomegalovirus and age. *J Med Virol*. 2012; 84: 1797–1802. doi: 10.1002/jmv.23397
- Szládek G, Juhász A, Kardos G, Szoke K, Major T, Sziklai I, Tar I, Márton I, Kónya J, Gergely L, Szarka K. High co-prevalence of genogroup 1 TT virus and human papillomavirus is associated with poor clinical outcome of laryngeal carcinoma. *J Clin Pathol* 2005; 58:402-405.
- Takahashi M, Asabe S, Gotanda Y, Kishimoto J, Tsuda F, Okamoto H. TT virus is distributed in various leukocyte subpopulations at distinct levels, with the highest viral load in granulocytes. *Biochem Biophys Res Commun*. 2002;290:242-248.

- Tokita H, Murai S, Kamitsukasa H, Yagura M, Harada H, Takahashi M, Okamoto H. Influence of TT virus infection on the thrombocytopenia of patients with chronic liver disease. *Hepatol Res.* 2001; 20:288-300.
- Ventura MT, Casciaro M, Gangemi S, Buquicchio R. Immunosenescence in aging: between immune cells depletion and cytokines up-regulation. *Clin Mol Allergy.* 2017;15:21. doi: [10.1186/s12948-017-0077-0](https://doi.org/10.1186/s12948-017-0077-0).
- Vignolini T, Macera L, Antonelli G, Pistello M, Maggi F, Giannecchini S. Investigation on torquetenovirus (TTV) microRNA transcriptome in vivo. *Virus Res.* 2016; 217:18-22. doi: [10.1016/j.virusres.2016.03.003](https://doi.org/10.1016/j.virusres.2016.03.003).
- Walton AH, Muenzer JT, Rasche D, Boomer JS, Sato B, Brownstein BH, Pachot A, Brooks TL, Deych E, Shannon WD, Green JM, Storch GA, Hotchkiss RS. Reactivation of multiple viruses in patients with sepsis. *PLoS One.* 2014; 9:e98819. doi: [10.1371/journal.pone.0098819](https://doi.org/10.1371/journal.pone.0098819). eCollection 2014.
- Westman G, Berglund D, Widén J, Ingelsson M, Korsgren O, Lannfelt L, Sehlin D, Lidehall AK, Eriksson BM. Increased inflammatory response in cytomegalovirus seropositive patients with Alzheimer's disease. *PLoS One.* 2014; 9:e96779. doi: [10.1371/journal.pone.0096779](https://doi.org/10.1371/journal.pone.0096779).
- Zhu M, John S, Berg M, Leonard WJ. Functional Association of Nmi with Stat5 and Stat1 in IL-2- and IFN  $\gamma$ -Mediated Signaling. *Cell.* 1999; 96:121-130.







Beta coefficient= -0.244  $p < 0.0001$




Beta coefficient= -0.269  $p < 0.05$




Males: Beta coefficient= 0.014 p=0.88  
Females: Beta coefficient= 0.206 p=0.009



Males: Beta coefficient= 0.178 p=0.014  
Females: Beta coefficient= 0.168 p=0.08




A



Model Term	Coefficient ►	Sig.	Importance
Intercept	1,547.914	.000	
Triglycerides	-5.172	.002	0.276
Monocytes	-90.758	.004	0.241
CRP	-287.906	.006	0.214
IL6	17.435	.023	0.147
Lymphocytes	-14.226	.037	0.122

B



Model Term	Coefficient ►	Sig.	Importance
AGE	589.612	.000	0.642
Albumin	-7,092.953	.009	0.160
Triglycerides	46.556	.015	0.142

## **Torquetenovirus (TTV) load predicts mortality in Italian elderly subjects**

Robertina Giacconi<sup>1\*</sup>, Fabrizio Maggi<sup>2°</sup>, Lisa Macera<sup>2</sup>, Mauro Pistello<sup>2</sup>, Mauro Provinciali<sup>1</sup>, Simone Giannecchini<sup>3</sup>, Francesco Martelli<sup>3</sup>, Pietro Giorgio Spezia<sup>2</sup>, Erminia Mariani<sup>4,5</sup>, [Roberta Galeazzi<sup>6</sup>](#), [Laura Costarelli<sup>6</sup>](#), Lorenzo Iovino<sup>7</sup>, Sara Galimberti<sup>7</sup>, Lorenzo Nisi<sup>1</sup>, Francesco Piacenza<sup>1</sup>, Marco Malavolta<sup>1</sup>

<sup>1</sup> Advanced Technology Center for Aging Research, Scientific and Technological Pole, Italian National Institute of Health and Science on Aging (INRCA), Ancona, Italy.

<sup>2</sup> Retrovirus Center and Virology Section, Department of Translational Research, University of Pisa, Pisa, Italy.

<sup>3</sup> Department of Experimental and Clinical Medicine, University of Florence, Florence, Italy.

<sup>4</sup> Laboratory of Immunorheumatology and Tissue Regeneration/RAMSES, Rizzoli Orthopaedic Institute, Bologna, Italy.

<sup>5</sup> Department of Medical and Surgical Sciences, University of Bologna, Bologna, Italy.

<sup>6</sup> [Clinical Laboratory & Molecular Diagnostics, INRCA-IRCCS, Ancona, Italy](#)

<sup>7</sup> Section of Hematology, Department of Clinical and Experimental Medicine, University of PisaPisa, Italy.

<sup>°</sup>Both authors contributed equally to this manuscript.

\*Address correspondence to this author at the Advanced Technology Center for Aging Research, IRCCS-INRCA, Via Birarelli 8, 60121, Ancona, Italy; Tel: +39-071-8004213; Fax +39-071-206791; E-mail [r.giacconi@inrca.it](mailto:r.giacconi@inrca.it)

**Table 1S Multivariate cox regression analysis of TTV DNA loads in PMNLs and all-cause mortality after adjustment for CMV IgG antibody titer in elderly subjects**

	<b>HR</b>	<b>CI 95%</b>	<b>P value</b>
<b>TTV <math>\geq</math> 4.0 DNA log copies/??g</b>	9.83	2.48-38.97	0.001
<b>Age</b>	1.06	0.98-1.14	0.125
<b>gender</b>			
<b>CMV IgG antibody titer (log values)</b>	187.11	0.63-55215	0.071
<b>Cu/Zn</b>	99.44	5.12-1928	0.002
<b>CRP</b>	0.70	0.44-1.13	0.151
<b>TG</b>	1.62	0.38-6.92	0.512
<b>TC</b>	0.21	0.007-6.67	0.380
<b>Albumin</b>	0.012	0.001-1.46	0.071
<b>Glycemia</b>	0.099	0.001-6.63	0.281

Cox regression model for all-cause mortality (N = 343) adjusted for age, gender, CMV IgG antibody titer, Cu/Zn (Copper Zinc ratio), CRP (C-reactive protein), TG, triglycerides, TC, total cholesterol, albumin and fasting glucose levels

**Table 2S Multivariate cox regression analysis of TTV DNA loads (continuous variable) in PMNLs and all-cause mortality in elderly subjects**

	<b>HR</b>	<b>CI 95%</b>	<b>P value</b>
<b>TTV DNA log copies/??g</b>	1.62	1.05-2.51	0.030
<b>Age</b>	1.10	1.02-1.18	0.006
<b>gender</b>	1.01	0.30-3.37	0.98
<b>Cu/Zn</b>	23.52	1.94-284.9	0.013
<b>CRP</b>	0.88	0.59-1.28	0.504
<b>TG</b>	1.84	0.50-6.76	0.359
<b>TC</b>	0.506	0.028-9.127	0.64
<b>Albumin</b>	0.003	0.00-0.285	0.012
<b>Glycemia</b>	0.061	0.002-1.998	0.116

Cox regression model for all-cause mortality (N = 379) TTV DNA copies was slightly but significantly associated with three year mortality after adjustment for age, gender, Cu/Zn

(Copper Zinc ratio), CRP (C-reactive protein), TG, triglycerides, TC, total cholesterol, albumin and fasting glucose levels

**Table 3S** TTV miRNAs expression in elderly participants

	Elderly subjects (n.103)	
	N. of positive patients (%)	Mean load
miRNA-t1a (TTV group 1)	0%	-
miRNA-t3b (TTV group 3)	28.2%	565±841
miRNA-tth8 (TTV group 5)	4.8%	720±624

A subgroup of subjects representative of the whole population were selected for the assay of TTV miRNA expression in plasma

TTV miRNA were expressed as copies per ng of total RNA extracted  $\pm$  standard deviation.


**Table 4S** Multiple linear regression model among miRNA-t3b of TTV group 3 and variables independently associated after Automatic linear modelling

Coefficients <sup>a</sup>						
Model		Unstandardized Coefficients		Standardized Coefficients	t	Sig.
		B	Std. Error	Beta		
1	(Constant)	6839,024	1652,913		4,138	,000
	Lntriglycerides	-972,889	248,722	-,487	-3,912	,000
	LnCRP	-317,369	78,388	-,577	-4,049	,000
	LnIL6	650,575	240,146	,326	2,709	,010
	LnMonocytes	-1258,752	283,372	-,557	-4,442	,000
	LnLymphocytes	-613,271	280,223	-,293	-2,189	,034

**Table 5S** Multiple linear regression model among TTV DNA loads in PMNL and variables independently associated after Automatic linear modelling


Coefficients <sup>a</sup>						
Model		Unstandardized Coefficients		Standardized Coefficients	t	Sig.
		B	Std. Error	Beta		
1	(Constant)	3,786	1,147		3,300	,001
	Age in years	,029	,007	,283	4,416	,000
	Lntriglycerides	-,071	,122	-,035	-,581	,561
	LnAlbumin	-1,905	,519	-,235	-3,669	,000

a. Dependent Variable: log10TTV




**Fig 1S Pearson correlation between TTV DNA loads in PMNL and telomere length in elderly subjects**

A negative correlation between TTV DNA loads (log-transformed data) and telomere length (Ln-transformed data) was observed ( $R=-0,219$   $P<0.05$ ) in a subgroup of 87 elderly subjects. However, Linear regression after adjusting for age showed no significant association (Beta coefficient:  $-0,093$   $p=0.41$ ). Telomere length was determined by flow-FISH methodology by using the Telomere PNA kit/FITC from Dako (Glostrup, Denmark) according manufacture's instruction. Viral loads were expressed as number of viral genomic copies per  $\mu\text{g}$  of genomic DNA extracted by PMNL (Copies/  $\mu\text{g}$ ).




**Fig 2S Pearson correlation between TTV DNA loads in PMNL and CMV IgG antibody titer in elderly subjects**

No significant correlation between TTV DNA loads (log-transformed data) and CMV IgG antibody titer (Log-transformed data) [Total population (N=343)  $R=0.034$ ,  $P=0.565$ ; Females  $R=-0.022$   $p=0.77$ ; Males  $R=0.122$   $P=0.189$ ]. Linear regression after adjusting for age confirmed that there was no significant association (Beta coefficient:0.005  $p=0.93$ ). Viral loads were expressed as number of viral genomic copies per  $\mu\text{g}$  of genomic DNA extracted by PMNL (Copies/  $\mu\text{g}$ ).




**Fig 3S** Pearson correlation between TTV DNA loads in PMNL, and Cu/Zn ratio in elderly subjects

Significant positive correlation between TTV DNA loads (log-transformed data) and Cu/Zn ratio (Ln-transformed data) (Total population  $R=0.115$ ,  $P<0.05$ ; Females  $R=0.169$   $p=0.019$ ; Males  $R=0.007$   $P=0.937$ ). However, Linear regression after adjusting for age showed no significant association (Beta coefficient:0.029  $p=0.53$ ). Viral loads were expressed as number of viral genomic copies per  $\mu$ g of genomic DNA extracted by PMNL (Copies/  $\mu$ g).




**Fig 4S** Pearson correlation between TTV DNA loads in PMNL, and IL6 in elderly subjects

Significant positive correlation between TTV DNA loads (log-transformed data) and IL6 (Ln-transformed data) ( $R=0.114$ ,  $P<0.05$ ). However, Linear regression after adjusting for age showed no significant association (Beta coefficient:0.093  $p=0.078$ ) Viral loads were expressed as number of viral genomic copies per  $\mu$ g of genomic DNA extracted by PMNL (Copies/  $\mu$ g).




**Fig 5S** Pearson correlation between TTV DNA loads in PMNL , and CRP in elderly subjects

No significant correlation between TTV DNA loads (log-transformed data) and CRP (Ln-transformed data) ( $R=-0.19$ ,  $P=0.735$ ) was observed. Viral loads were expressed as number of viral genomic copies per  $\mu$ g of genomic DNA extracted by PMNL (Copies/  $\mu$ g)




Ln Neutrophiles



Ln Lymphocytes

**Fig 6S** Pearson correlation between TTV DNA loads in PMNL , and leucocyte populations in elderly subjects

No Significant correlations between TTV DNA loads (log-transformed data) and neutrophils or lymphocytes (Ln-transformed data) ( $R = -0.054$ ,  $P = 0.376$ ;  $R = 0.019$ ,  $P = 0.751$ ) were found. Viral loads were expressed as number of viral genomic copies per  $\mu\text{g}$  of genomic DNA extracted by PMNL (Copies/  $\mu\text{g}$ )



**Fig 7S Pearson correlation between TTV DNA loads in PMNL and TTV DNA viremia in plasma samples from elderly subjects**

Significant positive correlation between TTV DNA loads in PMNL and plasma (log-transformed data) analyzed in 50 elderly subjects (N=50; 36 females and 14 males mean age:  $80.4 \pm 9.7$ ) ( $R=0.571$ ,  $P<0.0001$ ). 6/50 (12%) resulted TTV negative plasma samples. Viral loads were expressed as number of viral genomic copies per  $\mu\text{g}$  of genomic DNA extracted by PMNL (Copies/  $\mu\text{g}$ ) or per ml of plasma.