



ELSEVIER

Contents lists available at ScienceDirect

Annals of Physics

journal homepage: www.elsevier.com/locate/aop



Hawking radiation for a scalar field conformally coupled to an AdS black hole



P. Valtancoli*

Dipartimento di Fisica, Polo Scientifico Università di Firenze, Via G. Sansone 1, 50019 Sesto Fiorentino, Italy
INFN, Sezione di Firenze, Via G. Sansone 1, 50019 Sesto Fiorentino, Italy

ARTICLE INFO

Article history:

Received 15 June 2015

Accepted 5 August 2015

Available online 12 August 2015

Keywords:

Thermodynamics

Black holes

Heun equation

ABSTRACT

The decomposition in normal modes of a scalar field conformally coupled to an AdS black hole leads to a Heun equation with simple coefficients thanks to conformal invariance. By applying the Damour–Ruffini method we can relate the critical exponent of the radial part at the horizon surface to the Hawking radiation of scalar particles.

© 2015 Elsevier Inc. All rights reserved.

1. Introduction

An important application of quantum gravity has been the discover of the relation between black holes and thermodynamics. At large distance from the curvature singularity of the black hole, the gravitational effects are so weak that calculations can be based on the technique of quantum field theory on curved space.

In this way Hawking in 1974 proved that black holes can emit any kind of particle (transforming a pure state into a mixed state), similarly to the black-body radiation [1]. The Hawking radiation loses all the information about the black hole interior, apart from essential parameters like mass, angular momentum and charge (the so called no-hair theorem).

Afterwards many alternative methods have been proposed for a better understanding of this phenomenon:

* Correspondence to: Dipartimento di Fisica, Polo Scientifico Università di Firenze, Via G. Sansone 1, 50019 Sesto Fiorentino, Italy.

E-mail address: valtancoli@fi.infn.it.

(i) the tunnelling method; the Hawking radiation can be viewed as a tunnelling process, where the barrier is created by the tunnelling particle itself. To calculate this process, related to the imaginary part of the action, two methods have been developed, one known as null-geodesic method (Parikh and Wilczek [2]) and the other as Hamilton–Jacobi method (Angheben and others [3]).

(ii) the classical Damour–Ruffini method [4], that will be reviewed in the present article. To apply this method it is necessary studying the solutions of the Klein–Gordon equation in presence of a black hole, considered as a background.

Our contribution is deriving the exact solution of the Klein–Gordon equation for a scalar field conformally coupled to an *AdS* black hole in terms of known functions. In particular we have been to simplify the radial part in terms of a Heun function with elementary coefficients. An essential role for simplifying the solution is played by the conformally invariant coupling with the gravitational field.

In the *AdS* case we must avoid introducing an explicit mass term for the scalar field, which would break conformal invariance, because there is a spontaneous generation of a mass term from the coupling between the *AdS* curvature and the scalar field. Thanks to the Damour–Ruffini method one can verify that the Hawking radiation also arises in this case.

Moreover the exact solution of the wave function in presence of an *AdS* black hole may be useful for a better understanding of the gauge/gravity duality (the *AdS/CFT* correspondence) that in recent years has received a lot of attention in literature.

2. Properties of the *AdS* black hole

The *AdS* black hole metric can be expressed in the following form:

$$ds^2 = \frac{Q(r)}{r^2} dt^2 - r^2 \left[\frac{dr^2}{Q(r)} + d\theta^2 + \sin^2 \theta d\phi^2 \right] \quad (2.1)$$

where

$$Q(r) = \frac{|\Lambda|}{3} r \left(r^3 + \frac{3}{|\Lambda|} r - \frac{6m}{|\Lambda|} \right). \quad (2.2)$$

This metric satisfies the Einstein equations with negative cosmological constant outside the singularity $r = 0$ and in particular

$$R = -4|\Lambda|. \quad (2.3)$$

The non trivial roots of $Q(r)$, that are contained in the third degree polynomial equation, can be explicitly computed with the Cardano formula:

$$\begin{aligned} r_1 &= \frac{1}{\sqrt{|\Lambda|}} (\alpha - \alpha^{-1}) = \frac{6m}{1 + \alpha^2 + \alpha^{-2}} < 2m \\ r_2 &= -\frac{1}{2} r_1 + i \sqrt{\frac{3}{|\Lambda|}} \left(\frac{\alpha + \alpha^{-1}}{2} \right) \\ r_3 &= -\frac{1}{2} r_1 - i \sqrt{\frac{3}{|\Lambda|}} \left(\frac{\alpha + \alpha^{-1}}{2} \right) \end{aligned} \quad (2.4)$$

where

$$\alpha = (3m\sqrt{|\Lambda|} + \sqrt{1 + 9m^2|\Lambda|})^{\frac{1}{3}} > 1. \quad (2.5)$$

In the limit $\Lambda \rightarrow 0$ the complex roots r_2 and r_3 go to infinity and they decouple from the solution, while r_1 , the only positive real root, defines the event horizon.

The gravitational acceleration on the black hole horizon surface r_1 and the Hawking temperature can be computed in terms of the α parameter

$$\begin{aligned}
 k_1 &= \frac{Q'(r_1)}{2r_1^2} = \frac{|\Lambda|}{6} \frac{(r_1 - r_2)(r_1 - r_3)}{r_1} = \frac{1 + |\Lambda|r_1^2}{2r_1} \\
 &= \frac{1 + \alpha^4 + \alpha^{-4}}{12m} > \frac{1}{4m} \\
 T_1 &= \frac{k_1}{2\pi}.
 \end{aligned}
 \tag{2.6}$$

3. Conformal invariance

There are many ways to couple a scalar field to gravitation; they can be parameterized by ϵ in the following action:

$$S = \int d^Dx \sqrt{|g|} \frac{1}{2} (g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \epsilon R \phi^2).
 \tag{3.1}$$

The corresponding equation of motion is

$$(\square + \epsilon R)\phi = 0 \quad \square = |g|^{-\frac{1}{2}} \partial_\mu (|g|^{\frac{1}{2}} g^{\mu\nu} \partial_\nu).
 \tag{3.2}$$

Two cases are particularly relevant:

- (i) $\epsilon = 0$ minimal coupling;
- (ii) $\epsilon = \frac{D-2}{4(D-1)}$ conformal invariant coupling. In particular we will discuss in the following the case $\epsilon = \frac{1}{6}$ ($D = 4$), where the action (3.1) is invariant under the following conformal transformations

$$\begin{aligned}
 \tilde{g}_{\mu\nu} &= \Omega^2(x) g_{\mu\nu} \\
 \tilde{R} &= \Omega^{-2}(x) [R + 2(D-1)\square \ln \Omega + (D-1)(D-2) g^{\alpha\beta} \partial_\alpha (\ln \Omega) \partial_\beta (\ln \Omega)] \\
 \tilde{\phi} &= \Omega^{\frac{2-D}{2}} \phi.
 \end{aligned}
 \tag{3.3}$$

Note that in (3.1) we have omitted an explicit mass term to protect the conformal invariance of the action, which is physically important.

4. Equations of motion for the scalar field

The equation of motion for a scalar field in the background of an *AdS* black hole is given by (for the Kerr–Newman case see [5])

$$\begin{aligned}
 \frac{1}{\sqrt{|g|}} \partial_\mu (g^{\mu\nu} \sqrt{|g|} \partial_\nu) \psi + \frac{1}{6} R \psi &= 0 \\
 R &= -4|\Lambda|.
 \end{aligned}
 \tag{4.1}$$

By substituting the various components of the gravitational field (Eq. (2.1)) we obtain

$$\begin{aligned}
 \partial_r^2 \psi + (\partial_r \ln Q(r)) \partial_r \psi - \frac{r^4}{Q^2(r)} \partial_t^2 \psi + \\
 + \frac{1}{Q(r)} \left[\frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta) \psi + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \phi^2} \psi + \frac{2}{3} |\Lambda| r^2 \psi \right] &= 0.
 \end{aligned}
 \tag{4.2}$$

This equation can be resolved by separating the variables:

$$\begin{aligned} \psi &= R(r)S(\theta)e^{im\phi}e^{-i\omega t} \\ \partial_r^2 R(r) + (\partial_r \ln Q(r))\partial_r R(r) + \frac{\omega^2 r^4}{Q^2(r)}R(r) + \frac{1}{Q(r)}\left[\lambda + \frac{2}{3}|\Lambda|r^2\right]R(r) &= 0 \\ \frac{1}{\sin\theta}\partial_\theta(\sin\theta\partial_\theta)S - \frac{m^2}{\sin^2\theta}S &= \lambda S. \end{aligned} \tag{4.3}$$

The angular equation is equivalent to the angular momentum equation of quantum mechanics.

In this article we will discuss in detail the solution of the radial equation. We can map the two real solutions of the equation $Q(r) = 0$ into the points 0, 1 through the following transformation

$$z = \frac{r(r_1 - r_3)}{(r - r_3)r_1}. \tag{4.4}$$

The other two finite singularities of $Q(r)$ are mapped by the transformation (4.4) into the points

$$r_2 \rightarrow z = \xi = \frac{r_2(r_1 - r_3)}{(r_2 - r_3)r_1} \quad r_3 \rightarrow z = \infty \tag{4.5}$$

while the singularity at the infinity $r = \infty$ is mapped into the finite point

$$r = \infty \rightarrow z = \eta = \frac{r_1 - r_3}{r_1}. \tag{4.6}$$

Thanks to the multiple identity

$$\left(\frac{1}{r}, \frac{1}{r - r_1}, \frac{1}{r - r_2}, \frac{1}{r - r_3}\right) = \frac{z - \eta}{\eta r_3} \left(\frac{\eta}{z}, \frac{\eta - 1}{z - 1}, \frac{\eta - \xi}{z - \xi}, 1\right) \tag{4.7}$$

we can recast the radial equation in the z variable:

$$\begin{aligned} \frac{d^2}{dz^2}R(z) + P(z)\frac{d}{dz}R(z) + Q(z)R(z) &= 0 \\ P(z) &= \frac{1}{z} + \frac{1}{z - 1} + \frac{1}{z - \xi} - \frac{2}{z - \eta} \\ Q(z) &= \frac{\omega_1^2}{(z - 1)^2} + \frac{\omega_2^2}{(z - \xi)^2} + \frac{2\omega_1\omega_2}{(z - 1)(z - \xi)} \\ &+ \frac{2}{(z - \eta)^2} - \frac{1}{\eta(z - \eta)} - \frac{4}{\eta} \frac{1 + \xi - z}{(z - 1)(z - \xi)} + \frac{2}{(z - 1)(z - \xi)} \\ &+ \frac{1}{|\Lambda|r_1(r_2 - r_3)z(z - 1)(z - \xi)} \end{aligned} \tag{4.8}$$

where

$$\omega_1 = \frac{3\omega}{|\Lambda|\eta(r_1 - r_2)} \quad \omega_2 = -\xi\omega_1. \tag{4.9}$$

Apparently this equation depends on the singularity $z = \eta$ (which controls the behaviour at infinity in the r coordinate), but if we put this differential equation into the normal form, the whole dependence from $z - \eta$ is automatically removed, thanks to the conformal invariance of the coupling with the gravitational field (the original reference is [6] and afterwards [7–9]):

$$\begin{aligned} R(z) &= e^{-\frac{1}{2}\int P(z)dz}Z(z) \\ \frac{d^2}{dz^2}Z(z) + \tilde{Q}(z)Z(z) &= 0. \end{aligned} \tag{4.10}$$

where

$$\begin{aligned} \tilde{Q}(z) &= Q(z) - \frac{1}{2} \frac{d}{dz} P(z) - \frac{1}{4} P^2(z) \\ &= \frac{1}{4z^2} + \frac{1 + 4\omega_1^2}{4(z-1)^2} + \frac{1 + 4\omega_2^2}{4(z-\xi)^2} + \frac{4\omega_1\omega_2 - 1}{2(z-1)(z-\xi)} \\ &\quad + \left(\frac{1+\xi}{2} - \frac{\xi}{\eta} + \frac{3\lambda}{|\Lambda|r_1(r_2-r_3)} \right) \frac{1}{z(z-1)(z-\xi)}. \end{aligned} \tag{4.11}$$

This equation can be connected with the well known Heun equation

$$\frac{d^2}{dz^2} w(z) + P_H(z) \frac{d}{dz} w(z) + Q_H(z) w(z) = 0 \tag{4.12}$$

where

$$\begin{aligned} P_H(z) &= \frac{\gamma}{z} + \frac{\delta}{z-1} + \frac{\epsilon}{z-\xi} \\ Q_H(z) &= \frac{\alpha\beta z - q}{z(z-1)(z-\xi)}. \end{aligned} \tag{4.13}$$

After rewriting this equation in the normal form, we must identify

$$\begin{aligned} w(z) &= e^{-\frac{1}{2} \int P_H(z) dz} Z(z) \\ \tilde{Q}(z) &= \frac{\frac{\gamma}{2}(1-\frac{\gamma}{2})}{z^2} + \frac{\frac{\delta}{2}(1-\frac{\delta}{2})}{(z-1)^2} + \frac{\frac{\epsilon}{2}(1-\frac{\epsilon}{2})}{(z-\xi)^2} \\ &\quad + \frac{[\alpha\beta - \frac{1}{2}(\gamma\delta + \gamma\epsilon + \delta\epsilon)]}{(z-1)(z-\xi)} - \frac{q - \frac{1}{2}(\gamma\delta\xi + \gamma\epsilon)}{z(z-1)(z-\xi)}. \end{aligned} \tag{4.14}$$

By adding the Heun condition

$$\alpha + \beta = \gamma + \delta + \epsilon - 1 \tag{4.15}$$

we have the following table of identifications

$$\begin{aligned} \alpha &= \gamma = 1 \\ \beta &= 1 + 2i(\omega_1 + \omega_2) \\ \delta &= 1 + 2i\omega_1 \\ \epsilon &= 1 + 2i\omega_2 \\ q &= \frac{\xi}{\eta} - \frac{3\lambda}{|\Lambda|r_1(r_2-r_3)}. \end{aligned} \tag{4.16}$$

5. Hawking temperature

Let us recall that

$$R(z) = e^{-\frac{1}{2} \int P(z) dz} Z(z) = \frac{z-\eta}{\sqrt{z(z-1)(z-\xi)}} Z(z). \tag{5.1}$$

Let us note that the two solutions for $R(r)$ in our case behave as $1/r$ for large r and are both not normalizable (in and out states). The structure of the solution changes drastically in the unphysical region without conformal invariance, with some bizarre behaviour at infinity in the r coordinate.

Near the singularity $z = 1$ the behaviour of the wave function is (introducing the time factor)

$$\begin{aligned} Z(z) &= c_1(z-1)^{\frac{1}{2}+i\omega_1} + c_2(z-1)^{\frac{1}{2}-i\omega_1} \\ \psi(z) &= c_1 e^{-i\omega t} (z-1)^{i\omega_1} + c_2 e^{-i\omega t} (z-1)^{-i\omega_1}. \end{aligned} \tag{5.2}$$

We can relate the critical exponent ω_1 (Eq. (4.9))

$$\omega_1 = \frac{\omega}{2k_1} \tag{5.3}$$

to k_1 , the gravitational acceleration at the black hole horizon surface, and therefore to the Hawking temperature. At the horizon surface the ingoing and outgoing solutions are parameterized by

$$\begin{aligned} \psi_{in} &= e^{-i\omega t} (r - r_1)^{-i\frac{\omega}{2k_1}} \\ \psi_{out}(r > r_1) &= e^{-i\omega t} (r - r_1)^{i\frac{\omega}{2k_1}}. \end{aligned} \tag{5.4}$$

To simplify this discussion we can introduce the tortoise coordinate defined in general as

$$dr_* = \frac{r^2}{Q(r)} dr. \tag{5.5}$$

We can integrate this equation around $r \sim r_1$

$$r_* = \frac{1}{2k_1} \ln(r - r_1) \tag{5.6}$$

where the wave equation reduces to

$$\left(-\frac{\partial^2}{\partial t^2} + \frac{\partial^2}{\partial r_*^2} \right) \psi(r_*, t) = 0. \tag{5.7}$$

The ingoing and outgoing wave functions are therefore given by

$$\psi^{in} = e^{-i\omega(t+r_*)} \quad \psi^{out} = e^{-i\omega(t-r_*)}. \tag{5.8}$$

By changing from the problematic time coordinate to the well behaved Eddington coordinate $v = t + r_*$ the solutions discussed in (5.4) are re-obtained

$$\begin{aligned} \psi^{in} &= e^{-i\omega v} \\ \psi^{out} &= e^{-i\omega v} e^{2i\omega r_*} = e^{-i\omega v} (r - r_1)^{i\frac{\omega}{k_1}}. \end{aligned} \tag{5.9}$$

While the ingoing wave function is analytic the outgoing wave function has a logarithmic singularity at the horizon. We will use these solutions to investigate the Hawking radiation for scalar particles.

6. Damour–Ruffini method

To understand the decay rate it is necessary to build a damped part in the outgoing wave function (Damour–Ruffini method). The outgoing wave function is known only for $r > r_1$ therefore in the region outside the event horizon. A simple analytic continuation in the internal region gives:

$$\begin{aligned} (r - r_1) &\rightarrow e^{-i\pi} (r_1 - r) \\ \psi_{out}(r < r_1) &= e^{-i\omega v} (r_1 - r)^{i\frac{\omega}{k_1}} e^{\frac{\pi}{k_1} \omega}. \end{aligned} \tag{6.1}$$

Since the wave function is not analytic, the continuation produces a damping factor and the scattering probability of the scalar wave at the event horizon is given by

$$\Gamma_1 = \left| \frac{\psi_{out}(r > r_1)}{\psi_{out}(r < r_1)} \right|^2 = e^{-\frac{2\pi}{k_1} \omega}. \tag{6.2}$$

By introducing a normalization constant N_ω , the normalization condition of the wave function fixes the behaviour of N_ω :

$$|N_\omega|^2 = \frac{1}{e^{\frac{2\pi}{k_1} \omega} - 1} = \frac{1}{e^{\frac{\hbar\omega}{k_B T_1}} - 1}. \tag{6.3}$$

This formula implies that the Hawking radiation spectrum for the scalar particles has a thermal character analogous to the black-body spectrum with Hawking temperature T_1 .

7. Conclusions

In this paper we have presented the decomposition in normal modes of a scalar field conformally coupled to an *AdS* black hole. In particular the radial part leads to a differential equation with four finite singularity and the singularity at infinity. This scheme resembles the isomonodromy problem [9], however there is an important difference in our case. In fact one of the four finite singularities can be removed by simply choosing a conformal invariant coupling between the scalar field and the black hole. The resulting Heun equation (with only three finite singularities) has very simple coefficients and the critical exponent at the horizon singularity can be directly related to the Hawking radiation.

We expect that this study can be generalized to the case of integrable $2 + 1$ gravity, in which case black holes exist only for negative cosmological constant. Our study may have possible connections with the *AdS/CFT* correspondence and may be useful to study non-perturbative effects of quantum gravity.

References

- [1] S.W. Hawking, *Comm. Math. Phys.* 43 (1975) 199.
- [2] M.K. Parikh, F. Wilczek, *Phys. Rev. Lett.* 85 (2000) 5042. arXiv: [hep-th/9907001](#).
- [3] M. Angheben, M. Nadalini, L. Vanzo, S. Zerbini, *J. High Energy Phys.* 05 (2005) 014.
- [4] T. Damour, R. Ruffini, *Phys. Rev. D* 14 (1976) 332.
- [5] H.S. Vieira, V.B. Bezerra, C.R. Muniz, *Ann. Phys.* 350 (2014) 14. arXiv: [1401.5397v3](#) [gr-qc].
- [6] H. Suzuki, E. Takasugi, H. Umetsu, *Progr. Theoret. Phys.* 100 (1998) 491. [gr-qc/9805064](#).
- [7] M. Giammatteo, I.G. Moss, *Class. Quantum. Grav.* 22 (2005) 1803. [gr-qc/0502046](#).
- [8] D. Batic, H. Schmid, *J. Math. Phys.* 48 (2007) 042502. [gr-qc/0701064](#).
- [9] F. Novaes, B. Carneiro da Cunha, *J. High Energy Phys.* 07 (2014) 132. arXiv: [1404.5188v1](#) [hep-th].