



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

A direct approach to quaternionic manifolds

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

A direct approach to quaternionic manifolds / Gentili, Graziano; Gori, Anna; Sarfatti, Giulia.. - In: MATHEMATISCHE NACHRICHTEN. - ISSN 0025-584X. - ELETTRONICO. - 290:(2017), pp. 321-331. [10.1002/mana.201500489]

Availability:

The webpage <https://hdl.handle.net/2158/1042770> of the repository was last updated on 2021-03-07T11:31:26Z

Published version:

DOI: 10.1002/mana.201500489

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

A DIRECT APPROACH TO QUATERNIONIC MANIFOLDS

GRAZIANO GENTILI, ANNA GORI, AND GIULIA SARFATTI

ABSTRACT. The recent definition of slice regular function of several quaternionic variables suggests a new notion of quaternionic manifold. We give the definition of quaternionic regular manifold, as a space locally modeled on \mathbb{H}^n , in a slice regular sense. We exhibit some significant classes of examples, including manifolds which carry a quaternionic affine structure.

Mathematics Subject Classification (2010): 30G35, 53C15

keywords: Regular functions of quaternionic variables, quaternionic manifolds

1. INTRODUCTION

The definition of *slice regular* function of one quaternionic variable, given in [9], has originated a rich area of research that is producing a good quaternionic counterpart of the theory of holomorphic functions. In this theory - whose status was recently presented in [8] - many results hold in analogy with the complex case, while several properties show how deep the differences between the complex and the quaternionic setting can be. Slice regular functions play a fundamental role in surprising applications: one of them appears in differential geometry, where slice regular functions can be used for the classification of Orthogonal Complex Structures in certain subdomains of the space of quaternions, [6].

Let \mathbb{H} denote the skew field of quaternions and let $\mathbb{S} \subset \mathbb{H}$ denote the 2-dimensional sphere of purely imaginary quaternions, $\mathbb{S} = \{q \in \mathbb{H} : q^2 = -1\}$. Then \mathbb{H} is decomposed in “slices” as follows

$$\mathbb{H} = \bigcup_{I \in \mathbb{S}} \mathbb{R} + \mathbb{R}I$$

where each $L_I = \mathbb{R} + \mathbb{R}I$ is isomorphic to the complex plane \mathbb{C} . Slice regularity in one variable is then defined as follows (see [8]).

Definition 1.1. *Let Ω be a domain in \mathbb{H} and let $f : \Omega \rightarrow \mathbb{H}$ be a function. For all $I \in \mathbb{S}$, let us denote $L_I = \mathbb{R} + \mathbb{R}I$, $\Omega_I = \Omega \cap L_I$ and $f_I = f|_{\Omega_I}$. The function f is called (slice) regular if, for all $I \in \mathbb{S}$, the restriction f_I is holomorphic, i.e. the function $\bar{\partial}_I f : \Omega_I \rightarrow \mathbb{H}$ defined by*

$$\bar{\partial}_I f = \frac{1}{2} \left(\frac{\partial}{\partial x} + I \frac{\partial}{\partial y} \right) f_I$$

vanishes identically on Ω_I .

This project has been supported by G.N.S.A.G.A. of INdAM - Rome (Italy), by MIUR of the Italian Government (Research Projects: PRIN “Real and complex manifolds: geometry, topology and harmonic analysis” and FIRB “Geometric function theory and differential geometry”).

The class of slice regular functions shares with the one-Century-old family of Fueter regular functions (see [5]) the problem of not being closed under composition, but it has many nice properties: probably the most important is the fact of including the natural polynomials and power series of a quaternionic variable, [15].

The interest and the richness of the theory of holomorphic functions of several complex variables immediately encourages the natural search for a quaternionic analog. Such a search is also urged by the desire to define and study quaternionic manifolds. It turns out that, mainly due to the lack of commutativity, slice regular functions of several quaternionic variables are not easy to define. After different attempts, a recent and satisfactory approach is based on the use of *stem functions* (see Definition 2.1) introduced in [5] by Fueter himself. Indeed Fueter, after he gave his definition of quaternionic regularity in 1934, [5], envisaged a possible different approach. It was only very recently, with the paper by Ghiloni and Perotti [11], that this approach was fully understood and used to define slice regular functions of several quaternionic variables (see Definition 2.4). It is worthwhile noticing that, in one quaternionic variable, the definition of slice regularity made with the use of stem functions is a generalization of the direct definition that we have presented above (see [10]).

How to give a “direct” definition of quaternionic manifold is an old, well motivated, problem addressed by many authors. Indeed, several different definitions of quaternionic manifolds, as spaces locally modeled on \mathbb{H}^n , have been given, and have produced interesting theories. We like to mention here, in particular, the results by Kulkarni, [13], and Sommesse, [14]. Needless to say, the recently defined class of slice regular functions of several quaternionic variables encourages a further attempt to give a definition of quaternionic manifold. It is well known that the main difficulty that one encounters in doing such an attempt is the fact that in general slice regularity is not preserved by composition. For instance, in the one variable case, the only classes of slice regular functions preserved by compositions are the class of affine functions $q \mapsto qa + b$ with $a, b \in \mathbb{H}$, and the so called *slice preserving* functions, i.e. those slice regular functions $f : \Omega \rightarrow \mathbb{H}$ such that $f(\Omega_I) \subseteq \Omega_I$, for all $I \in \mathbb{S}$. As a consequence, natural examples of quaternionic regular manifolds in dimension one are: the quaternionic tori studied in [2] and the Hopf quaternionic manifolds considered in [14], whose transition functions are quaternionic affine; the classical quaternionic projective space \mathbb{HP}^1 , whose transition functions are slice preserving.

The aim of this paper is to give a definition of (slice) quaternionic regular manifold in higher dimension and provide some significant classes of examples.

Definition 1.2. *A differentiable manifold M of $4n$ real dimensions is a quaternionic regular manifold if it admits a differentiable atlas $\{(U_i, \varphi_i)\}_i$, $\varphi_i : U_i \rightarrow \mathbb{H}^n$, whose transition functions*

$$\varphi_{ij} := \varphi_j \circ \varphi_i^{-1} : \mathbb{H}^n \supseteq \varphi_i(U_i \cap U_j) \rightarrow \varphi_j(U_i \cap U_j) \subseteq \mathbb{H}^n,$$

$$\varphi_{ij} : (q_1, \dots, q_n) \mapsto (\varphi_{ij}^1(q_1, \dots, q_n), \dots, \varphi_{ij}^n(q_1, \dots, q_n))$$

are such that each component $\varphi_{ij}^k : \varphi_i(U_i \cap U_j) \rightarrow \mathbb{H}$ is a slice regular function of n quaternionic variables.

It turns out that Sommesse’s quaternionic manifolds, [14], including the quaternionic Iwasawa manifold, are quaternionic regular.

After presenting some preliminary results, we begin by proving, providing explicit coordinates, that, for all $n \in \mathbb{N}$, the quaternionic projective space $\mathbb{H}\mathbb{P}^n$ and the blow-up $Bl_0(\mathbb{H}^n)$ of \mathbb{H}^n at 0 are quaternionic regular manifolds.

We then identify an important class of quaternionic regular manifolds that, following what Kobayashi did in the complex case, [12], we call *affine quaternionic manifolds*, or *manifolds with an affine quaternionic structure*. Examples of these manifolds are constructed as quotients of \mathbb{H}^n with respect to the action of subgroups of affine transformations of \mathbb{H}^n that act freely and properly discontinuously. An exhaustive study of this class of manifolds will be the object of a forthcoming paper. Paper [4] by Díaz, Verjovsky and Vlacci is strictly related to this subject.

Then we pass to prove a few surgery-type results, concerning the connected sum of a quaternionic regular manifold M with $\mathbb{H}\mathbb{P}^n$, and the blow-up of M at a point $p \in M$. Namely, we prove the following result.

Theorem 1.3. *If M is a regular quaternionic manifold of quaternionic dimension n , then the connected sum of M and $\mathbb{H}\mathbb{P}^n$ is quaternionic diffeomorphic to the blow-up of M at a point $p \in M$. Moreover, both manifolds are quaternionic regular.*

We conclude the paper by noticing that the Grassmannians of p -dimensional \mathbb{H} -planes in \mathbb{H}^n are not quaternionic regular manifolds with respect to their natural atlases.

2. PRELIMINARIES ON REGULAR FUNCTIONS OF SEVERAL VARIABLES

We report here for the convenience of the reader the definition of regular function of several quaternionic variables. This notion was introduced by Ghiloni and Perotti in [11]. Let \mathbb{R}_n denote the real Clifford algebra of signature $(0, n)$ generated by e_1, \dots, e_n . Each element of \mathbb{R}_n is of the form $x = \sum_{K \in \mathcal{P}(n)} e_K x_K$ where $\mathcal{P}(n)$ is the powerset of $\{1, \dots, n\}$, $e_K = e_{k_1} \cdots e_{k_s}$ (with $k_1 < k_2 < \cdots < k_s$) are the basis element of \mathbb{R}_n (for $K = \emptyset$ one gets the unit of \mathbb{R}_n) and x_K are real numbers.

Definition 2.1. *Let D be an open subset of \mathbb{C}^n invariant with respect to complex conjugation in each variable z_1, \dots, z_n . A continuous function $F : D \rightarrow \mathbb{H} \otimes_{\mathbb{R}} \mathbb{R}_n$ of the form $F = \sum_{K \in \mathcal{P}(n)} e_K F_K$ is called a stem function if it is Clifford intrinsic, i.e. for any $K \in \mathcal{P}(n)$, $h \in \{1, \dots, n\}$ and $z = (z_1, \dots, z_n) \in D$ the components $F_K : D \rightarrow \mathbb{H}$ satisfy*

$$F_K(z_1, \dots, z_{h-1}, \bar{z}_h, z_{h+1}, \dots, z_n) = \begin{cases} F_K(z) & \text{if } h \notin K, \\ -F_K(z) & \text{if } h \in K. \end{cases}$$

Each stem function defined on $D \subseteq \mathbb{C}^n$ (open subset invariant by separate conjugation) induces a slice function defined on the *circular* subset of \mathbb{H}^n associated with D ,

$$\Omega_D = \{(q_1, \dots, q_n) \in \mathbb{H}^n \mid q_h = x_h + y_h J_h, J_h \in \mathbb{S}, \forall h = 1, \dots, n; (x_1 + y_1 i, \dots, x_n + y_n i) \in D\}.$$

Definition 2.2. *Let $F : D \rightarrow \mathbb{H} \otimes_{\mathbb{R}} \mathbb{R}_n$, with $F = \sum_{K \in \mathcal{P}(n)} e_K F_K$, be a stem function. We define the left slice function $\mathcal{I}(f) : \Omega_D \rightarrow \mathbb{H}$ induced by F as*

$$\mathcal{I}(F)(x_1 + y_1 J_1, \dots, x_n + y_n J_n) := \sum_{K \in \mathcal{P}(n)} J_K F_K(x_1 + y_1 i, \dots, x_n + y_n i)$$

where

$$J_K = \prod_{k \in K}^{\rightarrow} J_k = J_{k_1} \cdots J_{k_s}$$

is the ordered product.

In order to give the definition of *slice regular* functions of several quaternionic variables, we need to give a notion of holomorphicity for stem functions.

Definition 2.3. For any $h = 1, \dots, n$ define the complex structure \mathcal{J}_h on \mathbb{R}_n as

$$\mathcal{J}_h(e_K) := \begin{cases} -e_{K \setminus \{h\}} & \text{if } h \in K, \\ e_{K \cup \{h\}} & \text{if } h \notin K, \end{cases}$$

Notice that $\mathcal{J}_h^2 = -id_{\mathbb{R}_n}$, for any $h = 1, \dots, n$ so that \mathcal{J}_h defines an (almost) complex structure on \mathbb{R}_n . It is possible to extend any almost complex structure \mathcal{J}_h to $\mathbb{H} \otimes_{\mathbb{R}} \mathbb{R}_n$ by setting

$$\mathcal{J}_h(a \otimes x) = a \otimes \mathcal{J}_h(x)$$

for any $a \in \mathbb{H}$, $x \in \mathbb{R}_n$.

Definition 2.4. Let D be an open subset of \mathbb{C}^n invariant with respect to complex conjugation in each variable, let $F : D \rightarrow \mathbb{H} \otimes_{\mathbb{R}} \mathbb{R}_n$ be a stem function of class C^1 and let $f = \mathcal{I}(F) : \Omega_D \rightarrow \mathbb{H}$ be the induced slice function. F is called a *holomorphic stem function* if for any $h = 1, \dots, n$ and any fixed $z^0 = (z_1^0, \dots, z_n^0) \in D$, the function

$$F_h^{z^0} : D_h \rightarrow (\mathbb{H} \otimes_{\mathbb{R}} \mathbb{R}_n, \mathcal{J}_h), \quad z_h \mapsto F(z_1^0, \dots, z_{h-1}^0, z_h, z_{h+1}^0, \dots, z_n^0)$$

is holomorphic on a domain D_h of \mathbb{C} containing z_h . Equivalently, if

$$\bar{\partial}_h F := \frac{1}{2} \left(\frac{\partial F}{\partial x_h} + \mathcal{J}_h \frac{\partial F}{\partial y_h} \right) = 0$$

on D for every $h = 1, \dots, n$.

If F is holomorphic, the induced function $f = \mathcal{I}(F)$ is called a *left slice regular function* on Ω_D .

It is possible to give a “symmetric” definition, namely the definition of *right slice regular function*.

Definition 2.5. Let $F : D \rightarrow \mathbb{H} \otimes_{\mathbb{R}} \mathbb{R}_n$, with $F = \sum_{K \in \mathcal{P}(n)} e_K F_K$, be a stem function. We define the *right slice function* $\mathcal{I}^r(F) : \Omega_D \rightarrow \mathbb{H}$ induced by F as

$$\mathcal{I}^r(F)(x_1 + y_1 J_1, \dots, x_n + y_n J_n) := \sum_{K \in \mathcal{P}(n)} F_K(x_1 + y_1 i, \dots, x_n + y_n i) J_K$$

where

$$J_K = \prod_{k \in K}^{\rightarrow} J_k = J_{k_1} \cdots J_{k_s}$$

is the ordered product.

Right slice regularity can be expressed in terms of the family of (almost) complex structures (extended as before to $\mathbb{H} \otimes_{\mathbb{R}} \mathbb{R}_n$) defined as follows.

Definition 2.6. For any $h = 1, \dots, n$ define the complex structure \mathcal{J}_h^r on \mathbb{R}_n as

$$\mathcal{J}_h^r(e_K) := -(\mathcal{J}_h(e_K^c))^c,$$

where $x \mapsto x^c$ denotes the Clifford conjugation in \mathbb{R}_n .

Definition 2.7. Let D be an open subset of \mathbb{C}^n invariant with respect to complex conjugation in each variable, let $F : D \rightarrow \mathbb{H} \otimes_{\mathbb{R}} \mathbb{R}_n$ be a stem function of class C^1 and let $f = \mathcal{I}^r(F) : \Omega_D \rightarrow \mathbb{H}$ be the induced right slice function. F is called a right holomorphic stem function if

$$\bar{\partial}_h^r F := \frac{1}{2} \left(\frac{\partial F}{\partial x_h} + \mathcal{J}_h^r \frac{\partial F}{\partial y_h} \right) = 0$$

on D for every $h = 1, \dots, n$.

If F is right holomorphic, the induced right slice function $f = \mathcal{I}^r(F)$ is called a right slice regular function on Ω_D .

Example 2.8. Let us collect here some significant examples, referring to [11] for an extensive description. We point out that the ordering of the variables is important for regularity:

- the function $(q_1, q_2) \mapsto q_1^{-1} q_2$ is left slice regular on $\mathbb{H} \setminus \{0\} \times \mathbb{H}$;
- the function $(q_1, q_2) \mapsto q_2 q_1$ is right slice regular on $\mathbb{H} \times \mathbb{H}$;
- the function $(q_1, q_2, q_3) \mapsto q_2 q_1 q_3$ is neither right nor left slice regular.

Definition 2.9. Let $f = (f_1, f_2, \dots, f_m) : \Omega_D \rightarrow \mathbb{H}^m$ be a differentiable function. We say that f is slice regular if, and only if, each component f_i is either a right or a left slice regular function of n quaternionic variables.

3. QUATERNIONIC REGULAR MANIFOLDS

In this section, we give the announced definition of quaternionic regular manifold and we exhibit several examples.

Definition 3.1. A differentiable manifold M of $4n$ real dimensions is a quaternionic regular manifold if it admits a differentiable atlas $\{(U_i, \varphi_i)\}_i$, $\varphi_i : U_i \rightarrow \mathbb{H}^n$, whose transition functions

$$\varphi_{ij} := \varphi_j \circ \varphi_i^{-1} : \mathbb{H}^n \supseteq \varphi_i(U_i \cap U_j) \rightarrow \varphi_j(U_i \cap U_j) \subseteq \mathbb{H}^n,$$

$$\varphi_{ij} : (q_1, \dots, q_n) \mapsto (\varphi_{ij}^1(q_1, \dots, q_n), \dots, \varphi_{ij}^n(q_1, \dots, q_n))$$

are slice regular functions of n quaternionic variables. Such an atlas is called a regular atlas.

Definition 3.2. Let M and N be two quaternionic regular manifolds with regular atlases $\{(U_i, \varphi_i)\}_i$ and $\{(V_j, \psi_j)\}_j$ respectively. A differentiable function $f : M \rightarrow N$ is called quaternionic regular at a point $p \in M$ if, and only if, there exist two systems of local coordinates $\{(U_{i_0}, \varphi_{i_0})\}$ for p , and $\{(V_{j_0}, \psi_{j_0})\}$ for $f(p)$, so that $\psi_{j_0} \circ f \circ \varphi_{i_0}^{-1}$ is slice regular. The function f is called quaternionic regular if it is quaternionic regular at all $p \in M$.

Notice that the quaternionic regular functions between manifolds, that appear in the examples considered in this paper, have the property that their expressions are slice regular in all local coordinates of the given regular atlases. However one cannot expect that this holds in general, since composition of slice regular functions does not always maintain regularity.

3.1. Quaternionic projective spaces. Of course \mathbb{H}^n is a quaternionic regular manifold. If $(q_1, \dots, q_{n+1}) \in \mathbb{H}^{n+1} \setminus \{0\}$, then $[q_1, \dots, q_{n+1}]$ denotes the (right) vector line $\{(q_1 \lambda, \dots, q_{n+1} \lambda) \in \mathbb{H}^{n+1} : \lambda \in \mathbb{H}\}$ of \mathbb{H}^{n+1} . As usual $\mathbb{H}\mathbb{P}^n$ denotes the set of (right) vector lines in \mathbb{H}^{n+1} . It is easy to see that the natural system of coordinates of a quaternionic projective space endows $\mathbb{H}\mathbb{P}^n$ with a structure of quaternionic regular manifold: the usual transition functions are easily seen to be slice regular.

3.2. The blow-up of \mathbb{H}^n . In order to prove that the blow-up of \mathbb{H}^n is quaternionic regular, we use the natural construction of a structure of differentiable manifold on the blow-up of \mathbb{H}^n at a point. To begin with, consider the projection onto the quotient space

$$\varphi : \mathbb{H}^n \setminus \{0\} \rightarrow \mathbb{HP}^{n-1}$$

defined, with classical notation, by

$$\varphi(q_1, \dots, q_n) = [q_1, \dots, q_n],$$

and set $\Gamma_\varphi \subset \mathbb{H}^n \times \mathbb{HP}^{n-1}$ to be the graph of the map φ , i.e. the set

$$\Gamma_\varphi = \{((q_1, \dots, q_n), [a_1, \dots, a_n]) \in (\mathbb{H}^n \setminus \{0\}) \times \mathbb{HP}^{n-1} : (q_1, \dots, q_n) \in [a_1, \dots, a_n]\}.$$

Definition 3.3. *The subset $Bl_0(\mathbb{H}^n) = \Gamma_\varphi \cup (\{0\} \times \mathbb{HP}^{n-1})$ of the Cartesian product $\mathbb{H}^n \times \mathbb{HP}^{n-1}$, endowed with the natural induced topology, is called the blow-up of \mathbb{H}^n at 0. The subset $\{0\} \times \mathbb{HP}^{n-1}$ is called the exceptional set of $Bl_0(\mathbb{H}^n)$.*

The blow-up of \mathbb{H}^n at 0 is a Hausdorff, paracompact and connected topological space, that can be described as follows:

Proposition 3.4. *The blow-up $Bl_0(\mathbb{H}^n)$ of \mathbb{H}^n at 0 can be globally described as the set*

$$A = \{((q_1, \dots, q_n), [a_1, \dots, a_n]) \in \mathbb{H}^n \times \mathbb{HP}^{n-1} : (q_1, \dots, q_n) \text{ is in the line } [a_1, \dots, a_n]\}.$$

Proof. The point $0 \in \mathbb{H}^n$ belongs to all quaternionic 1-dimensional subspaces of \mathbb{H}^n , and hence $(\{0\} \times \mathbb{HP}^{n-1}) \in A$. Moreover, for $(q_1, \dots, q_n) \neq 0$, we have that (q_1, \dots, q_n) belongs to the quaternionic 1-dimensional subspace $[q_1, \dots, q_n]$ of \mathbb{HP}^{n-1} . As a consequence Γ_φ is also contained in A . There are no other possible elements of A , which consequently coincides with $Bl_0(\mathbb{H}^n)$. \square

We now state a couple of technical lemmas, which use the notion of Dieudonné determinant of 2×2 matrices with quaternionic entries. As it is known, the *Dieudonné determinant*, defined as

$$(3.1) \quad \det_{\mathbb{H}} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \sqrt{|a|^2|d|^2 + |c|^2|b|^2 - \operatorname{Re}(c\bar{a}b\bar{d})}$$

generalizes the notion of determinant to the case of quaternionic matrices (see, e.g. [3]). Notice that a straightforward extension of the usual definition of determinant to matrices with quaternionic entries does not maintain the key properties that one expects from the determinant. For example, for a, c, λ quaternions, the two columns of the matrix

$$\begin{pmatrix} a & a\lambda \\ c & c\lambda \end{pmatrix}$$

are \mathbb{H} -linearly dependent, but the Cayley determinant of the same matrix (see e.g. [1]) $ac\lambda - ca\lambda$ does not vanish when $ac \neq ca$ and $\lambda \neq 0$. Moreover, if $ac \neq ca$, then for $\lambda = c^{-1}a^{-1}ca \neq 1$ we have that the two columns of the matrix

$$\begin{pmatrix} a & a \\ c & c\lambda \end{pmatrix}$$

are \mathbb{H} -linearly independent but $ac\lambda - ca = 0$.

Lemma 3.5. *The Dieudonné determinant $\det_{\mathbb{H}}$ has the following properties: For any $\lambda, \mu \in \mathbb{H}$ and any matrix $X = \begin{pmatrix} x & y \\ z & t \end{pmatrix} \in M(2, \mathbb{H})$ we have that*

$$\text{i) } \det_{\mathbb{H}} \begin{pmatrix} x & y\lambda \\ z & t\lambda \end{pmatrix} = \det_{\mathbb{H}} \begin{pmatrix} x\lambda & y \\ z\lambda & t \end{pmatrix} = |\lambda| \det_{\mathbb{H}} \begin{pmatrix} x & y \\ z & t \end{pmatrix}$$

$$\text{ii) } \det_{\mathbb{H}} \begin{pmatrix} \mu x & \mu y \\ z & t \end{pmatrix} = \det_{\mathbb{H}} \begin{pmatrix} x & y \\ \mu z & \mu t \end{pmatrix} = |\mu| \det_{\mathbb{H}} \begin{pmatrix} x & y \\ z & t \end{pmatrix}$$

iii) *If the matrix Y is obtained from the matrix X by: (a) substituting to a row the sum of the two rows, or (b) substituting to a column the sum of the two columns, then $\det_{\mathbb{H}}(X) = \det_{\mathbb{H}}(Y)$.*

Proof. A direct substitution and computation show the assertions (see also [3]). \square

It is natural at this point to search for the inverse of a 2×2 quaternionic matrix having nonvanishing Dieudonné determinant (see [3]). If the matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in M(2, \mathbb{H})$ is right-invertible, then $a \neq 0$ or $b \neq 0$. If $a \neq 0$, then the right inverse of A is

$$(3.2) \quad \begin{bmatrix} x & y \\ t & z \end{bmatrix} = \begin{bmatrix} a^{-1} + a^{-1}b(d - ca^{-1}b)^{-1}ca^{-1} & -a^{-1}b(d - ca^{-1}b)^{-1} \\ -(d - ca^{-1}b)^{-1}ca^{-1} & (d - ca^{-1}b)^{-1} \end{bmatrix}.$$

If instead we are in the case $b \neq 0$ then, the right inverse has the form

$$(3.3) \quad \begin{bmatrix} x & y \\ t & z \end{bmatrix} = \begin{bmatrix} -(c - db^{-1}a)^{-1}db^{-1} & (c - db^{-1}a)^{-1} \\ b^{-1} + b^{-1}a(c - db^{-1}a)^{-1}db^{-1} & -b^{-1}a(c - db^{-1}a)^{-1} \end{bmatrix}.$$

As one may expect, when $ab \neq 0$ then the two forms (3.2) and (3.3) of the inverse of A do coincide. If $abcd \neq 0$ then the inverse matrix of A assumes an even nicer form,

$$(3.4) \quad \begin{bmatrix} x & y \\ t & z \end{bmatrix} = \begin{bmatrix} (a - bd^{-1}c)^{-1} & (c - db^{-1}a)^{-1} \\ (b - ac^{-1}d)^{-1} & (d - ca^{-1}b)^{-1} \end{bmatrix}$$

which allows a Cramer-type rule to solve 2×2 linear systems with quaternionic coefficients (see again [3]).

Lemma 3.6. *Let (q_1, \dots, q_n) and (a_1, \dots, a_n) be elements of \mathbb{H}^n , with $(a_1, \dots, a_n) \neq 0$. Then there exists $\lambda \in \mathbb{H}$ such that $(q_1, \dots, q_n) = (a_1, \dots, a_n)\lambda$ if, and only if, the following system of $\binom{n}{2}$ equations is satisfied:*

$$\det_{\mathbb{H}} \begin{pmatrix} q_i & a_i \\ q_j & a_j \end{pmatrix} = 0$$

for all $1 \leq i < j \leq n$.

Proof. The proof is a consequence of Lemma 3.5. In fact if there is $\lambda \in \mathbb{H}$ such that $(a_1, \dots, a_n)\lambda = (q_1, \dots, q_n)$, then by Lemma 3.5 iii), for all $1 \leq i < j \leq n$, we get

$$\det_{\mathbb{H}} \begin{pmatrix} q_i & a_i \\ q_j & a_j \end{pmatrix} = \det_{\mathbb{H}} \begin{pmatrix} a_i\lambda & a_i \\ a_j\lambda & a_j \end{pmatrix} = |\lambda| \det_{\mathbb{H}} \begin{pmatrix} a_i & a_i \\ a_j & a_j \end{pmatrix} = |\lambda| \det_{\mathbb{H}} \begin{pmatrix} 0 & a_i \\ 0 & a_j \end{pmatrix} = 0.$$

Conversely, if for all $1 \leq i < j \leq n$,

$$\det_{\mathbb{H}} \begin{pmatrix} q_i & a_i \\ q_j & a_j \end{pmatrix} = 0,$$

then in particular (re-ordering the entries if necessary) we have $a_1 \neq 0$ and

$$(3.5) \quad \det_{\mathbb{H}} \begin{pmatrix} q_1 & a_1 \\ q_j & a_j \end{pmatrix} = 0$$

for all $1 < j \leq n$. If $\lambda \in \mathbb{H}$ is such that $q_1 = -a_1\lambda$, then again by Lemma 3.5

$$(3.6) \quad 0 = \det_{\mathbb{H}} \begin{pmatrix} q_1 & a_1 \\ q_j & a_j \end{pmatrix} = \det_{\mathbb{H}} \begin{pmatrix} q_1 & a_1\lambda \\ q_j & a_j\lambda \end{pmatrix}$$

$$(3.7) \quad = \det_{\mathbb{H}} \begin{pmatrix} 0 & a_1\lambda \\ q_j + a_j\lambda & a_j\lambda \end{pmatrix} = \sqrt{|a_1\lambda|^2|q_j + a_j\lambda|^2}$$

and hence either $\lambda = 0$ (and hence $q_1 = 0$) and $q_j = 0$, or $q_j + a_j\lambda = 0$ (for all $1 < j \leq n$). In any case $q_j = -a_j\lambda$ for all $1 < j \leq n$. We then conclude that $(q_1, \dots, q_n) = (a_1, \dots, a_n)\lambda$. \square

We can now prove

Theorem 3.7. *The blow-up $Bl_0(\mathbb{H}^n)$ of \mathbb{H}^n at 0 is a quaternionic regular manifold of real dimension $4n$. Moreover the exceptional set $\{0\} \times \mathbb{H}\mathbb{P}^{n-1}$ is a quaternionic submanifold of real dimension $4n - 4$.*

Proof. To begin with notice that the two natural projections

$$\begin{aligned} \pi_1 &: Bl_0(\mathbb{H}^n) \rightarrow \mathbb{H}^n & \pi_1((q_1, \dots, q_n), [a_1, \dots, a_n]) &= (q_1, \dots, q_n) \\ \pi_2 &: Bl_0(\mathbb{H}^n) \rightarrow \mathbb{H}\mathbb{P}^{n-1} & \pi_2((q_1, \dots, q_n), [a_1, \dots, a_n]) &= [a_1, \dots, a_n] \end{aligned}$$

are continuous maps from $Bl_0(\mathbb{H}^n)$ to \mathbb{H}^n and $\mathbb{H}\mathbb{P}^{n-1}$, respectively. If we use homogeneous coordinates $[a] = [a_1, \dots, a_n]$ in $\mathbb{H}\mathbb{P}^{n-1}$ and set $q = (q_1, \dots, q_n) \in \mathbb{H}^n$ then, using the Dieudonné determinant (3.1), we can write

$$Bl_0(\mathbb{H}^n) = \{(q, [a]) \in \mathbb{H}^n \times \mathbb{H}\mathbb{P}^{n-1} : |q_i|^2|a_j|^2 + |q_j|^2|a_i|^2 - \operatorname{Re}(q_j\bar{q}_ia_i\bar{a}_j) = 0, 1 \leq i < j \leq n\}.$$

On $\mathbb{H}\mathbb{P}^{n-1}$ we use the n natural systems of coordinates $\{(U_i, p_i)\}_{i=1}^n$ where $U_i = \{[a] \in \mathbb{H}\mathbb{P}^{n-1} : a_i \neq 0\}$ and where $p_i : U_i \rightarrow \mathbb{H}^{n-1}$ is such that

$$(3.8) \quad p_i([a_1, \dots, a_{i-1}, a_i, a_{i+1}, \dots, a_n]) = (a_1a_i^{-1}, \dots, a_{i-1}a_i^{-1}, a_{i+1}a_i^{-1}, \dots, a_na_i^{-1})$$

In turn, a $4n$ dimensional, differentiable atlas $\{(V_i, \varphi_i)\}_{i=1}^n$ can be constructed on $Bl_0(\mathbb{H}^n)$ by considering the n open sets $V_i = \pi_2^{-1}(U_i)$ that cover all of $Bl_0(\mathbb{H}^n)$ and, for each $i = 1, \dots, n$, the homeomorphism of V_i onto \mathbb{H}^n defined by

$$\varphi_i(q, [a]) = (a_1a_i^{-1}, \dots, a_{i-1}a_i^{-1}, q_i, a_{i+1}a_i^{-1}, \dots, a_na_i^{-1})$$

whose inverse is

$$\varphi_i^{-1}(b_1, \dots, b_n) = ((b_1b_i, \dots, b_{i-1}b_i, b_i, b_{i+1}b_i, \dots, b_nb_i), [(b_1, \dots, b_{i-1}, 1, b_{i+1}, \dots, b_n)])$$

At this point, for $1 \leq i \neq j \leq n$ we can easily compute the change of coordinates

$$\begin{aligned} \varphi_j \circ \varphi_i^{-1}(b_1, \dots, b_n) &= \varphi_j((b_1b_i, \dots, b_{i-1}b_i, b_i, b_{i+1}b_i, \dots, b_nb_i), [(b_1, \dots, b_{i-1}, 1, b_{i+1}, \dots, b_n)]) \\ &= (b_1b_j^{-1}, \dots, b_{i-1}b_j^{-1}, b_j^{-1}, b_{i+1}b_j^{-1}, \dots, b_{j-1}b_j^{-1}, b_jb_i, b_{j+1}b_j^{-1}, \dots, b_nb_j^{-1}) \end{aligned}$$

that is clearly a diffeomorphism. To prove that π_1 is differentiable it is enough to compute explicitly the map $\pi_1 \circ \varphi_i^{-1} : \mathbb{H}^n \rightarrow \mathbb{H}^n$

$$(3.9) \quad \pi_1 \circ \varphi_i^{-1}(b_1, \dots, b_n) = (b_1 b_i, \dots, b_{i-1} b_i, b_i, b_{i+1} b_i, \dots, b_n b_i)$$

and see that this map is differentiable for every $i = 1, \dots, n$. Analogously, the fact that the map $p_i \circ \pi_2 \circ \varphi_i^{-1} : \mathbb{H}^n \rightarrow \mathbb{H}^{n-1}$

$$p_i \circ \pi_2 \circ \varphi_i^{-1}(b_1, \dots, b_n) = (b_1, \dots, b_{i-1}, b_{i+1}, \dots, b_n)$$

is differentiable for every $i = 1, \dots, n$ proves that π_2 is differentiable. The transition functions $\varphi_j \circ \varphi_i^{-1}$ have components which are right or left slice regular, thus showing that $Bl_0(\mathbb{H}^n)$ is a quaternionic regular manifold.

To prove that the exceptional set is a quaternionic regular submanifold of real dimension $4n - 4$, we can notice that, for $i = 1, \dots, n$, the intersection $(\{0\} \times \mathbb{H}^{n-1}) \cap V_i$ is mapped by φ_i to the \mathbb{H} -hyperplane $b_i = 0$, which has real dimension $4n - 4$. \square

As in the case of $Bl_0(\mathbb{H}^n)$, we can define the blow-up $Bl_0(B^n)$ of the closed unit ball $B^n = \overline{\mathbb{B}}^n$ of \mathbb{H}^n at the point 0. The restriction to $Bl_0(B^n)$ of the charts of the atlas of $Bl_0(\mathbb{H}^n)$ define on $Bl_0(B^n)$ a natural structure of regular quaternionic manifold (with boundary).

3.3. Quaternionic affine manifolds. The Dieudonné determinant, that we have encountered in dimension 2, has been defined for any $n \times n$ quaternionic matrix $A \in M(n, \mathbb{H})$ (see, e.g. [7]). It can be used, in the usual fashion, to define the group of quaternionic $n \times n$ invertible matrices $GL(n, \mathbb{H})$. For $Q = (q_1, q_2, \dots, q_n) \in \mathbb{H}^n$, we can define the group of all quaternionic affine transformations

$$\mathcal{A}(n, \mathbb{H}) = \{Q \mapsto QA + B : A \in GL(n, \mathbb{H}), B = (b_1, b_2, \dots, b_n) \in \mathbb{H}^n\}$$

which is included in the class of slice regular functions. In complete analogy with what Kobayashi does in the complex case, [12], we give the following:

Definition 3.8. *A differentiable manifold M of $4n$ real dimensions has a quaternionic affine structure if it admits a differentiable atlas whose transition functions are restrictions of quaternionic affine functions of $\mathcal{A}(n, \mathbb{H})$.*

In particular, differentiable manifolds endowed with a quaternionic affine structure are quaternionic regular. This can be used to construct a large class of quaternionic regular manifolds; indeed, for any subgroup $\Gamma \subset \mathcal{A}(n, \mathbb{H})$ which acts freely and properly discontinuously on \mathbb{H}^n , the quotient space

$$M = \mathbb{H}^n / \Gamma$$

admits an atlas whose transition functions are slice regular belonging to $\Gamma \subset \mathcal{A}(n, \mathbb{H})$, and hence has a quaternionic affine structure. It is worthwhile noticing that the *quaternionic manifolds* studied by Sommese in [14] all admit a quaternionic affine structure, and hence many significant examples can be found in his paper.

3.4. The connected sum of a quaternionic regular manifold M and $\mathbb{H}\mathbb{P}^n$. In what follows, the elements of the space \mathbb{H}^{n+1} are denoted by (w, q_1, \dots, q_n) , and we set $|q|^2 = q_1\bar{q}_1 + \dots + q_n\bar{q}_n = |q_1|^2 + \dots + |q_n|^2$. We consider the quaternionic regular manifold $\mathbb{H}\mathbb{P}^n$, and the map

$$h : (\mathbb{H}\mathbb{P}^n \setminus [1, 0, \dots, 0]) \rightarrow \mathbb{H}^n$$

defined by

$$h([w, q_1, \dots, q_n]) = \left(\frac{q_1}{|q|^2} \bar{w}, \dots, \frac{q_n}{|q|^2} \bar{w} \right)$$

Let us notice that h is well defined, since

$$\begin{aligned} h([w\lambda, q_1\lambda, \dots, q_n\lambda]) &= \left(\frac{q_1\lambda}{|q|^2|\lambda|^2} \bar{\lambda}\bar{w}, \dots, \frac{q_n\lambda}{|q|^2|\lambda|^2} \bar{\lambda}\bar{w} \right) \\ &= \left(\frac{q_1}{|q|^2} \bar{w}, \dots, \frac{q_n}{|q|^2} \bar{w} \right) = h([w, q_1, \dots, q_n]), \end{aligned}$$

for all non zero quaternions λ . It is straightforward to prove that h is differentiable, and we have

Proposition 3.9. *The map $H : (\mathbb{H}\mathbb{P}^n \setminus [1, 0, \dots, 0]) \rightarrow Bl_0(\mathbb{H}^n)$ defined by*

$$(3.10) \quad H([w, q_1, \dots, q_n]) = \left(\left(\frac{q_1}{|q|^2} \bar{w}, \dots, \frac{q_n}{|q|^2} \bar{w} \right), \left[\frac{q_1}{|q|^2}, \dots, \frac{q_n}{|q|^2} \right] \right)$$

is a diffeomorphism.

Proof. Since there exists $i \in \{1, \dots, n\}$ such that $q_i \neq 0$, then we can use the charts U_{i+1} and V_i , (see the proof of Theorem 3.7) and write the map H as

$$\begin{aligned} (w, q_1, \dots, q_{i-1}, q_{i+1}, \dots, q_n) &\mapsto \\ \mapsto (w, q_1, \dots, q_{i-1}, 1, q_{i+1}, \dots, q_n) &\mapsto (q_1, \dots, q_{i-1}, \frac{\bar{w}}{|q|^2}, q_{i+1}, \dots, q_n). \end{aligned}$$

As a consequence, the map H is differentiable. The inverse map

$$H^{-1} : Bl_0(\mathbb{H}^n) \rightarrow (\mathbb{H}\mathbb{P}^n \setminus [1, 0, \dots, 0])$$

can be expressed as

$$H^{-1}((b_1u, \dots, b_nu), [b_1, \dots, b_n]) = [\bar{u}, \frac{b_1}{|b|^2}, \dots, \frac{b_n}{|b|^2}]$$

and, in coordinates on $\varphi_i(V_i)$, $i = 1, \dots, n$,

$$\begin{aligned} (b_1, \dots, b_{i-1}, u, b_{i+1}, \dots, b_n) &\mapsto \\ \mapsto ((b_1u, \dots, b_{i-1}u, u, b_{i+1}u, \dots, b_nu), [b_1, \dots, b_{i-1}, 1, b_{i+1}, \dots, b_n]) & \\ \mapsto [\bar{u}, \frac{b_1}{|b|^2}, \dots, \frac{b_{i-1}}{|b|^2}, \frac{1}{|b|^2}, \frac{b_{i+1}}{|b|^2}, \dots, \frac{b_n}{|b|^2}] & \\ \mapsto (\bar{u}|b|^2, b_1, \dots, b_n). & \end{aligned}$$

Therefore H^{-1} is differentiable and H is a diffeomorphism. \square

By simply restricting the map H , and with obvious notations, we obtain the following

Corollary 3.10. *The restriction of the map*

$$H([w, q_1, \dots, q_n]) = ((\frac{q_1}{|q|^2}\bar{w}, \dots, \frac{q_n}{|q|^2}\bar{w}), [\frac{q_1}{|q|^2}, \dots, \frac{q_n}{|q|^2}])$$

to $(\mathbb{H}\mathbb{P}^n \setminus [1 \times \mathbb{B}^n])$ establishes a diffeomorphism between $(\mathbb{H}\mathbb{P}^n \setminus [1 \times \mathbb{B}^n])$ and $Bl_0(B^n)$.

In analogy with the real and complex cases, one can define the blow-up $Bl_p(M)$ of a quaternionic regular manifold M of real dimension $4n$, at a point p , as the manifold obtained from M by substituting the point p with the set of all quaternionic (right) vector lines of the tangent space $T_p M$ (i.e. with the $\mathbb{H}\mathbb{P}^{n-1}$ obtained as the quaternionic projective space over $T_p M \cong \mathbb{H}^n$). After recalling that blowing-up at a point is a local operation we are able to prove that

Theorem 3.11. *Let M be a real $4n$ -dimensional differentiable manifold. The connected sum of M and $\mathbb{H}\mathbb{P}^n$ is diffeomorphic to the blow-up of M at a point $p \in M$.*

Proof. As it is well known, to construct the connected sum described in the statement we can start by considering, on one side, a chart (U, φ) of M such that $\varphi(p) = 0$ and that $\varphi(U) \supset \mathbb{B}^n$. In this way $\varphi^{-1}|_{\partial\mathbb{B}^n} : \partial\mathbb{B}^n \rightarrow \varphi^{-1}(\partial\mathbb{B}^n)$ is the “glueing diffeomorphism” of $M \setminus \varphi^{-1}(\mathbb{B}^n)$ and B^n along the boundaries $\varphi^{-1}(\partial\mathbb{B}^n)$ and $\partial\mathbb{B}^n$. We need now to consider $(\mathbb{H}\mathbb{P}^n \setminus [1 \times \mathbb{B}^n])$, which - by Corollary 3.10 - is a manifold with boundary diffeomorphic to $Bl_0(B^n)$. The projection map π_1 defined in Theorem 3.7 induces a diffeomorphism between $(Bl_0(B^n) \setminus \{0\} \times \mathbb{H}\mathbb{P}^{n-1})$ and $B^n \setminus \{0\}$, whose restriction

$$\pi_1|_{\partial Bl_0(B^n)} : \partial Bl_0(B^n) \rightarrow \partial\mathbb{B}^n$$

can be used to construct the glueing diffeomorphism

$$\varphi^{-1} \circ \pi_1|_{\partial Bl_0(B^n)} : \partial Bl_0(B^n) \rightarrow \varphi^{-1}(\partial\mathbb{B}^n)$$

of the connected sum $M \# \mathbb{H}\mathbb{P}^n$. □

3.5. The blow-up of a quaternionic regular manifold at a point. It is easy to prove the following consequence of Theorem 3.11.

Theorem 3.12. *If M is a regular quaternionic manifold of quaternionic dimension n , then the connected sum of M and $\mathbb{H}\mathbb{P}^n$ is quaternionic diffeomorphic to the blow-up of M at a point $p \in M$. Moreover, both manifolds are quaternionic regular.*

Proof. It is enough to show that gluing the boundary of $Bl_0(B^n)$ with that of $M \setminus \varphi^{-1}(\mathbb{B}^n)$ we obtain a regular quaternionic manifold. To this aim, we can find an open neighborhood V such that $Bl_0((1+\epsilon)B^n) \setminus \{0 \times \mathbb{H}\mathbb{P}^{n-1}\} \subseteq V \subseteq Bl_0(\mathbb{H}^n) \setminus \{0 \times \mathbb{H}\mathbb{P}^{n-1}\}$ and consider the map $\psi : V \rightarrow M$ defined as in the preceding proof by $\varphi^{-1} \circ \pi_1$. Indeed, using the chart φ_i , ($i = 1, \dots, n$), of $Bl_0(\mathbb{H}^n)$, see (3.9), and the chart φ of M we obtain that $\varphi \circ \psi \circ \varphi_i^{-1}(q_1, \dots, q_n) = \varphi \circ \varphi^{-1} \circ \pi_1 \circ \varphi_i^{-1}(q_1, \dots, q_n) = (q_1 q_i, \dots, q_{i-1} q_i, q_i, q_{i+1} q_i, \dots, q_n q_i)$ is a regular diffeomorphism between open sets of \mathbb{H}^n . Therefore $(\varphi^{-1} \circ \pi_1 \circ \varphi_i)^{-1}$ is a compatible chart for the quaternionic regular atlas of the connected sum of M and $\mathbb{H}\mathbb{P}^n$. The assertion follows. □

For the sake of completeness, we recall here that the issue of orientation is important when using the connected sum; for example, when considering the sum of two copies of $\mathbb{H}\mathbb{P}^n$, one of the copies has to be endowed with the reverse orientation with respect to the other. For

this reason, to keep trace of the different orientation of the two copies, the connected sum is classically denoted by $\mathbb{H}\mathbb{P}^n \# \overline{\mathbb{H}\mathbb{P}^n}$.

Corollary 3.13. *The connected sum $\mathbb{H}\mathbb{P}^n \# \overline{\mathbb{H}\mathbb{P}^n}$ is quaternionic diffeomorphic to the blow-up of $\mathbb{H}\mathbb{P}^n$ at a point $p \in \mathbb{H}\mathbb{P}^n$.*

3.6. Quaternionic Grassmannians are not quaternionic regular manifolds. A natural generalization of the quaternionic projective space $\mathbb{H}\mathbb{P}^n$, $n \in \mathbb{N}$, is the quaternionic Grassmannian $Gr_{n,p}(\mathbb{H})$, $p < n \in \mathbb{N}$. One may expect that the natural transition functions that one uses to define the quaternionic Grassmannian of p -planes in \mathbb{H}^n are slice regular functions. This is not the case: in fact even the Grassmannian $Gr_{2,4}(\mathbb{H})$ of 2-planes in \mathbb{H}^4 is not a quaternionic regular manifold with respect to the atlas of natural charts that one constructs. To see this, consider the following transition function obtained when the Dieudonné determinant of the

matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in M(2, \mathbb{H})$ is non zero and $abcd \neq 0$ (see (3.4)):

$$\begin{aligned} \begin{bmatrix} a & b \\ c & d \end{bmatrix} &\mapsto \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ a & b \\ c & d \end{bmatrix} \mapsto \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ a & b \\ c & d \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \\ &= \begin{bmatrix} (a - bd^{-1}c)^{-1} & (c - db^{-1}a)^{-1} \\ (b - ac^{-1}d)^{-1} & (d - ca^{-1}b)^{-1} \\ 1 & 0 \\ 0 & 1 \end{bmatrix} \mapsto \begin{bmatrix} (a - bd^{-1}c)^{-1} & (c - db^{-1}a)^{-1} \\ (b - ac^{-1}d)^{-1} & (d - ca^{-1}b)^{-1} \end{bmatrix}. \end{aligned}$$

In view of Definition 2.9, the function

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \mapsto \begin{bmatrix} (a - bd^{-1}c)^{-1} & (c - db^{-1}a)^{-1} \\ (b - ac^{-1}d)^{-1} & (d - ca^{-1}b)^{-1} \end{bmatrix}$$

is not slice regular.

Acknowledgments. The authors wish to thank Alessandro Perotti for useful discussions.

REFERENCES

- [1] H. Aslaksen, *Quaternionic determinants*, Math. Intelligencer **18** (1996), 57-65.
- [2] C. Bisi, G. Gentili, *On quaternionic Tori and their Moduli Space* Preprint 2015.
- [3] C. Bisi, G. Gentili, *Möbius transformations and the Poincaré distance in the quaternionic setting*, Indiana Univ. Math. J., **58** (2009), 2729–2764.
- [4] J. P. Díaz, A. Verjovsky, F. Vlacci, *Quaternionic Kleinian modular groups and arithmetic hyperbolic orbifolds over the quaternions*, arXiv:1503.07214.
- [5] R. Fueter, *Die Funktionentheorie der Differentialgleichungen $\Delta u = 0$ und $\Delta \Delta u$ mit vier reellen Variablen*, Comment. Math. Helv., **7** (1934), 307-330.
- [6] G. Gentili, S. Salamon, C. Stoppato, *Twistor transforms of quaternionic functions and orthogonal complex structures*, J. Eur. Math. Soc., **16** (2014), 2323-2353.
- [7] J. Dieudonné, *Les déterminants sur un corps non commutatif*, Bull. Soc. Math. France, **71** (1943), 27-45.
- [8] G. Gentili, C. Stoppato, D. C. Struppa, *Regular Functions of a Quaternionic Variable*, Springer Monographs in Mathematics, Springer, Berlin-Heidelberg, 2013.
- [9] G. Gentili, D. Struppa, *A new theory of regular functions of a quaternionic variable*, Adv. Math., **216**, 279–301 (2007).
- [10] R. Ghiloni, A. Perotti, *Slice regular functions on real alternative algebras* Adv. Math **226** (2011) 1662–1691.

- [11] R. Ghiloni, A. Perotti, *Slice regular functions of several Clifford variables* in 9th International Conference on Mathematical Problems in Engineering, Aerospace and Sciences : ICNPAA 2012 : Vienna, Austria.
- [12] S. Kobayashi, T. Ochiai, *Holomorphic projective structures on compact complex surfaces*, Math. Ann., **249** (1980), 75-94.
- [13] R. S. Kulkarni, *On the principle of uniformization*, J. Diff. Geom., **13** (1978), 109-138.
- [14] A. J. Sommese, *Quaternionic manifolds*, Math. Ann., **212** (1975), 191-214.
- [15] C. Stoppato, *A new series expansion for slice regular functions*, Adv. Math., **231** (2012), 1401-1416.

DIPARTIMENTO DI MATEMATICA E INFORMATICA “U. DINI”, UNIVERSITÀ DI FIRENZE, 50134 FIRENZE, ITALY

E-mail address: `gentili@math.unifi.it`

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI MILANO, VIA SALDINI 50, 20133 MILANO, ITALY

E-mail address: `anna.gori@unimi.it`

ISTITUTO NAZIONALE DI ALTA MATEMATICA “F. SEVERI”, CITTÀ UNIVERSITARIA, PIAZZALE ALDO MORO 5, 00185 ROMA, ITALY AND INSTITUT DE MATHÉMATIQUES DE JUSSIEU, UNIVERSITÉ PIERRE ET MARIE CURIE, 4, PLACE JUSSIEU, F-75252 PARIS, FRANCE

E-mail address: `sarfatti@math.unifi.it`