



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Nutrigenomics of extra-virgin olive oil: A review

Questa è la versione Preprint (Submitted version) della seguente pubblicazione:

Original Citation:

Nutrigenomics of extra-virgin olive oil: A review / Piroddi, Marta; Albini, Adriana; Fabiani, Roberto; Giovannelli, Lisa; Luceri, Cristina; Natella, Fausta; Rosignoli, Patrizia; Rossi, Teresa; Taticchi, Agnese; Servili, Maurizio; Galli, Francesco. - In: BIOFACTORS. - ISSN 0951-6433. - STAMPA. - 43:(2017), pp. 17-41. [10.1002/biof.1318]

Availability:

This version is available at: 2158/1056521 since: 2021-03-02T11:04:44Z

Published version:

DOI: 10.1002/biof.1318

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

Conformità alle politiche dell'editore / Compliance to publisher's policies

Questa versione della pubblicazione è conforme a quanto richiesto dalle politiche dell'editore in materia di copyright.

This version of the publication conforms to the publisher's copyright policies.

(Article begins on next page)



Nutrigenomics of extra-virgin olive oil: a review.

Journal:	<i>BioFactors</i>
Manuscript ID	BIOF-16-0113.R1
Wiley - Manuscript type:	Review Article
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Piroddi, Marta; University of Perugia, Department of Pharmaceutical Sciences - Clinical and Nutritional Biochemistry lab Albini, Adriana; IRCCS MultiMedica, Scientific and Technology Pole Fabiani, Roberto; University of Perugia, Department of Chemistry, Biology and Biotechnologies Giovannelli, L.; University of Firenze, NEUROFARBA – Section of Famacology and Toxicology Luceri, Cristina; University of Firenze, NEUROFARBA – Section of Famacology and Toxicology Natella, F.; CREA, Consiglio per la Ricerca in Agricoltura e l'Analisi dell'Economia Agraria, CREA-NUT, Food and Nutrition Research Centre Rosignoli, Patrizia; University of Perugia, Department of Chemistry, Biology and Biotechnologies Rossi, Teresa; IRCCS "Tecnologie Avanzate e Modelli Assistenziali in Oncologia" Arcispedale S. Maria Nuova, Research and Statistics, Department Taticchi, Agnese; University of Perugia, Department of Agricultural, Food and Environmental Sciences Servili, Maurizio; University of Perugia, Department of Agricultural, Food and Environmental Sciences Galli, Francesco; University of Perugia, Department of Internal Medicine - Clinical and Nutritional Biochemistry lab</p>
Keywords:	olive oil, extravirgin olive oil, polyphenols, vitamin E, nutrigenomics

SCHOLARONE™
Manuscripts

Nutrigenomics of extra-virgin olive oil: a review.

Marta Piroddi ^{1§}, Adriana Albini ², Roberto Fabiani ³, Lisa Giovannelli ⁴, Cristina Luceri ⁴,
Fausta Natella ⁵, Patrizia Rosignoli ³, Teresa Rossi ⁶, Agnese Taticchi ^{7§}, Maurizio Servili ⁷
and Francesco Galli ^{1*}

¹ Nutrition and clinical biochemistry lab Dept of Pharmaceutical Sciences, University of Perugia Italy

² Scientific and Technology Pole, IRCCS MultiMedica, Milan, Italy

³ Department of Chemistry, Biology and Biotechnologies, University of Perugia, Italy.

⁴ NEUROFARBA – Section of Pharmacology and Toxicology, University of Firenze, Italy

⁵ CREA-NUT, Consiglio per la Ricerca in Agricoltura e l'Analisi dell'Economia Agraria, Food and Nutrition Research Centre, Via Ardeatina 546, 00178 Roma, Italy

⁶ Research and Statistics, Department, IRCCS “Tecnologie Avanzate e Modelli Assistenziali in Oncologia” Arcispedale S. Maria Nuova, Reggio Emilia, Italy.

⁷ Department of Agricultural, Food and Environmental Sciences, University of Perugia, Via S. Costanzo, 06126 Perugia, Italy

§ These authors contributed equally to the work.

* Corresponding author at. Present address:

Francesco Galli, PhD

Laboratory of Biochemistry and Nutrition,

Department of Pharmaceutical Sciences,

University of Perugia,

Via del Giochetto, IT-06126, Perugia, Italy.

E-mail addresses:

francesco.galli@unipg.it,

roosters2110@gmail.com (F. Galli).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

Table of contents

1. Introduction

2. Virgin Olive Oil (VOO) bioactives

3. Nutrigenomics of VOO

4. Toward a hormetic role of extra virgin olive oil (EVOO) phenolic compounds

5. Functional components of olive oil as anti-aging agents: modulation of the miRNome

6. Chemopreventive effects of olive oil polyphenols: molecular and cellular mechanisms.

7. Olive oil polyphenols and their effect on tumor vascularization and progression

8. Signaling and gene response of vitamin E: a focus on the main fat-soluble vitamin in olives and olive oil

9. Conclusions and perspectives

Abstract

Nutrigenomics data on the functional components of olive oil are still sparse, but the available literature on this subject is increasing. Olive oil is the main source of fat and a key health-promoting component of the Mediterranean diet with proposed positive effects on genes involved in the pathobiology of most prevalent age- and lifestyle-related human conditions such as cancer cardiovascular disease and neurodegeneration. Other effects on the regulation of health-promoting genes have been identified for bioactive components of olives and olive leaves. Omics technologies are offering unique opportunities to identify nutritional and health biomarkers associated with these gene responses and to use them with a personalized and even predictive approach, which is a main breakthrough in modern medicine and nutrition. Gene regulation properties of the functional components of olive oil, such as oleic acid, biophenols and vitamin E, point to a role for these molecules as natural homeostatic and even hormetic factors for an application as cytoprotection and early prevention agents in conditions of premature and pathologic aging. Even therapeutic applications can be foreseen in conditions of chronic inflammation, and particularly in cancer, which will be discussed in detail in this review paper as major clinical target of olive oil and its functional components.

Word count: 192

Keywords:

olive oil;

extravirgin olive oil;

polyphenols;

vitamin E;

nutrigenomics.

1. Introduction

Olive Oil (OO) in its production variants of virgin and extra virgin OO (VOO and EVOO, respectively), is universally recognized as a symbol, and the major source of fat, of the Mediterranean diet [1]. In all the traditional forms of this diet found in the Mediterranean basin, VOO, the OO obtained directly from olives and solely by mechanical means, is proposed as main health-promoting component with effects that include a reduced risk of morbidity and mortality for cancer, neurodegenerative diseases such as Parkinson's and Alzheimer's Disease, metabolic syndrome and cardio-cerebro-vascular events [2; 3; 4; 5; 6].

Gene modulation by VOO combined with the other components of Mediterranean diet have been investigated to provide a mechanistic rationale to such a positive clinical outcome (recently reviewed in [7; 8]). Solid evidence was obtained on converging effects of VOO and Mediterranean diet on the homeostatic control of genes having a role in immune-inflammatory pathways, vessel protection and blood pressure control, metabolic regulation and detoxification of reactive species. The actual molecular players of these nutrigenomic effects have been tentatively identified in animal models and humans [7; 9; 10] and the available evidence is strong enough to consider VOO a natural functional food. Besides having a high content of monounsaturated fatty acids (MUFAs), VOO contains a number of "bioactives", such as biophenols (Figure 1) and vitamin E (Figure 2), the latter being the main fat-soluble vitamin of this oil (Section 8), the pattern of which shows huge variability in olives and VOO products available for human consumption [11].

The impact of the accumulated evidence on this edible oil on health and nutrition policies of different regions has been huge. Recommendations of World Health Organization for a healthy diet include VOO as a source of unsaturated fats that should be preferred to more saturated ones, found for instance in fatty meat and dairy products [12], which is in agreement with previous health claims on oleic acid of the European Food Safety Authority (EFSA) [13] and Food and Drug Administration (FDA) [14] (Section 2). Further recognition of the importance of functional components in VOO has been provided by the EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA), concerning polyphenols, and vitamin E [15; 16].

Nutrigenomics encompassing high-throughput omics technologies, such as transcriptomics, proteomics, metabolomics, interactomics and fluxomics, and their implementation with the latest bioinformatics tools, is now available to characterize the molecular markers of these claims. In fact, a growing body of evidence is accumulating on the identification of genes and

1
2
3 1 metabolic responses that link the functional properties of VOO bioactives with the nutritional
4 2 and health-promoting activity of the bioactive molecules found in this edible oil. These
5 3 aspects are discussed in this review paper, which is aimed at providing an updated description
6 4 of the most recently identified gene-bioactive functional interactions produced during the
7 5 consumption of VOO.
8 6
9 7

10 8

11 9 **2. Virgin Olive Oil bioactives**

12 10

13 11 The nutritional and healthy values as well as the sensory and biological properties of VOO
14 12 have been ascribed to the presence of bioactive components such as Monounsaturated and
15 13 Polyunsaturated Fatty Acids (MUFAs and PUFAs), squalene, phytosterols, triterpenic acids
16 14 and dialcohols, pigments, tocopherols and polyphenols.

17 15 Initially, the health-promoting effects of VOO have been attributed to its high MUFAs
18 16 amount. Among them, the oleic acid (18:1 ω -9), representing 49% to 83% of the total FA in
19 17 VOO, is supposed to be the most important one from a healthy point of view [17]. In fact, it
20 18 exerts high efficiency in the modulation of gastrointestinal and metabolic functions and of
21 19 extrinsic cardiovascular risk factors. The ameliorative effect of oleic acid in olive oil is
22 20 thought to occur via modifications to plasma lipid and lipoprotein patterns and levels. Cell
23 21 membrane composition and fluidity, inhibition of coagulation, improvement in glucose
24 22 homeostasis and blood pressure, and attenuation of inflammation and oxidative states in
25 23 fasting conditions, have also been described to be affected. Other important functions
26 24 associated to the oleic acid consumption include the a better control of the secretory activity
27 25 of pancreas and liver (bile secretion) and an improved protection of the gastric mucosa by a
28 26 reduced secretion of hydrochloric acid that helps constraining the risk of gastric-duodenal
29 27 ulcers [18]. However, the most convincing evidence in medicine on the health-promoting
30 28 activity of this MUFA was obtained in cardiovascular prevention trials. This evidence has
31 29 resulted in the formulation of the health claims introduced above and now appearing on olive
32 30 oil labels. More in detail, the FDA in 2004 issued “the benefits on the risk of coronary heart
33 31 disease of eating about two tablespoons (i.e. 23g) of VOO daily, due to the MUFAs (oleic
34 32 acid) in olive oil” [14]. The European Authority EFSA has moved in the same direction with
35 33 the following sentence in a recent opinion: “Replacing saturated fats in the diet with
36 34 unsaturated fats contributes to the maintenance of normal blood cholesterol levels. Oleic acid
37 35 is an unsaturated fat” [13].
38 36
39 37
40 38
41 39
42 40
43 41
44 42
45 43
46 44
47 45
48 46
49 47
50 48
51 49
52 50
53 51
54 52
55 53
56 54
57 55
58 56
59 57
60 58

1
2
3 1 On the other hand, the PUFA linoleic acid (18:2 ω -6) and linolenic acid (18:3 ω -3),
4 2 respectively, as essential FA, are indispensable components of the cell structure and are also
5 3 fundamental for the development of the brain and retina, especially during the growth [19]. In
6 4 humans they are biosynthetic precursors of other long chain unsaturated fatty acids, namely
7 5 arachidonic acid (20:4 n-6) and eicosapentaenoic acid (20:5 n-3), which are involved in
8 6 eicosanoid metabolism, thereby regulating important functions of inflammatory leukocytes,
9 7 platelets and vascular cells. Furthermore, most of the nutrition guidelines agree in considering
10 8 the VOO ratios PUFA/SFA and ω -3/ ω -6 as the best occurring in natural fats. Some
11 9 parameters, such as the area of production, altitude, climate, fruit variety, and stage of
12 10 maturity of the fruit can greatly affect the FA composition of virgin olive oil. It is generally
13 11 accepted that cooler areas produce oil with higher monounsaturated content than warmer
14 12 climates [20].

15 13 Current epidemiological and experimental studies strongly support the fact that the beneficial
16 14 effects of VOO are also due to its minor bioactive components. Among them, the squalene,
17 15 besides its well known anticancer properties, shows several biological activities, with the
18 16 antioxidant one being similar to that of trans retinol [21; 22]. The squalene content in olive oil
19 17 is especially high (up to 0.7% (7 mg/g)) when compared to other oils and human dietary fats.
20 18 Moreover, squalene plays a key role as intermediate metabolite in cholesterol metabolism. In
21 19 vivo and in vitro studies have shown that it regulates the absorption, synthesis, esterification
22 20 and elimination of cholesterol [23] by stimulating acyl-coenzyme A. At the same time it
23 21 reduces cholesterol, thereby increasing the efficiency of statins and reduces the UV-induced
24 22 DNA damage thus preventing human skin photo-aging [24]. Squalene is also an important
25 23 intermediate in the biosynthetic pathway of sterols in both plants and animals [25]. The
26 24 phytosterols represent a major fraction of unsaponifiables molecules in VOO (ranging from
27 25 80 to 260 mg/100g of VOO), mainly represented by β -sitosterol ($\geq 93.0\%$ of total sterols). In
28 26 vivo phytosterols, and particularly β -sitosterol, are effective in reducing the concentrations of
29 27 total and LDL cholesterol, and in stimulating the apoptotic signaling of prostate cancer cells;
30 28 moreover, these sterols are used in the natural treatment of benign prostatic hyperplasia [26;
31 29 27]. The beneficial effect of phytosterols is obtained only with a daily intake of at least 0.8 g
32 30 of plant sterols/stanols, according to the Claim of EFSA named "The sterols / stanols ratio
33 31 contributes to the maintenance of normal levels of blood cholesterol" [16].

34 32 The triterpenes are bioactive molecules found in olive skin and in the leaves of olive trees.
35 33 Hydroxyl pentacyclic triterpene acids (HPTA) (oleanolic and maslinic acid) and dialcohols
36 34 (uvaol and erythrodiol) are responsible for several biological activities attributed to VOO,

1
2
3 1 such as anti-inflammatory, hepatoprotective, anticancer, antiviral, anti-HIV, anti-microbial,
4 2 antifungal, anti-diabetic, gastroprotective and anti-hyperlipemia [28; 29; 30; 31]. Recent
5 3 findings demonstrated also different neuroprotective effects exerted by maslinic and oleanolic
6 4 acid [32; 33; 34]. However, their concentration in VOO is very weak, ranging between 17 and
7 5 344 mg/Kg for oleanolic acid, 19 and 250 mg/Kg for maslinic acid and traces of ursolic, while
8 6 it is significantly higher in crude olive pomace oil. According to some authors, the main
9 7 factors causing variability in the HPTA concentration are the oil free acidity olive variety,
10 8 olive ripeness, and oil extraction system [35].

11 9 Furthermore, VOO contains a considerable amount of pigments (chlorophylls and
12 10 carotenoids). Consistent clinical evidence has been obtained on the antioxidant activity of
13 11 carotenoids as well as on other molecular effects that were associated with the prevention or
14 12 amelioration of serious human ailments [24; 36]; these include cancer and cardiovascular
15 13 disease, and skin and eye disorders. In the latter, carotenoids enhance the optical density of
16 14 macular pigments and protects against the formation of age-related cataracts.. Their health-
17 15 promoting properties are mostly due to carotenes (e.g. β -carotene) and xanthophylls (e.g.
18 16 lutein). In particular, carotenes (precursors of vitamin A) are proposed to quench the singlet
19 17 oxygen, a reactive intermediate of the molecular oxygen formed during the process of light-
20 18 induced oxidation (photo oxidation) of the biomolecules. Lutein shows higher efficiency in
21 19 protecting cellular membranes against lipid peroxidation and in preventing oxidative damage
22 20 to the retina [24].

23 21 Tocopherols (vitamin E) are presented in VOO essentially as α -tocopherol ($\geq 90\%$ of
24 22 tocopherols in EVOO) i.e. the main form of this vitamin E also found in human tissues [37]
25 23 (Section 8). This is one of the most important lipophilic antioxidants found in nature [38; 39]
26 24 and its role in preventing lipid peroxidation of cellular membranes and lipoproteins [40] has
27 25 been recognized in the recent health claim released from EFSA: "Vitamin E helps to protect
28 26 cells from oxidative stress" [16]. Moreover, both the redox-dependent and -independent
29 27 properties of this vitamin have been demonstrated to influence the expression of homeostatic
30 28 genes that protect tissues from oxidative and inflammatory processes associated with aging,
31 29 degenerative diseases and cancer [37]. Noteworthy, the levels of α -tocopherol in different
32 30 VOO products show marked variability depending on pedoclimatic factors and agronomic
33 31 practices, such as the area of origin, the cultivar and the stage of fruit ripening [20; 41]. The
34 32 data obtained assessing 430 samples of EVOO have showed a range of variability between 23
35 33 and 751 mg/kg [17].

1
2
3 1 Besides the compounds described above, it is well known that biophenols are the most
4 2 represented bioactive molecules of VOO. There are more than 100 different biophenols
5 3 reported in olive products (fruit, oil, leaves, and waste) [42]. The composition in terms of
6 4 phenolic compounds is different in olive fruit, oil, leaves, and waste. Additionally agronomic
7 5 (varieties, ripeness, and agro-climatic aspects) and technological factors (variables of
8 6 mechanical extraction process and storage conditions) have a significant impact on the
9 7 biophenol composition of olives, VOOs and by-products [43].

10 8 The main biophenols occurring in olives include the phenyl alcohols hydroxytyrosol (HT),
11 9 and tyrosol, and the secoiridoids oleuropein and ligstroside showed in Figure 1. Other forms
12 10 include verbascoside, lignans, and flavonoids (rutin and glycosides of luteolin and apigenin).
13 11 In VOO the main classes of phenols are phenolic acids, phenolic alcohols, flavonoids,
14 12 secoiridoids (as aglycon derivatives) and lignans. Secoiridoids are characterized by the
15 13 presence in their molecules of elenolic acid (EA) or its derivative forms (Figure 1); they occur
16 14 only in plants belonging to the family of Oleaceae, which includes *Olea europaea* L., thus the
17 15 only natural food sources are table olives and VOO. The most abundant secoiridoids in VOO
18 16 are the dialdehydic form of decarboxymethyl-EA linked to HT or tyrosol (3,4-DHPEA-EDA
19 17 and p-HPEA-EDA, respectively), an isomer of oleuropein aglycone (3,4-DHPEA-EA), and
20 18 the ligstroside aglycone (p-HPEA-EA) (Figure 1), found and characterized for the first time
21 19 by Montedoro et al. [44]. These substances are aglycone derivatives of secoiridoid glucosides
22 20 contained in the olive fruit, originating during the oil mechanical extraction process, by the
23 21 hydrolysis of oleuropein, demethyloleuropein, and ligstroside through the activity of
24 22 endogenous β -glucosidases [45]. The composition in those compounds may be extremely
25 23 variable due to the combination of several factors including agronomical, technological and
26 24 storage factors. Most of the variables involved in such modifications, in fact, have been
27 25 widely investigated during the last 35 years. As an example about the variability in the
28 26 concentration in VOO, in 210 oil samples obtained in industrial plants, average values of the
29 27 prevalent phenolic alcohols, phenolic acids and secoiridoids was in total 352.4 mg/Kg (133.5
30 28 and 950.5 mg/Kg were the lower and the upper quintile, respectively) [45].

31 29 Being responsible for the bitter and pungent sensory attributes of VOO, the secoiridoid
32 30 derivatives in VOO represent a rare case of the healthy value of a food product which is
33 31 directly perceptible by means of sensory stimuli. All the scientific evidences regarding the
34 32 role of those compounds in the prevention of several diseases have contributed in increasing
35 33 consumer awareness about the positive correlation between VOO bitterness and pungency
36 34 and its good quality [46].

1
2
3 1 VOO phenolics may exert beneficial effects as a consequence of their antioxidant,
4 2 antimicrobial and anti-inflammatory activities. Observational and epidemiological studies
5 3 demonstrated the efficacy of the VOO's phenolic compounds on the prevention of chronic
6 4 and inflammatory diseases such as cardio-cerebro-vascular disease and cancer, which is
7 5 consistent with the results of clinical trials that have confirmed the importance of phenolic
8 6 compounds cardiovascular risk protection afforded by VOO (recently reviewed in [45; 47; 48;
9 7 49]). In 2011 the EFSA agency released a health claim [15; 16] concerning the recognition of
10 8 the effectiveness of the ingestion of VOO phenols (HT and its derivatives, 5 mg/day per 20 g
11 9 of VOO) in protecting LDL from oxidation. This is the sole example between the available
12 10 health claims that identified in the phenotype of LDL cholesterol oxidation an underlying
13 11 event in cardiovascular risk to be targeted with a specific dose of a natural functional
14 12 ingredient of food.

15 13 Besides these effects on LDL protection, other molecular mechanisms by which the phenolic
16 14 fraction of VOO could protect human tissues from the pathogenic cues of chronic and
17 15 degenerative diseases have been tentatively identified. These mechanisms presented in detail
18 16 in the next sections, are proposed to include effects on other aspects of lipid metabolism such
19 17 as on HDL levels, the anti-atherogenic fraction of cholesterol [50], as well as on oxidative
20 18 stress and inflammatory parameters (Section 3 and 4), platelet function [51] and activity of
21 19 the fibrinolytic factors PAI-1 and FVII [52], and on endothelial parameters such as the release
22 20 and pro-oxidant effects of the vasodilating gas NO [53], blood cell adhesiveness [54] and
23 21 angiogenic activity that is an important target in chemoprevention and therapy of different
24 22 cancers (Section 7) together with the control of DNA damage as an early event in
25 23 carcinogenesis [48], and with specific effects on signal transduction and gene regulation
26 24 pathways that control proliferation, invasiveness and apoptotic death of cancer cells (Section
27 25 6).

28 26 The possibility that biophenols of VOO and from other food items may reach key molecular
29 27 targets of human cells and tissues to produce such an impressive series of effects, depends on
30 28 the metabolism and bioavailability features of the active forms of these compounds. Data
31 29 regarding the metabolism of VOO phenolics in humans are very limited, and contrasting
32 30 results have been obtained regarding the amounts and forms in which they are present in
33 31 plasma and excreted in urine (reviewed in [55]). These aspects further discussed below
34 32 (Section 4) have been assessed in some human trials that conclusively demonstrated how
35 33 VOO phenolics are resistant to the acidic conditions of the stomach and thus are readily
36 34 absorbed in humans (55–60 % for HT and oleuropein aglycone) [56]. However, blood

1 kinetics were found to vary among the different VOO biophenols. Oleuropein is rapidly
2 absorbed after oral administration with a maximum plasma concentration occurring 2 h after
3 administration [57]. Tyrosol and HT are the products of the metabolic transformation of this
4 and other VOO phenolics such as ligstroside aglycone, which are absorbed in a dose-
5 dependent manner in humans [57]. HT appears in plasma minutes after oral administration,
6 with maximal concentrations observed in 5–10 min and then the renal clearance produces a
7 rapid drop of circulating levels within the first hour [58].

8 Notwithstanding, the free form of HT, that is commonly investigated in cellular tests, is
9 almost completely undetectable in plasma and urine, being > 95% in the conjugated form,
10 mainly condensed with glucuronyl residues, while methylconjugates are a minor form of the
11 molecule excreted in human urine [57; 59]. The same is for oleuropein [60; 61]. Therefore,
12 HT should reach tissues mainly as conjugated form.

13 If the biological activity of HT or other biophenols could be attributed to endogenous
14 metabolites is still matter of investigation and preliminary data on antioxidant properties of
15 HT have provided conflicting results in literature (reviewed in [62]). Other options may
16 include the role of cellular esterase enzymes in the local metabolism and bioactivation of
17 derivatized forms that may have great relevance in the pathophysiology of GI tract, mainly in
18 gut and hepatic tissue. These aspects are worth of further and more accurate investigation also
19 considering the limits imposed by the metabolism of VOO bioactives in animal models and
20 particularly in rodents when compared to humans (reviewed in [55]). Genomics and
21 molecular data on the absorption and biotransformation of VOO biophenols in humans are
22 also elusive and this has fuelled speculation and a biased interpretation of mechanisms that
23 lay behind the biological effects of these molecules.

24 Olive Oil vitamin E (Section 8), similarly to dietary fatty acids, is readily absorbed and
25 delivered to liver and then to systemic lipid pools, through the pre-hepatic and post-hepatic
26 lipoprotein metabolism.

27 28 **3. Nutrigenomics of VOO.**

29
30 In the last few years, the impressive growth of omics technologies has offered the opportunity
31 to deepen the knowledge on the molecular and metabolic effects of VOO and its functional
32 components. High-throughput transcriptomic and metabolomic profiling of VOO
33 administered alone or in combination with Mediterranean diet have been implemented in
34 animal models and humans in the frame of healthy or pathological conditions. Nutrigenomics

1 of biophenols has also been investigated separately from other components of VOO.
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 of biophenols has also been investigated separately from other components of VOO.
2 Transcriptomic fingerprints from a series of these studies have been recently compared in the
3 elegant review paper of Konstantinidou et al. [7] revealing key molecular targets of VOO in
4 the area of prevention and management of CVD and other inflammatory and age-related
5 disorders. Among the gene transcripts investigated in peripheral blood cells, few of these
6 were identified as common molecular signs of the intervention with VOO phenolic
7 compounds or with the Mediterranean diet combined to VOO. The following pathways
8 associated with coronary artery disease [63] appear to harbour most of the genes identified in
9 transcriptomic studies on VOO: oxidoreductase activity (JUN), hydroxymethylglutaryl-CoA
10 reductase activity (HMB-CoA), adipocytokine receptor signaling pathway (ADIPOQ,
11 GLUT4, NFkB, TNF- α), VEGF signaling pathway (COX2), hematopoietic cell lineage
12 (CD14), and cytokine–cytokine receptor interaction (CCL5, LEP, IL6, IL8R, IL7R, IL1B,
13 TNF- α , IFN γ).

14 Most significant changes toward a protective mode were observed in atherosclerosis,
15 inflammation, and oxidative stress-related genes such as MCP, IL7R, IFN γ , TNF α and the β -
16 adrenergic receptor B2. Monocyte chemoattractant protein-1 (MCP1), also known as CCL2,
17 chemokine C-Cmotif ligand 2, is modulated by the VOO polyphenols within and out of the
18 context of the Mediterranean diet, while TNF- α gene was a point of transcriptional
19 convergence between the Mediterranean diet and VOO intake. MCP1 is a crucial chemokine
20 responsible for the recruitment of monocytes to inflammatory lesions in the vasculature and
21 its decreased expression is a convincing evidence of the anti-inflammatory effect of VOO. In
22 association with other inflammatory mediators, it plays a fundamental role in monocyte
23 chemotaxis to sites of injury and infection; the levels of this protein have been shown to
24 increase in conditions of chronic inflammation and accelerated aging, such as rheumatoid
25 arthritis or lupus [64] and chronic kidney disease [65]. TNF- α , one of the earliest
26 inflammatory cytokines generated during monocyte activation, was downregulated together
27 with interferon gamma (IFN γ) and interleukin-7 receptor (IL7R) in some intervention studies
28 with diets rich in VOO polyphenols [66; 67]. TNF- α sustains the production of late
29 inflammatory cytokines, such as IL-6 [68]. These control the activation of cascades of genes
30 associated with the synthesis of acute phase proteins, endothelial cell activation, metabolism
31 and stress response of tissues.

32 Molecular and cellular effects of oleic acid (C18:1, n-9) have been extensively characterized
33 as this MUFA is relatively abundant in food and is at the same time a product of the cellular
34 biosynthesis of FA, a process activated in response to dietary carbohydrates [69]. ChRBP

1 transcription factor is essential to control the series of genes that implement the biosynthetic
2 steps of this metabolism, i.e. the “de novo lipogenesis” (DNL), starting from the molecular
3 precursor in this polymerization process, e.g. acetyl-CoA. A critical step in oleic acid
4 biosynthesis downstream of palmitic acid (C16:0) formation, the major product of DNL, and
5 elongation to stearic acid (C18:0), is the $\Delta 9$ desaturation which is catalyzed by the enzyme
6 protein Stearoyl-Coenzyme A desaturase [70]. This gene may represent a major player of the
7 lipotoxicity that an increased DNL may generate in cellular systems through the stressogenic
8 activity of palmitic acid, a pro-oxidant and pro-apoptotic agent with proposed pathogenic
9 roles in non-alcoholic steatohepatitis [71]. The capability to synthesize oleic acid at the
10 cellular level is thus indicative of an efficient lipid metabolism that promotes the FA
11 catabolism through the β -oxidation pathway. Accordingly, both gene transcription and
12 lipidomic data demonstrate that oleic acid exerts much less toxicity than SFA, such as
13 palmitic and stearic acid, when assessed in murine and human hepatocytes [70] and in β -cells,
14 the insulin-secreting cellular component of the endocrine pancreas [71; 72]. The same
15 findings have been reported in other cellular models outside of the gastrointestinal tract such
16 as cardiomyocytes [73]. Moreover, in these studies oleic acid and other unsaturated species
17 have been convincingly demonstrated to be protective against the lipotoxicity of palmitic
18 acid.

19 HT, one of the most active biophenols of VOO, has been extensively investigated in a number
20 of in vitro and in vivo studies focused on anti-cancer, anti-inflammatory and cardio-
21 prevention effects (reviewed in [62; 74]). Biochemistry, pharmacokinetics, and toxicology
22 data obtained on HT as minor component of VOO or pure molecule of natural or synthetic
23 origin, point to a use of this biophenol as a potential drug for the chemoprevention of highly
24 prevalent chronic and inflammatory diseases. In a recent study by Giordano et al. [9], mice
25 fed with a diet rich in HT (0.03 % w/w) showed a significant modification of a series of
26 glutathione-related genes in the adipose organ, one of the target organs in cardiometabolic
27 prevention [75]. Among the responding genes the microsomal and cytosolic glutathione S-
28 transferase, selenium and non-selenium glutathione-peroxidases, forms 1 and 7, respectively,
29 and γ -glutamyltransferase 5 were found; this in vivo transcriptional effect of the adipose
30 tissue could produce functional interactions with other components of the detoxification and
31 antioxidant protection system such as superoxide dismutase 1, 2 and 3, and some members of
32 the cytochrome P450 family of genes, namely CYP1A1, 1A2 and 2E1. HT was also observed
33 to influence the redox status of the tripeptide GSH in cultured adipocytes, i.e. the GSH/GSSG
34 ratio, which points to a role of this functional component of VOO in the homeostatic control

1
2
3 1 of the cellular redox at the interface between the master regulators of transcriptional and
4 2 metabolic pathways involved in the stress adaptation response, such as the Nrf2 transcription
5 3 factor, and the pentose phosphate pathway (reviewed in [76]). Altogether these effects are
6 4 crucial to preserve the endocrine and metabolic functions of this organ also including its
7 5 capability to control the compensatory and regenerative processes associated with the
8 6 differentiation of adipose stem cells [75]. Obesity can lead to develop pathologic phenotypes
9 7 in which chronic inflammation can impair these functions of the adipose tissue thus
10 8 increasing the risk of cardiometabolic events and that of other age-related disorders [77].
11 9 Olive oil bioactives are among the dietary phytochemicals that have been described to
12 10 influence the control of “inflammagens” [77; 78]. In a recent comparative study among OO
13 11 polyphenols, Richard et al. [79] described HT as the most effective inhibitor of inflammatory
14 12 pathways that stimulate the production of NO, the eicosanoid PGE₂, and cytokines such as
15 13 IL-1 α , IL-1 β , IL-6, IL-12, TNF- α , and the chemokines CXCL10/IP-10, CCL2/MPC-1.
16 14 Corresponding effects of inhibition were observed in the gene expression of the inducible
17 15 nitric oxide synthase (iNOS), IL-1 α , CXCL10/IP-10, MIP-1 β , matrix metalloproteinase-9,
18 16 and prostaglandin E₂ synthase. The inhibition of COX-2 and iNOS genes, responsible for the
19 17 transcriptional control of TNF- α , was confirmed in other studies in human monocyte-
20 18 macrophages [80; 81; 82] and in vivo in a model of inflammatory response in rats in which
21 19 the inhibitory effects of HT on COX-1 and -2 have been reported to have the same potential
22 20 of the non-steroidal anti-inflammatory drugs ibuprofen and celecoxib [83]. Also the control of
23 21 NF- κ B transcription factor is proposed to play a major role in the anti-inflammatory gene
24 22 response to HT [84]. Moreover, at nutritionally relevant concentrations, HT was shown to
25 23 have additive effects with other VOO components such as OA, in preventing gene defects
26 24 associated with the metabolic defect of inflamed adipocytes, a major event in metabolic
27 25 syndrome and cardiovascular risk [85]. HT and OA synergize in preventing the
28 26 downregulative effect of TNF- α on gene expression and secretion of adiponectin, a
29 27 cardioprotective hormone of the adipose tissue. This effect was caused by this inflammatory
30 28 cytokine through JNK (stress kinase)-mediated suppression of PPAR γ activity.
31 29 The activity of HT on stress-activated kinases, such as JNK, and transcription factors, such as
32 30 NF- κ B and PPAR γ , also suggests effects of this biophenol on the regulation of cell cycle and
33 31 apoptotic pathways, which have sure relevance in the chemo-prevention and possibly
34 32 hormetic role of this compound (Section 4) as well as on anticancer properties described
35 33 below in Sections 6 and 7. Similarly to oleuropein and other related polyphenols, HT
36 34 possesses cytotoxic activity and depending on the experimental model it has been described to

1
2
3 1 inhibit either initiation or promotion/progression phases of carcinogenesis. In fact, this
4 2 molecule can prevent the DNA damage induced by different genotoxic molecules of pre-
5 3 tumoral models and, at the same time, it was found to arrest cell cycle thus inhibiting
6 4 proliferation and inducing apoptosis in different tumors cell lines (Section 6).

7 5 Signaling and gene regulation effects of HT and oleuropein have been consistently
8 6 demonstrated to depend on redox-dependent properties of these molecules that include the
9 7 stimulation of hydrogen peroxide production both at the extracellular and intracellular level
10 8 [86; 87]. However, redox-independent processes could also have a role in the pro-apoptotic
11 9 activity of HT documented on different tumour cells [86].

12 10 According with a redox-dependent mechanism of action within the
13 11 cytoprotection/chemoprevention function of HT, the treatment of cells with this biophenol
14 12 was found to activate a series of antioxidant and detoxification genes, including heme
15 13 oxygenase-1 (15-fold upregulation), glutaredoxin (1.65) and glutathione peroxidase (1.53)
16 14 [88]. These are typical Nrf2 transcription factor-dependent genes that collectively produce the
17 15 adaption response of cells to oxidative stress and, more in general, to noxious stimuli deriving
18 16 from the exposure to electrophiles and lipophilic xenobiotics [76; 89] also including natural
19 17 and food-derived biophenols [90]. Nrf2 operates the transcriptional control of these genes in
20 18 concert with other elements. In the case of the in vitro anti-cancer effects of HT, changes in
21 19 the expression of the transcription factors STAT3, STAT6, SMAD7 and ETS-1 as well as of
22 20 the telomerase subunit TERT have been described. Trans-regulation effects between these
23 21 regulatory elements may occur by means of functional interactions with components of the
24 22 redox sensing and signaling platform of the cell. Glutathione S-transferase P was recently
25 23 observed to represent a protein hub for a redox-sensitive protein interaction network that
26 24 coordinates the transcriptional activity of Nrf2 and STAT3 with the signaling of stress
27 25 kinases, detoxification and redox-regulating enzymes and cell cycle checkpoints [76; 89].
28 26 Such a regulatory network is one of the cellular interactomes with important role in the
29 27 crosstalk between complex cellular responses at the interface between inflammatory pathways
30 28 and oxidative stress [76] that surely deserve further investigation as far as hormetic effects of
31 29 HT and other VOO bioactives may have on human tissues (Section 4).

32 30 Extensive in vitro investigation on anticancer activity (reviewed in [55; 62]) clearly
33 31 demonstrates that HT affects the regulation of genes associated with the arrest of cell cycle
34 32 during G0/G1 or G2/M transitions, which results in cell senescence and activation of the
35 33 canonical (mitochondrial-dependent) pathway of apoptotic cell death. The same signaling and
36 34 cell cycle regulation activity is reported for another functional biocomponent of VOO, e.g.

1
2
3 1 vitamin E, even if the desmethyl and tocotrienol forms of this vitamin seem to possess higher
4 2 activity (reviewed in [91; 92; 93]) when compared with the main form present in VOO,
5 3 namely α -tocopherol (Section 8).
6 4
7 5

6 *Metabolomics, proteomics and epigenetics of VOO bioactives*

7 6 Efforts have been made to identify metabolomics fingerprints of VOO administration to
8 7 humans, alone or in the context of the beneficial effects that the Mediterranean diet is
9 8 proposed to have on human health. The effect of the Mediterranean diet supplemented with
10 9 either EVOO (MD + EVOO) in nondiabetic subjects on the ¹H-NMR urinary metabolome
11 10 was recently investigated in one of the PREDIMED intervention trials with a follow-up of 1
12 11 and 3 years [94]. Potential metabolome biomarker discriminating MD + EVOO from
13 12 baseline and from a low-fat diet (LFD) were concerning the metabolism of carbohydrates (3-
14 13 hydroxybutyrate, citrate, and cis-aconitate), creatine, creatinine, amino acids (proline, N-
15 14 acetylglutamine, glycine, branched-chain amino acids, and derived metabolites), lipids (oleic
16 15 and suberic acids), and microbial cometabolites (phenylacetylglutamine and p-cresol).
17 16 Hippurate, trimethylamine-N-oxide, histidine and derivatives (methylhistidines, carnosine, and
18 17 anserine), and xanthosine were predominant after the administration of the LFD.

19 18 Metabolomics investigations were recently extended to plasma metabolites that may provide
20 19 biomarker and mechanistic cues of the influence of EVOO and MD on cardiovascular risk
21 20 (recently reviewed in [95]). Although still very speculative in nature, such efforts led to
22 21 tentatively identify specific candidates of a targeted investigation that include branched-chain
23 22 and aromatic amino acids, the glutamine-to-glutamate ratio, some short- to medium-chain
24 23 acylcarnitines, gut flora metabolites (choline, betaine, and trimethylamine N-oxide), urea
25 24 cycle metabolites (citrulline and ornithine) and specific lipid subclasses. These candidates
26 25 together with a large number of untargeted metabolites are now under investigation to further
27 26 examine the effects of VOO and other food interventions within the MD dietary pattern on
28 27 CVD risk.

29 28 Next-generation omics are now available in several laboratories, which should help to expand
30 29 and even increase the scientific level of this research. This technology and its extensive
31 30 application in the next years are expected to boost to development of protocols for the
32 31 personalized evaluation of nutritional and health outcomes of dietary patterns and functional
33 32 ingredients with a predictive power that was unimaginable before. These omics approaches
34 33 include the investigation of epigenetic drift (epigenetic modifications as they occur as a direct
35 34 function with age) of blood cells, and its ancillary elements, including platelets, secreted

1
2
3 1 microvesicles (MVs), and microRNA (miRNA) [96]. These are reflective of various diseases
4 2 as well as of lifestyle changes, making them extremely sensitive biomarkers of human health
5 3 and nutrition interventions. Animal studies have ascertained these aspects in the case of VOO
6 4 biophenols. Recently, the effects of these molecules on miRNA and gene modulation have
7 5 been investigated in the mouse brain [97] and findings are compatible with positive regulatory
8 6 effects on neuronal function and synaptic plasticity, leading to improve cognitive, motor and
9 7 emotional behaviour in aged animals treated with these bioactives (this theme is further
10 8 discussed in Section 5).

11 9 Dietary FA also influence miRNA expression of adult rats and to a certain extent of the
12 10 offspring of mothers that consumed different dietary sources of fat including VOO and these
13 11 changes appear to have effects on lipid composition and metabolism of the liver [98]. These
14 12 results point to a major impact of VOO components on epigenetic players that warrant further
15 13 investigation and translation into human clinical trials.

16 14 The impact of VOO phenolics on proteomic profiles of urine and blood was also investigated.
17 15 A recent randomized controlled trial confirmed the positive effect of VOO on a series of
18 16 urinary proteomic biomarkers associated with the cardiovascular risk in healthy volunteers,
19 17 but this effect was independent from the polyphenol content of the administered VOO [99]. In
20 18 the same study, the urinary proteome was not associated with other types of risk such as that
21 19 for chronic kidney disease or diabetes.

22 20 The effects of VOO phenolic compounds on a targeted blood proteome of
23 21 hypercholesterolemic patients, that of HDL apoproteins, was investigated in a recent double-
24 22 blind randomized controlled trial [100]. The comparison with VOO enriched in its phenolics
25 23 or complemented with thyme polyphenols showed minor differences in the quantitative
26 24 changes produced by the reference treatment, i.e. that with VOO (25 ml/day for 3 weeks) thus
27 25 suggesting minor effects on HDL proteome of VOO biophenols. Of the 127 HDL-associated
28 26 proteins identified in the proteomic profiling of these patients, 15 were differently expressed
29 27 after the three VOO interventions compared to baseline, with some quantitative changes
30 28 specific to each treatment. These changes suggest a cardioprotective impact on the HDL
31 29 proteome with the up-regulation of proteins related to cholesterol homeostasis, protection
32 30 against lipoprotein oxidation and blood coagulation while down-regulated proteins are
33 31 implicated in acute-phase response, lipid transport, and immune response. Pathway analysis
34 32 revealed the involvement of these changes in the control of LXR/RXR nuclear receptor, acute
35 33 phase response, and atherosclerosis.

1
2
3 1 The effect of VOO on blood and urinary proteome biomarkers is obviously the result of
4 2 complex effects on tissues and particularly on the liver proteome. Unluckily, these aspects
5 3 have not been investigated in human liver, but in a recent animal study a proteomic approach
6 4 was used to characterize molecular targets of the prevention effects of VOO on fibrotic liver
7 5 damage during chronic exposure to the free radical generating agent CCL4 [101]. In this
8 6 experimental model of inflammation and oxidative stress, the administration of EVOO
9 7 improved histology and molecular markers of lipid peroxidation and fibrosis, with effects on
10 8 the liver expression of proteins associated with antioxidant protection, cellular detoxification,
11 9 and the intermediary metabolism. In detail, the proteins that increased in association with the
12 10 liver protection effect of VOO were: thioredoxin domain-containing protein 12,
13 11 peroxiredoxin-1, thiosulphate sulphurtransferase, calcium-binding protein 1, Annexin A2 and
14 12 heat shock cognate 71 kDa protein. Conversely, COQ9, cAMP-dependent protein kinase type
15 13 I-alpha regulatory subunit, phenylalanine hydroxylase and glycerate kinase were decreased.
16 14 These results point to a major effect of VOO on the liver proteome and possibly on other
17 15 proteomes, such as those of blood plasma and other biological fluids, that may have both
18 16 casual and causal association with metabolic risk factors of age- and lifestyle-dependent
19 17 diseases of the liver.
20 18

21 19 **4. Toward a hormetic role of EVOO phenolic compounds**

22 20 Many biological effects of VOO phenolics (VOOPs) introduced above have been ascribed to
23 21 their antioxidant activity that has been largely demonstrated in vitro [102] and seems to be
24 22 retained in vivo [103; 104]. However, a crucial element in determining VOOPs antioxidant
25 23 activity in vivo is represented by their bioavailability from diet (i.e. to achieve a biological
26 24 effect in a specific tissue or organ, VOOPs should be present in the active form in that tissue
27 25 or organ).

28 26 Several supplementation studies in human and animal models demonstrated that VOOPs are
29 27 rapidly absorbed and undergo to first-pass intestinal/hepatic metabolism [57; 105; 106; 107].
30 28 This process leads to the formation of sulphate and glucuronide conjugates to such an extent
31 29 that the free forms are almost undetectable in body fluids (around 98% of Tyr and HT in
32 30 plasma and urine are present in conjugated forms [108]. The conjugation and the resulting
33 31 loss of the OH groups decreases the radical scavenging activity of Ps, completely modifying
34 32 the antioxidant nature of the parent molecules. Actually, the few studies considering the
35 33 antioxidant activity of VOOPs metabolites demonstrated a general drop in their radical
36 34 scavenging activity. Using a LDL oxidation test and DPPH assay, Khymenets[66]

1 demonstrated that VOOP metabolites (obtained by chemical synthesis) did not possess any
2 significant antioxidant activity when used at physiological concentration. In particular, the
3 glucuronides of Tyr, OH-Tyr and HVA were not able to inhibit Cu-catalyzed LDL oxidation,
4 while the parent molecule OH-Tyr had strong antioxidant effect. Tuck et al. [109] observed
5 contrasting results, but they used VOOPs metabolites purified from human urines; in
6 particular, they observed that Homovannillic Acid and OH-Tyr-3-O-Glucuronide had a high
7 radical scavenging activity (DPPH test), while the sulfate conjugate of OH-Tyr did not
8 possess any radical scavenging activity. Finally, Deiana demonstrated that low concentration
9 of 3 different OH-Tyr glucuronide (obtained by chemical synthesis) were able to slightly
10 protect renal cells against H₂O₂ induced membrane oxidative damage, but they did not exert
11 any significant protection against H₂O₂ induced cellular injury [110].

12 Moreover, due to their low bioavailability, the conjugated forms are present in plasma at very
13 low concentration. In fact, the maximum concentration reached in human plasma by HT
14 metabolites after a single dose supplementation does not exceed 5 µM (Table 1), and similar
15 results have been obtained also after longer supplementation [111]. This concentration
16 appears quite low especially if we compare it to those of the main plasma antioxidants (Table
17 2), with which they should compete.

18 Inside tissues and cells the concentration of PCs is even lower than that present in plasma, and
19 this is quite critical as molecular components of cells and tissues should be key targets of the
20 antioxidant action. Unfortunately, it is very difficult to evaluate the presence of PCs in human
21 tissues and, to the best of our knowledge, there are no human studies on VOOP accumulation
22 in tissues and also animal studies are very limited. Serra et al. [112] quantified VOOP
23 metabolites in different tissues of rats supplemented with a high dose of VOOPs (3 g/kg of
24 phenolic extract from olive cake containing 100 mg PC/g extract). They observed that VOOPs
25 were distributed through the blood stream almost everywhere in the body, however PCs were
26 present especially in their conjugated forms and the tissue concentration was quite low
27 (nmol/g tissue). In a recent paper, tissue uptake of HT was studied in rats after the
28 supplementation of a refined oil containing HT in a dose compatible with human dietary
29 intake (1 mg HT/Kg). The concentration of HT metabolites recovered in rat tissues was even
30 lower (pmol/g tissue) than that previously observed [113]. Finally, in the study of Rublio et
31 al. [114] VOOP metabolites were detected in rat red blood cell (RBC) after an oral
32 administration of a VOO phenolic extract (1.5 g extract/Kg by intragastric gavage). The
33 maximum RBC concentration reached by HT and its metabolites was about 200 nM.
34 Considering the high efficiency of our cellular antioxidant defence system, providing

1
2
3 1 cytosolic GSH at 1-10 mM concentration, such small amount of VOOP metabolites in cells
4 2 should not play a direct influence on the cellular redox.

5 3 To sum up, the bioavailability of these compounds is quite low and the attained
6 4 concentrations (in plasma and tissue) after ingestion appear to be too low to compete with and
7 5 impact on the endogenous antioxidant system and the redox homeostasis of tissues. Therefore,
8 6 a direct antioxidant action in vivo seems quite improbable. However, several supplementation
9 7 studies demonstrated that VOO or VOOP consumption induces an increase antioxidant
10 8 capacity of human plasma [104; 115; 116; 117] and prevents oxidative damage [103].

11 9 To explain this apparent contradiction several hypotheses have been made. First of all, it has
12 10 been proposed that the conjugated metabolites may act as storage forms and that parent
13 11 molecules could be freed from their glucuronide and/or sulphate conjugates in tissues. A
14 12 recent animal study suggested that the de-conjugation process could really occur in vivo; in
15 13 this study, the content of OH-Tyr and OH-Tyr metabolites was measured in rat RBCs before
16 14 and after an oral administration of olive extract. The results show a decrease in the level of
17 15 conjugated forms and a parallel increase of the parent molecule concentrations in RBCs
18 16 suggesting that the OH-Tyr conjugated forms could be hydrolysed intracellularly [114].

19 17 Incorporation of VOOPs into lipoproteins could represent another mechanism through which
20 18 they could prevent the oxidative damage. Several human studies demonstrate that VOO
21 19 consumption increase the ex vivo LDL resistance against oxidation (Table 3). On this basis,
22 20 the EFSA released a claim concerning the benefits of olive oil polyphenols (5 mg/d) for the
23 21 protection of LDL from oxidation [15]. But, if at least 10 μ M HT concentration is needed to
24 22 lower LDL levels, as demonstrated in in vitro studies [106], how the low attained plasma
25 23 concentrations of this PC can explain such results? The answer could be found in the
26 24 formation of molecular interactions between VOOPs and the LDL particle; these interactions
27 25 can determine a much higher local concentration that will be able to protect LDL from
28 26 oxidation. Unfortunately, just few studies have assessed physical and chemical aspects of
29 27 VOOP incorporation into LDL (Table 3), and further results are needed to explore the
30 28 possibility that this incorporation represents a prerequisite for the inhibition of LDL
31 29 oxidation. Because, VOOPs-LDL interaction appears to be transient (time-coinciding with
32 30 the plasma absorption peak) and strongly affected by dialysis procedure (90 % of the HT
33 31 present in LDL disappears within the first 10 minutes of dialysis) [118], the selection of the
32 32 right time point and analytical method used for LDL preparation should be considered as key
33 33 aspects in future human studies.

1
2
3 1 Finally, also it can be hypothesized that the systemic antioxidant effect (increase of
4 2 antioxidant capacity and prevention of oxidative damage) of VOOPs could be the result of an
5 3 indirect effect, which may lead to improve the endogenous antioxidant defence essentially
6 4 through transcriptional effects on phase II drug metabolizing genes [90]. By its own nature,
7 5 antioxidants (PCs among them) are prone to oxidation and their oxidation products if not
8 6 redox cycled can be toxic and then operate as hormetic compounds, activating adaptive
9 7 cellular stress response pathways [119]. Therefore, the oxidized forms of VOOP could induce
10 8 a temporary oxidative stress and cause the activation of signalling pathway that in turn trigger
11 9 the up-regulation of endogenous antioxidant and detoxification enzymes, which may lead to
12 10 higher level of protection against the long-term damaging effects of oxidative stress.

13 11 As introduced above (Section 3), one of the most important pathways of cellular hormesis is
14 12 the Nrf2/antioxidant response element (ARE) system. Nrf2 regulates the expression of phase
15 13 II detoxification and antioxidant genes in response to harmful stimuli and food bioactives
16 14 [90]. Emerging evidence indicates that VOOPs can activate the Nrf2 pathway both in cellular
17 15 models [120; 121; 122] and in vivo [123]. Finally, several human studies demonstrated that
18 16 the consumption of VOO or VOOPs activates this endogenous antioxidant response system
19 17 increasing enzyme activities such as plasma GPX [124; 125] and erythrocyte CAT [126], and
20 18 the levels of GSH [127] and its redox control [116] in blood plasma.

21 19 Then we can speculate that at the low doses reached in vivo VOOPs can activate
22 20 cytoprotective pathways, not acting directly as free radical scavenger, but as a sort of Nrf2-
23 21 targeted “early warning signal” (positive hormetic effect).

22 23 **5. Functional components of olive oil as anti-aging agents: modulation of the miRNome.**

24
25 Epidemiological studies indicate that olive oil consumption is associated with reduced
26 27 mortality and improved cognitive function in elderly subjects at high cardiovascular risk [128;
28 29 129]. Clinical trials have shown that the assumption of these compounds, either in VOO or as
30 31 extracts, induced a decrease in a series of inflammation and oxidative stress parameters in the
32 32 blood [130]. As many of these parameters are associated with cardiovascular risk, and
33 33 considering that cardiovascular pathologies are the first death cause in the over-65 population,
it can be hypothesized that the protection from this particular risk is one of the main
mechanisms through which VOO phenols can reduce both mortality and some age-associated
pathologies.

1
2
3 1 Preclinical animal studies have confirmed the protective effects of olive oil phenols on
4 2 cognitive function, both in normal and in accelerated aging rodent models [131], showing also
5 3 positive effects on motor function and emotional behaviors related to anxiety [132; 133]. HT,
6 4 tyrosol, their sulfate metabolites and oleuropein have been found in the brain after
7 5 administration of VOO extracts to laboratory animals, although in smaller amount as
8 6 compared to other organs [112]. Thus, it is possible that these behavioral actions are also due
9 7 to a direct interaction with the brain tissue.

10 8 Moreover, both in vitro and in vivo evidences indicate that VOO phenolics have the ability to
11 9 modulate cellular pathways that are relevant to the aging process, such as those related to cell
12 10 protection and survival, energy metabolism, and the inflammation process [134].

13 11 While part of these actions can be due to direct interaction with proteins, they can also be
14 12 ascribed to the modulation of gene expression. DNA microarray-based studies have shown
15 13 that a large number of genes is modulated in the aging heart and brain of mice, and that
16 14 dietary interventions have the ability to counteract these changes [135]. Very few works have
17 15 addressed the modulation of gene expression by VOO components during aging. Bayram et
18 16 al. [123] have reported an induction of genes related to antioxidant defenses and longevity in
19 17 SAMP8 mice, a widely used model of accelerated senescence, treated for 4.5 months with a
20 18 diet rich in VOO phenols (10% extra virgin olive oil, EVOO).

21 19 The role of miRNAs, the small non-coding RNAs, in modulating gene expression at the post-
22 20 transcriptional level has gained increasing importance during the last decade, and it has been
23 21 shown that miRNA regulation might also play an important role in the shaping of age-related
24 22 mRNA changes [136]. Furthermore, a dietary intervention, namely calorie restriction, known
25 23 to exert anti-aging actions, has been shown to induce the down-regulation of miR-30e and
26 24 miR-34a, up-regulated in the aging brain [137].

27 25 Recently, it has been shown that olive oil phenols were able to modulate both the gene and
28 26 miRNA expression profiles evaluated by the microarray technique in mice treated from age
29 27 10 to 16 months with an EVOO naturally rich in phenols (H-EVOO) and these changes were
30 28 associated with reduced age-related decline of motor coordination and contextual memory
31 29 [97]. At the end of the treatment, most of the genes modulated by aging resulted to be down-
32 30 regulated and restricted to the cerebral cortex. Compared to L-EVOO (the same oil deprived
33 31 of phenols), the treatment with H-EVOO was instead associated with a significant up-
34 32 regulation of genes associated with synaptic plasticity and with motor and cognitive behavior,
35 33 such as Notch1, BMPs, NGFR, GLP1R and CRT3. The agrin pathway was also
36 34 significantly modulated.

1
2
3 1 Opposite to genes, miRNAs in the cerebral cortex were mostly up-regulated by aging, and the
4 2 treatment with olive oil phenols modified the miRNA profile in aging mice so that it resulted
5 3 very similar to the young mice profile. In particular, 63 miRNAs were significantly down-
6 4 regulated by the H-EVOO treatment [97].

7 5 Further, some of the H-EVOO-modulated miRNAs were found to target genes associated
8 6 with synaptic plasticity and neuronal function protection, whose expression was also modified
9 7 by the treatment. A computational analysis on miRNAs modified according to the changes of
10 8 their respective target genes allowed to identify a further restricted list of 14 age-modulated
11 9 miRNAs, and a partially overlapped list of 6 treatment-modulated miRNAs, shown in Table 4
12 10 (Giovannelli, unpublished data). Among the age-modulated miRNAs, those with the top
13 11 scores were miR-681, -709, -706, all reduced in aging, and -30a-5p, which was up-regulated
14 12 in older mice. Of the 6 treatment-modulated miRNAs, all down-regulated in the H-EVOO
15 13 group, 5 were also up-regulated in aging: miR-30a-5p, -434-5p, -369-5p, -451 and -126-3p.
16 14 At the top ranking score was miR-30a-5p, predicted to control a large numbers of genes
17 15 involved in several pathways, among which axon guidance, ubiquitin-mediated proteolysis,
18 16 regulation of actin cytoskeleton and long-term potentiation. MiR-126, a well-studied miRNA
19 17 in vascular biology, has been found increased in colon-derived myofibroblasts cells upon in
20 18 vitro treatment with wine-derived polyphenols, and demonstrated to be associated to reduced
21 19 expression of inflammatory genes [138]. No association with aging or polyphenols has been
22 20 previously described for miR-434, -369 and -451.

23 21 Among the miRNAs consistently reported to be associated with the aging process in the brain,
24 22 mir-30, mir-34 and mir-124 have been reported to be up-regulated in aged animals, and these
25 23 changes were associated to learning dysfunctions in animal models and in Alzheimer's
26 24 disease [139]. As these and other miRNAs can be down-regulated by dietary interventions,
27 25 the modulation of the miRNome, and of specific miRNAs, by natural compounds might
28 26 represent a novel mechanism of action for nutraceutical compounds, and become part of a
29 27 neuroprotective strategy in the prevention of age-related pathologies.

30 28 Recently, much interest has focused on circulating miRNAs, as potential biomarkers that can
31 29 be measured in plasma by simple PCR techniques. In a future perspective, specific miRNAs
32 30 might serve as markers in intervention trials in elderly humans, to evaluate the extent of age-
33 31 associated dysfunctions and the efficacy of anti-aging treatments.

34 32 In conclusion, olive oil phenols are able to counteract age-induced alterations in brain
35 33 function, and the treatment has proved effective also in animals that were fully adult at the

1
2
3 1 beginning of the intervention. These changes can be related to a complex modulation of gene
4 and miRNA expression patterns.

5 2
6 3 Finally, it is worth mentioning the recent interest in evaluating the effects of exogenous plant-
7 derived miRNAs [140]. Exogenous miRNAs from animal and plant dietary sources can be
8 transported to human blood and tissues through exosomes [141; 142] and the possibility of
9 cross-kingdom gene regulation has been raised. mi-RNAs have been detected with next
10 generation sequencing in olives [143], but up to date the bioavailability of these miRNAs
11 ingested with olives or VOO remains unexplored [144]. Future work is needed to clarify
12 whether part of the beneficial effects of olive oil can also be attributed to such olive-derived
13 miRNAs.

14 11 15 12 **6. Chemoprevention effects of olive oil polyphenols: molecular and cellular mechanisms.**

16 13
17 14 In the last few years, VOOPs have received growing attention because of their healthy
18 properties including the chemopreventive activity. Several processes are essential for cancer
19 development: DNA damage, sustained proliferation and insensitivity to antigrowth signals,
20 evasion of apoptosis, sustained angiogenesis, tissue invasion, metastatization and
21 inflammation [145]. Some VOOPs are capable to affect many if not all these processes thus
22 interfering with the carcinogenic process. This chemopreventive activity of VOOPs is the
23 result of specific gene regulation effects some of which are now identified and are described
24 here in this section.

25 15
26 16 Oxidative DNA damage plays a central role in both the stages of cancer initiation and
27 promotion/progression. The protection effect of different VOOPs against the H₂O₂-induced
28 DNA damage has been investigated in HL60 human lymphoblasts and in peripheral blood
29 mononuclear cells (PBMC). HT (3,4-DHPEA) significantly reduced the extent of DNA
30 damage at concentrations as low as 1 μ M, as evaluated by the single cell gel electrophoresis
31 (SCGE or Comet assay) [146]. Other compounds structurally related to HT showed the same
32 effect on the H₂O₂-induced DNA damage, but the potency of the different compounds
33 changed. In particular, tyrosol (p-HPEA), a phenol compound lacking of the ortho-hydroxyl
34 group on the phenol ring, had less effect than 3,4-DHPEA [146]. These VOOPs were also
35 able to prevent the oxidative DNA damage induced in human lymphocytes by the co-culture
36 with monocyte-macrophages stimulated with PMA, an ex vivo model that mimics the
37 pathophysiology of oxidative stress of an inflammatory lesion [147]. In this experimental
38 system, tyrosol was more effective than HT in preventing oxidative DNA damage [146].

1
2
3 1 Similar results were obtained in Jurkat cells exposed to continuously generated H₂O₂ and
4 2 treated with phenolic extracts from OO and olive mill waste water as well as with isolated
5 3 compounds (HT and caffeic acid) [148]. Additional studies have demonstrated that
6 4 pretreatment of HeLa cells with VOO phenolic extract also prevents DNA damage induced by
7 5 H₂O₂ [149]. The effect of VOO was tested also in vivo. Quiles et al. tested the effects of
8 6 feeding male Wistar rats with diets containing different sources of fat, such as VOO and
9 7 sunflower oil (SO) [150]. Lower levels of DNA double-strand breaks in peripheral blood
10 8 lymphocytes were found in young animals fed on VOO, which reached around one half of the
11 9 damage found in SO treated animals. The same measurements were carried out in aged rats
12 10 showing that the age-related increase of DNA double-strand breaks was significantly lower in
13 11 rats fed a diet containing VOO [150]. Another study from the same group showed the higher
14 12 efficacy of VOO compared to SO in decrease deletions of the mitochondrial genes ND1 and
15 13 ND4 of rat liver [151]. These results strongly suggest that VOOPs may prevent one of the
16 14 main steps in the initiation phase of carcinogenesis, i.e. the oxidative lesion of DNA, a
17 15 process that is responsible for oncogene activation [152].

18 16 Regarding the effects of VOOPs on cancer progression it was found that 3,4-DHPEA reduces
19 17 the proliferation of different human tumour cell lines derived from colon (HCT116, SW480),
20 18 prostate (PC3 and LnCap), breast (MDA and MCF-7) [86] and leukemia (HL60) [153].
21 19 Among the solid tumour cells those derived from breast were the most sensitive to 3,4-
22 20 DHPEA treatment followed by colon and prostate cells. The greatest inhibition of
23 21 proliferation rate was found in HL60 cells (IC₅₀ = 75µM) and this response was associated
24 22 with an increased apoptotic cell death. On the other hand, p-HPEA failed to inhibit cell
25 23 growth and to induce apoptosis in the same experimental conditions [153]. Two secoiridoid
26 24 derivatives of 3,4-DHPEA and pHPEA linked to the dialdehydic form of elenoic acid (EDA)
27 25 and relatively abundant in olive oil were also investigated. 3,4-DHPEA-EDA and pHPEA-
28 26 EDA were able to inhibit the proliferation of HL60 cells and pHPEA-EDA was more potent
29 27 than 3,4-DHPEA-EDA to suppress cell growth and to induce apoptosis in these cells [154].
30 28 Worth of note, the secoiridoid derivatives of 3,4-DHPEA and pHPEA were more efficient
31 29 than phenolics in producing these cellular responses of leukemia cells. In particular, the
32 30 apparent IC₅₀ calculated on cell proliferation for pHPEA-EDA was 10µM while pHPEA
33 31 was ineffective also at 100µM. When pHPEA and 3,4-DHPEA were combined with the
34 32 dialdehydic form of elenoic acid, the physicochemical and structural properties of the
35 33 resulting compounds undergo to major modifications [154]. The molecular and mechanistic
36 34 aspects of the cellular response to 3,4-DHPEA have been investigated in HL60 cells. The

1
2
3 1 expression of some cyclins, kinases cycline dependent (CDK) and inhibitors of kinases
4 2 cycline dependent (CDKi), are under the influence of this molecule, e.g. 3,4-DHPEA or HT
5 3 [155], and demonstrate changes in cell cycle checkpoints that are consistent with a block of
6 4 the cycle in G0/G1 phase. Most relevant changes observed at the proteomic and
7 5 transcriptomic level were an increased expression of cyclin D3 and of the CDKi p21 and p27.
8 6 At the same time, CDK6 decreased and the cyclins E, B1, and A remained unchanged
9 7 together with p15 levels. These findings are suggestive of the inhibitory activity that 3,4-
10 8 DHPEA may exert on Rb phosphorylation, a key pathway in the control of genes responsible
11 9 for DNA synthesis and G1-S transition [156].

12 10 3,4-DHPEA-targeted checkpoints of cell cycle, such as p21 and p27, also provide regulation
13 11 effects on the cellular differentiation process [157]. Accordingly, 3,4-DHPEA similarly to the
14 12 positive controls DMSO and ATRA, induced the granulocytic differentiation of HL60 cells
15 13 [155].

16 14 The same or even more potent activity of 3,4-DHPEA on these anti-cancer effects associated
17 15 with a modified regulation of cell cycle were also observed for other VOO bioactives, such as
18 16 pinoresinol. This is one of the simplest lignans present in olives and VOO, which was found
19 17 to exert a potent anti-proliferative activity both in HL60 cells (IC50% 8 μ M) and in the
20 18 multidrug resistant variant HL60R (IC50% 32 μ M). Again, this effect was associated with a
21 19 block of cell cycle in G0/G1 phase, up-regulation of the CDK inhibitor p21 and induction of
22 20 cellular differentiation [158].

23 22 *Redox-dependent properties of anticancer VOOPs*

24 23 Underlying molecular events in the anticancer function of main VOOPs, such as tyrosol and
25 24 HT, have been reported to include effects on the redox homeostasis of cancer cells (see
26 25 Section 4). Mechanisms of the redox sensing for these molecules may include a direct
27 26 interaction with cellular proteins and lipids at the level of plasmalemma or even
28 27 intracellularly for the VOOP forms with cellular bioavailability. Both the redox-dependent and
29 28 independent interactions with the cellular components are markedly influenced by the
30 29 structural and physico-chemical features of VOOP compounds. For instance, different
31 30 reaction rates with radical probes have been reported for phenolics that show very close
32 31 structural similarity [102] and lipophilic properties of biophenols influences the possibility to
33 32 modify the physical properties of the lipid bilayer or even permeate cellular membranes to a
34 33 relevant extent thus reaching intracellular interactors/sensors. The latter is for instance the
35 34 case of oleocanthal [159], an anti-inflammatory fat-soluble biophenol first identified in VOO

1
2
3 1 by the group of Montedoro and Servili [44]. As far as, the redox properties of 3,4-DHPEA are
4 2 concerned, H₂O₂ was generated and released in the culture medium of cancer cells that
5 3 responded to the anti-proliferative and pro-apoptotic activity of this molecule (16). Steady-
6 4 state levels of H₂O₂ were influenced by the presence of serum and pyruvate in the culture
7 5 medium, and from the ability of cells to remove this oxidant agent. Worth of note, other
8 6 VOOPs structurally related to HT, such as p-HPEA, do not stimulate the production of H₂O₂
9 7 in the culture medium and the apoptotic death of HL60 cells [160].

10 8 Recently, the attention of different groups has been focused on the effects of 3,4-DHPEA on
11 9 different inflammatory pathways either in association with cancerogenesis or other chronic
12 10 diseases such as arthritis, pathologic obesity and atherosclerosis (reviewed in [161; 162]). In
13 11 this regard, it was demonstrated that 3,4-DHPEA reduced the production of superoxide anions
14 12 (O₂⁻), prostaglandin E₂ (PGE₂), and the expression of cyclooxygenase-2 (COX-2) in
15 13 freshly-isolated human monocytes [80]. Similar effects were previously reported in murine
16 14 and human macrophage cell lines [79; 163; 164].

17 15 The decreased generation of O₂⁻ suggests that 3,4-DHPEA can inhibit the activity of the
18 16 enzyme NADPH-oxidase, a membrane protein complex expressed in several cellular systems
19 17 involved in the production of reactive oxygen species at the inflammatory site [165]. This
20 18 effect on O₂⁻ production was also observed with other OOPs with an efficacy that changed
21 19 according with the following order of magnitude: pHPEA-EDA \approx 3,4-DHPEA > 3,4-
22 20 DHPEA-EDA > p-HPEA [80]. In LPS-stimulated human monocytes, 3,4-DHPEA
23 21 significantly decreased PGE₂ production and release in the culture medium. PGE₂ is an
24 22 inflammatory mediator involved in angiogenesis, proliferation, invasion and apoptosis [166;
25 23 167]. This 3,4-DHPEA dependent decrease of PGE₂ secretion resulted from the
26 24 downregulation of COX-2 gene [80]. In the same experimental system, 3,4-DHPEA
27 25 stimulated the production of TNF- α , one of the earliest pro-inflammatory cytokine with key
28 26 role in immune cell activation and cancer mechanisms [168; 169]. The addition of exogenous
29 27 PGE₂ to the cell culture medium hampered this effect, pointing to a PGE₂-mediated control
30 28 of 3,4-DHPEA on TNF- α gene expression and cytokine secretion. Pharmacologic activation
31 29 of adenylate cyclase by Forskolin and the consequent increment of intracellular cAMP,
32 30 reduced the LPS-stimulated secretion of TNF- α in 3,4-DHPEA-treated monocytes [82].
33 31 Therefore, in close functional similarity with non steroidal anti-inflammatory drugs
34 32 (NSAIDs), such as celecoxib and rofecoxib (19), 3,4-DHPEA behaves as a natural anti-
35 33 inflammatory agent decreasing COX-2 activity and PGE₂ production, and increasing the
36 34 secretion of TNF- α in LPS-activated human monocytes.

1
2
3 1 The in vivo anticancer activity of oleuropein, the precursor of olive oil secoiridoids, has been
4 2 also investigated and convincing pre-clinical in vitro findings have been obtained in several
5 3 models of hormone-sensitive cancers (reviewed in [170]). In animal models oleuropein has
6 4 been described to prevent the colitis-associated molecular changes reducing the risk of
7 5 developing colorectal cancer [171]. Recent studies in a MCF-7 breast carcinoma xenograft
8 6 generated in ovariectomised nude mice, demonstrated that an oleuropein-enriched diet
9 7 compared to the baseline control diet decreases the tumour growth rate and significantly
10 8 prevents the formation of both peripulmonary and parenchymal lung metastases [172]. This
11 9 earliest in vivo evidence on the chemopreventive activity of this VOO compound in breast
12 10 cancer is worth of further clinical investigation.

11 12 **7. Olive oil polyphenols and their effect on tumor vascularization and progression**

13
14 14 Angiogenesis is the formation of new blood vessels from the pre-existing vasculature [173;
15 15 174]. It is a physiologic process regulated through the fine balance between stimulatory
16 16 factors, such as Vascular Endothelial Growth Factor (VEGF), basic Fibroblast Growth Factor
17 17 (bFGF), Placental Growth Factor (PlGF), inhibitory factors as Tissue Inhibitor of
18 18 MetalloProteinase (TIMP), Angiopietin-2 (Ang-2), Interferon- α (IFN- α), Thrombospondin
19 19 family of genes (TSP) and other molecular players with accessory pro-angiogenic properties,
20 20 such as endothelial Nitric Oxide Synthase (eNOS) and Cyclooxygenase-2 (COX-2). This
21 21 process could shift to a pathological condition when feedback mechanisms are deficient or
22 22 completely lost, a condition promoted by the tumor microenvironment, that is the cellular
23 23 environment surrounding the tumor, including blood vessels, immune cells, fibroblasts and
24 24 signaling molecules [175; 176]. Tumor and microenvironment are closely related and interact
25 25 constantly by the releasing of extracellular signals that promote tumor angiogenesis and pro-
26 26 tumoral stimuli. Indeed, tumor cells can produce chemokines and cytokines that stimulate
27 27 angiogenesis in terms of growth, progression, and dissemination; these events are grouped
28 28 together under the definition of angiogenic switch, a process that allows the tumor
29 29 transformation from a microscopic lesion with a low malignant potential to a high aggressive
30 30 mass favouring malignancy and metastatization.

31 31 The newly-formed tumor vasculature is structurally and functionally abnormal and it differs
32 32 from the normal vasculature by means of several phenotypic aspects. Blood vessels are leaky,
33 33 tortuous, dilated and show a disorganized pattern of interconnection that witnesses the loss of
34 34 the vascular architecture. The endothelial cells show an aberrant morphology, pericytes (cells

1
2
3 1 that provide support for the endothelial cells and are able to regulate blood flow) are loosely
4 2 attached or absent, and the basement membrane is often abnormal. These structural
5 3 abnormalities contribute to spatial and temporal heterogeneity in tumor blood flow.

6 4 Under physiological situations, such as in wound healing, angiogenesis is closely related with
7 5 the mobilization of inflammatory cells that is a transient process. However, in certain
8 6 pathological conditions, such as in cancer, a continuous recruitment of inflammatory cells is
9 7 observed, which are also a source of ROS. This tight connection between the inflammation-
10 8 dependent generation of ROS and angiogenesis is believed to play a leading role during
11 9 tumorigenesis, from angiogenic switch to metastatization [177].

12 10 Considering the relevant role of angiogenesis in tumor progression, several studies have
13 11 focused on the prevention of the angiogenic switch, e.g. angioprevention [173], which is the
14 12 inhibition of angiogenesis using chemopreventive drugs targeted to key players of this
15 13 process. Accordingly, angiopreventive compounds interrupt autocrine or paracrine stimuli
16 14 generated by the cancer cells or the tumor microenvironment that in turn influence specific
17 15 endothelial pathways, such as inhibition of VEGF-pathway, c-Met (Hepatocyte Growth
18 16 Factor Receptor) axis and COX-2 signaling.

19 17 For instance, Terzuoli [178] demonstrated that DPE (2-(3,4-dihydroxyphenyl)-ethanol), a
20 18 polyphenol present in virgin olive oil, exerts its effects on colon cancer cell growth and
21 19 angiogenesis through the inhibition of Hypoxia Inducible Factor-1alpha (HIF-1 α) pathway.
22 20 HIF-1 α stimulates the migration of mature endothelial cells increasing vascularization of
23 21 hypoxic areas in the tumor microenvironment and this response stimulates the activation of
24 22 microsomal prostaglandin-E synthase-1 and VEGF pathways [179]. In consideration of the
25 23 close relationship between angiogenesis and inflammation in the tumor microenvironment.
26 24 Scoditti et al. [180] investigated the protective role of oleuropein and HT in endothelial
27 25 models after the induction of an inflammatory state with the pro-angiogenic factor phorbol
28 26 12-myristate 13-acetate-PMA. The Authors demonstrated that these polyphenols suppress
29 27 inflammatory angiogenesis attenuating COX-2 expression and the production and release of
30 28 Matrix MetalloProteinase-9 (MMP-9).

31 29 HT, oleic acid and taxifolin have also been studied for their anti-angiogenic potential in
32 30 endothelial cells [181]. Lamy and colleagues demonstrated that these polyphenols present in
33 31 olive oil inhibit angiogenesis downregulating the phosphorylation of VEGF receptor-2 and the
34 32 downstream pathways [182].

35 33 The auto-phosphorylation of different tyrosine residues is linked to a specific effect on
36 34 endothelial cells, for examples, Tyr951 is essential for endothelial cell migration, whereas

1
2
3 1 Tyr1059 is required for VEGF-induced MAPK pathway activation that leads to cell
4 2 proliferation. Secoiridoid derivatives, in particular (-)-oleocanthal, have been assessed for
5 3 their capability to promote chemopreventive effects. In two different studies (-)-oleocanthal
6 4 was demonstrated to exert inhibitory effects towards c-Met 8;9 [183; 184] ; c-Met is a proto-
7 5 oncogene encoding the heterodimeric tyrosine kinase receptor of Hepatocyte Growth Factor
8 6 (HGF) that has been associated with several cellular mechanism, such as cell proliferation,
9 7 epithelial-to-mesenchymal transition (EMT), invasion and metastasis. Furthermore, the
10 8 HGF/c-Met axis represents an additional VEGF-independent mechanism of tumor
11 9 angiogenesis, and its inhibition correlates with repression of angiogenesis.

12 10 Certain anti-angiogenic agents transiently "normalize" the abnormal structure and function of
13 11 tumor vasculature to make them more efficient for oxygen and drug delivery [185]. Drugs
14 12 that are capable to induce vascular normalization can modulate hypoxia increasing the
15 13 efficacy of conventional therapies [185]. Palmieri et al. [186] studied the anoxic environment
16 14 surrounding tumor cells and the correlation of antioxidant activity by olive phenols. They
17 15 demonstrated that polyphenols decreased the anoxia-associated levels of iNOS, COX-2, TNF-
18 16 α , MMP-2/-9 and restored that of the TIMP-1 leading to inhibition of vascular injury and
19 17 correlated hemodynamic alteration (vascular normalization hypothesis [187]).

20 18 All these studies are focused on the chemoprevention properties of VOOPs as anti-angiogenic
21 19 agents that delay or impede the neovascularization by the tumor cells and the tumor
22 20 microenvironment. Early work by some of us demonstrated the anti-angiogenic potential of a
23 21 phenol-rich olive mill wastewater purified extract, an aqueous byproduct of VOO production
24 22 process. We found that this phenol extract inhibits the angiogenic process, lowering the index
25 23 of proliferation, migration/invasion and network formation of endothelial cells both in vitro
26 24 and in vivo [188]. These results further demonstrate the chemoprevention potential of these
27 25 natural functional agents that deserve further exploration in preclinical and clinical protocols
28 26 of therapy for tumor relapse and progression also to face the problem of the severe collateral
29 27 effects that the available anti-angiogenesis drugs are known to produce.

30 **8. Signaling and gene response of vitamin E: a focus on the main fat-soluble vitamin in** 31 **olives and olive oil.**

32
33 33 Vitamin E was discovered in 1922 by Evans and Bishop as a factor essential for rat fertility
34 34 [189]. Nowadays, this term is used to identify a family of eight tocopherols produced in

1 photosynthetic organisms [190], 4 tocopherols (TOH) and 4 tocotrienols (T3) (Figure 2).
2 These molecules share as common structural component a chroman group with a hydroxyl
3 moiety in position 6 and a 13-carbon atom side chain in position 2. This chain has phytyl-
4 derived or geranylgeranyl-derived (or isoprenoid) structure in the subfamily of tocopherols
5 and tocotrienols, respectively. In each of these two subfamilies the four vitamers identified
6 with Greek letters, from alpha to delta, have different methylation patterns of the chroman
7 ring with alpha as the fully methylated (trimethylated) and delta as the desmethyl
8 (monomethylated) form. Both the degree of methylation and the presence of unsaturations in
9 the side chain confer specific biological, physico-chemical and analytical characteristics to the
10 different forms of vitamin E [191].

11 Tocopherols are considerably more widespread in the plant kingdom than tocotrienols [192]
12 and are found in leaves, seeds, roots, tuber, fruits, stems, hypocotyls and cotyledons of higher
13 plants with very heterogeneous concentrations and relative composition of the vitamers [190;
14 192]. In general, α -tocopherol is the main form of tocopherol in photosynthetic tissues. Seeds
15 generally accumulate ten to twenty times more tocopherols than leaves and in some cases,
16 such as Arabidopsis, soy or sunflower seeds, the desmethyl forms of tocopherol largely
17 predominate over the content of α -tocopherol. The presence of tocotrienols in photosynthetic
18 tissues is relatively rare but various tocotrienols can be present in significant amounts in
19 monocot seeds [193].

20 As introduced in Section 2, α -TOH accounts for more than 90% of the vitamin E found in
21 EVOO with key role in the antioxidant protection of other lipid components of this edible oil
22 [20; 41]. Absolute levels of this vitamin however vary substantially in different oil products;
23 in a recent comparative analysis on 430 samples of EVOO showed levels of this vitamin
24 between 23 and 751 mg/kg [17].

25 α -TOH also represents the main form of vitamin E of the human organism (reviewed in [194;
26 195]) and this could be the consequence of a series of physicochemical and biological
27 properties, which are unique among this family of fat-soluble molecules. Convincing
28 evidence was obtained on the fact that α -TOH provides optimal structural interactions with
29 membrane phospholipids. A differential distribution with respect to cholesterol and ceramides
30 has been described to occur on cellular membranes and this may help to regulate fluidity and
31 the functional properties of the different areas of the plasmalemma [34; 196]. Reduction
32 potential and lipid-lipid interaction properties of α -tocopherol are also distinctive of a role for
33 this molecule as the main non-enzymatic antioxidant of cellular membranes. The dynamic
34 interaction of α -TOH with the allylic moieties of unsaturated phospholipids strategically

1
2
3 1 influences the antioxidant protection and the physical stability of the membrane. α -TOH is by
4 2 far the most important hydrogen donating species of biological membranes and in general of
5 3 macromolecular complexes such as the lipoprotein particles [40] and other lipid structures
6 4 even in plant organisms [190]. This is a key and early event in the radical chain-breaking
7 5 activity that non-enzymatic fat-soluble antioxidants embedded in the lipid structure may
8 6 promote. A further requisite for chain breakers is to generate a corresponding radical with
9 7 higher degree of stability with respect to the lipid hydroperoxyl radical formed during free
10 8 radical damage. This non-enzymatic step is a one-electron reaction that in the end promotes
11 9 the formation of a lipid hydroperoxide for a subsequent two-electron step of enzymatic
12 10 reduction that restores the lipid structure of PUFA at the site of free radical attack. In the cell
13 11 membrane this late step is catalysed by the peroxidase activity of enzyme proteins such as the
14 12 selenium-dependent glutathione peroxidase 4 [197]; other glutathione- or thioredoxin-
15 13 dependent enzymatic reactions take place in the different compartments of cells and tissues to
16 14 implement this antioxidant protection of lipid environments, the 1-Cys Peroxiredoxin 6
17 15 reaction is reported as one of these antioxidant enzymes (reviewed in [198]).

18 16 As far as the radical species produced during H-donation reactions of tocopherols concern,
19 17 tocopheryl radicals and their quinone derivatives, such as tocopheryl quinone (TQ), show
20 18 different degree of stability and cytotoxicity. Desmethyl quinone derivatives generate much
21 19 greater toxicity than α -tocopheryl quinone [40; 199]. Such a potentially harmful activity is
22 20 based on peculiar signaling and gene modulation effects that may help to explain the
23 21 beneficial roles so far reported for the desmethyl forms of vitamin E [200; 201]. The same
24 22 type of considerations has been made on the quinone analogues of VOO phenolics [202]. The
25 23 mitochondrial-dependent pro-apoptotic activity of α -tocotrienol in HER2-neu positive breast
26 24 cancer cells and the corresponding in vivo anticancer activity are a specific example of these
27 25 properties [92; 203]. Molecular players of this signaling include among the others,
28 26 phospholipase A2, PKC isoforms and their corresponding phosphatases, pro-survival and
29 27 stress-related MAPKs, and the regulatory components of cell cycle (reviewed in [37; 204]).

30 28 The lower chemoprevention potential of α -TOH suggests a physiological regulatory activity
31 29 in lipid/phosphorylative signalling of cells which is another potential reason for its
32 30 preferential selection and bioavailability to tissues. The physiological role of α -TOH in
33 31 cellular signaling and functional control of biological processes distinct from other forms of
34 32 vitamin E can also be deduced from the gene response of immune cells. In fact, in vivo a high
35 33 intake of α -TOH sustains the gene response of stimulated T-cells thus suggesting immune-
36 34 modulation effects useful to promote host defence and the inflammatory response of tissues

1 [205], a response that was also observed in other experimental models and in humans [206].
2 On the contrary, γ -TOH assessed under the same experimental conditions, repressed T-cell
3 inflammatory genes with higher efficacy than α -TOH, pointing to an anti-inflammatory role
4 for this desmethyl form of vitamin E [205].

5 Emerging evidence demonstrates that α -TOH may exert some of its roles by means of the
6 produce some of its biological roles useful to

7 Therefore, available data are sufficient to propose roles for α -TOH as key cytoprotection
8 factor with homeostatic effects that spread virtually to all the cellular components and
9 biological fluids [207]. Such an important series of functions justifies the definition for this
10 molecule as an “essential factor” also representing a valid reason for the hepatic metabolism
11 to preferentially retain α -TOH and distribute it with the lipoprotein particles to blood and then
12 to peripheral tissues (reviewed in [194; 195]). In fact, its levels in plasma are by far the
13 highest among the fat-soluble molecules reported to possess antioxidant activity with levels
14 that vary depending on gene characteristics and dietary habits from 20 to 60 μ M. Plasma α -
15 TOH is approximately 10 to 20-fold less abundant than α -TOH; δ -TOH shows even lower
16 levels, usually below 0.5 μ M.

17 The selective liver uptake and distribution of α -TOH is under the control of the α -Tocopherol
18 Transfer Protein (α -TTP), a cytosolic protein whose gene defect causes a severe and even
19 fatal form of spinocerebellar ataxia associated with vitamin E deficiency, also known with the
20 term of AVED (OMIM #277460; [208]) that can be successfully treated by vitamin E therapy
21 [209; 210].

22 *Nutritional considerations*

23 Regardless of the variability among VOO products, the intake of α -TOH with this oil within a
24 Mediterranean diet food pattern is of nutritional relevance and this can become even more
25 relevant if the quality of VOO is the highest in terms of vitamin E levels. The amount of
26 tocopherols in other edible oils not common in the Mediterranean diet is between 0.31 and
27 39.2 mg/100 g for α -TOH, 0.92 and 64.9 for γ -TOH, 2.8 and 20.38 for δ -TOH [211]. These
28 composition data support a much higher intake of γ -TOH than α -TOH in regions that
29 consume soy and corn oil as main source of fat in their diets, such as US and Australia [194].
30 High intakes of tocotrienols are found in Asiatic regions cause of the sustained consumption
31 of palm oil or other characteristic oil products such as annatto oil [91].

32 Non-alpha forms and tocotrienols have been reported to promote beneficial effects such as
33 cancer prevention and cholesterol lowering activity [91; 194], but so far the most convincing
34

1 literature on nutritional and health-promoting effects of vitamin E was produced on α -TOH
2 [40]. Along with antioxidant and gene regulation properties reported above for this vitamer, it
3 is reported to have higher biological activity in different tests used to comparatively assess the
4 nutritional properties of the forms of this vitamin such as the rat foetal resorption, H_2O_2 -
5 induced red blood cell haemolysis and prevention of muscular degeneration in animal models
6 of vitamin E deficiency. Accordingly, nutritional recommendations for non-alpha forms of
7 vitamin E are expressed in terms of α -TOH equivalents (one equivalent is defined by the
8 biological activity of 1 mg of RRR- α -TOH in the resorption-gestation test, i.e. β -TOH should
9 be multiplied by 0.5, γ -TOH by 0.1, and α -T3 by 0.3 [212])

10 These aspects altogether support the definition of α -TOH as the actual molecule that provides
11 vitamin E activity to humans and that one on which nutritional recommendations should
12 focus. Although the debate on nutritional requirements and optimal intake of this vitamin is
13 still open [37], the European Food Safety Authority (EFSA) panel on Dietetic Products,
14 Nutrition, and Allergies recently assessed the intakes in healthy populations with no apparent
15 deficiency for α -tocopherol in the EU and used these data to define the Adequate Intakes
16 (AIs) for this vitamin [213]. These were set at 13 mg/day for men and 11 mg/day for women,
17 and at 6 and 9 mg/day for children aged 1 to 3 years and 3 to 10 years, respectively. A value
18 of 5 mg/day was proposed for infants aged 7–11 months and no evidence for an increased
19 intake was reported for pregnant or lactating women. However, recent studies suggest that
20 vitamin E requirements could increase from 15 to 25 mg/d or more, depending on the intake
21 of polyunsaturated fatty acids (PUFA); consequently, additional needs of vitamin E could be
22 considered at least for some individuals [214] and other adjustments on recommended intakes
23 are needed in case of some diseases (reviewed in [37]).

24 From these data it is obvious that a significant proportion of these AI values can be met with
25 few tablespoons of EVOO rich in vitamin E. Therefore, a further reason for nutritionists to
26 recommend EVOO as a health-promoting food is that this can represent a significant dietary
27 source of α -TOH.

28 29 *Emerging aspects of vitamin E biology and function*

30 If on one hand the hepatic metabolism of vitamin E provides a preferential route for α -TOH
31 to be selected and delivered to tissues, on the other hand it may represent a site of biological
32 activation for this molecule as it is for other fat-soluble vitamins such as vitamin A and D
33 (reviewed in [37]). As discussed above, α -TTP and possibly other binding proteins of liver
34 cells select the quota of α -TOH to be retained in the human organism thus directing the

1
2
3 1 excess of this molecule together with the non-alpha forms to hepatic catabolism and excretion
4 2 [37]. In the last years, the metabolites produced during the hepatic catabolism of vitamin E
5 3 have been identified to regulate gene expression and signaling pathways with biological
6 4 effects that are specific or superior than those produced by the vitamin precursor [215; 216;
7 5 217; 218; 219]. The metabolism of vitamin E is initiated by the enzymatic activity of a
8 6 tocopherol ω -oxidase that produces the long-chain metabolite (LCM) 13'-hydroxy-6-
9 7 hydroxychromanol (13'-OH) (Figure 2) [220; 221]. This early metabolite found at low
10 8 nanomolar levels in human serum [191; 217] is further oxidized by the ω -oxidase activity of
11 9 alcohol or aldehyde dehydrogenase enzymes to the corresponding 13'-COOH derivative that
12 10 is the substrate of the β -oxidation like shortage of the side chain to form the final
13 11 carboxyethylchroman metabolite found in plasma and then of urine and bile as main end-
14 12 products of vitamin E catabolism in animal organisms [195; 222; 223]. Convincing evidence
15 13 was obtained on the identification of a form of the cytochrome P450 family of enzymes as the
16 14 main tocopherol ω -oxidase of liver cells, namely the CYP4F2 isoenzyme [221; 224; 225].
17 15 This isoenzyme may represent a point of convergence between the metabolism of vitamin E
18 16 and that of middle and long-chain fatty acids with roles in the inflammatory signalling of cells
19 17 [226]. These include the PUFA arachidonic acid and the eicosanoid derivatives prostaglandin
20 18 A1 and E1, some lipotoxins, the 5- and 12-hydroxyeicosatetraenoic acid (HETE) forms, and
21 19 particularly leukotriene B4, a potent mediator of inflammation. Indeed CYP4F2 gene encodes
22 20 for the leukotriene B4 ω -oxidase 1 enzyme protein and, together with CYP4A11, it is the
23 21 main 20-HETE-synthesizing enzymes in humans, a bioactivation process of arachidonic acid
24 22 substrate with prominent role in the control of renal processes that regulate ion absorption,
25 23 blood flow, vascularization, and thus blood pressure of rodents and possibly humans
26 24 (reviewed in [227]). According with this hypothesis of a functional convergence between the
27 25 metabolism of vitamin E and that of inflammatory lipids, the supplementation of mononuclear
28 26 leukocytes with vitamin E has been described to control inflammatory pathways that include
29 27 NF κ B transcription factor and the enzymes iNOS and COX-2, as well as lipoxygenase
30 28 isoenzymes (reviewed in [37; 228] and references therein). Among the products of this
31 29 metabolism, LCMs of α -TOH and γ -TOH, appear to produce regulatory effects on these
32 30 inflammatory processes that are more potent than those promoted by their vitamin precursor
33 31 forms. Effects on these metabolites [229] and to a lower extent of CEHC metabolites [230;
34 32 231] were also reported on genes responsible of cell cycle regulation. More recently LCMs
35 33 have been found to act in a feed-forward mechanism with the dietary fatty acids oleate and
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 linoleate to control the expression of CYP4F2 gene in human hepatocytes, a regulatory
2 response that is under the transcriptional influence of PPAR γ nuclear receptor [215].
3 Other bioactive metabolites of α -TOH include α -TOH phosphate, the enzymatic product of
4 an unknown cellular kinase that has been identified in tissues and biological fluids [204].
5 The effect of EVOO rich in α -TOH on the circulating level of LCMs in healthy volunteers
6 and fatty liver patients is under investigation (Piroddi et al., unpublished data).
7

For Peer Review

9. Conclusions and perspectives

Olives and OO are a source of bioactive phytonutrients the additive and synergistic effects of which are reported to exert unique and powerful health-promoting effects in antagonism with risk factors for inflammatory and age-dependent conditions such as cardiovascular disease, tumorigenesis and cancer progression, and the cognitive decline of neurodegenerative diseases. Some of these effects have been confirmed in randomized clinical trial and are now the subject of nutritional and health claims endorsed by the institutional authorities in Europe and US. From a nutritional perspective, VOO is a precious source of the MUFA oleic acid and with the average intake of the traditional Mediterranean diet, products with sustained content of α -tocopherol can provide a significant part of, if not all, the adequate intake for this vitamin. These components of VOO are claimed to help regulating the metabolism and antioxidant protection of lipoprotein particles and cellular membranes, thereby leading to a better control of inflammatory and endothelial risk factors for cardiovascular and cerebrovascular disease such as insulin resistance and an increased generation and progression rate of atherosclerotic lesions. Under the form of EVOO, it provides a mixture of biophenols with biological roles that further expand these effects. Antioxidant, anti-inflammatory and anti-cancer have been among the most investigated ones in vitro, in animal models and even in humans.

Health-promoting and nutritional properties of VOO and its functional molecules are now supported by a first set of data coming from nutrigenomics investigations that evaluated this oil alone or within the Mediterranean diet food pattern. These include few but significant gene transcription, metabolomics and proteomics studies carried out in the last few years. The resulting bunch of data tentatively defined a molecular signature for the positive effects of VOO bioactives; biological targets and mechanisms of action are now better understood and available for experimental validation and even clinical application; molecular and cellular targets identified in the nutrigenomics of VOO biophenols are suggestive of specific effects on disease mechanisms. These include receptors, signaling kinases and transcription factors associated with cellular stress and inflammation, lipoprotein damage and endothelial function and more in general with pathways responsible for cell cycle regulation and metabolism that include mitochondrial function and signaling, ER stress, DNA damage, and the response to growth factors, cytokines and hormones (mainly associated with insulin resistance). Efforts have been made to characterize the molecular effects of the individual components and particularly of phenolic compounds such as tyrosol, HT, oleuropein and ligstroside

1
2
3 1 derivatives, hydrocarbons (as squalene), sterols (as β -sitosterol) carotenoids and
4 2 phospholipids. Olive oil phenolics are the most characterized species and available omics data
5 3 clearly confirm the potential for the bioavailable form of these molecules as homeostatic
6 4 factors of cells of the GI tract, particularly of liver and pancreas, as well as of inflammatory
7 5 and vascular cells at the systemic level. Gene expression data are consistent with a role as
8 6 anti-inflammatory and immunomodulating molecules, and recent evidence clearly
9 7 demonstrates for some of these molecules a potent activity as hormetic agents with effects
10 8 on antioxidant and detoxification genes and thus on the control of the cellular redox. Specific
11 9 responses to VOOPs such as HT, oleuropein and oleocanthal have been observed on cell
12 10 cycle regulation and anticancer genes.

13 11 On this ground, VOO-derived nutraceuticals or functional ingredients for food fortification or
14 12 formulation of nutritional supplements, have been developed with proposed use in
15 13 chemoprevention of chronic inflammation and age-related diseases, particularly cancer and
16 14 CVD. These applications however need more extensive evaluation as far as nutritional and
17 15 health effects concern, and such an important evaluation should be implemented at the
18 16 individual level. The perspective of using the available molecular data to predict and even
19 17 assess the gene response to VOO bioactive in protocols of personalized nutrition and
20 18 medicine is very attractive and worth of further investigation. Omics technology is going to
21 19 make this perspective real in the near future.
22 20

1
2
3 **1** **Legends to figures.**
4
5 **2**
6 **3** **Figure 1. Structures of main secoiridoid derivatives and phenyl alcohols of VOO. From**
7 **[102].**
8 **4**
9 **5** **Figure 2. Structures of tocopherols, tocotrienols and their earlier hepatic long-chain**
10 **6** **metabolites.**
11
12 **7**
13
14 **8**
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1. Structures of main secoiridoid derivatives and phenyl alcohols of VOO. From [102].

Figure 2. Structures of tocopherols, tocotrienols and their earlier hepatic long-chain metabolites.

For Peer Review

References

1. Willett, W. C., Sacks, F., Trichopoulou, A., Drescher, G., Ferro-Luzzi, A., Helsing, E. and Trichopoulos, D. (1995) Mediterranean diet pyramid: a cultural model for healthy eating. *Am J Clin Nutr* **61**, 1402S-1406S.
2. Trichopoulou, A., Costacou, T., Bamia, C. and Trichopoulos, D. (2003) Adherence to a Mediterranean diet and survival in a Greek population. *N Engl J Med* **348**, 2599-608.
3. Sofi, F., Macchi, C., Abbate, R., Gensini, G. F. and Casini, A. (2014) Mediterranean diet and health status: an updated meta-analysis and a proposal for a literature-based adherence score. *Public Health Nutr* **17**, 2769-82.
4. Estruch, R., Ros, E., Salas-Salvado, J., Covas, M. I., Corella, D., Aros, F., Gomez-Gracia, E., Ruiz-Gutierrez, V., Fiol, M., Lapetra, J., Lamuela-Raventos, R. M., Serra-Majem, L., Pinto, X., Basora, J., Munoz, M. A., Sorli, J. V., Martinez, J. A. and Martinez-Gonzalez, M. A. (2013) Primary prevention of cardiovascular disease with a Mediterranean diet. *N Engl J Med* **368**, 1279-90.
5. Willett, W. C. (2006) The Mediterranean diet: science and practice. *Public Health Nutr* **9**, 105-10.
6. Turner, R., Etienne, N., Alonso, M. G., de Pascual-Teresa, S., Minihane, A. M., Weinberg, P. D. and Rimbach, G. (2005) Antioxidant and anti-atherogenic activities of olive oil phenolics. *Int J Vitam Nutr Res* **75**, 61-70.
7. Konstantinidou, V., Covas, M. I., Sola, R. and Fito, M. (2013) Up-to date knowledge on the in vivo transcriptomic effect of the Mediterranean diet in humans. *Mol Nutr Food Res* **57**, 772-83.
8. Castaner, O., Corella, D., Covas, M. I., Sorli, J. V., Subirana, I., Flores-Mateo, G., Nonell, L., Bullo, M., de la Torre, R., Portoles, O. and Fito, M. (2013) In vivo transcriptomic profile after a Mediterranean diet in high-cardiovascular risk patients: a randomized controlled trial. *Am J Clin Nutr* **98**, 845-53.
9. Giordano, E., Davalos, A. and Visioli, F. (2014) Chronic hydroxytyrosol feeding modulates glutathione-mediated oxido-reduction pathways in adipose tissue: a nutrigenomic study. *Nutr Metab Cardiovasc Dis* **24**, 1144-50.
10. Martin-Pelaez, S., Castaner, O., Konstantinidou, V., Subirana, I., Munoz-Aguayo, D., Blanchart, G., Gaixas, S., de la Torre, R., Farre, M., Saez, G. T., Nyyssonen, K., Zunft, H. J., Covas, M. I. and Fito, M. (2015) Effect of olive oil phenolic compounds on the expression of blood pressure-related genes in healthy individuals. *Eur J Nutr*.
11. Bayram, B., Esatbeyoglu, T., Schulze, N., Ozcelik, B., Frank, J. and Rimbach, G. (2012) Comprehensive analysis of polyphenols in 55 extra virgin olive oils by HPLC-ECD and their correlation with antioxidant activities. *Plant Foods Hum Nutr* **67**, 326-36.
12. WHO (2015) Healthy diet - Fact sheet N°394. World Health Organization.
13. EFSA Panel on Dietetic Products, N. a. A. N. (2011a) Scientific Opinion on the substantiation of health claims related to oleic acid intended to replace saturated fatty acids (SFAs) in foods or diets and maintenance of normal blood LDL-cholesterol concentrations (ID 673, 728, 729, 1302, 4334) and maintenance of normal (fasting) blood concentrations of triglycerides (ID 673, 4334) pursuant to Article 13(1) of Regulation (EC) No 1924/2006. *EFSA Journal* **9**, 2043.
14. FDA (2004) Monounsaturated fatty acids from olive oil and coronary heart disease - Health Claim Petition Docket No. 2003Q-0559. Vol. No. 2003Q-0559, Rome, Italy.
15. EFSA Panel on Dietetic Products, N. a. A. N. (2011) Scientific opinion on the substantiation of health claims related to polyphenols in olive and protection of LDL particles from oxidative damage (ID 1333, 1638, 1639, 1696, 2865), maintenance of normal blood HDL-cholesterol concentrations (ID 1639), maintenance of normal blood pressure (ID 3781),

- “anti-inflammatory properties” (ID 1882), “contributes to the upper respiratory tract health” (ID 3468), “can help to maintain a normal function of gastrointestinal tract” (3779), and “contributes to body defences against external agents” (ID 3467) pursuant to Article 13(1) of Regulation (EC) No 1924/2006. *EFSA Journal* **9**, 2033
16. European Union (2012) Commission Regulation (EU) No 432/2012 of 16 May 2012 establishing a list of permitted health claims made on foods, other than those referring to the reduction of disease risk and to children’s development and health. *Off. J. Eur. Union* **L136**, 1-40.
 17. Servili, M., Sordini, B., Esposto, S., Urbani, S., Veneziani, G., Di Maio, I., Selvaggini, R. and Taticchi, A. (2013) Biological Activities of Phenolic Compounds of Extra Virgin Olive Oil. *Antioxidants (Basel)* **3**, 1-23.
 18. Bermudez, B., Lopez, S., Ortega, A., Varela, L. M., Pacheco, Y. M., Abia, R. and Muriana, F. J. (2011) Oleic acid in olive oil: from a metabolic framework toward a clinical perspective. *Curr Pharm Des* **17**, 831-43.
 19. Viola, P. and Viola, M. (2009) Virgin olive oil as a fundamental nutritional component and skin protector. *Clin Dermatol* **27**, 159-65.
 20. Inglese, P., Famiani, F., Galvano, F., Servili, M., Esposto, S. and Urbani, S. (2010) Factors Affecting Extra-Virgin Olive Oil Composition. In *Horticultural Reviews*, Vol. 38 Janick, J. ed. John Wiley & Sons, Inc., Hoboken, NJ, USA.
 21. Warleta, F., Campos, M., Allouche, Y., Sanchez-Quesada, C., Ruiz-Mora, J., Beltran, G. and Gaforio, J. J. (2010) Squalene protects against oxidative DNA damage in MCF10A human mammary epithelial cells but not in MCF7 and MDA-MB-231 human breast cancer cells. *Food Chem Toxicol* **48**, 1092-100.
 22. Sanchez-Fidalgo, S., Villegas, I., Rosillo, M. A., Aparicio-Soto, M. and de la Lastra, C. A. (2015) Dietary squalene supplementation improves DSS-induced acute colitis by downregulating p38 MAPK and NFκB signaling pathways. *Mol Nutr Food Res* **59**, 284-92.
 23. Naziri, E., Consonni, R. and Tsimidou, M. Z. (2014) Squalene oxidation products: Monitoring the formation, characterisation and pro-oxidant activity. *European Journal of Lipid Science and Technology* **116**, 1400-1411.
 24. Lopez, S., Bermudez, B., Montserrat-de la Paz, S., Jaramillo, S., Varela, L. M., Ortega-Gomez, A., Abia, R. and Muriana, F. J. (2014) Membrane composition and dynamics: a target of bioactive virgin olive oil constituents. *Biochim Biophys Acta* **1838**, 1638-56.
 25. Newmark, H. L. (1997) Squalene, olive oil, and cancer risk: a review and hypothesis. *Cancer Epidemiol Biomarkers Prev* **6**, 1101-3.
 26. Woyengo, T. A., Ramprasath, V. R. and Jones, P. J. (2009) Anticancer effects of phytosterols. *Eur J Clin Nutr* **63**, 813-20.
 27. Awad, A. B. and Fink, C. S. (2000) Phytosterols as anticancer dietary components: evidence and mechanism of action. *J Nutr* **130**, 2127-30.
 28. Sanchez-Quesada, C., Lopez-Biedma, A., Warleta, F., Campos, M., Beltran, G. and Gaforio, J. J. (2013) Bioactive properties of the main triterpenes found in olives, virgin olive oil, and leaves of *Olea europaea*. *J Agric Food Chem* **61**, 12173-82.
 29. Kalogeropoulos, N. and Tsimidou, M. Z. (2014) Antioxidants in Greek Virgin Olive Oils. *Antioxidants (Basel)* **3**, 387-413.
 30. Habib, L., Jraij, A., Khreich, N., Charcosset, C. and Greige-Gerges, H. (2015) Effect of Erythrodiol, A Natural Pentacyclic Triterpene from Olive Oil, on the Lipid Membrane Properties. *J Membr Biol* **248**, 1079-87.
 31. Rodriguez-Morato, J., Xicotá, L., Fito, M., Farre, M., Dierssen, M. and de la Torre, R. (2015) Potential role of olive oil phenolic compounds in the prevention of neurodegenerative diseases. *Molecules* **20**, 4655-80.

- 1
2
3 1 32. Qian, Y., Guan, T., Tang, X., Huang, L., Huang, M., Li, Y. and Sun, H. (2011) Maslinic acid, a natural triterpenoid compound from *Olea europaea*, protects cortical neurons against oxygen-glucose deprivation-induced injury. *Eur J Pharmacol* **670**, 148-53.
- 4 2
5 3
6 4 33. Qian, Y., Guan, T., Tang, X., Huang, L., Huang, M., Li, Y., Sun, H., Yu, R. and Zhang, F. (2011) Astrocytic glutamate transporter-dependent neuroprotection against glutamate toxicity: an in vitro study of maslinic acid. *Eur J Pharmacol* **651**, 59-65.
- 7 5
8 6
9 7 34. Marquardt, D., Williams, J. A., Kucerka, N., Atkinson, J., Wassall, S. R., Katsaras, J. and Harroun, T. A. (2013) Tocopherol activity correlates with its location in a membrane: a new perspective on the antioxidant vitamin E. *J Am Chem Soc* **135**, 7523-33.
- 10 8
11 9
12 10 35. Perez-Camino, M. C. and Cert, A. (1999) Quantitative determination of hydroxy pentacyclic triterpene acids in vegetable oils. *J Agric Food Chem* **47**, 1558-62.
- 13 11
14 12 36. Moyano, M. J., Heredia, F. J. and Meléndez-Martínez, A. J. (2010) The Color of Olive Oils: The Pigments and Their Likely Health Benefits and Visual and Instrumental Methods of Analysis. *Comprehensive Reviews in Food Science and Food Safety* **9**, 278-291.
- 15 13
16 14 37. Galli, F., Birringer, M., Lorkowski, S., Frank, J., Azzi, A., Traber, M. G., Eggersdorfer, M., Ozer, K. N., Cook-Mills, J. and Cruciani, G. (2016) Vitamin E: emerging aspects and new directions. *Free Radic Biol Med* (Submitted).
- 17 15
18 16
19 17 38. Ghanbari, R., Anwar, F., Alkharfy, K. M., Gilani, A. H. and Saari, N. (2012) Valuable nutrients and functional bioactives in different parts of olive (*Olea europaea* L.)-a review. *Int J Mol Sci* **13**, 3291-340.
- 20 18
21 19
22 20 39. Tsimidou, M. Z. (2012) Virgin Olive Oil (VOO) and Other Olive Tree Products as Sources of α -Tocopherol. Updating and Perspective. In *Tocopherol: Sources, Uses and Health Benefits*, Catala, A. ed, pp. 1-21. Nova Science Publisher, New York, NY, USA.
- 23 21
24 22 40. Traber, M. G. and Atkinson, J. (2007) Vitamin E, antioxidant and nothing more. *Free Radic Biol Med* **43**, 4-15.
- 25 23
26 24 41. EL Riachy, M. E., Priego-Capote, F., León, L., Rallo, L. and Luque de Castro, M. D. (2011) Hydrophilic antioxidants of virgin olive oil. Part 1: Hydrophilic phenols: A key factor for virgin olive oil quality. *European Journal of Lipid Science and Technology* **113**, 678-691.
- 27 25
28 26 42. Obied, H. K., Bedgood, D. R., Jr., Prenzler, P. D. and Robards, K. (2007) Chemical screening of olive biophenol extracts by hyphenated liquid chromatography. *Anal Chim Acta* **603**, 176-89.
- 29 27
30 28 43. Servili, M., Esposto, S., Taticchi, A., Urbani, S., Di Maio, I., Sordini, B. and Selvaggini, R. (2012) The Effect of Diverse Agricultural and Technological Factors on Olive Oil Quality and Yield. Paper presented at the VII International Symposium on Olive Growing, San Juan, Argentina, 2012.
- 31 29
32 30 44. Montedoro, G., Servili, M., Badioli, M., Selvaggini, R., Miniati, E. and Macchioni, A. (1993) Simple and hydrolyzable compounds in virgin olive oil. 3. Spectroscopic characterizations of the secoiridoid derivatives. *J. Agric. Food Chem.* **41**, 2228-2234.
- 33 31
34 32 45. Servili, M., Selvaggini, R., Esposto, S., Taticchi, A., Montedoro, G. and Morozzi, G. (2004) Health and sensory properties of virgin olive oil hydrophilic phenols: agronomic and technological aspects of production that affect their occurrence in the oil. *J Chromatogr A* **1054**, 113-27.
- 35 33
36 34 46. Taticchi, A., Esposto, S. and Servili, M. (2013) The basis of sensory properties of virgin olive oil. In *Olive Oil Sensory Science*, Monteleone, E. and Langstaff, S. eds, pp. 33-54. John Wiley & Sons, Ltd.
- 37 35
38 36 47. Visioli, F. (2012) Olive oil phenolics: where do we stand? Where should we go? *J Sci Food Agric* **92**, 2017-9.
- 39 37
40 38 48. Casaburi, I., Puoci, F., Chimento, A., Sirianni, R., Ruggiero, C., Avena, P. and Pezzi, V. (2013) Potential of olive oil phenols as chemopreventive and therapeutic agents against cancer: a review of in vitro studies. *Mol Nutr Food Res* **57**, 71-83.
- 41 39
42 40
43 41
44 42
45 43
46 44
47 45
48 46
49 47
50 48

- 1
2
3 1 49. Covas, M. I., de la Torre, R. and Fito, M. (2015) Virgin olive oil: a key food for
4 2 cardiovascular risk protection. *Br J Nutr* **113 Suppl 2**, S19-28.
- 5 3 50. Covas, M. I., de la Torre, K., Farre-Albaladejo, M., Kaikkonen, J., Fito, M., Lopez-Sabater,
6 4 C., Pujadas-Bastardes, M. A., Joglar, J., Weinbrenner, T., Lamuela-Raventos, R. M. and de
7 5 la Torre, R. (2006) Postprandial LDL phenolic content and LDL oxidation are modulated by
8 6 olive oil phenolic compounds in humans. *Free Radic Biol Med* **40**, 608-16.
- 9 7 51. Dell'Agli, M., Maschi, O., Galli, G. V., Fagnani, R., Dal Cero, E., Caruso, D. and Bosisio,
10 8 E. (2008) Inhibition of platelet aggregation by olive oil phenols via cAMP-
11 9 phosphodiesterase. *Br J Nutr* **99**, 945-51.
- 12 10 52. Delgado-Lista, J., Lopez-Miranda, J., Cortes, B., Perez-Martinez, P., Lozano, A., Gomez-
13 11 Luna, R., Gomez, P., Gomez, M. J., Criado, J., Fuentes, F. and Perez-Jimenez, F. (2008)
14 12 Chronic dietary fat intake modifies the postprandial response of hemostatic markers to a
15 13 single fatty test meal. *Am J Clin Nutr* **87**, 317-22.
- 16 14 53. Esposito, K., Marfella, R., Ciotola, M., Di Palo, C., Giugliano, F., Giugliano, G.,
17 15 D'Armiento, M., D'Andrea, F. and Giugliano, D. (2004) Effect of a mediterranean-style diet
18 16 on endothelial dysfunction and markers of vascular inflammation in the metabolic
19 17 syndrome: a randomized trial. *JAMA* **292**, 1440-6.
- 20 18 54. Manna, C., Napoli, D., Cacciapuoti, G., Porcelli, M. and Zappia, V. (2009) Olive oil
21 19 phenolic compounds inhibit homocysteine-induced endothelial cell adhesion regardless of
22 20 their different antioxidant activity. *J Agric Food Chem* **57**, 3478-82.
- 23 21 55. Bulotta, S., Oliverio, M., Russo, D. and Procopio, A. (2013) Biological Activity of
24 22 Oleuropein and Its Derivatives. In *Natural Products, Chapter: , Publisher: , Editors: , pp.*,
25 23 Ramawat, K. G. and Merillon, J. M. eds, pp. 3605-3638. Springer-Verlag Berlin Heidelberg.
- 26 24 56. Vissers, M. N., Zock, P. L., Roodenburg, A. J., Leenen, R. and Katan, M. B. (2002) Olive
27 25 oil phenols are absorbed in humans. *J Nutr* **132**, 409-17.
- 28 26 57. Visioli, F., Galli, C., Bornet, F., Mattei, A., Patelli, R., Galli, G. and Caruso, D. (2000)
29 27 Olive oil phenolics are dose-dependently absorbed in humans. *FEBS Lett* **468**, 159-60.
- 30 28 58. Visioli, F., Bellomo, G. and Galli, C. (1998) Free radical-scavenging properties of olive oil
31 29 polyphenols. *Biochem Biophys Res Commun* **247**, 60-4.
- 32 30 59. Miro-Casas, E., Covas, M. I., Farre, M., Fito, M., Ortuno, J., Weinbrenner, T., Roset, P. and
33 31 de la Torre, R. (2003) Hydroxytyrosol disposition in humans. *Clin Chem* **49**, 945-52.
- 34 32 60. Tan, H. W., Tuck, K. L., Stupans, I. and Hayball, P. J. (2003) Simultaneous determination
35 33 of oleuropein and hydroxytyrosol in rat plasma using liquid chromatography with
36 34 fluorescence detection. *J Chromatogr B Analyt Technol Biomed Life Sci* **785**, 187-91.
- 37 35 61. Del Boccio, P., Di Deo, A., De Curtis, A., Celli, N., Iacoviello, L. and Rotilio, D. (2003)
38 36 Liquid chromatography-tandem mass spectrometry analysis of oleuropein and its metabolite
39 37 hydroxytyrosol in rat plasma and urine after oral administration. *J Chromatogr B Analyt
40 38 Technol Biomed Life Sci* **785**, 47-56.
- 41 39 62. Vilaplana-Perez, C., Aunon, D., Garcia-Flores, L. A. and Gil-Izquierdo, A. (2014)
42 40 Hydroxytyrosol and potential uses in cardiovascular diseases, cancer, and AIDS. *Front Nutr*
43 41 **1**, 18.
- 44 42 63. Zhang, L., Li, X., Tai, J., Li, W. and Chen, L. (2012) Predicting candidate genes based on
45 43 combined network topological features: a case study in coronary artery disease. *PLoS One* **7**,
46 44 e39542.
- 47 45 64. Niu, J. and Kolattukudy, P. E. (2009) Role of MCP-1 in cardiovascular disease: molecular
48 46 mechanisms and clinical implications. *Clin Sci (Lond)* **117**, 95-109.
- 49 47 65. Galli, F., Benedetti, S., Floridi, A., Canestrari, F., Piroddi, M., Buoncristiani, E. and
50 48 Buoncristiani, U. (2005) Glycoxidation and inflammatory markers in patients on treatment
51 49 with PMMA-based protein-leaking dialyzers. *Kidney Int* **67**, 750-9.
- 52 50 66. Konstantinidou, V., Covas, M. I., Munoz-Aguayo, D., Khymenets, O., de la Torre, R., Saez,
53 51 G., Tormos Mdel, C., Toledo, E., Marti, A., Ruiz-Gutierrez, V., Ruiz Mendez, M. V. and

- 1
2
3 1 Fito, M. (2010) In vivo nutrigenomic effects of virgin olive oil polyphenols within the frame
4 2 of the Mediterranean diet: a randomized controlled trial. *FASEB J* **24**, 2546-57.
- 5 3 67. Castaner, O., Covas, M. I., Khymenets, O., Nyssonen, K., Konstantinidou, V., Zunft, H. F.,
6 4 de la Torre, R., Munoz-Aguayo, D., Vila, J. and Fito, M. (2012) Protection of LDL from
7 5 oxidation by olive oil polyphenols is associated with a downregulation of CD40-ligand
8 6 expression and its downstream products in vivo in humans. *Am J Clin Nutr* **95**, 1238-44.
- 9 7 68. McInnes, I. B. and Schett, G. (2007) Cytokines in the pathogenesis of rheumatoid arthritis.
10 8 *Nat Rev Immunol* **7**, 429-42.
- 11 9 69. Sae-Lee, C., Moolsuwan, K., Chan, L. and Pongvarin, N. (2016) ChREBP Regulates Itself
12 10 and Metabolic Genes Implicated in Lipid Accumulation in beta-Cell Line. *PLoS One* **11**,
13 11 e0147411.
- 14 12 70. Li, Z. Z., Berk, M., McIntyre, T. M. and Feldstein, A. E. (2009) Hepatic lipid partitioning
15 13 and liver damage in nonalcoholic fatty liver disease: role of stearyl-CoA desaturase. *J Biol*
16 14 *Chem* **284**, 5637-44.
- 17 15 71. Listenberger, L. L., Han, X., Lewis, S. E., Cases, S., Farese, R. V., Jr., Ory, D. S. and
18 16 Schaffer, J. E. (2003) Triglyceride accumulation protects against fatty acid-induced
19 17 lipotoxicity. *Proc Natl Acad Sci U S A* **100**, 3077-82.
- 20 18 72. Plotz, T., Hartmann, M., Lenzen, S. and Elsner, M. (2016) The role of lipid droplet
21 19 formation in the protection of unsaturated fatty acids against palmitic acid induced
22 20 lipotoxicity to rat insulin-producing cells. *Nutr Metab (Lond)* **13**, 16.
- 23 21 73. Bosma, M., Dapito, D. H., Drosatos-Tampakaki, Z., Huiping-Son, N., Huang, L. S.,
24 22 Kersten, S., Drosatos, K. and Goldberg, I. J. (2014) Sequestration of fatty acids in
25 23 triglycerides prevents endoplasmic reticulum stress in an in vitro model of cardiomyocyte
26 24 lipotoxicity. *Biochim Biophys Acta* **1841**, 1648-55.
- 27 25 74. Granados-Principal, S., Quiles, J. L., Ramirez-Tortosa, C. L., Sanchez-Rovira, P. and
28 26 Ramirez-Tortosa, M. C. (2010) Hydroxytyrosol: from laboratory investigations to future
29 27 clinical trials. *Nutr Rev* **68**, 191-206.
- 30 28 75. Murdolo, G., Bartolini, D., Tortoioli, C., Piroddi, M., Iuliano, L. and Galli, F. (2013)
31 29 Lipokines and oxysterols: novel adipose-derived lipid hormones linking adipose dysfunction
32 30 and insulin resistance. *Free Radic Biol Med* **65**, 811-20.
- 33 31 76. Bartolini, D. and Galli, F. (2016) The functional interactome of GSTP: A regulatory
34 32 biomolecular network at the interface with the Nrf2 adaption response to oxidative stress. *J*
35 33 *Chromatogr B Analyt Technol Biomed Life Sci* **1019**, 29-44.
- 36 34 77. Davinelli, S., Maes, M., Corbi, G., Zarrelli, A., Willcox, D. C. and Scapagnini, G. (2016)
37 35 Dietary phytochemicals and neuro-inflammation: from mechanistic insights to translational
38 36 challenges. *Immun Ageing* **13**, 16.
- 39 37 78. Ostan, R., Lanzarini, C., Pini, E., Scurti, M., Vianello, D., Bertarelli, C., Fabbri, C., Izzi, M.,
40 38 Palmas, G., Biondi, F., Martucci, M., Bellavista, E., Salvioli, S., Capri, M., Franceschi, C.
41 39 and Santoro, A. (2015) Inflammation and cancer: a challenge for the Mediterranean diet.
42 40 *Nutrients* **7**, 2589-621.
- 43 41 79. Richard, N., Arnold, S., Hoeller, U., Kilpert, C., Wertz, K. and Schwager, J. (2011)
44 42 Hydroxytyrosol is the major anti-inflammatory compound in aqueous olive extracts and
45 43 impairs cytokine and chemokine production in macrophages. *Planta Med* **77**, 1890-7.
- 46 44 80. Rosignoli, P., Fuccelli, R., Fabiani, R., Servili, M. and Morozzi, G. (2013) Effect of olive oil
47 45 phenols on the production of inflammatory mediators in freshly isolated human monocytes.
48 46 *J Nutr Biochem* **24**, 1513-9.
- 49 47 81. Zhang, X., Cao, J., Jiang, L. and Zhong, L. (2009) Suppressive effects of hydroxytyrosol on
50 48 oxidative stress and nuclear Factor-kappaB activation in THP-1 cells. *Biol Pharm Bull* **32**,
51 49 578-82.

- 1
2
3 1 82. Fuccelli, R., Fabiani, R., Sepporta, M. V. and Rosignoli, P. (2015) The hydroxytyrosol-
4 2 dependent increase of TNF-alpha in LPS-activated human monocytes is mediated by PGE2
5 3 and adenylate cyclase activation. *Toxicol In Vitro* **29**, 933-7.
- 6 4 83. Procopio, A., Alcaro, S., Nardi, M., Oliverio, M., Ortuso, F., Sacchetta, P., Pieragostino, D.
7 5 and Sindona, G. (2009) Synthesis, biological evaluation, and molecular modeling of
8 6 oleuropein and its semisynthetic derivatives as cyclooxygenase inhibitors. *J Agric Food*
9 7 *Chem* **57**, 11161-7.
- 10 8 84. Killeen, M. J., Linder, M., Pontoniere, P. and Crea, R. (2014) NF-kappabeta signaling and
11 9 chronic inflammatory diseases: exploring the potential of natural products to drive new
12 10 therapeutic opportunities. *Drug Discov Today* **19**, 373-8.
- 13 11 85. Scoditti, E., Massaro, M., Carluccio, M. A., Pellegrino, M., Wabitsch, M., Calabriso, N.,
14 12 Storelli, C. and De Caterina, R. (2015) Additive regulation of adiponectin expression by the
15 13 mediterranean diet olive oil components oleic Acid and hydroxytyrosol in human
16 14 adipocytes. *PLoS One* **10**, e0128218.
- 17 15 86. Fabiani, R., Sepporta, M. V., Rosignoli, P., De Bartolomeo, A., Crescimanno, M. and
18 16 Morozzi, G. (2012) Anti-proliferative and pro-apoptotic activities of hydroxytyrosol on
19 17 different tumour cells: the role of extracellular production of hydrogen peroxide. *Eur J Nutr*
20 18 **51**, 455-64.
- 21 19 87. Odiatou, E. M., Skaltsounis, A. L. and Constantinou, A. I. (2013) Identification of the
22 20 factors responsible for the in vitro pro-oxidant and cytotoxic activities of the olive
23 21 polyphenols oleuropein and hydroxytyrosol. *Cancer Lett* **330**, 113-21.
- 24 22 88. Rafehi, H., Smith, A. J., Balcerczyk, A., Ziemann, M., Ooi, J., Loveridge, S. J., Baker, E.
25 23 K., El-Osta, A. and Karagiannis, T. C. (2012) Investigation into the biological properties of
26 24 the olive polyphenol, hydroxytyrosol: mechanistic insights by genome-wide mRNA-Seq
27 25 analysis. *Genes Nutr* **7**, 343-55.
- 28 26 89. Bartolini, D., Comodi, J., Piroddi, M., Incipini, L., Sancineto, L., Santi, C. and Galli, F.
29 27 (2015) Glutathione S-transferase pi expression regulates the Nrf2-dependent response to
30 28 hormetic diselenides. *Free Radic Biol Med* **88**, 466-80.
- 31 29 90. Galli, F. (2007) Interactions of polyphenolic compounds with drug disposition and
32 30 metabolism. *Curr Drug Metab* **8**, 830-8.
- 33 31 91. Viola, V., Pilolli, F., Piroddi, M., Pierpaoli, E., Orlando, F., Provinciali, M., Betti, M.,
34 32 Mazzini, F. and Galli, F. (2012) Why tocotrienols work better: insights into the in vitro anti-
35 33 cancer mechanism of vitamin E. *Genes Nutr* **7**, 29-41.
- 36 34 92. Pierpaoli, E., Viola, V., Barucca, A., Orlando, F., Galli, F. and Provinciali, M. (2013) Effect
37 35 of annatto-tocotrienols supplementation on the development of mammary tumors in HER-
38 36 2/neu transgenic mice. *Carcinogenesis* **34**, 1352-60.
- 39 37 93. Mazzini, F., Betti, M., Canonico, B., Netscher, T., Luchetti, F., Papa, S. and Galli, F. (2010)
40 38 Anticancer activity of vitamin E-derived compounds in murine C6 glioma cells.
41 39 *ChemMedChem* **5**, 540-3.
- 42 40 94. Vazquez-Fresno, R., Llorach, R., Urpi-Sarda, M., Lupianez-Barbero, A., Estruch, R.,
43 41 Corella, D., Fito, M., Aros, F., Ruiz-Canela, M., Salas-Salvado, J. and Andres-Lacueva, C.
44 42 (2015) Metabolomic pattern analysis after mediterranean diet intervention in a nondiabetic
45 43 population: a 1- and 3-year follow-up in the PREDIMED study. *J Proteome Res* **14**, 531-40.
- 46 44 95. Martinez-Gonzalez, M. A., Ruiz-Canela, M., Hruby, A., Liang, L., Trichopoulou, A. and
47 45 Hu, F. B. (2016) Intervention Trials with the Mediterranean Diet in Cardiovascular
48 46 Prevention: Understanding Potential Mechanisms through Metabolomic Profiling. *J Nutr.*
- 49 47 96. Wallace, R. G., Twomey, L. C., Custaud, M. A., Moyna, N., Cummins, P. M., Mangone, M.
50 48 and Murphy, R. P. (2016) Potential Diagnostic and Prognostic Biomarkers of Epigenetic
51 49 Drift within the Cardiovascular Compartment. *Biomed Res Int* **2016**, 2465763.

- 1
2
3 1 97. Luceri, C., Bigagli, E., Pitozzi, V. and Giovannelli, L. (2015) A nutrigenomics approach for
4 2 the study of anti-aging interventions: olive oil phenols and the modulation of gene and
5 3 microRNA expression profiles in mouse brain. *Eur J Nutr*.
- 6 4 98. Casas-Agustench, P., Fernandes, F. S., Tavares do Carmo, M. G., Visioli, F., Herrera, E. and
7 5 Davalos, A. (2015) Consumption of distinct dietary lipids during early pregnancy
8 6 differentially modulates the expression of microRNAs in mothers and offspring. *PLoS One*
9 7 **10**, e0117858.
- 10 8 99. Silva, S., Bronze, M. R., Figueira, M. E., Siwy, J., Mischak, H., Combet, E. and Mullen, W.
11 9 (2015) Impact of a 6-wk olive oil supplementation in healthy adults on urinary proteomic
12 10 biomarkers of coronary artery disease, chronic kidney disease, and diabetes (types 1 and 2):
13 11 a randomized, parallel, controlled, double-blind study. *Am J Clin Nutr* **101**, 44-54.
- 14 12 100. Pedret, A., Catalan, U., Fernandez-Castillejo, S., Farras, M., Valls, R. M., Rubio, L., Canela,
15 13 N., Aragonés, G., Romeu, M., Castaner, O., de la Torre, R., Covas, M. I., Fito, M., Motilva,
16 14 M. J. and Sola, R. (2015) Impact of Virgin Olive Oil and Phenol-Enriched Virgin Olive Oils
17 15 on the HDL Proteome in Hypercholesterolemic Subjects: A Double Blind, Randomized,
18 16 Controlled, Cross-Over Clinical Trial (VOHF Study). *PLoS One* **10**, e0129160.
- 19 17 101. Wang, H., Sit, W. H., Tipoe, G. L. and Wan, J. M. (2014) Differential protective effects of
20 18 extra virgin olive oil and corn oil in liver injury: a proteomic study. *Food Chem Toxicol* **74**,
21 19 131-8.
- 22 20 102. Servili, M., Esposto, S., Fabiani, R., Urbani, S., Taticchi, A., Mariucci, F., Selvaggini, R.
23 21 and Montedoro, G. F. (2009) Phenolic compounds in olive oil: antioxidant, health and
24 22 organoleptic activities according to their chemical structure. *Inflammopharmacology* **17**, 76-
25 23 84.
- 26 24 103. Salvini, S., Sera, F., Caruso, D., Giovannelli, L., Visioli, F., Saieva, C., Masala, G., Ceroti,
27 25 M., Giovacchini, V., Pitozzi, V., Galli, C., Romani, A., Mulinacci, N., Bortolomeazzi, R.,
28 26 Dolara, P. and Palli, D. (2006) Daily consumption of a high-phenol extra-virgin olive oil
29 27 reduces oxidative DNA damage in postmenopausal women. *Br J Nutr* **95**, 742-51.
- 30 28 104. Visioli, F., Caruso, D., Grande, S., Bosisio, R., Villa, M., Galli, G., Sirtori, C. and Galli, C.
31 29 (2005) Virgin Olive Oil Study (VOLOS): vasoprotective potential of extra virgin olive oil in
32 30 mildly dyslipidemic patients. *Eur J Nutr* **44**, 121-7.
- 33 31 105. Miro Casas, E., Farre Albadalejo, M., Covas Planells, M. I., Fito Colomer, M., Lamuela
34 32 Raventos, R. M. and de la Torre Fornell, R. (2001) Tyrosol bioavailability in humans after
35 33 ingestion of virgin olive oil. *Clin Chem* **47**, 341-3.
- 36 34 106. Visioli, F., Bellomo, G., Montedoro, G. and Galli, C. (1995) Low density lipoprotein
37 35 oxidation is inhibited in vitro by olive oil constituents. *Atherosclerosis* **117**, 25-32.
- 38 36 107. Weinbrenner, T., Fito, M., Farre Albaladejo, M., Saez, G. T., Rijken, P., Tormos, C.,
39 37 Coolen, S., De La Torre, R. and Covas, M. I. (2004) Bioavailability of phenolic compounds
40 38 from olive oil and oxidative/antioxidant status at postprandial state in healthy humans.
41 39 *Drugs Exp Clin Res* **30**, 207-12.
- 42 40 108. de la Torre, R. (2008) Bioavailability of olive oil phenolic compounds in humans.
43 41 *Inflammopharmacology* **16**, 245-7.
- 44 42 109. Tuck, K. L., Hayball, P. J. and Stupans, I. (2002) Structural characterization of the
45 43 metabolites of hydroxytyrosol, the principal phenolic component in olive oil, in rats. *J Agric*
46 44 *Food Chem* **50**, 2404-9.
- 47 45 110. Deiana, M., Incani, A., Rosa, A., Atzeri, A., Loru, D., Cabboi, B., Paola Melis, M., Lucas,
48 46 R., Morales, J. C. and Assunta Dessi, M. (2011) Hydroxytyrosol glucuronides protect renal
49 47 tubular epithelial cells against H₂O₂ induced oxidative damage. *Chem Biol Interact* **193**,
50 48 232-9.
- 51 49 111. Oliveras-Lopez, M.-J., Berna, G., Jurado-Ruiz, E., Lopez-Garcia de la Serrana, H. and
52 50 Martin, F. (2014) Consumption of extra-virgin olive oil rich in phenolic compounds has

- beneficial antioxidant effects in healthy human adults. *Journal of Functional Foods* **10**, 475-484.
112. Serra, A., Rubio, L., Borrás, X., Macià, A., Romero, M. P. and Motilva, M. J. (2012) Distribution of olive oil phenolic compounds in rat tissues after administration of a phenolic extract from olive cake. *Mol Nutr Food Res* **56**, 486-96.
113. Lopez de las Hazas, M. C., Rubio, L., Kotronoulas, A., de la Torre, R., Sola, R. and Motilva, M. J. (2015) Dose effect on the uptake and accumulation of hydroxytyrosol and its metabolites in target tissues in rats. *Mol Nutr Food Res* **59**, 1395-9.
114. Rubio, L., Serra, A., Macià, A., Pinol, C., Romero, M.-P. and Motilva, M.-J. (2014) In vivo distribution and deconjugation of hydroxytyrosol phase II metabolites in red blood cells: A potential new target for hydroxytyrosol. *Journal of Functional Foods* **10**, 139-143.
115. Bonanome, A., Pagnan, A., Caruso, D., Toia, A., Xamin, A., Fedeli, E., Berra, B., Zamburlini, A., Ursini, F. and Galli, G. (2000) Evidence of postprandial absorption of olive oil phenols in humans. *Nutr Metab Cardiovasc Dis* **10**, 111-20.
116. Covas, M. I., Nyssonen, K., Poulsen, H. E., Kaikkonen, J., Zunft, H. J., Kiesewetter, H., Gaddi, A., de la Torre, R., Mursu, J., Baumler, H., Nascetti, S., Salonen, J. T., Fito, M., Virtanen, J., Marrugat, J. and Group, E. S. (2006) The effect of polyphenols in olive oil on heart disease risk factors: a randomized trial. *Ann Intern Med* **145**, 333-41.
117. Razquin, C., Martínez, J. A., Martínez-González, M. A., Mitjavila, M. T., Estruch, R. and Martí, A. (2009) A 3 years follow-up of a Mediterranean diet rich in virgin olive oil is associated with high plasma antioxidant capacity and reduced body weight gain. *Eur J Clin Nutr* **63**, 1387-93.
118. Gonzalez-Santiago, M., Fonolla, J. and Lopez-Huertas, E. (2010) Human absorption of a supplement containing purified hydroxytyrosol, a natural antioxidant from olive oil, and evidence for its transient association with low-density lipoproteins. *Pharmacol Res* **61**, 364-70.
119. Forman, H. J., Davies, K. J. and Ursini, F. (2014) How do nutritional antioxidants really work: nucleophilic tone and para-hormesis versus free radical scavenging in vivo. *Free Radic Biol Med* **66**, 24-35.
120. Martín, M. A., Ramos, S., Granado-Serrano, A. B., Rodríguez-Ramiro, I., Trujillo, M., Bravo, L. and Goya, L. (2010) Hydroxytyrosol induces antioxidant/detoxicant enzymes and Nrf2 translocation via extracellular regulated kinases and phosphatidylinositol-3-kinase/protein kinase B pathways in HepG2 cells. *Mol Nutr Food Res* **54**, 956-66.
121. Zou, X., Feng, Z., Li, Y., Wang, Y., Wertz, K., Weber, P., Fu, Y. and Liu, J. (2012) Stimulation of GSH synthesis to prevent oxidative stress-induced apoptosis by hydroxytyrosol in human retinal pigment epithelial cells: activation of Nrf2 and JNK-p62/SQSTM1 pathways. *J Nutr Biochem* **23**, 994-1006.
122. Parzonko, A., Czerwinska, M. E., Kiss, A. K. and Naruszewicz, M. (2013) Oleuropein and oleacein may restore biological functions of endothelial progenitor cells impaired by angiotensin II via activation of Nrf2/heme oxygenase-1 pathway. *Phytomedicine* **20**, 1088-94.
123. Bayram, B., Ozcelik, B., Grimm, S., Roeder, T., Schrader, C., Ernst, I. M., Wagner, A. E., Grune, T., Frank, J. and Rimbach, G. (2012) A diet rich in olive oil phenolics reduces oxidative stress in the heart of SAMP8 mice by induction of Nrf2-dependent gene expression. *Rejuvenation Res* **15**, 71-81.
124. Weinbrenner, T., Fito, M., de la Torre, R., Saez, G. T., Rijken, P., Tormos, C., Coolen, S., Albaladejo, M. F., Abanades, S., Schroder, H., Marrugat, J. and Covas, M. I. (2004) Olive oils high in phenolic compounds modulate oxidative/antioxidative status in men. *J Nutr* **134**, 2314-21.
125. Fito, M., Cladellas, M., de la Torre, R., Martí, J., Alcantara, M., Pujadas-Bastardes, M., Marrugat, J., Bruguera, J., Lopez-Sabater, M. C., Vila, J., Covas, M. I. and members of the,

- 1
2
3 1 S. I. (2005) Antioxidant effect of virgin olive oil in patients with stable coronary heart
4 2 disease: a randomized, crossover, controlled, clinical trial. *Atherosclerosis* **181**, 149-58.
- 5 3 126. Oliveras-Lopez, M. J., Molina, J. J., Mir, M. V., Rey, E. F., Martin, F. and de la Serrana, H.
6 4 L. (2013) Extra virgin olive oil (EVOO) consumption and antioxidant status in healthy
7 5 institutionalized elderly humans. *Arch Gerontol Geriatr* **57**, 234-42.
- 8 6 127. Visioli, F., Wolfram, R., Richard, D., Abdullah, M. I. and Crea, R. (2009) Olive phenolics
9 7 increase glutathione levels in healthy volunteers. *J Agric Food Chem* **57**, 1793-6.
- 10 8 128. Buckland, G., Mayen, A. L., Agudo, A., Travier, N., Navarro, C., Huerta, J. M., Chirlaque,
11 9 M. D., Barricarte, A., Ardanaz, E., Moreno-Iribas, C., Marin, P., Quiros, J. R., Redondo, M.-
12 10 L., Amiano, P., Dorronsoro, M., Arriola, L., Molina, E., Sanchez, M.-J. and Gonzalez, C. A.
13 11 (2012) Olive oil intake and mortality within the Spanish population (EPIC-Spain). *The*
14 12 *American journal of clinical nutrition* **96**, 142-9.
- 15 13 129. Valls-Pedret, C., Lamuela-Raventos, R. M., Medina-Remon, A., Quintana, M., Corella, D.,
16 14 Pinto, X., Martinez-Gonzalez, M. A., Estruch, R. and Ros, E. (2012) Polyphenol-rich foods
17 15 in the Mediterranean diet are associated with better cognitive function in elderly subjects at
18 16 high cardiovascular risk. *Journal of Alzheimer's disease : JAD* **29**, 773-82.
- 19 17 130. Martin-Pelaez, S., Covas, M. I., Fito, M., Kusar, A. and Pravst, I. (2013) Health effects of
20 18 olive oil polyphenols: recent advances and possibilities for the use of health claims.
21 19 *Molecular nutrition & food research* **57**, 760-71.
- 22 20 131. Farr, S. A., Price, T. O., Dominguez, L. J., Motisi, A., Saiano, F., Niehoff, M. L., Morley, J.
23 21 E., Banks, W. A., Ercal, N. and Barbagallo, M. (2012) Extra virgin olive oil improves
24 22 learning and memory in SAMP8 mice. *Journal of Alzheimer's disease : JAD* **28**, 81-92.
- 25 23 132. Pitozzi, V., Jacomelli, M., Zaid, M., Luceri, C., Bigagli, E., Lodovici, M., Ghelardini, C.,
26 24 Vivoli, E., Norcini, M., Gianfriddo, M., Esposto, S., Servili, M., Morozzi, G., Baldi, E.,
27 25 Bucherelli, C., Dolara, P. and Giovannelli, L. (2010) Effects of dietary extra-virgin olive oil
28 26 on behaviour and brain biochemical parameters in ageing rats. *The British journal of*
29 27 *nutrition* **103**, 1674-83.
- 30 28 133. Pitozzi, V., Jacomelli, M., Catelan, D., Servili, M., Taticchi, A., Biggeri, A., Dolara, P. and
31 29 Giovannelli, L. (2012) Long-term dietary extra-virgin olive oil rich in polyphenols reverses
32 30 age-related dysfunctions in motor coordination and contextual memory in mice: role of
33 31 oxidative stress. *Rejuvenation research* **15**, 601-12.
- 34 32 134. Giovannelli, L. (2013) Beneficial effects of olive oil phenols on the aging process:
35 33 Experimental evidence and possible mechanisms of action. *Nutrition and Aging* **1**, 24.
- 36 34 135. Park, S.-K., Kim, K., Page, G. P., Allison, D. B., Weindruch, R. and Prolla, T. A. (2009)
37 35 Gene expression profiling of aging in multiple mouse strains: identification of aging
38 36 biomarkers and impact of dietary antioxidants. *Aging cell* **8**, 484-95.
- 39 37 136. Somel, M., Guo, S., Fu, N., Yan, Z., Hu, H. Y., Xu, Y., Yuan, Y., Ning, Z., Hu, Y., Menzel,
40 38 C., Hu, H., Lachmann, M., Zeng, R., Chen, W. and Khaitovich, P. (2010) MicroRNA,
41 39 mRNA, and protein expression link development and aging in human and macaque brain.
42 40 *Genome research* **20**, 1207-18.
- 43 41 137. Khanna, A., Muthusamy, S., Liang, R., Sarojini, H. and Wang, E. (2011) Gain of survival
44 42 signaling by down-regulation of three key miRNAs in brain of calorie-restricted mice. *Aging*
45 43 **3**, 223-36.
- 46 44 138. Angel-Morales, G., Noratto, G. and Mertens-Talcott, S. (2012) Red wine polyphenolics
47 45 reduce the expression of inflammation markers in human colon-derived CCD-18Co
48 46 myofibroblast cells: potential role of microRNA-126. *Food & function* **3**, 745-52.
- 49 47 139. Zovoilis, A., Agbemenyah, H. Y., Agis-Balboa, R. C., Stilling, R. M., Edbauer, D., Rao, P.,
50 48 Farinelli, L., Delalle, I., Schmitt, A., Falkai, P., Bahari-Javan, S., Burkhardt, S.,
51 49 Sananbenesi, F. and Fischer, A. (2011) microRNA-34c is a novel target to treat dementias.
52 50 *The EMBO journal* **30**, 4299-308.

- 1
2
3 1 140. Wagner, A. E., Piegholdt, S., Ferraro, M., Pallauf, K. and Rimbach, G. (2015) Food derived
4 2 microRNAs. *Food Funct* **6**, 714-8.
- 5 3 141. Baier, S. R., Nguyen, C., Xie, F., Wood, J. R. and Zempleni, J. (2014) MicroRNAs are
6 4 absorbed in biologically meaningful amounts from nutritionally relevant doses of cow milk
7 5 and affect gene expression in peripheral blood mononuclear cells, HEK-293 kidney cell
8 6 cultures, and mouse livers. *The Journal of nutrition* **144**, 1495-500.
- 9 7 142. Zhang, L., Hou, D., Chen, X., Li, D., Zhu, L., Zhang, Y., Li, J., Bian, Z., Liang, X., Cai, X.,
10 8 Yin, Y., Wang, C., Zhang, T., Zhu, D., Zhang, D., Xu, J., Chen, Q., Ba, Y., Liu, J., Wang,
11 9 Q., Chen, J., Wang, J., Wang, M., Zhang, Q., Zhang, J., Zen, K. and Zhang, C. Y. (2012)
12 10 Exogenous plant MIR168a specifically targets mammalian LDLRAP1: evidence of cross-
13 11 kingdom regulation by microRNA. *Cell Res* **22**, 107-26.
- 14 12 143. Donaire, L., Pedrola, L., Rosa Rde, L. and Llave, C. (2011) High-throughput sequencing of
15 13 RNA silencing-associated small RNAs in olive (*Olea europaea* L.). *PLoS One* **6**, e27916.
- 16 14 144. Mico, V., Martin, R., Lasuncion, M. A., Ordovas, J. M. and Daimiel, L. (2016)
17 15 Unsuccessful Detection of Plant MicroRNAs in Beer, Extra Virgin Olive Oil and Human
18 16 Plasma After an Acute Ingestion of Extra Virgin Olive Oil. *Plant Foods Hum Nutr* **71**, 102-
19 17 8.
- 20 18 145. Hanahan, D. and Weinberg, R. A. (2011) Hallmarks of cancer: the next generation. *Cell*
21 19 **144**, 646-74.
- 22 20 146. Fabiani, R., Rosignoli, P., De Bartolomeo, A., Fuccelli, R., Servili, M., Montedoro, G. F.
23 21 and Morozzi, G. (2008) Oxidative DNA damage is prevented by extracts of olive oil,
24 22 hydroxytyrosol, and other olive phenolic compounds in human blood mononuclear cells and
25 23 HL60 cells. *J Nutr* **138**, 1411-6.
- 26 24 147. Fabiani, R., De Bartolomeo, A., Rosignoli, P. and Morozzi, G. (2001) Antioxidants prevent
27 25 the lymphocyte DNA damage induced by PMA-stimulated monocytes. *Nutr Cancer* **39**,
28 26 284-91.
- 29 27 148. Nosis, L., Doulias, P. T., Aligiannis, N., Bazios, D., Agalias, A., Galaris, D. and Mitakou,
30 28 S. (2005) DNA protecting and genotoxic effects of olive oil related components in cells
31 29 exposed to hydrogen peroxide. *Free Radic Res* **39**, 787-95.
- 32 30 149. Erol, O., Arda, N. and Erdem, G. (2012) Phenols of virgin olive oil protects nuclear DNA
33 31 against oxidative damage in HeLa cells. *Food Chem Toxicol* **50**, 3475-9.
- 34 32 150. Quiles, J. L., Ochoa, J. J., Ramirez-Tortosa, C., Battino, M., Huertas, J. R., Martin, Y. and
35 33 Mataix, J. (2004) Dietary fat type (virgin olive vs. sunflower oils) affects age-related
36 34 changes in DNA double-strand-breaks, antioxidant capacity and blood lipids in rats. *Exp*
37 35 *Gerontol* **39**, 1189-98.
- 38 36 151. Quiles, J. L., Ochoa, J. J., Ramirez-Tortosa, M. C., Huertas, J. R. and Mataix, J. (2006) Age-
39 37 related mitochondrial DNA deletion in rat liver depends on dietary fat unsaturation. *J*
40 38 *Gerontol A Biol Sci Med Sci* **61**, 107-14.
- 41 39 152. Besaratinia, A. and Pfeifer, G. P. (2006) Investigating human cancer etiology by DNA
42 40 lesion footprinting and mutagenicity analysis. *Carcinogenesis* **27**, 1526-37.
- 43 41 153. Fabiani, R., De Bartolomeo, A., Rosignoli, P., Servili, M., Montedoro, G. F. and Morozzi,
44 42 G. (2002) Cancer chemoprevention by hydroxytyrosol isolated from virgin olive oil through
45 43 G1 cell cycle arrest and apoptosis. *Eur J Cancer Prev* **11**, 351-8.
- 46 44 154. Fabiani, R., De Bartolomeo, A., Rosignoli, P., Servili, M., Selvaggini, R., Montedoro, G. F.,
47 45 Di Saverio, C. and Morozzi, G. (2006) Virgin olive oil phenols inhibit proliferation of
48 46 human promyelocytic leukemia cells (HL60) by inducing apoptosis and differentiation. *J*
49 47 *Nutr* **136**, 614-9.
- 50 48 155. Fabiani, R., Rosignoli, P., De Bartolomeo, A., Fuccelli, R. and Morozzi, G. (2008)
51 49 Inhibition of cell cycle progression by hydroxytyrosol is associated with upregulation of
52 50 cyclin-dependent protein kinase inhibitors p21(WAF1/Cip1) and p27(Kip1) and with
53 51 induction of differentiation in HL60 cells. *J Nutr* **138**, 42-8.

- 1
2
3 1 156. Wikman, H. and Kettunen, E. (2006) Regulation of the G1/S phase of the cell cycle and
4 2 alterations in the RB pathway in human lung cancer. *Expert Rev Anticancer Ther* **6**, 515-30.
5 3 157. Dotto, G. P. (2000) p21(WAF1/Cip1): more than a break to the cell cycle? *Biochim Biophys*
6 4 *Acta* **1471**, M43-56.
7 5 158. Sepporta, M. V., Mazza, T., Morozzi, G. and Fabiani, R. (2013) Pinoreosinol inhibits
8 6 proliferation and induces differentiation on human HL60 leukemia cells. *Nutr Cancer* **65**,
9 7 1208-18.
10 8 159. Cassiano, C., Casapullo, A., Tosco, A., Monti, M. C. and Riccio, R. (2015) In Cell
11 9 Interactome of Oleocanthal, an Extra Virgin Olive Oil Bioactive Component. *Nat Prod*
12 10 *Commun* **10**, 1013-6.
13 11 160. Fabiani, R., Fuccelli, R., Pieravanti, F., De Bartolomeo, A. and Morozzi, G. (2009)
14 12 Production of hydrogen peroxide is responsible for the induction of apoptosis by
15 13 hydroxytyrosol on HL60 cells. *Mol Nutr Food Res* **53**, 887-96.
16 14 161. Bernini, R., Gilardini Montani, M. S., Merendino, N., Romani, A. and Velotti, F. (2015)
17 15 Hydroxytyrosol-Derived Compounds: A Basis for the Creation of New Pharmacological
18 16 Agents for Cancer Prevention and Therapy. *J Med Chem* **58**, 9089-107.
19 17 162. Hu, T., He, X. W., Jiang, J. G. and Xu, X. L. (2014) Hydroxytyrosol and its potential
20 18 therapeutic effects. *J Agric Food Chem* **62**, 1449-55.
21 19 163. Maiuri, M. C., De Stefano, D., Di Meglio, P., Irace, C., Savarese, M., Sacchi, R., Cinelli, M.
22 20 P. and Carnuccio, R. (2005) Hydroxytyrosol, a phenolic compound from virgin olive oil,
23 21 prevents macrophage activation. *Naunyn Schmiedebergs Arch Pharmacol* **371**, 457-65.
24 22 164. Zhang, X., Cao, J. and Zhong, L. (2009) Hydroxytyrosol inhibits pro-inflammatory
25 23 cytokines, iNOS, and COX-2 expression in human monocytic cells. *Naunyn Schmiedebergs*
26 24 *Arch Pharmacol* **379**, 581-6.
27 25 165. Sumimoto, H. (2008) Structure, regulation and evolution of Nox-family NADPH oxidases
28 26 that produce reactive oxygen species. *FEBS J* **275**, 3249-77.
29 27 166. Kurtova, A. V., Xiao, J., Mo, Q., Pazhanisamy, S., Krasnow, R., Lerner, S. P., Chen, F.,
30 28 Roh, T. T., Lay, E., Ho, P. L. and Chan, K. S. (2015) Blocking PGE2-induced tumour
31 29 repopulation abrogates bladder cancer chemoresistance. *Nature* **517**, 209-13.
32 30 167. Nakanishi, M. and Rosenberg, D. W. (2013) Multifaceted roles of PGE2 in inflammation
33 31 and cancer. *Semin Immunopathol* **35**, 123-37.
34 32 168. Sasi, S. P., Yan, X., Enderling, H., Park, D., Gilbert, H. Y., Curry, C., Coleman, C., Hlatky,
35 33 L., Qin, G., Kishore, R. and Goukassian, D. A. (2012) Breaking the 'harmony' of TNF-alpha
36 34 signaling for cancer treatment. *Oncogene* **31**, 4117-27.
37 35 169. Lees, C. W., Ironside, J., Wallace, W. A. and Satsangi, J. (2008) Resolution of non-small-
38 36 cell lung cancer after withdrawal of anti-TNF therapy. *N Engl J Med* **359**, 320-1.
39 37 170. Fayyaz, S., Aydin, T., Cakir, A., Gasparri, M. L., Panici, P. B. and Farooqi, A. A. (2016)
40 38 Oleuropein mediated targeting of signaling network in cancer. *Curr Top Med Chem*.
41 39 171. Giner, E., Recio, M. C., Rios, J. L., Cerda-Nicolas, J. M. and Giner, R. M. (2016)
42 40 Chemopreventive effect of oleuropein in colitis-associated colorectal cancer in c57bl/6
43 41 mice. *Mol Nutr Food Res* **60**, 242-55.
44 42 172. Sepporta, M. V., Fuccelli, R., Rosignoli, P., Ricci, G., Servili, M., Morozzi, G. and Fabiani,
45 43 R. (2014) Oleuropein inhibits tumour growth and metastases dissemination in
46 44 ovariectomised nude mice with MCF-7 human breast tumour xenografts. *J Functional*
47 45 *Foods* **8**, 269-273.
48 46 173. Albin, A., Tosetti, F., Li, V. W., Noonan, D. M. and Li, W. W. (2012) Cancer prevention
49 47 by targeting angiogenesis. *Nat Rev Clin Oncol* **9**, 498-509.
50 48 174. Tosetti, F., Noonan, D. M. and Albin, A. (2009) Metabolic regulation and redox activity as
51 49 mechanisms for angioprevention by dietary phytochemicals. *Int J Cancer* **125**, 1997-2003.
52
53
54
55
56
57
58
59
60

- 1
2
3 1 175. Bruno, A., Ferlazzo, G., Albini, A. and Noonan, D. M. (2014) A think tank of
4 2 TINK/TANKs: tumor-infiltrating/tumor-associated natural killer cells in tumor progression
5 3 and angiogenesis. *J Natl Cancer Inst* **106**, dju200.
- 6 4 176. Bruno, A., Pagani, A., Magnani, E., Rossi, T., Noonan, D. M., Cantelmo, A. R. and Albini,
7 5 A. (2014) Inflammatory angiogenesis and the tumor microenvironment as targets for cancer
8 6 therapy and prevention. *Cancer Treat Res* **159**, 401-26.
- 9 7 177. Liou, G. Y. and Storz, P. (2010) Reactive oxygen species in cancer. *Free Radic Res* **44**, 479-
10 8 96.
- 11 9 178. Terzuoli, E., Donnini, S., Giachetti, A., Iniguez, M. A., Fresno, M., Melillo, G. and Ziche,
12 10 M. (2010) Inhibition of hypoxia inducible factor-1alpha by dihydroxyphenylethanol, a
13 11 product from olive oil, blocks microsomal prostaglandin-E synthase-1/vascular endothelial
14 12 growth factor expression and reduces tumor angiogenesis. *Clin Cancer Res* **16**, 4207-16.
- 15 13 179. Movafagh, S., Crook, S. and Vo, K. (2015) Regulation of hypoxia-inducible factor-1a by
16 14 reactive oxygen species: new developments in an old debate. *J Cell Biochem* **116**, 696-703.
- 17 15 180. Scoditti, E., Calabriso, N., Massaro, M., Pellegrino, M., Storelli, C., Martines, G., De
18 16 Caterina, R. and Carluccio, M. A. (2012) Mediterranean diet polyphenols reduce
19 17 inflammatory angiogenesis through MMP-9 and COX-2 inhibition in human vascular
20 18 endothelial cells: a potentially protective mechanism in atherosclerotic vascular disease and
21 19 cancer. *Arch Biochem Biophys* **527**, 81-9.
- 22 20 181. Lamy, S., Ouanouki, A., Beliveau, R. and Desrosiers, R. R. (2014) Olive oil compounds
23 21 inhibit vascular endothelial growth factor receptor-2 phosphorylation. *Exp Cell Res* **322**, 89-
24 22 98.
- 25 23 182. Lamy, S., Akla, N., Ouanouki, A., Lord-Dufour, S. and Beliveau, R. (2012) Diet-derived
26 24 polyphenols inhibit angiogenesis by modulating the interleukin-6/STAT3 pathway. *Exp Cell*
27 25 *Res* **318**, 1586-96.
- 28 26 183. Akl, M. R., Ayoub, N. M., Mohyeldin, M. M., Busnena, B. A., Foudah, A. I., Liu, Y. Y. and
29 27 Sayed, K. A. (2014) Olive phenolics as c-Met inhibitors: (-)-Oleocanthal attenuates cell
30 28 proliferation, invasiveness, and tumor growth in breast cancer models. *PLoS One* **9**, e97622.
- 31 29 184. Elnagar, A. Y., Sylvester, P. W. and El Sayed, K. A. (2011) (-)-Oleocanthal as a c-Met
32 30 inhibitor for the control of metastatic breast and prostate cancers. *Planta Med* **77**, 1013-9.
- 33 31 185. Jain, R. K. (2013) Normalizing tumor microenvironment to treat cancer: bench to bedside to
34 32 biomarkers. *J Clin Oncol* **31**, 2205-18.
- 35 33 186. Palmieri, D., Aliakbarian, B., Casazza, A. A., Ferrari, N., Spinella, G., Pane, B., Cafueri, G.,
36 34 Perego, P. and Palombo, D. (2012) Effects of polyphenol extract from olive pomace on
37 35 anoxia-induced endothelial dysfunction. *Microvasc Res* **83**, 281-9.
- 38 36 187. Goel, S., Wong, A. H. and Jain, R. K. (2012) Vascular normalization as a therapeutic
39 37 strategy for malignant and nonmalignant disease. *Cold Spring Harb Perspect Med* **2**,
40 38 a006486.
- 41 39 188. Rossi, T., Bassani, B., Gallo, C., Maramotti, S., Noonan, D., Albini, A. and Bruno, A.
42 40 (2015) Effect of a Purified Extract of Olive Mill Waste water on Endothelial Cell
43 41 Proliferation, Apoptosis, Migration and Capillary-Like Structure in vitro and in vivo. . *J*
44 42 *Bioanal Biomed* **S12:006**.
- 45 43 189. Evans, H. M. and Bishop, K. S. (1922) On the Existence of a Hitherto Unrecognized Dietary
46 44 Factor Essential for Reproduction. *Science* **56**, 650-651.
- 47 45 190. Mene-Saffrane, L. and DellaPenna, D. (2010) Biosynthesis, regulation and functions of
48 46 tocochromanols in plants. *Plant Physiol Biochem* **48**, 301-9.
- 49 47 191. Torquato, P., Ripa, O., Giusepponi, D., Galarini, R., Bartolini, D., Wallert, M., Pellegrino,
50 48 R., Cruciani, G., Lorkowski, S., Birringer, M., Mazzini, F. and Galli, F. (2016) Analytical
51 49 strategies to assess the functional metabolome of vitamin E. *J Pharm Biomed Anal*.

- 1
2
3 1 192. Horvath, G., Wessjohann, L., Bigirimana, J., Jansen, M., Guisez, Y., Caubergs, R. and
4 2 Horemans, N. (2006) Differential distribution of tocopherols and tocotrienols in
5 3 photosynthetic and non-photosynthetic tissues. *Phytochemistry* **67**, 1185-95.
- 6 4 193. Cahoon, E. B., Hall, S. E., Ripp, K. G., Ganzke, T. S., Hitz, W. D. and Coughlan, S. J.
7 5 (2003) Metabolic redesign of vitamin E biosynthesis in plants for tocotrienol production and
8 6 increased antioxidant content. *Nat Biotechnol* **21**, 1082-7.
- 9 7 194. Hensley, K., Benaksas, E. J., Bolli, R., Comp, P., Grammas, P., Hamdheydari, L., Mou, S.,
10 8 Pye, Q. N., Stoddard, M. F., Wallis, G., Williamson, K. S., West, M., Wechter, W. J. and
11 9 Floyd, R. A. (2004) New perspectives on vitamin E: gamma-tocopherol and
12 10 carboxyethylhydroxychroman metabolites in biology and medicine. *Free Radic Biol Med*
13 11 **36**, 1-15.
- 14 12 195. Galli, F., Cristina Polidori, M., Stahl, W., Mecocci, P. and Kelly, F. J. (2007) Vitamin E
15 13 biotransformation in humans. *Vitam Horm* **76**, 263-80.
- 16 14 196. Atkinson, J., Harroun, T., Wassall, S. R., Stillwell, W. and Katsaras, J. (2010) The location
17 15 and behavior of alpha-tocopherol in membranes. *Mol Nutr Food Res* **54**, 641-51.
- 18 16 197. Brigelius-Flohe, R. and Maiorino, M. (2013) Glutathione peroxidases. *Biochim Biophys*
19 17 *Acta* **1830**, 3289-303.
- 20 18 198. Bartolini, D., Piroddi, M., Tidei, C., Giovagnoli, S., Pietrella, D., Manevich, Y., Tew, K. D.,
21 19 Giustarini, D., Rossi, R., Townsend, D. M., Santi, C. and Galli, F. (2015) Reaction kinetics
22 20 and targeting to cellular glutathione S-transferase of the glutathione peroxidase mimetic
23 21 PhSeZnCl and its D,L-poly lactide microparticle formulation. *Free Radic Biol Med* **78**, 56-
24 22 65.
- 25 23 199. Cornwell, D. G., Williams, M. V., Wani, A. A., Wani, G., Shen, E. and Jones, K. H. (2002)
26 24 Mutagenicity of tocopheryl quinones: evolutionary advantage of selective accumulation of
27 25 dietary alpha-tocopherol. *Nutr Cancer* **43**, 111-8.
- 28 26 200. Ogawa, Y., Saito, Y., Nishio, K., Yoshida, Y., Ashida, H. and Niki, E. (2008) Gamma-
29 27 tocopheryl quinone, not alpha-tocopheryl quinone, induces adaptive response through up-
30 28 regulation of cellular glutathione and cysteine availability via activation of ATF4. *Free*
31 29 *Radic Res* **42**, 674-87.
- 32 30 201. Cornwell, D. G. and Ma, J. (2007) Studies in vitamin E: biochemistry and molecular biology
33 31 of tocopherol quinones. *Vitam Horm* **76**, 99-134.
- 34 32 202. Cornwell, D. G. and Ma, J. (2008) Nutritional benefit of olive oil: the biological effects of
35 33 hydroxytyrosol and its arylating quinone adducts. *J Agric Food Chem* **56**, 8774-86.
- 36 34 203. Viola, V., Ciffolilli, S., Legnaioli, S., Piroddi, M., Betti, M., Mazzini, F., Pierpaoli, E.,
37 35 Provinciali, M. and Galli, F. (2013) Mitochondrial-dependent anticancer activity of delta-
38 36 tocotrienol and its synthetic derivatives in HER-2/neu overexpressing breast
39 37 adenocarcinoma cells. *Biofactors* **39**, 485-93.
- 40 38 204. Zingg, J. M., Libinaki, R., Lai, C. Q., Meydani, M., Gianello, R., Ogru, E. and Azzi, A.
41 39 (2010) Modulation of gene expression by alpha-tocopherol and alpha-tocopheryl phosphate
42 40 in THP-1 monocytes. *Free Radic Biol Med*.
- 43 41 205. Zingg, J. M., Han, S. N., Pang, E., Meydani, M., Meydani, S. N. and Azzi, A. (2013) In vivo
44 42 regulation of gene transcription by alpha- and gamma-tocopherol in murine T lymphocytes.
45 43 *Arch Biochem Biophys* **538**, 111-9.
- 46 44 206. Molano, A. and Meydani, S. N. (2012) Vitamin E, signalosomes and gene expression in T
47 45 cells. *Mol Aspects Med* **33**, 55-62.
- 48 46 207. Galli, F. and Azzi, A. (2010) Present trends in vitamin E research. *Biofactors* **36**, 33-42.
- 49 47 208. Cavalier, L., Ouahchi, K., Kayden, H. J., Di Donato, S., Reutenauer, L., Mandel, J. L. and
50 48 Koenig, M. (1998) Ataxia with isolated vitamin E deficiency: heterogeneity of mutations
51 49 and phenotypic variability in a large number of families. *Am J Hum Genet* **62**, 301-10.

- 1
2
3 1 209. Gohil, K., Vasu, V. T. and Cross, C. E. (2010) Dietary alpha-tocopherol and neuromuscular
4 2 health: search for optimal dose and molecular mechanisms continues! *Mol Nutr Food Res*
5 3 **54**, 693-709.
- 6 4 210. Muller, D. P. (2010) Vitamin E and neurological function. *Mol Nutr Food Res* **54**, 710-8.
- 7 5 211. Jiang, Q. (2014) Natural forms of vitamin E: metabolism, antioxidant, and anti-
8 6 inflammatory activities and their role in disease prevention and therapy. *Free Radic Biol*
9 7 *Med* **72**, 76-90.
- 10 8 212. FAO/WHO, e. c. r.-N. D. (2002) Human Vitamin and Mineral Requirements: Vitamin E
11 9 (chapter 9). In *Training Materials for Agricultural Planning*, WHO, F. ed. FAO & WHO,
12 10 Bangkok, Thailand.
- 13 11 213. EFSA Panel on Dietetic Products, N., and Allergies (NDA) (2015) Scientific Opinion on
14 12 Dietary Reference Values or vitamin E as α -tocopherol. *EFSA Journal* **13**, 4149.
- 15 13 214. Raederstorff, D., Wyss, A., Calder, P. C., Weber, P. and Eggersdorfer, M. (2015) Vitamin E
16 14 function and requirements in relation to PUFA. *Br J Nutr* **114**, 1113-22.
- 17 15 215. Torquato, P., Bartolini, D., Giusepponi, D., Saluti, G., Russo, A., Barola, C., Birringer, M.,
18 16 Galarini, R. and Galli, F. (2016) alpha-13'-OH is the main product of alpha-tocopherol
19 17 metabolism and influences CYP4F2 and PPARgamma gene expression in HepG2 human
20 18 hepatocarcinoma cells. *Free Radic Biol Med* **in press**.
- 21 19 216. Wallert, M., Schmolz, L., Koeberle, A., Krauth, V., Gleis, M., Galli, F., Werz, O., Birringer,
22 20 M. and Lorkowski, S. (2015) alpha-Tocopherol long-chain metabolite alpha-13'-COOH
23 21 affects the inflammatory response of lipopolysaccharide-activated murine RAW264.7
24 22 macrophages. *Mol Nutr Food Res* **59**, 1524-34.
- 25 23 217. Cifollilli, S., Wallert, M., Bartolini, D., Krauth, V., Werz, O., Piroddi, M., Sebastiani, B.,
26 24 Torquato, P., Lorkowski, S., Birringer, M. and Galli, F. (2015) Human serum determination
27 25 and in vitro anti-inflammatory activity of the vitamin E metabolite alpha-(13'-hydroxy)-6-
28 26 hydroxychroman. *Free Radic Biol Med* **89**, 952-962.
- 29 27 218. Wallert, M., Schmolz, L., Galli, F., Birringer, M. and Lorkowski, S. (2014) Regulatory
30 28 metabolites of vitamin E and their putative relevance for atherogenesis. *Redox Biol* **2**, 495-
31 29 503.
- 32 30 219. Wallert, M., Mosig, S., Rennert, K., Funke, H., Ristow, M., Pellegrino, R. M., Cruciani, G.,
33 31 Galli, F., Lorkowski, S. and Birringer, M. (2014) Long-chain metabolites of alpha-
34 32 tocopherol occur in human serum and inhibit macrophage foam cell formation in vitro. *Free*
35 33 *Radic Biol Med* **68**, 43-51.
- 36 34 220. Birringer, M., Pfluger, P., Kluth, D., Landes, N. and Brigelius-Flohe, R. (2002) Identities
37 35 and differences in the metabolism of tocotrienols and tocopherols in HepG2 cells. *J Nutr*
38 36 **132**, 3113-8.
- 39 37 221. Sontag, T. J. and Parker, R. S. (2002) Cytochrome P450 omega-hydroxylase pathway of
40 38 tocopherol catabolism. Novel mechanism of regulation of vitamin E status. *J Biol Chem*
41 39 **277**, 25290-6.
- 42 40 222. Galli, F., Lee, R., Dunster, C. and Kelly, F. J. (2002) Gas chromatography mass
43 41 spectrometry analysis of carboxyethyl-hydroxychroman metabolites of alpha- and gamma-
44 42 tocopherol in human plasma. *Free Radic Biol Med* **32**, 333-40.
- 45 43 223. Stahl, W., Graf, P., Brigelius-Flohe, R., Wechter, W. and Sies, H. (1999) Quantification of
46 44 the alpha- and gamma-tocopherol metabolites 2,5,7, 8-tetramethyl-2-(2'-carboxyethyl)-6-
47 45 hydroxychroman and 2,7, 8-trimethyl-2-(2'-carboxyethyl)-6-hydroxychroman in human
48 46 serum. *Anal Biochem* **275**, 254-9.
- 49 47 224. Sontag, T. J. and Parker, R. S. (2007) Influence of major structural features of tocopherols
50 48 and tocotrienols on their omega-oxidation by tocopherol-omega-hydroxylase. *J Lipid Res*
51 49 **48**, 1090-8.

- 1
2
3 1 225. Bardowell, S. A., Duan, F., Manor, D., Swanson, J. E. and Parker, R. S. (2012) Disruption
4 2 of mouse cytochrome p450 4f14 (Cyp4f14 gene) causes severe perturbations in vitamin E
5 3 metabolism. *J Biol Chem* **287**, 26077-86.
- 6 4 226. Kalsotra, A. and Strobel, H. W. (2006) Cytochrome P450 4F subfamily: at the crossroads of
7 5 eicosanoid and drug metabolism. *Pharmacol Ther* **112**, 589-611.
- 8 6 227. Hoopes, S. L., Garcia, V., Edin, M. L., Schwartzman, M. L. and Zeldin, D. C. (2015)
9 7 Vascular actions of 20-HETE. *Prostaglandins Other Lipid Mediat* **120**, 9-16.
- 10 8 228. Schmolz, L., Birringer, M., Lorkowski, S. and Wallert, M. (2016) Complexity of vitamin E
11 9 metabolism. *World J Biol Chem* **7**, 14-43.
- 12 10 229. Birringer, M., Lington, D., Vertuani, S., Manfredini, S., Scharlau, D., Gleib, M. and Ristow,
13 11 M. (2010) Proapoptotic effects of long-chain vitamin E metabolites in HepG2 cells are
14 12 mediated by oxidative stress. *Free Radic Biol Med* **49**, 1315-22.
- 15 13 230. Galli, F., Stabile, A. M., Betti, M., Conte, C., Pistilli, A., Rende, M., Floridi, A. and Azzi, A.
16 14 (2004) The effect of alpha- and gamma-tocopherol and their carboxyethyl hydroxychroman
17 15 metabolites on prostate cancer cell proliferation. *Arch Biochem Biophys* **423**, 97-102.
- 18 16 231. Conte, C., Floridi, A., Aisa, C., Piroddi, M. and Galli, F. (2004) Gamma-tocotrienol
19 17 metabolism and antiproliferative effect in prostate cancer cells. *Ann N Y Acad Sci* **1031**,
20 18 391-4.
21 19
22 20
23 21
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgments

This manuscript was planned and prepared with the collaboration of the Scientific committee of the 2° Workshop PROS.IT “Cluster Agrifood Nazionale - Area Nutrizione e Salute”, Perugia, Italy, 20 of April, 2015.

This work was supported by the grant program of the Italian Ministry of University and Research (MIUR), National Technology Agrifood Cluster, Health and Nutrition area, PROS.IT project (CTN01 00230 413096)

AA was supported by the Fattoria La Violla di Gianni, Antonio e Bandino Lo Franco – SAS.

We acknowledge Miss Jenny Severi for her support in the organization of this manuscript, part of which was included in her dissertation for the Master Degree in Food Science and Human Nutrition at the University of Perugia, July 2015.

Conflict of interest statement

The authors have no conflict of interest to declare.

1
2
3 Table 2. Main plasma antioxidants
45
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	μM
Uric Acid	200-400
Ascorbic Acid	50-100
α -tocopherol	20-40
Protein thiols	400-500
Low MW thiols	0.1-20

From: [129]

For Peer Review

Table 3. Human Intervention Studies with OOPs: effects on LDL Oxidation

Supplement	Type of study	Control	n subject	Dose of VOO PC	Effect	LDL PC incorporation	ref
EVOO	Chronic 3 weeks	Sunflower oil	10	na	Mild (not signific) effect on ex vivo LDL oxidation	na	[130]
EVOO (50g/day)	Chronic 4 weeks	Refined oil (no PCs)	14	na	No difference between treatment	No	[112]
Fortified olive oil (47 g)	Acute 0-2 hrs	Placebo	12	31 mg	No difference vs control	na	[131]
EVOO (69 g/day)	Chronic 3 weeks	Phenol-poor EVOO	49	21 mg vs 3 mg	No difference between treatment Fasting blood	na	[132]
EVOO	Chronic 4 weeks	Olive Oil	10 hyperlipidemic		Reduction of ex vivo LDL oxidation	na	[133]
HPC OO (25 ml/day)	Chronic 4 days	LPC OO MPC OO	12	11 mg vs 3 vs 0.2	Reduced oxLDL in plasma (dose dependent effect)	na	[121]
VOO (25 ml/day)	Chronic 3 weeks	Refined oil (no PC) Common oil (medium PC)	30	4 mg	Reduction of ex vivo LDL oxidation Reduced oxLDL in plasma (dose dependent effect)	na	[134]
HPC OO (25 ml/day)	Chronic 3 weeks	LPC OO MPC OO	182	8 mg vs 4 vs 0.1	Reduced oxLDL in plasma (dose dependent effect)	na	[135]
HPC OO (40 ml)	Acute 0-6 hrs	LPC OO MPC OO	12	13 mg vs 6 vs 0.1	Reduced oxLDL in plasma vs LPC OO	Yes	[48]
VOO (25 ml/day)	Chronic 3 weeks	Refined oil (no PC) Common oil (medium PC)	33	21 mg vs 0	Reduction of ex vivo LDL oxidation Reduced oxLDL in plasma (dose dependent effect)	Yes	[136]
HT in H2O (2.5mg/kg)	Acute 0-2 hrs	Time 0	10	2.5mg/kg	No difference respect time 0	Yes but measured 10/20 min after supplementation	[115]
VOO (25 ml/day)	Chronic 3 weeks	Refined oil (no PC)	36	16 mg vs 0	Reduced oxLDL in plasma	Yes as conjugated	[137]
HPC OO (25 ml/day)	Chronic 3 weeks	LPC OO	18	8 mg vs 0	Reduced oxLDL in plasma	na	[65]
HPC OO (25 ml/day)	Chronic 3 weeks	LPC OO	25	8 mg vs 0	Decrease LDL concentration Decrease of number of small LDL Reduction of ex vivo LDL oxidation	na	[138]

na: not analyzed/available

Table 4. miRNAs modified by aging and/or treatment with olive oil phenols in concordance with the changes of the respective target genes.

Aging-modified miRNAs	Treatment-modified miRNAs
mmu-miR-681	mmu-miR-30a-5p*
mmu-miR-709	mmu-miR-484
mmu-miR-706	mmu-miR-434-5p*
mmu-miR-30a-5p*	mmu-miR-369-5p*
mmu-miR-129-5p	mmu-miR-451*
mmu-miR-434-3p	
mmu-miR-380-3p	
mmu-miR-30a-3p	
mmu-miR-434-5p*	
mmu-miR-433-3p	
mmu-miR-451*	
mmu-miR-720	
mmu-miR-126-3p*	
mmu-miR-369-5p*	

C57Bl mice were treated from age 10 to 16 months with an extra-virgin olive oil naturally rich in phenols (n=9), or with an extra-virgin olive oil deprived of phenols (n=9, control group), and miRNA expression was evaluated in the cerebral cortex through microarray. Age-related changes in miRNAs expression were obtained by comparing the control group of aged mice with a group of young animals (age 4 months, n=6). Treatment effects were evaluated comparing the two groups of aged mice. * miRNA present in both the lists.




Figure 1

Figure 1. Structures of main secoiridoid derivatives and phenyl alcohols of VOO. From [102].

251x188mm (96 x 96 DPI)




Figure 2.

Vitamin E isomers and metabolites	R ₁	R ₂	R ₃ [*]
α	CH ₃	CH ₃] OH OOH
β	CH ₃	H	
γ	H	CH ₃	
δ	H	H	

* Long-chain metabolites:
13'-Hydroxychromanol
13'-Carboxychromanol

Figure 2. Structures of tocopherols, tocotrienols and their earlier hepatic long-chain metabolites.

244x179mm (96 x 96 DPI)