



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

A clinical-molecular prognostic model to predict survival in patients with post polycythemia vera and post essential thrombocythemia

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

A clinical-molecular prognostic model to predict survival in patients with post polycythemia vera and post essential thrombocythemia myelofibrosis / Passamonti, F.; Giorgino, T.; Mora, B.; Guglielmelli, P.; Rumi, E.; Maffioli, M.; Rambaldi, A.; Caramella, M.; Komrokji, R.; Gotlib, J.; Kiladjian, J. J.; Cervantes, F.; Devos, T.; Palandri, F.; De Stefano, V.; Ruggeri, M.; Silver, R. T.; Benevolo, G.; Albano, F.; Caramazza, D.; Merli, M.;

Availability:

The webpage <https://hdl.handle.net/2158/1106622> of the repository was last updated on 2020-10-16T11:02:53Z

Published version:

DOI: 10.1038/leu.2017.169

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)



A clinical-molecular prognostic model to predict survival in patients with post polycythemia vera and post essential thrombocythemia myelofibrosis

F Passamonti, T Giorgino, B Mora, P Guglielmelli, E Rumi, M Maffioli, A Rambaldi, M Caramella, R Komrokji, J Gotlib, J J Kiladjan, F Cervantes, T Devos, F Palandri, V De Stefano, M Ruggeri, R T Silver, G Benevolo, F Albano, D Caramazza, M Merli, D Pietra, R Casalone, G Rotunno, T Barbui, M Cazzola, A M Vannucchi

Cite this article as: F Passamonti, T Giorgino, B Mora, P Guglielmelli, E Rumi, M Maffioli, A Rambaldi, M Caramella, R Komrokji, J Gotlib, J J Kiladjan, F Cervantes, T Devos, F Palandri, V De Stefano, M Ruggeri, R T Silver, G Benevolo, F Albano, D Caramazza, M Merli, D Pietra, R Casalone, G Rotunno, T Barbui, M Cazzola, A M Vannucchi, A clinical-molecular prognostic model to predict survival in patients with post polycythemia vera and post essential thrombocythemia myelofibrosis, *Leukemia* accepted article preview 31 May 2017; doi: [10.1038/leu.2017.169](https://doi.org/10.1038/leu.2017.169).

This is a PDF file of an unedited peer-reviewed manuscript that has been accepted for publication. NPG are providing this early version of the manuscript as a service to our customers. The manuscript will undergo copyediting, typesetting and a proof review before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers apply.

Received 8 February 2017; revised 27 April 2017; accepted 18 May 2017;
Accepted article preview online 31 May 2017

A clinical-molecular prognostic model to predict survival in patients with post polycythemia vera and post essential thrombocythemia myelofibrosis

Francesco Passamonti,^{1*} Toni Giorgino,^{2*} Barbara Mora,¹ Paola Guglielmelli,³ Elisa Rumi,⁴ Margherita Maffioli,¹ Alessandro Rambaldi,⁵ Marianna Caramella,⁶ Rami Komrokji,⁷ Jason Gotlib,⁸ Jean Jacques Kiladjan,⁹ Francisco Cervantes,¹⁰ Timothy Devos,¹¹ Francesca Palandri,¹² Valerio De Stefano,¹³ Marco Ruggeri,¹⁴ Richard T. Silver,¹⁵ Giulia Benevolo,¹⁶ Francesco Albano,¹⁷ Domenica Caramazza,¹ Michele Merli,¹ Daniela Pietra,⁴ Rosario Casalone,¹⁸ Giada Rotunno,³ Tiziano Barbui,¹⁹ Mario Cazzola^{4§} and Alessandro Maria Vannucchi.^{3§}

* Equal contribution §co-senior Authors

¹Hematology, Department of Medicine and Surgery, University of Insubria, Ospedale di Circolo, ASST Sette Laghi, Varese, Italy; ²Institute of Neurosciences, National Research Council of Italy, Padova, Italy; ³CRIMM-Centro Ricerca e Innovazione delle Malattie Mieloproliferative, Department of Experimental and Clinical Medicine, Azienda ospedaliera-Universitaria Careggi, University of Florence, Florence, Italy; ⁴Department of Hematology Oncology, Fondazione IRCCS Policlinico San Matteo, Università di Pavia, Pavia, Italy; ⁵University of Milan, Hematology and BMT Unit, ASST Papa Giovanni XXIII, Bergamo, Italy; ⁶Ospedale Niguarda Cà Granda, Milano, Italy; ⁷Moffitt Cancer Center, Tampa, FL, United States; ⁸Stanford University, Palo Alto, CA, United States; ⁹Hôpital Saint-Louis et Université Paris Diderot, Paris, France; ¹⁰Hospital Clínic, IDIBAPS, University of Barcelona, Barcelona, Spain; ¹¹Department of Hematology, University Hospitals Leuven and Laboratory of Experimental Transplantation, Department of Microbiology and Immunology, KU Leuven, Leuven, Belgium; ¹²Policlinico S. Orsola-Malpighi, Bologna, Italy; ¹³Università Cattolica del Sacro Cuore, Roma, Italy; ¹⁴Ospedale S. Bortolo, Vicenza, Italy; ¹⁵Weill Cornell Medical College, New York, United States; ¹⁶SC Hematology, A.O. Città della Salute e della Scienza, Turin Italy; ¹⁷Università di Bari, Bari, Italy; ¹⁸Cytogenetics and Medical Genetics Laboratory, Ospedale di Circolo, ASST Sette Laghi, Varese, Italy; ¹⁹FROM Research Foundation, ASST Papa Giovanni XXIII, Bergamo, Italy.

Running title: Post-PV and post-ET myelofibrosis

Contact information for the corresponding author: Francesco Passamonti, MD, Hematology, Dipartimento di Medicina e Chirurgia, University of Insubria, Via Guicciardini - 21100 Varese, Italy. Tel/Fax: +39 0332 393 648. E-mail francesco.passamonti@uninsubria.it

Abstract

Polycythemia vera (PV) and essential thrombocythemia (ET) are myeloproliferative neoplasms with variable risk of evolution into post-PV and post-ET myelofibrosis, from now on referred to as secondary myelofibrosis (SMF). No specific tools have been defined for risk stratification in SMF. To develop a prognostic model for predicting survival, we studied 685 *JAK2*, *CALR*, and *MPL* annotated patients with SMF. Median survival of the whole cohort was 9.3 years (95% CI: 8-not reached-NR-). Through penalized Cox regressions we identified negative predictors of survival and according to beta risk coefficients we assigned 2 points to hemoglobin level <11 g/dL, to circulating blasts $\geq 3\%$, and to *CALR*-unmutated genotype, 1 point to platelet count <150 x 10⁹/L and to constitutional symptoms, and 0.15 points to any year of age. MYSEC-PM (Myelofibrosis Secondary to PV and ET-Prognostic Model) allocated SMF patients into four risk categories with different survival ($P < 0.0001$): low (median survival NR; 133 patients), intermediate-1 (9.3 years, 95% CI: 8.1-NR; 245 patients), intermediate-2 (4.4 years, 95% CI: 3.2-7.9; 126 patients), and high risk (2 years, 95% CI: 1.7-3.9; 75 patients). Finally, we found that the MYSEC-PM represents the most appropriate tool for SMF decision-making to be used in clinical and trial settings.

INTRODUCTION

Polycythemia vera (PV), essential thrombocythemia (ET) and primary myelofibrosis (PMF) are the classical BCR/ABL1-negative myeloproliferative neoplasms (MPN).^{1, 2} The two more indolent diseases, PV and ET, nevertheless, can progress to secondary myelofibrosis (SMF), named post-PV (PPV) MF and post-ET (PET) MF,³ and to blast phase (BP),⁴ that result in worsening survival.⁵

Because of the lack of information on SMF, PMF and SMF are considered similar. The IPSS (International Prognostic Scoring System)⁶ and its time-dependent variants (Dynamic IPSS –DIPSS and DIPSS-plus)^{7, 8} are often used to predict survival and to plan therapy for SMF patients. However, these models have been developed in patients with PMF and are suboptimal to predict survival in SMF.⁹⁻¹¹ Recently, having acquired the prognostic implication of phenotype driver mutations and of additional mutations, the prognostication in MPN is moving towards integrated clinical-molecular models.¹²⁻¹⁶

The MYSEC (MYelofibrosis SEcondary to PV and ET) project recently disclosed genotype-phenotype associations in the largest cohort of SMF patients published to date, including 685 patients.¹⁷ We found that at presentation *JAK2*-mutated patients had higher white blood cell count and greater splenomegaly than *CALR*-mutated patients and that *CALR* type 1/type 1-like and *CALR* type 2/type 2-like were similar in terms of clinical presentation and outcome. Blast phase incidence was higher in *JAK2*-mutated PET MF and TN patients (triple negative, i.e. without *JAK2*, *MPL*, *CALR* mutations) when compared with *CALR*-mutated patients.

In this study, based on the MYSEC database, we developed an integrated clinical-molecular model to predict survival of SMF. We call this the MYSEC-PM (Myelofibrosis Secondary to PV and ET-Prognostic Model).

SUBJECTS AND METHODS***Study population***

This study includes 781 patients collected from 16 international centers (Table 1S). All patients have demographic, clinical, and hematologic data at diagnosis and an adequate follow-up. No differences in disease presentation (white blood cell count, hemoglobin level, platelet count) were observed among centers applying the Kruskal Wallis and pairwise Wilcoxon rank-sum tests. Driver mutation status was requested as secondary objective and available in 685 patients. Diagnoses of PPV MF and PET MF were performed between 1981 to 2015 and were locally reviewed according to the International Working Group on Myeloproliferative Neoplasm Research and Treatment (IWG-MRT) criteria.³ Evolution to BP was defined when leukemic blast cells were more than 20%, according to the World Health Organization (WHO) criteria.¹⁸ The study was approved by the Institutional Review Board of each Institution and conducted in accordance with the principles of the Declaration of Helsinki.

Statistical analysis

Descriptive summaries are reported as median and range for continuous covariates, and count and relative frequency for categorical ones. Continuous baseline values were compared via non-parametric Mann-Whitney U tests; categorical feature counts were compared with Fisher's exact tests. Time-to-event analyses were performed via Kaplan-Meier curves, using log-rank tests for comparisons and semi-parametric Cox models for regression. Events were assumed to be death for any cause (censored at last follow-up or at the time of stem cell transplant), thrombosis and leukemia. P values <.05 (2-tailed) were considered significant. To test impact on survival, we first performed an exploratory univariate analysis developing Cox regression models considering each covariate separately. To account for possible nonlinear effects, restricted cubic spline with 3 nodes were considered for continuous predictors. To select a parsimonious set of covariates on which to base the prediction algorithm, we fitted regularized regression models according to the least absolute shrinkage selection operator (LASSO) method, entering all the available discretized covariates. The selected value of the regularization parameter was $\lambda = 0.053$. LASSO fits a sequence of models with varying degrees of penalization in order to shrink less-relevant coefficients to zero, thus effectively performing a variable selection.¹⁹ The performance of the models was evaluated with 10-fold cross-validation; the highest shrinkage factor providing

performance within one standard deviation of the optimal cross-validated one was selected.²⁰ The ability of the final score to discriminate survival was verified via Harrell's concordance index C and its cross-validation. Statistical analyses were performed using R version 3.3.2.

RESULTS

Presenting features at diagnosis of SMF, comparison of PET MF and PPV MF

We developed the analyses on 685 (333 PET MF, 352 PPV MF) SMF patients with phenotype driver mutations available. Demographics and clinical features of patients at onset of SMF are shown in Table 1. Patients with PPV MF were older, had higher values of white blood cells and hemoglobin, larger spleen size and lower platelet count than those with PET MF. Pearson pairwise test demonstrated that at diagnosis patients with PPV MF had significantly higher frequency of constitutional symptoms, abnormal karyotype and prior thrombosis than those with PET MF. A significantly higher number of PPV MF patients had received cytoreductive treatments (231 with PET MF, 287 PPV MF, $P < .001$).

Events occurring after diagnosis of SMF

Incidence rates of events are reported in Table 2. For their calculation we took into account death and stem cell transplant as competing risks with thrombosis and leukemia. In detail, thrombotic events occurred in 67 SMF (12%; 29 PET MF and 38 PPV MF), blast phase in 52 SMF (7.5%, 30 PET MF and 22 PPV MF) and death in 169 SMF (25%, 69 PET MF and 100 PPV MF). Cause of death was known in 136 of the 169 patients who died: non-clonal disease progression in 52 (38%), blast phase in 43 (32%), second malignancy in 10 (7%), infection in 12 (9%), heart failure in 11 (8%), vascular complications in seven (5%), and other in one (1%). Median survival was 14.5 years (95% CI: 8-NR) in PET MF and 8.1 years (95% CI: 7.2-10.1) in PPV MF, with a borderline difference (Supplemental Figure 1, log-rank test, $P = .051$).

Analysis of survival and identification of risk factors

Median survival of SMF was 9.3 years (95% CI: 8-NR), as illustrated in Figure 1. To ascertain whether SMF survival has increased over calendar years, we performed a Cox regression including calendar year of diagnosis (as a linear covariate), correcting for IPSS risk category. We found that the trend of survival was not significantly changed ($P = .064$).

Univariate Cox Proportional Hazards regression showed that advanced age, male gender, lower hemoglobin level, greater white blood cell count, lower platelet count, higher circulating blast count, bone marrow fibrosis grade 3 vs. grade 2, presence of constitutional symptoms (fever, weight loss, night sweats), history of thrombosis before SMF, longer time from ET/PV to SMF negatively affected survival (maximum P values = .004). Interestingly, a normal karyotype was associated with longer survival ($P = .001$), but, as cytogenetic data were available in only 340 patients (49%), we excluded this variable from the statistical analysis. Conversely, type of diagnosis (PET MF, PPV MF), centers, spleen and liver size were neutral for survival.

Analysis of cutpoints of continuous variables indicated marked differences for patients with white blood cell count higher than $25 \times 10^9/L$, hemoglobin value lower than 11 g/dL, platelet count lower than $150 \times 10^9/L$, circulating blast equal to or higher than 3% and time to SMF greater than 10 years ($P < .0001$ each). An exploratory multi-class regression showed that HRs (hazard ratios) for *CALR*-unmutated genotypes (i.e. *JAK2*-mutated, *MPL*-mutated and triple negative) had overlapping confidence intervals, and significantly different from *CALR*-mutated genotype ($P = .003$), thus determining a binary category (*CALR*-mutated vs. *CALR*-unmutated) for genotype. Multivariate models consistently showed age at diagnosis to be an important predictor for survival ($P < 0.0001$). In order to minimize information loss on this covariate, we retained age at diagnosis as a continuous covariate.

We then selected the significant covariates employing a LASSO Cox regression. Six covariates remained with non-null coefficients: advanced age, hemoglobin level below 11 g/dL, platelet count below $150 \times 10^9/L$, circulating blasts equal to or higher than 3%, *CALR*-unmutated genotype, presence of constitutional symptoms. We generated a final Cox regression model incorporating the identified covariates (Table 3). All coefficients remained highly significant ($P < .003$); a test for Schönfeld residuals revealed no deviations from the proportional hazards assumption, except for a minor departure for constitutional symptoms.²¹

Development of the prognostic model

All factors shown in Table 3 were therefore included in the MYSEC-PM. To simplify the application of the risk score, we rounded the risk coefficients as risk points (Table 3). Namely, we allocated two points to hemoglobin level below 11 g/dL, to circulating blasts equal to or higher than 3% and to *CALR*-unmutated genotype, one point to platelet count lower than $150 \times 10^9/L$ and to the presence of constitutional symptoms. Age-related risk was rescaled accordingly, yielding approximately 0.15 points per year.

We thus recoded the MYSEC-PM into four categories of adequate size by pooling consecutive score values. The resulting risk categories were: low-risk (score less than 11, 133 patients), intermediate-1 risk (score equal to or higher than 11 and lower than 14, 245 patients), intermediate-2 risk (score equal to or higher than 14 and less than 16, 126 patients) and high risk (score equal to or higher than 16, 75 patients). Survival was significantly different among the risk groups (Figure 2A, log-rank test $P < 10^{-6}$). Median survival was not reached in the low risk, 9.3 years (95% CI: 8.1-NR) in the intermediate-1 risk, 4.4 (95% CI: 3.2-7.9) in the intermediate-2 risk and 2 years (95% CI: 1.7-3.9) in the high risk category. Additional Figure 2 shows survival compared to year-, age- and sex-matched U.S. population. Taking low risk as reference, the estimated average HR for intermediate-1 risk was 3.6 (95% CI: 1.8-7.2), for intermediate-2 risk was 10.6 (95% CI: 5.3-21.1) and that for high risk was 29.1 (95% CI: 14.1-59.8). When used to assign patients to the four discrete risk categories, the test retained very good predictivity (cross-validated C statistics 0.79) and calibration.

We validated the MYSEC-PM risk score via the internal 40-fold cross-validation of Harrell's concordance index, or C-statistic; the procedure re-trains the model multiple times on random resamples of the original data, aggregating the corresponding values obtained for C. The resulting cross-validated value of the C-statistics was $C = 0.78$ (cf. $C = 0.79$ of the full data set), confirming the validity of the model.

How to use the prognostic model in clinical practice: the MYSEC PM nomogram

Given the hybrid nature (continuous age, discrete points) of the risk prediction model, we provide a discrete/continuous nomogram (Figure 2B) to interpolate the final score and assess the individual patient's risk in an easy manner. The MYSEC PM nomogram provides an at-a-glance

diagram to combine the effect of age (continuous) and other covariates, at the same time providing color-coded read-outs on the resulting risk category. To calculate the MYSEC-PM doctors have to: 1) collect information on non-age prognostic variables (hemoglobin value, platelet count, circulating blast counts, genotype, constitutional symptoms), thus refer to Table 3 to assign the points and calculate their sum (score); 2) collect patient's age; 3) use the nomogram (Figure 2B) to locate the combination of score (read on the vertical axis) and age (on the horizontal axis) – the color at the location indicates the final risk category, 3) estimate the individual survival on the Kaplan Mayer curve (Figure 2A). To further illustrate and expedite the use of the score, the nomogram is also made available as an interactive web application for desktop and mobile use (available online at https://mysec.shinyapps.io/prognostic_model/).

Mutation status distribution in the MYSEC-PM risk categories

Supplemental Figure 3 describes the distribution of the phenotype driver mutations in the four MYSEC-PM risk groups. Of interest, *CALR* mutations were absent in high risk patients.

Comparison of the MYSEC-PM with the IPSS

We compared the quality on the risk stratification of the MYSEC-PM prognostic model with respect to the previously used IPSS (developed in PMF). The MYSEC-PM risk categories had higher predictive values than IPSS both in the original data set ($C = 0.79$ for MYSEC-PM and $C = 0.70$ for IPSS) and the 40-fold validation ($C = 0.78$ for MYSEC-PM and $C = 0.71$ for IPSS). The same conclusion has been obtained considering Akaike information criterion (AIC) values for the two models, which amount to 1416 and 1485 respectively for MYSEC-PM and IPSS (preferable models have lower AIC values). In summary, the predictive power of the MYSEC-PM is very high, and significantly better than the IPSS model.

In addition, we also checked whether it was possible, for the sake of simplicity, to integrate the IPSS model with genotype, by fitting a Cox model including the IPSS risk factors augmented by the “not-*CALR*” covariate. In this analysis, leukocyte count $> 25 \times 10^9/L$ ceases to be a significant predictor, suggesting that its adoption would require further changes to scores and thresholds with respect to the original IPSS.

DISCUSSION

Diagnosis of SMF is based on the IWG-MRT criteria, established in 2008: an antecedent WHO-based diagnosis of PV or ET including appropriate mutations and a bone marrow fibrosis above grade 1 are the two main criteria.³ The molecular anatomy of PV and ET has changed from 2008, leading to the new WHO classification in 2016.¹ By enriching the MYSEC database with the phenotype driver mutations of the *JAK2*, *CALR* and *MPL* genes,^{1,2} we provide a molecularly updated diagnosis of PV and ET and consequently of SMF. Concerning the accompanying mutations of MF,¹ no impact on SMF survival has been demonstrated,²² differently from their effect in PMF.¹² The assessment of bone marrow myelofibrosis requires bone marrow biopsy. Our study is representative of real-life in Europe and the United States: doctors perform bone marrow biopsy when they suspect disease evolution, an approach that remains a mainstay in recent recommendations.²³ Of note, the MYSEC database showed that the longer the span between PV/ET diagnosis and SMF, the worse the survival. This suggests to carefully monitor PV/ET patients in order to identify SMF evolution earlier, especially if disease-modifying treatments may be envisaged. In our series, we cannot completely exclude that some ET are prefibrotic/early MF (WHO, 2016).¹

The MYSEC study also characterized clinical phenotype and events of SMF. PPV MF and PET MF had substantial differences in clinical presentation, with a more “proliferative” phenotype in PPV MF, a pattern that is confirmed by the higher rate of PPV MF patients receiving cytoreductive agents. Of interest, the incidence of thrombosis ranged from 2.4 to 3.1 / 100 patients-year in PET MF and PPV MF, respectively, and accounted for 5% of deaths. These data clearly indicate that the risk of vascular complications is still significant in SMF. Perhaps, thromboprophylaxis should be considered in SMF, if not contraindicated because of a bleeding history or a low platelet count.

The median survival in SMF was 9.3 years without significant differences between PPV MF and PET MF. The MYSEC dataset did not disclose any change of SMF survival over calendar years of diagnosis. This seems to suggest that treatment strategies have not changed the disease history yet. Modern approach to myelofibrosis treatment includes the use of JAK inhibition and allogeneic stem cell transplantation (ASCT).²⁴ In PMF, we demonstrated that ruxolitinib might modify life expectancy in higher risk categories²⁵ with some criticisms²⁶⁻²⁸ and that ASCT improves survival in

higher risk categories, with the opposite effect in low risk patients,²⁹ when matched with a cohort of conventionally treated individuals. To date, no information is available on survival effect of these strategies in SMF.

Concerning current risk stratification of patients with SMF, the IPSS⁶ and DIPSS⁸ prognostic models are used in clinical practice²³ as well as in clinical trials.³⁰⁻³⁴ Introducing the MYSEC-PM instead of the IPSS model in patients with SMF will provide strong advantages: 1) model development in the correct setting of patients: IPSS/DIPSS were generated in PMF and not SMF, and, as a consequence, their application outside that setting is arbitrary and not data-supported; 2) the integrated clinical-molecular MYSEC-PM provides an excellent discrimination of survival (C=0.79), much better than the clinical-based IPSS; 3) a tentative integrated approach we did combining IPSS risk factors with mutational status failed in SMF.

Advanced age, hemoglobin level below 11 g/dL, platelet count below $150 \times 10^9/L$, circulating blast cells equal to or greater than 3%, *CALR*-unmutated genotype and the presence of constitutional symptoms are the risk factors composing the MYSEC-PM. Advanced age, anemia, circulating blast cells and the presence of constitutional symptoms are both components of the MYSEC-PM and the IPSS model,⁶ and advanced age and constitutional symptoms also stratify patients at the time of ASCT for survival.³⁵ This indicates a role of these factors in myelofibrosis survival prediction in general.

Myelofibrosis is an age-related disease and advanced age is the most powerful prognostic factor for survival prediction. This is not surprising from a biological standpoint as hematopoietic stem cells are modified during aging influencing disease development and eventually favoring clonal hematopoiesis with selection of mutated cells.³⁶ It is noteworthy that the most frequently involved age-related somatic mutations (*DNMT3A*, *TET2*, *ASXL1*, and *JAK2*)³⁷ are also implicated in myelofibrosis development.¹²

The extended study of the three phenotype driver mutations helped to recognize the favorable impact on survival of *CALR* mutations,¹⁷ and in this latter analysis *CALR*-unmutated genotypes (*JAK2*-mutated, *MPL*-mutated, triple negativity) are associated with a worse survival in multivariable analysis. The association of *CALR* mutations with a benign outcome in SMF, also

highlighted by the absence of *CALR*-mutated patients within the MYSEC-PM high-risk group, remains to be determined. Although all phenotype driver mutations activate the JAK/STAT pathway, subtle changes in the activation mechanism have been described among mutants.³⁸ The molecular profiling of SMF patients allows the MYSEC-PM to improve risk stratification in SMF, as demonstrated by the superior accuracy in survival prediction of MYSEC-PM over IPSS.

The MYSEC-PM identifies four risk categories with different survival: median survival was not reached in the low risk, 9.3 years in the intermediate-1 risk, 4.5 years in the intermediate-2 risk and 2 years in the high risk category. This information may be directly translated into clinical practice to personalize treatment options. Young and fit patients with intermediate-2 and high risk disease can be considered candidates for ASCT on the basis of the European LeukemiaNet recommendations,²³ which give an indication for ASCT in MF patients with a life expectancy below five years. On the opposite, patients at low risk have an indolent disease and a more conservative approach seems reasonable. Patients at intermediate-1 risk should be discussed on an individual basis in SMF. Ruxolitinib can be offered on the basis of the national indication/reimbursement rules since it has been intensively studied in SMF patients with intermediate and high risk disease according to clinical-based prognostic models.^{31, 32} Concerning investigative clinical trials, the use of MYSEC-PM in the selection of SMF patients may help in the identification of patients at higher risk who may be candidates for new treatment strategies or at lower risk who may be candidates for preventive approaches targeting disease progression/survival.

In conclusion, the MYSEC-PM is an integrated clinical-molecular prognostic model uniquely developed in SMF patients with a superior accuracy over IPSS. This clearly indicates that the MYSEC-PM is appropriate to make a clinical decision or design new clinical trials for patients with SMF.

Acknowledgments: This work was supported by a grant from the Associazione Italiana per la Ricerca sul Cancro (AIRC; Milano, Italy), Special Program Molecular Clinical Oncology 5x1000 to AIRC-Gruppo Italiano Malattie Mieloproliferative (AGIMM) project #1005. A complete list of AGIMM investigators is available at <http://www.progettoagimm.it>. P.G. also received funding by AIRC IG2014-15967 and by the Ministero della Salute (project code GR-2011-02352109). The Varese group was also supported by grants from the Fondazione Matarelli (Milano, Italy), Fondazione Rusconi (Varese, Italy) and AIL Varese ONLUS. M.C and F.P. were supported by a grant from the Fondazione Regionale Ricerca Biomedica (FRRB), Regione Lombardia. R.T.S. was supported in part by the Cancer Research and Treatment Fund, Inc. , New York, NY.

Conflict-of-interest disclosure: No conflict of interest to disclose for this study by all Authors.

Accepted manuscript

References

1. Arber DA, Orazi A, Hasserjian R, Thiele J, Borowitz MJ, Le Beau MM, *et al.* The 2016 revision to the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia. *Blood* 2016 Apr 11.
2. Passamonti F, Maffioli M. Update from the latest WHO classification of MPNs: a user's manual. *Hematology Am Soc Hematol Educ Program* 2016 Dec 02; **2016**(1): 534-542.
3. Barosi G, Mesa RA, Thiele J, Cervantes F, Campbell PJ, Verstovsek S, *et al.* Proposed criteria for the diagnosis of post-polycythemia vera and post-essential thrombocythemia myelofibrosis: a consensus statement from the International Working Group for Myelofibrosis Research and Treatment. *Leukemia* 2008 Feb; **22**(2): 437-438.
4. Passamonti F, Rumi E, Arcaini L, Castagnola C, Lunghi M, Bernasconi P, *et al.* Leukemic transformation of polycythemia vera: a single center study of 23 patients. *Cancer* 2005 Sep 1; **104**(5): 1032-1036.
5. Passamonti F, Rumi E, Caramella M, Elena C, Arcaini L, Boveri E, *et al.* A dynamic prognostic model to predict survival in post-polycythemia vera myelofibrosis. *Blood* 2008 Apr 1; **111**(7): 3383-3387.
6. Cervantes F, Dupriez B, Pereira A, Passamonti F, Reilly JT, Morra E, *et al.* New prognostic scoring system for primary myelofibrosis based on a study of the International Working Group for Myelofibrosis Research and Treatment. *Blood* 2009 Mar 26; **113**(13): 2895-2901.
7. Gangat N, Caramazza D, Vaidya R, George G, Begna K, Schwager S, *et al.* DIPSS plus: a refined Dynamic International Prognostic Scoring System for primary myelofibrosis that incorporates prognostic information from karyotype, platelet count, and transfusion status. *J Clin Oncol* 2011 Feb 1; **29**(4): 392-397.
8. Passamonti F, Cervantes F, Vannucchi AM, Morra E, Rumi E, Pereira A, *et al.* A dynamic prognostic model to predict survival in primary myelofibrosis: a study by the IWG-MRT (International Working Group for Myeloproliferative Neoplasms Research and Treatment). *Blood* 2010 Mar 4; **115**(9): 1703-1708.
9. Hernandez-Boluda JC, Pereira A, Gomez M, Boque C, Ferrer-Marin F, Raya JM, *et al.* The International Prognostic Scoring System does not accurately discriminate different risk categories in patients with post-essential thrombocythemia and post-polycythemia vera myelofibrosis. *Haematologica* 2014 Apr; **99**(4): e55-57.
10. Gowin K, Coakley M, Kosiorek H, Mesa R. Discrepancies of applying primary myelofibrosis prognostic scores for patients with post polycythemia vera/ essential thrombocytosis myelofibrosis. *Haematologica* 2016 Jun 27.
11. Chen M, Xu ZF, Xu JQ, Li B, Zhang PH, Qin TJ, *et al.* [Analysis of prognostic factors in Chinese patients with post-polycythemia vera myelofibrosis and post-essential thrombocythemia myelofibrosis]. *Zhonghua Xue Ye Xue Za Zhi* 2016 Oct 14; **37**(10): 876-880.

12. Vannucchi AM, Lasho TL, Guglielmelli P, Biamonte F, Pardanani A, Pereira A, *et al.* Mutations and prognosis in primary myelofibrosis. *Leukemia* 2013 Sep; **27**(9): 1861-1869.
13. Tefferi A, Lasho TL, Finke CM, Knudson RA, Ketterling R, Hanson CH, *et al.* CALR vs JAK2 vs MPL-mutated or triple-negative myelofibrosis: clinical, cytogenetic and molecular comparisons. *Leukemia* 2014 Jul; **28**(7): 1472-1477.
14. Wassie E, Finke C, Gangat N, Lasho TL, Pardanani A, Hanson CA, *et al.* A compendium of cytogenetic abnormalities in myelofibrosis: molecular and phenotypic correlates in 826 patients. *Br J Haematol* 2015 Apr; **169**(1): 71-76.
15. Rumi E, Pietra D, Pascutto C, Guglielmelli P, Martinez-Trillos A, Casetti I, *et al.* Clinical effect of driver mutations of JAK2, CALR, or MPL in primary myelofibrosis. *Blood* 2014 Aug 14; **124**(7): 1062-1069.
16. Rozovski U, Verstovsek S, Manshour T, Dembitz V, Bozinovic K, Newberry K, *et al.* An accurate, simple prognostic model consisting of age, JAK2, CALR, and MPL mutation status for patients with primary myelofibrosis. *Haematologica* 2016 Sep 29.
17. Passamonti F, Mora B, Giorgino T, Guglielmelli P, Cazzola M, Maffioli M, *et al.* Driver mutations' effect in secondary myelofibrosis: An international multicenter study based on 781 patients. *Leukemia* 2016 Nov 25.
18. Vardiman JW, Thiele J, Arber DA, Brunning RD, Borowitz MJ, Porwit A, *et al.* The 2008 revision of the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia: rationale and important changes. *Blood* 2009 Jul 30; **114**(5): 937-951.
19. Pavlou M, Ambler G, Seaman SR, Guttman O, Elliott P, King M, *et al.* How to develop a more accurate risk prediction model when there are few events. *BMJ* 2015 Aug 11; **351**: h3868.
20. Friedman J, Hastie T, Tibshirani R. Regularization Paths for Generalized Linear Models via Coordinate Descent. *J Stat Softw* 2010; **33**(1): 1-22.
21. Grambsch PM, Therneau TM. Proportional Hazards Tests and Diagnostics Based on Weighted Residuals. *Biometrika* 1994 Sep; **81**(3): 515-526.
22. Rotunno G, Pacilli A, Artusi V, Rumi E, Maffioli M, Delaini F, *et al.* Epidemiology and clinical relevance of mutations in post-polycythemia vera and post-essential thrombocythemia myelofibrosis. A study on 359 patients of the AGIMM group. *Am J Hematol* 2016 Apr 2.
23. Barbui T, Barosi G, Birgegard G, Cervantes F, Finazzi G, Griesshammer M, *et al.* Philadelphia-negative classical myeloproliferative neoplasms: critical concepts and management recommendations from European LeukemiaNet. *J Clin Oncol* 2011 Feb 20; **29**(6): 761-770.

24. Mesa RA, Passamonti F. Individualizing Care for Patients With Myeloproliferative Neoplasms: Integrating Genetics, Evolving Therapies, and Patient-Specific Disease Burden. *Am Soc Clin Oncol Educ Book* 2016; **35**: e324-335.
25. Passamonti F, Maffioli M, Cervantes F, Vannucchi AM, Morra E, Barbui T, *et al.* Impact of ruxolitinib on the natural history of primary myelofibrosis: a comparison of the DIPSS and the COMFORT-2 cohorts. *Blood* 2014 Mar 20; **123**(12): 1833-1835.
26. Barosi G, Zhang MJ, Peter Gale R. Does ruxolitinib improve survival of persons with MPN-associated myelofibrosis? Should it? *Leukemia* 2014 Jul 16.
27. Passamonti F, Vannucchi AM, Cervantes F, Harrison C, Morra E, Kantarjian H, *et al.* Ruxolitinib and survival improvement in patients with myelofibrosis. *Leukemia* 2015 Mar; **29**(3): 739-740.
28. Cervantes F, Pereira A. Does ruxolitinib prolong the survival of patients with myelofibrosis? *Blood* 2016 Dec 28.
29. Kroger N, Giorgino T, Scott BL, Ditschkowski M, Alchalby H, Cervantes F, *et al.* Impact of allogeneic stem cell transplantation on survival of patients less than 65 years of age with primary myelofibrosis. *Blood* 2015 May 21; **125**(21): 3347-3350; quiz 3364.
30. Pardanani A, Laborde RR, Lasho TL, Finke C, Begna K, Al-Kali A, *et al.* Safety and efficacy of CYT387, a JAK1 and JAK2 inhibitor, in myelofibrosis. *Leukemia* 2013 Jun; **27**(6): 1322-1327.
31. Verstovsek S, Mesa RA, Gotlib J, Levy RS, Gupta V, DiPersio JF, *et al.* A double-blind, placebo-controlled trial of ruxolitinib for myelofibrosis. *N Engl J Med* 2012 Mar 1; **366**(9): 799-807.
32. Harrison C, Kiladjian JJ, Al-Ali HK, Gisslinger H, Waltzman R, Stalbovskaya V, *et al.* JAK inhibition with ruxolitinib versus best available therapy for myelofibrosis. *N Engl J Med* 2012 Mar 1; **366**(9): 787-798.
33. Pardanani A, Harrison C, Cortes JE, Cervantes F, Mesa RA, Milligan D, *et al.* Safety and Efficacy of Fedratinib in Patients With Primary or Secondary Myelofibrosis: A Randomized Clinical Trial. *JAMA Oncol* 2015 Aug 1; **1**(5): 643-651.
34. Tefferi A, Al-Ali HK, Barosi G, Devos T, Gisslinger H, Jiang Q, *et al.* A randomized study of pomalidomide vs placebo in persons with myeloproliferative neoplasm-associated myelofibrosis and RBC-transfusion dependence. *Leukemia* 2016 Nov 18.
35. Alchalby H, Yunus DR, Zabelina T, Kobbe G, Holler E, Bornhauser M, *et al.* Risk models predicting survival after reduced-intensity transplantation for myelofibrosis. *Br J Haematol* 2012 Apr; **157**(1): 75-85.
36. Marty C, Lacout C, Droin N, Le Couedic JP, Ribrag V, Solary E, *et al.* A role for reactive oxygen species in JAK2(V617F) myeloproliferative neoplasm progression. *Leukemia* 2013 Nov; **27**(11): 2187-2195.

37. Jaiswal S, Fontanillas P, Flannick J, Manning A, Grauman PV, Mar BG, *et al.* Age-Related Clonal Hematopoiesis Associated with Adverse Outcomes. *New Engl J Med* 2014 Dec 25; **371**(26): 2488-2498.
38. Vainchenker W, Constantinescu SN, Plo I. Recent advances in understanding myelofibrosis and essential thrombocythemia. *F1000Res* 2016; **5**.

Accepted manuscript

Legend to Figures

Figure 1. Estimate of survival in 685 patients with post polycythemia vera and post essential thrombocythemia myelofibrosis. This represents the estimate of survival of the entire cohort of patients with secondary myelofibrosis.

Figure 2. The MYSEC-PM. (A) The MYSEC-PM estimate of survival in patients with secondary myelofibrosis molecularly annotated for *JAK2*, *CALR*, *MPL* mutations. Risk factors and relative points composing the MYSEC-PM are patient's age (0.15 per patient's year of age), hemoglobin level below 11 g/dL (2 points), platelet count lower than $150 \times 10^9/L$ (1 point), circulating blasts equal to or higher than 3% (2 points), presence of constitutional symptoms (1 point) and *CALR*-unmutated genotype (2 points). The final risk category is to be calculated with the MYSEC-PM nomogram (Figure 2B). The four risk categories are: low-risk (median survival not reached; 133 patients), intermediate-1 risk (median survival 9.3 years, 95% CI: 8.0-NR; 245 patients), intermediate-2 risk (median survival 4.5, 95% CI: 3.2-7.9; 126 patients) and high risk (median survival 2.0 years, 95% CI: 1.7-3.9; 75 patients) (2). (B) The MYSEC-PM nomogram. The MYSEC PM nomogram visually assigns the MYSEC-PM risk category starting from the non-age prognostic variables (vertical axis) and the patient's age (horizontal axis) illustrated in Table 3. To determine the risk category of an individual patient with hemoglobin value of 10 g/dL and circulating blast of 6%, for example, follow the horizontal line, starting from the non-age-parameter-sum of 4 on the vertical axis (see Table 3 for points) to the age of the patient and record the color at that point. If the patient is 40 years old, the 4-line and the vertical 40-year line cross in the green field, corresponding to the low risk category, while if the patient is 70 years old, the 4-line and the vertical 70-year line cross in the violet field, corresponding to the intermediate-2 risk category.

Table 1. Hematological and clinical features of 685 patients with post essential thrombocythemia and post polycythemia vera myelofibrosis.

	SMF (n = 685)	PET MF (n = 333)	PPV MF (n = 352)	P value
Age, median (range), years	64 (25-96)	64 (25-93)	65 (34-96)	.02
Age older than 65 years, n. (%)	360 (53)	146 (44)	179 (51)	.08
Follow-up, median (range), years	3.0 (0.6-27.3)	3.1 (0.6-17.4),	2.9 (0.6-27.3)	.88
Time to SMF, years (range)	10.7 (0.6-41.4)	10.3 (0.7-34.8)	11.1 (0.6-41.4)	.36
History of cancer, n. (%)	87 (13)	36 (11)	51 (15)	.16
History of thrombosis, n. (%)	171 (26)	70 (22)	101 (29)	.03
Male gender, n. (%)	356 (52)	165 (50)	191 (54)	.23
WBC, median (range), x10 ⁹ /L	10.2 (1.1-98.4)	7.8 (1.1-97.3)	13.5 (1.7-98.4)	< .001
Hb, median (range), g/dL	11 (5-15.7)	10.7 (5-15.4)	12 (6.8-15.7)	< .001
PLT, median (range), x 10 ⁹ /L	336 (15-1908)	379 (40-1908)	294 (15-1689)	< .001
Circulating blast 3% or more (%)	55 (9)	24 (8)	31 (10)	.43
Spleen size,* median (range)	7 (0-34)	4 (0-27)	10 (0-34)	< .001
Constitutional symptoms, n. (%)	285 (44)	113 (37)	172 (51)	< .001
Normal karyotype,** n. (%)	223 (66)	118 (73)	105 (59)	.005
Favorable karyotype,** n. (%)	283 (87)	138 (88)	145 (85)	.58
<i>JAK2</i> (V617F)	534 (78)	181 (54)	352(100)	< .001
<i>CALR</i>	102 (15)	102 (31)	-	
<i>MPL</i>	30 (4)	30 (9)	-	
Triple negative	19 (3)	19 (6)	-	

SMF: secondary myelofibrosis; PET MF: post essential thrombocythemia myelofibrosis; PPV MF: post polycythemia vera myelofibrosis; WBC: white blood cell count; Hb: hemoglobin level; PLT: platelet count.

*palpable from the left costal margin

**Karyotype was available in 340 patients

Table 2. Incidence of events during the follow-up of 685 patients with post essential thrombocythemia and post polycythemia vera myelofibrosis.


Incidence	PET MF	PPV MF	P value
/100 patients-year (95% CI)	(n = 333)	(n = 352)	
Thrombosis	2.4 (1.6-3.4)	3.1 (2.2-4.3)	.2
Blast phase	2.3 (1.6-3.4)	1.6 (1-2.5)	.2
Mortality	5.5 (4.3-7)	7.4 (6-8.9)	.06



Accepted manuscript

Table 3. Results of the multivariable analysis to define predictors of inferior survival in 685 molecularly annotated patients with post essential thrombocythemia and post polycythemia vera myelofibrosis.

Covariates	HR	95% CI	P value	Risk coefficient	Points assigned in
				Beta	the MYSEC-PM
Age at diagnosis of SMF	1.07	1.05-1.09	<.0001	0.068	0.15
Hemoglobin < 11 g/dL	2.3	1.6-3.3	<.0001	0.8	2
Platelet < 150 x10 ⁹ /L	1.7	1.2-2.5	.006	0.5	1
Circulating blast cells ≥ 3%	2.9	1.8-4.8	<.0001	1.1	2
<i>CALR</i> -unmutated genotype	2.6	1.2-5.3	.001	0.9	2
Constitutional symptoms	1.5	1.0-2.0	.03	0.4	1

HR: Hazard Ratio; CI: confidence interval





© 2017 Macmillan Publishers Limited. All rights reserved.

A

B