

Slab-derived melt involvement in petrogenesis of the high-Nb basalts and magnesian andesites-dacites from NE Iran

Parham Ahmadi (1), Mohammad Reza Ghorbani (1), Massimo Coltorti (2), Takeshi Kuritani (3), Yue Cai (4), Anna Maria Fioretti (5), Eleonora Braschi (6), Pier Paolo Giacomoni (2), Shahrouz Babazadeh (1), Sandro Conticelli (6,7)

(1) Department of Geology, Tarbiat Modares University, Tehran 114115-175, Iran, (2) Department of Earth Sciences, Ferrara University, Via Saragat 1, 44100 Ferrara, Italy, (3) Department of Earth and Planetary Sciences, Graduate School of Science, Hokkaido University, (4) Lamont Doherty Earth Observatory at the Columbia University, 61 Route 9W, Palisades NY 10964, USA, (5) CNR, Istituto di Geoscienze e Georisorse, sezione di Padova, Corso Garibaldi 37, I-35137 Padova, Italy, (6) CNR, Istituto di Geoscienze e Georisorse, sezione di Firenze, Via Giorgio La Pira, 4,I-50121, Firenze, Italy, (7) Dipartimento di Scienze della Terra, Università degli Studi di Firenze, Via Giorgio La Pira, 4,I-50121, Firenze, Italy

Abstract

Tertiary volcanic rocks in NW Firoozeh region, also known as Meshkan triangular structural unit, represent an episode of Tertiary post-collisional magmatism in NE Iran. The volcanic suite is made of Magnesian andesites to dacites associated with some high-Nb basalts. Despite a rather restricted range of emplacement age, variable from 24.1 to 21.7 Ma obtained on the basis of ^{40}Ar - ^{39}Ar dating. Major and trace element data show that Firoozeh volcanic rocks were derived from two distinct parental magmas. The dominant high magnesian magmatic series constitutes a wide spectrum of volcanic rocks from andesite to dacite. The magnesian andesites-dacites are subalkaline and characterized by rather high MgO, Ni, Cr and Sr/Y ratio. High Sr/Y ratio of the magnesian andesites-dacites as well as their Sr-Nd isotopic composition support the notion that the magnesian andesites were originated by interaction of slab derived melts with overlying mantle wedge. Fractional crystallization of an amphibole and plagioclase mineral assemblage is found responsible for evolutionary path from the magnesian andesite to more evolved rocks. In contrast to the subalkaline magnesian andesites-dacites, the high Nb basalts are sodic alkaline rocks and show silica-undersaturated degree. The high-Nb basalts are enriched in Nb, and a wide range of incompatible trace elements that include LILE, LREE. Almost identical Sr-Nd isotopic composition of the magnesian andesites-dacites and high-Nb basalts imply that the two magmatic series has probably shared a common mantle source. Generation of parental magmas of the two magmatic series are attributed to the asthenospheric upwelling and subsequent partial melting of mantle metasomatized by slab-derived melts.

Key words: slab melt, High-Nb basalt, $^{40}\text{Ar}/^{39}\text{Ar}$, High-Mg andesite