



UNIVERSITÀ  
DEGLI STUDI  
FIRENZE

## FLORE

# Repository istituzionale dell'Università degli Studi di Firenze

### **A new real-time loop-mediated isothermal amplification (LAMP) assay to rapidly detect *Phytophthora ramorum* and *P. lateralis* invasive**

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

*Original Citation:*

A new real-time loop-mediated isothermal amplification (LAMP) assay to rapidly detect *Phytophthora ramorum* and *P. lateralis* invasive plant pathogens / C. Aglietti; A. Santini; P. Bartolini; P. Capretti; N. Luchi. - STAMPA. - (2018), pp. 74-74. ( XXIV National Congress Italian Phytopathological Society (SIPaV)5-7 September 2018).

*Availability:*

The webpage <https://hdl.handle.net/2158/1135343> of the repository was last updated on 2018-09-20T09:39:14Z

*Publisher:*

G. Romanazzi, L. Landi, S. Murolo, E. Feliziani, V. Mancini, L. Rubino

*Terms of use:*

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

*Publisher copyright claim:*

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

**A new real-time loop-mediated isothermal amplification (LAMP) assay to rapidly detect *Phytophthora ramorum* and *P. lateralis* invasive plant pathogens**

**C. Aglietti<sup>1</sup>, A. Santini<sup>2</sup>, P. Bartolini<sup>2</sup>, P. Capretti<sup>1</sup>, N. Luchi<sup>2</sup>**

*Department of Agrifood Production and Environmental Sciences, University of Florence (DISPAA), Piazzale delle Cascine 28, I-50144 Firenze, Italy; <sup>2</sup> Institute for Sustainable Plant Protection - National Research Council (IPSP-CNR), Via Madonna del Piano 10, I-50019 Sesto Fiorentino, Firenze, Italy. E-mail: nicola.luchi@ipsp.cnr.it*

An effective framework for early warning and rapid response is a crucial element to mitigate or prevent the ecological and economic impacts of invasive plant pathogens. Molecular detection of these pathogens is now preferred for its greater sensitivity and specificity respect to classical tools. These analyses are generally PCR-based and require a well-equipped lab. Significant advantages in terms of prompt response to threats might be obtained by using an on-site detection tool. A real-time monitoring LAMP-based method has been optimized for a rapid and sensitive detection of *P. ramorum* and *P. lateralis*, two quarantine pathogens that cause mortality on forest trees and ornamental plants in Europe and North America, by using the Genie<sup>®</sup> II system. The assay was able to recognize each pathogen with a high level of specificity and sensitivity also in plant tissues (to 4 pg  $\mu\text{l}^{-1}$  of DNA) in only 30 minutes. Great simplicity, sensitivity, specificity, and minimum required equipment make the LAMP assay ideal for application in the field and for routine plant testing at ports of entry. The use of portable and handled instruments allows a fast analysis of the collected sample reducing the diagnosis time and may have implications for disease management and for the control of *P. lateralis* and *P. ramorum*.