



FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Role of TRPA1 receptors in skin inflammation induced by volatile chemical irritants in mice

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Role of TRPA1 receptors in skin inflammation induced by volatile chemical irritants in mice / Noroes M.M.; Santos L.G.; Gavioli E.C.; de Paula Soares Rachetti V.; Otuki M.F.; de Almeida Cabrini D.; da Silveira Prudente A.; Oliveira J.R.J.M.; de Carvalho Goncalves M.; Ferreira J.; Preti D.; De Logu F.; Nassini R.; Andre E.. - In: EUROPEAN JOURNAL OF PHARMACOLOGY. - ISSN 0014-2999. - ELETTRONICO. - 858:(2019), pp.

Availability:

The webpage https://hdl.handle.net/2158/1163838 of the repository was last updated on 2019-07-19T16:03:36Z

Published version: DOI: 10.1016/j.ejphar.2019.172460

Terms of use: Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The abovementioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

Contents lists available at ScienceDirect



European Journal of Pharmacology



journal homepage: www.elsevier.com/locate/ejphar

Full length article

Role of TRPA1 receptors in skin inflammation induced by volatile chemical irritants in mice



Maíra Macedo Norões^c, Larissa Gonzaga Santos^b, Elaine Cristina Gavioli^a, Vanessa de Paula Soares Rachetti^a, Michel Fleith Otuki^b, Daniela de Almeida Cabrini^b, Arthur da Silveira Prudente^{b,f}, Janiana Raíza Jentsch Matias Oliveira^b, Muryel de Carvalho Gonçalves^c, Juliano Ferreira^c, Delia Preti^d, Francesco De Logu^e, Romina Nassini^e, Eunice André^{b,*}

^a Department of Biophysics and Pharmacology, Federal University of Rio Grande do Norte, Campus Universitário UFRN – Lagoa Nova, Natal, RN, Brazil

^b Department of Pharmacology, Federal University of Paraná, Centro Politécnico – Jardim das Américas, Curitiba, PR, Brazil

^c Department of Pharmacology, Federal University of Santa Catarina, Campus Universitário, Córrego Grande, Florianópolis, SC, Brazil

^d Department of Chemical and Pharmaceutical Sciences, University of Ferrara, Ferrara, Italy

^e Department of Health Sciences, Section of Clinical Pharmacology and Oncology, University of Florence, Florence, Italy

^fFederal University of Latin American Integration, Avenida Silvio Américo Sasdelli, Bairro Itaipu A, Foz do Iguaçu, PR, Brazil

ARTICLE INFO

Keywords: Transient receptor potential ankyrin 1 Inflammatory skin disease Chemical irritants Irritant contact dermatitis

ABSTRACT

Contact dermatitis is a very common inflammatory reaction in the skin, causing not only aesthetic problems but also loss functionality at work. The molecular mechanisms of contact dermatitis induced by chemical irritants are still unclear. Considering that transient receptor potential channels (TRP) may induce neurogenic inflammation and the exacerbation of inflammatory responses, here we investigated the role of transient receptor potential channel ankyrin type-1 (TRPA1) in skin inflammation evoked by chemical irritants. Ear oedema and nociceptive responses elicited by the topical application of xylene and toluene were measured in Swiss mice, wild type and TRPA1 knockout (Trpa1^{-/-}) C57BL/6 mice. Histological analyses were performed in mice subjected to the ear oedema assay. Topical application of xylene and toluene in the mouse ear induced an edematogenic response (0.113 \pm 0.008 mm and 0.067 \pm 0.011 mm), compared to vehicle (0.008 \pm 0.008 mm), assessed by ear thickness measurements and histological analyses. These responses were prevented by topical pretreatment with a selective TRPA1 antagonist, HC-030031 (% inhibition: xylene 36.8 \pm 9.4% and toluene 50.7 \pm 11.0%), and by the genetic deletion of TRPA1 ((% inhibition: xylene 66.6 \pm 16.7% and toluene $75 \pm 0\%$). In addition, the topical application of xylene and toluene to the mouse paw elicited nociceptive responses, which were significantly reduced by oral treatment with HC-030031 ((% of inhibition: 84.9 \pm 1.3% and 27.1 \pm 8.0%, respectively); nociceptive responses were almost completely abolished in Trpa1^{-/-}mice. Our data suggest that the activation of TRPA1 could be involved in some of the symptoms of irritant-mediated contact dermatitis, such as oedema, pain and neurogenic inflammation.

1. Introduction

Nowadays, due to increasing exposure to chemical agents and environmental pollutants, the number of people developing inflammatory reactions in the skin has increased. Most of these are clinically described as contact dermatitis, which can cause not only aesthetic problems, but also loss functionality at work (Kaplan et al., 2012). Epidemiological studies show that contact dermatitis accounts for 50% of all occupational diseases in the United States, and is the fourth leading

cause of work-related disability (Cao and Taylor, 2008; Ngo and Maibach, 2010). The incidence varies according to the risk of exposure of the population to chemical agents, and is higher in chemical industry workers, household workers, painters and hairdressers, among others (Sasseville, 2008).

Both types of contact dermatitis, i.e. irritant-mediated and allergic, are caused by dermal exposure to chemical agents found in detergents, solvents, insecticides, beauty and house cleaning products (Cao and Taylor, 2008). While the allergic reaction results from an immune

https://doi.org/10.1016/j.ejphar.2019.172460 Received 7 March 2019; Received in revised form 13 June 2019; Accepted 14 June 2019 Available online 20 June 2019 0014-2999/ © 2019 Elsevier B.V. All rights reserved.

^{*} Corresponding author. Department of Pharmacology, Federal University of Paraná, Curitiba, PR, 81531-980, Brazil. *E-mail address:* eunice.andre@ufpr.br (E. André).

response triggered in the epidermis and dermis after a previous sensitisation (Cao and Taylor, 2008), irritant-mediated contact dermatitis is caused by chemical agents that directly irritate the skin (Sasseville, 2008; Ngo and Maibach, 2010). Currently, a family of channels known as transient receptor potential channels (TRPs) are considered to be molecular integrators of a large number of noxious chemical stimuli (Boonen et al., 2017). This receptor family has been studied with great interest in several pathologies including skin inflammation (for a review, see Caterina and Pang, 2016). Among these receptors, we highlight the transient receptor potential channels ankyrin type-1 (TRPA1).

TRPA1 channels are activated by pungent chemicals such as horseradish, mustard and cinnamon oils, temperatures less than or equal to 17 °C, endogenous inflammatory mediators, as well as two α.βunsaturated aldehydes (acrolein and crotonaldehyde) generated from the combustion process of cigarette smoking and petrochemical fuels (Bautista et al., 2006; Andrè et al., 2008; Nilius et al., 2012). TRPA1 is expressed in primary sensory fibers and non-neuronal cells such as keratinocytes (Atoyan et al., 2009; Bíró and Kovács, 2009). Some studies have demonstrated that TRPA1 is involved in inflammatory skin symptoms such as erythema, pain and heat hyperalgesia (Silva et al., 2011; Nilius et al., 2012; Brederson et al., 2013). For example, chemicals used in the manufacturing sector, i.e. xylene and formaldehyde, when topically applied to the skin or injected via the intraplantar route, respectively, result in nociception in mice via TRPV1 and TRPA1 activation (McNamara et al., 2007; Sándor et al., 2009; Tian et al., 2009). In addition, it has also been reported that toluene diisocyanate, a hazardous irritant used in paints and foams, causes sensory nerve activation in vitro via TRPA1 gating (Taylor-Clark et al., 2009). However, whether the activation of TRPs, in particular TRPA1, is essential to the development of inflammatory processes, pain and other symptoms observed during irritative contact dermatitis caused by these volatile organic compounds remains unclear. In this context, we investigated the role of TRPA1 in the cutaneous inflammatory responses evoked by chemical irritants such as xylene and toluene after topical application to the skin.

2. Material and methods

2.1. Animals

Experiments were performed on adult male Swiss mice (25-30 g) and wild type $(Trpa1^{+/+})$ or Trpa1-deficient $(Trpa1^{-/-}B6; 129P-Trpa1tm1Kykw/J)$ mice generated in heterozygous animals on a C57BL/6 background (Jackson Laboratories, Bar Harbor, Maine, USA). Animals were maintained in standard housing conditions (12 h light/ dark cycle, lights on at 6 a.m., food and water ad libitum in a climatically controlled environment). The experimental procedures were performed according to the ARRIVE guidelines for reporting experiments involving animals (Kilkenny et al., 2013) and with the standards set in the directives for animal care in Brazilian law n°11.794/2008. All protocols employed have been approved by the Ethics Committee of the Federal University of Rio Grande do Norte (Protocol n°: 023/2010) and Federal University of Paraná (Protocol nº: 698), except for the experiments with $Trpa1^{+/+}$ or $Trpa1^{-/-}$ mice. These were conducted at the University of Florence (research permit #204/2012-B) in conformity with the European Communities Council (ECC) guidelines for animal care procedures and Italian legislative application (DL 116/92) of the ECC directive (86/609/EEC).

2.2. Drugs and reagents

The TRPA1 antagonist 1,2,3,6-Tetrahydro-1,3-dimethyl-N-[4-(1-methylethyl)phenyl]-2,6-dioxo-7H-purine-7-acetamide, 2-(1,3-Dimethyl-2,6dioxo-1,2,3,6-tetrahydro-7H-purin-7-yl)-N-(4-isopropylphenyl)acetamide (HC-030031) was synthesised and purified as previously described (Andrè et al., 2008) and the structure of the compound was assigned by H-NMR and mass spectroscopy (final purity > 98%) determined by RP-HPLC. Toluene and xylene were purchased from Sigma (St. Louis, MO, USA). All drugs used topically were dissolved in acetone immediately prior to use. HC-030031 was prepared in 0.9% NaCl containing 5% dimethylsulphoxide plus 5% Tween 80 for intragastric administration. The solutions were diluted on the day of the experiment immediately before use.

2.3. Volatile chemical irritant-induced ear oedema

Oedema induced by the topical application of chemical irritants was measured as an increase in ear thickness. The chemical irritants were pipetted directly onto the ears of the mice. Ear thickness was assessed with an electronic digital micrometer (Starrett Series 734) applied near the tip of the ear distal to the cartilaginous ridges; the difference in thickness between the basal measurement and after challenge was calculated (Silva et al., 2011). To minimise variation, a single investigator performed the measurements and the values were recorded in mm. Ear thickness was measured before and until 120 min after the topical application of xylene (100%/20 μ /ear), toluene (100%/20 μ /ear) or their vehicle (acetone, 20 μ /ear) onto the right ear.

2.4. Involvement of the TRPA1 receptor in the edematogenic response induced by volatile chemical irritants

Separate groups of animals were topically pretreated with the TRPA1 selective antagonist HC-030031 (300 nmol/20 µl/ear) (Silva et al., 2011) or its vehicle (acetone, 20 µl/ear) 15 min before challenge with the organic compounds. Ear thickness was measured before and 30 min after the topical application of xylene (100%/20 µl/ear), to-luene (100%/20 µl/ear) or vehicle (acetone, 20 µl/ear) to the right ear. In another set of experiments, oedema induced by the topical application of organic compounds in wild type and *Trpa1*^{-/-}mice was evaluated until 120 min after exposure.

2.5. Histology

Histological changes were evaluated in ear samples from mice pretreated with the topical TRPA1 antagonist HC-030031 (300 nmol/ 20 μ /ear) or vehicle in *Trpa1* wild type and *Trpa1^{-/-}*mice, challenged with the organic compounds. Thirty min after the induction of oedema by xylene or toluene, mice were euthanised with thiopental (100 mg/kg, intraperitoneal) and ear tissues were removed and fixed in alfac solution (16:2:1 mixture of 80% ethanol, 40% formaldehyde and glacial acetic acid). Each sample was embedded in paraffin wax, sectioned at 5 μ m and stained with haematoxylin-eosin. A representative area was selected for qualitative light microscopic analysis of the inflammatory cellular response with a 40 × objective lens (Recio et al., 2000). These analysis were carried out by using imagej program. To minimise sources of bias, the investigator analysed the specimens blindly.

2.6. Measurement of behavioural nociceptive responses

In this study, we also investigated whether xylene or toluene could induce acute nociception after applying irritant compounds directly to mouse skin. Briefly, animals were firstly individually placed in transparent glass chambers for 1 h of acclimation. After the adaptation period, xylene (30, 50, 100%/20 µl/paw) or toluene (30, 50, 100%/20 µl/paw) was topically applied (using a Gilson P100 pipette) onto the plantar surface of the right hind paw. Immediately after the topical application, mice were gently introduced into the chamber for the evaluation of the acute nociceptive response. The amount of time spent licking the exposed hind paw for 5 min after chemical irritant application was recorded with a chronometer and was considered a measure of acute nociceptive behaviour.

2.7. Involvement of the TRPA1 receptor in volatile chemical irritant-induced nociceptive responses

To assess the role of the TRPA1 receptor in the nociceptive response, mice received a single oral dose of a TRPA1 antagonist HC-030031 (100 mg/kg; intragastric) (Eid et al., 2008) or vehicle. Sixty min later, xylene (50%/20 µl) or toluene (50%/20 µl) was applied to the right hind paw and the index of the nociceptive response was evaluated for a period of 5 min. In another set of experiments, xylene (50%/20 µl) or toluene (50%/20 µl) was applied to the right hind paw of wild type and $Trpa1^{-/-}$ mice and the nociceptive response was evaluated for a period of 5 min.

2.8. Statistical analysis

Data are reported as the mean \pm standard error of the mean (S.E.M.) of at least 4 animals per group. The statistical analysis was assessed by the unpaired two-tailed Student's t-test (to evaluate statistical significance between two groups) or by repeated measures two-way analysis of variance (ANOVA) followed by Bonferroni's multiple comparison post hoc test. These evaluations were carried out using GraphPad Prism 5 (GraphPad Software, San Diego, CA, USA). Values of $P \leq 0.05$ were considered to be statistically significant.

3. Results

3.1. Effect of topical of xylene or toluene application on ear oedema

Topical application of xylene (100%/20 μ /ear) and toluene (100%/20 μ /ear) induced ear oedema in Swiss mice. The peak of the edematogenic response was observed at 30 min (0.113 \pm 0.008 mm and 0.067 \pm 0.011 mm, respectively) when compared to vehicle (0.008 \pm 0.008 mm) (Fig. 1A, B).

3.2. Effect of pharmacological blockade and genetic deletion of TRPA1 on ear oedema induced by xylene and toluene

In Swiss mice, ear oedema evoked by xylene (100%/20 µl/ear) and toluene (100%/20 µl/ear) was reduced (36.8 ± 9.4% and 50.7 ± 11.0%, respectively) by pretreatment (local application) with the selective TRPA1 antagonist HC-030031 (300 nmol/20 µl/ear, 15 min prior) (Fig. 2A, B). In addition, in *Trpa1^{-/-}*mice, the ear oedema induced by xylene and toluene was strongly reduced (66.6 ± 16.7% and 75 ± 0%, respectively) (calculated on the peak of 30 min) when compared with wild type mice (Fig. 3A, B).

3.3. Histological findings

Cross-sections of mouse ears collected 30 min after being xylene or



Fig. 1. Ear oedema induced by volatile chemical irritants. Time-course of ear oedema induced by topical application xylene (100%/20 µl/ear) and toluene (100%/20 µl/ear) in mice. The effects of these chemical irritants were measured 30, 60, 90 and 120 min after the administration of chemical irritants and were expressed as the change in ear thickness. Each point represents the mean \pm S.E.M. for 8 animals. . *P < 0.05 vs. vehicle (Student's t-test).



Fig. 2. Effect of TRPA1 receptor antagonist on the edematogenic responses induced by volatile chemical irritants. HC-030031 topical pretreatment was performed 15 min before the topical application of xylene (100%/20 µl/ear; A) or toluene (100%/20 µl/ear; B). Ear oedema was measured 30 min after administration of each chemical irritant and were expressed as the change in ear thickness. Each column represents the mean \pm S.E.M. for 8 animals. *P < 0.05 vs. vehicle plus xylene or toluene (Veh₁) (one-way ANOVA followed by Bonferroni's test).



Fig. 3. Ear oedema induced by topical application of volatile chemical irritants in wild type (*Trpa1*^{+/+}) or *TRPA1*-deficient mice (*Trpa1*^{-/-}). Ear oedema was measured 30, 60, 90 and 120 min after the topical administration of xylene (100%/20 µl/ear; A) or toluene (100%/20 µl/ear; B) and are expressed as the change in ear thickness. Each point represents the mean \pm S.E.M. for at least 5 animals. *P < 0.05 vs. *Trpa1*^{+/+} (Student's t-test).

toluene challenge showed differences in the thickness of the histological structures, when compared to vehicle-treated mice (Fig. 4A,B,D,E). Observations performed with the $40 \times$ objective revealed an edematogenic response, mainly between the skin and the cartilage. In contrast, pretreatment with the TRPA1 antagonist HC-030031 prevented the ear oedema induced by both chemical irritants (Fig. 4 C,F).

Increased mouse ear thickness was observed in wild type mice after a single topical application of the chemical irritants (Figs. 5–6). However, no histological alterations were observed in $Trpa1^{-/-}$ mice challenged with xylene (Fig. 5) and toluene (Fig. 6).

3.4. Nociceptive response induced by xylene or toluene application to the mouse paw

The topical application of xylene (50 and 100%/20 μ l/paw) or toluene (50 and 100%/20 μ l/paw) on the plantar surface of the right hind paw of Swiss mice induced licking reactions in a dose-dependent manner, characteristic of a painful sensation (Fig. 7A, B).

3.5. Effects of pharmacological and genetic blockade of the TRPA1 receptor on xylene or toluene-induced nociceptive responses

Xylene (50%/20 µl/paw) and toluene (50%/20 µl/paw) evoked nociceptive responses were significantly reduced (% of inhibition: 84.9 \pm 1.3% and 27.1 \pm 8.0%, respectively) by oral pretreatment with the TRPA1 antagonist HC-030031 (100 mg/kg, 60 min prior the chemical application) in Swiss mice (Fig. 8A, B). Furthermore, the nociceptive response elicited by the topical application of xylene (50%/20



Fig. 4. Representative light microphotograph and graphic of the effects of TRPA1 antagonist on xylene and toluene-induced ear edema in mice. A thin section was obtained from ear tissue of mice pretreated with HC-030031 (300 nmol/20 µl/ear) or vehicle (acetone, 20 µl/ear) 30 min after the topical application of xylene (100%/20 µl/ear), toluene (100%/20 µl/ear) or vehicle (Veh, acetone, 20 µl/ear) (hematoxylineosin staining) x40. (A) Veh- Veh; (B) Veh-Xylene; (C) HC-030031-Xylene; (D) Veh-Toluene; (E) HC-030031-Toluene; (F) Representative graphic of light microphotograph of the effects of TRPA1 antagonist on xylene and toluene-induced ear edema in mice. Each column represents the mean \pm S.E.M. for 8 animals. *P < 0.05 vs. xylene or toluene (one-way ANOVA followed by Bonferroni's test).



 $Trpa1^{+/+}$ $Trpa1^{+/+}$

Fig. 5. Representative light microphotograph and graph indicating the effect of genetic deletion TRPA1 receptor on xylene-induced ear oedema in mice. Thin sections were obtained from the ear tissue of mice *Trpa1^{+/+}* and *Trpa^{-/-}*30 min after the topical application of xylene or vehicle (Veh, 20 µl/ear) (haematox-ylin-eosin staining × 40). *P < 0.05 vs. vehicle (veh), §P < 0.05 vs. xylene, (one-way ANOVA followed by Bonferroni's test).

 μ l/paw; 32.0 \pm 5.1 s) and toluene (50%/20 μ l/paw; 17.4 \pm 2.3 s) was suppressed in mice with genetic deletion of TRPA1 (0.3 \pm 0.1 s and 2.4 \pm 0.8 s, respectively) as compared to wild type mice (Fig. 9A, B).

4. Discussion

Exposure to noxious chemical compounds may occur throughout multiple routes, including dermal contact. Xylene and toluene are common pollutants in the plastic, chemical and leather industries. Skin contact with chemical substances may result in local diseases such as irritant contact dermatitis, the most common type of occupational skin disorder, or allergic dermatitis (Sasseville, 2008). The present study shows that the topical application of chemical irritants, i.e. xylene and toluene, induces robust and long-lasting ear oedema in mice.

By focusing on the role played by TRPA1 signalling in the mediation of skin inflammation induced by xylene and toluene, here we show through ear thickness measurements and histological analyses that pharmacological and genetic blockade of TRPA1 significantly inhibited xylene- and toluene-induced ear oedema. Thus, these findings suggest that some volatile irritant compounds, after contact with the skin, promote vascular reactions mediated by TRPA1 receptor signalling. The role of TRPA1 in skin inflammation has been previously reported by our

Fig. 6. Representative light microphotograph and graph indicating the effect of genetic deletion TRPA1 receptor on toluene-induced ear oedema in mice. Thin sections were obtained from ear tissue of mice $Trpa1^{+/+}$ and $Trpa1^{-/-}$ 30 min after the topical application of toluene or vehicle (Veh, acetone, 20 µl/ear) (haematoxylin-eosin staining ×40). *P < 0.05 vs. vehicle (veh), §P < 0.05 vs. toluene, (one-way ANOVA followed by Bonferroni's test).



Fig. 7. Nociceptive responses induced by volatile chemical irritants. Nociceptive dose-response curves induced by topical application of xylene (30–100%/20 µl/paw; A) or toluene (30–100%/20 µl/paw; B) in the right hind paw of mice. Nociception time was assessed by recording the time spent evoking nociceptive responses during 5 min of observation immediately after exposure to chemical irritants. Each point represents the mean \pm S.E.M. for 8 animals per group. *P < 0.05 compared to the xylene 30% or toluene 30% group (one-way ANOVA followed by Bonferroni's test).



Fig. 8. Effects of TRPA1 antagonist on the nociceptive responses produced by volatile chemical irritants. Effects of oral pretreatment with HC-030031 on the nociceptive responses induced by the topical administration of xylene (50%/20 μ //ear; A) or toluene (50%/20 μ //ear; B) in the hind paw of mice. Nociception time was assessed by recording the time spent evoking nociceptive responses during 5 min of observation immediately after exposure to chemical irritants. Each column represents the mean \pm S.E.M. for at least 8 animals. *P < 0.05 vs. vehicle plus xylene or toluene (Veh₁) (Student's t-test).



Fig. 9. Nociceptive responses induced by the topical application of volatile chemical irritants in wild type $(Trpa1^{+/+})$ or *TRPA1*-deficient mice $(Trpa1^{-/-})$. Nociception time was measured during 5 min after the topical administration of xylene (50%/20 µl/paw; A) or toluene (50%/20 µl/paw; B) in the hind paw of mice. Each point represents the mean \pm S.E.M. for 5 animals. *P < 0.05 vs. TRPA1 wild type mice $(Trpa1^{+/+})$ (Student's t-test).

research group (Silva et al., 2011). We showed that the topical administration to the mouse ear of the TRPA1 receptor agonist cinnamaldehyde promoted inflammatory responses with oedema formation, which was inhibited by pre-exposure to the TRPA1 antagonist HC-030031 (Silva et al., 2011). To reinforce the role of TRPA1 in skin disorders, it has been demonstrated that this channel is expressed in many non-neuronal cell types, such as mast cells, keratinocytes (Bíró and Kovács, 2009) and melanocytes (Bellono et al., 2013), and in sensory nerves, where it works as a nociceptive sensor (Atoyan et al., 2009).

It was interesting to note that, besides oedema and erythema, other relevant clinical manifestations mentioned by patients after dermal exposure to chemical irritants are pain sensations and itching at the site of injury (Ngo and Maibach, 2010). We also observed that xylene and toluene contact with the skin induced an acute nociceptive response. In fact, the topical application of xylene and toluene directly to the mouse paw was sufficient to elicit nociceptive responses, as assessed by an increased amount of time spent licking the treated paw. Taken together, the sensory and vascular manifestations observed after the application of xylene and toluene to the ear and paw of mice mimic the main symptoms of irritant contact dermatitis.

In the present study, we also observed that both pretreatment with the TRPA1 antagonist HC-030031 and genetic deletion of the TRPA1 receptor significantly inhibited xylene-and toluene-induced nociceptive responses when topically applied to the mouse paw. A growing body of evidence supports a role for TRPA1 in mediating pain and pruritic responses caused by several exogenous noxious stimuli (Oh et al., 2013; Wilson et al., 2013; Chen and Hackos, 2015). However, Sándor et al. (2009) employed a similar method to that used in this study and reported that xylene-induced nociception is mediated by the TRPV1 receptor signalling. These authors showed that pharmacological and genetic blockade of TRPV1 partially prevented xylene-induced nociceptive responses (Sándor et al., 2009).

It is interesting to note that a close relationship between TRPA1 and TRPV1 has already been reported. Both TRPV1 and TRPA1 are co-expressed in a large subset of sensory nerves and, using transient co-expression of these two receptors in Chinese hamster ovary cells, Staruschenko et al. (2010) demonstrated a direct interaction between TRPV1 and TRPA1, resulting in heterotetramer formation by these two ion channels. Recently, it has also been demonstrated that Ca²⁺ influx through TRPV1 opening may increase the sensitivity of the TRPA1 channel (Hsu and Lee, 2015). Therefore, it is possible that both TRPA1 and TRPV1, working synergistically, can mediate the cutaneous inflammatory response caused by some volatile irritant chemical compounds.

5. Conclusion

In conclusion, we have shown that the activation of TRPA1 could be involved in some of the symptoms of irritant contact dermatitis, such as pain, neurogenic inflammation and possibly itching induced by the topical exposure of some volatile chemical irritants. The present data also support a role for TRPA1 antagonists as innovative drugs for topical pharmacological intervention during skin inflammation. However, other studies are required to further clarify the role of other TRP receptors in inflammatory skin reactions.

Ethical approval and consent to participate

All protocols employed have been approved by the Ethics Committee of the Federal University of Rio Grande do Norte (Protocol n°: 023/2010) and the Federal University of Paraná (Protocol n°: 698), except for the experiments with Trpa1 + / + or Trpa1 - / - mice. These were conducted at the University of Florence (research permit #204/ 2012-B) in conformity with the European Communities Council (ECC) guidelines for animal care procedures and the Italian legislative application (DL 116/92) of the ECC directive (86/609/EEC).

Consent for publication

Not applicable.

Competing interests

The authors have no conflicts of interest to disclose.

Funding

This work was supported by CNPq (grant number 305058/2018-5) and Araucaria Foundation, Brazil (grant number 37556.413.40855.12042013).

CRediT authorship contribution statement

Maíra Macedo Norões: Methodology, Data curation. Larissa Gonzaga Santos: Methodology, Data curation. Elaine Cristina Gavioli: Formal analysis. Vanessa de Paula Soares Rachetti: Formal analysis. Michel Fleith Otuki: Writing - review & editing. Daniela de Almeida Cabrini: Writing - review & editing. Arthur da Silveira Prudente: Methodology, Data curation. Janiana Raíza Jentsch Matias Oliveira: Writing - original draft. Muryel de Carvalho Gonçalves: Methodology, Data curation. Juliano Ferreira: Conceptualization, Supervision, Funding acquisition. Delia Preti: Conceptualization, Supervision, Funding acquisition. Francesco De Logu: Conceptualization, Supervision, Funding acquisition. Romina Nassini: Conceptualization, Supervision, Funding acquisition. Eunice André: Conceptualization, Supervision, Funding acquisition.

Acknowledgements

J.R.J.M. de Oliveira is a PhD student in Pharmacology and thanks CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico, Brazil) for the fellowship support.

References

- André, E., Campi, B., Materazzi, S., Trevisani, M., Amadesi, S., Massi, D., Creminon, C., Vaksman, N., Nassini, R., Civelli, M., Baraldi, P.G., Poole, D.P., Bunnett, N.W., Geppetti, P., Patacchini, R., 2008. Cigarette smoke-induced neurogenic inflammation is mediated by $\alpha_i\beta$ -unsaturated aldehydes and the TRPA1 receptor in rodents. J. Clin. Investig. https://doi.org/10.1172/JCI34886.
- Atoyan, R., Shander, D., Botchkareva, N.V., 2009. Non-neuronal expression of transient receptor potential type A1 (TRPA1) in human skin. J. Investig. Dermatol. https://doi. org/10.1038/jid.2009.58.
- Bautista, D.M., Jordt, S.E., Nikai, T., Tsuruda, P.R., Read, A.J., Poblete, J., Yamoah, E.N., Basbaum, A.I., Julius, D., 2006. TRPA1 mediates the inflammatory actions of environmental irritants and proalgesic agents. Cell. https://doi.org/10.1016/j.cell. 2006.02.023.
- Bellono, N.W., Kammel, I.G., Zimmerman, A.L., Oancea, E., 2013. UV light phototransduction activates transient receptor potential A1 ion channels in human melanocytes. Proc. Natl. Acad. Sci. https://doi.org/10.1073/pnas.1215555110.
- Bíró, T., Kovács, L., 2009. An ice-Cold TR(i)P to skin biology: the role of TRPA1 in human epidermal keratinocytes. J. Investig. Dermatol. https://doi.org/10.1038/jid.2009. 179
- Boonen, B., Startek, J.B., Talavera, K., 2017. Chemical activation of sensory TRP channels. Top. Med. Chem. https://doi.org/10.1007/7355_2015_98.
- Brederson, J.D., Kym, P.R., Szallasi, A., 2013. Targeting TRP channels for pain relief. Eur. J. Pharmacol. https://doi.org/10.1016/j.ejphar.2013.03.003.
- Cao, L.Y., Taylor, J.S., 2008. Contact dermatitis and related disorders. In: ACP Medicine. Section 2. Dermatology. BC Decker, Hamilton.
- Caterina, M.J., Pang, Z., 2016. TRP channels in skin biology and pathophysiology. Pharmaceuticals. https://doi.org/10.3390/ph9040077.
- Chen, J., Hackos, D.H., 2015. TRPA1 as a drug target promise and challenges. Naunyn-Schmiedeberg's Arch. Pharmacol. https://doi.org/10.1007/s00210-015-1088-3.
- Eid, S.R., Crown, E.D., Moore, E.L., Liang, H.A., Choong, K.C., Dima, S., Henze, D.A.,

Kane, S.A., Urban, M.O., 2008. HC-030031, a TRPA1 selective antagonist, attenuates inflammatory- and neuropathy-induced mechanical hypersensitivity. Mol. Pain. https://doi.org/10.1186/1744-8069-4-48.

- Hsu, C.-C., Lee, L.-Y., 2015. Role of calcium ions in the positive interaction between TRPA1 and TRPV1 channels in bronchopulmonary sensory neurons. J. Appl. Physiol. https://doi.org/10.1152/japplphysiol.00043.2015.
- Kaplan, D.H., Igyártó, B.Z., Gaspari, A.A., 2012. Early immune events in the induction of allergic contact dermatitis. Nat. Rev. Immunol. https://doi.org/10.1038/nri3150.
- Kilkenny, C., Browne, W.J., Cuthill, I.C., Emerson, M., Altman, D.G., 2013. Improving Bioscience Research Reporting: the Arrive Guidelines for Reporting Animal Research. Animals. https://doi.org/10.3390/ani4010035.
- McNamara, C.R., Mandel-Brehm, J., Bautista, D.M., Siemens, J., Deranian, K.L., Zhao, M., Hayward, N.J., Chong, J.A., Julius, D., Moran, M.M., Fanger, C.M., 2007. TRPA1 mediates formalin-induced pain. Proc. Natl. Acad. Sci. https://doi.org/10.1073/ pnas.0705924104.
- Ngo, M.A., Maibach, H.I., 2010. Dermatotoxicology: historical perspective and advances. Toxicol. Appl. Pharmacol.
- Nilius, B., Appendino, G., Owsianik, G., 2012. The transient receptor potential channel TRPA1: from gene to pathophysiology. Pflugers Arch. Eur. J. Physiol. https://doi. org/10.1007/s00424-012-1158-z.
- Oh, M.-H., Oh, S.Y., Lu, J., Lou, H., Myers, A.C., Zhu, Z., Zheng, T., 2013. TRPA1-Dependent pruritus in IL-13-induced chronic atopic dermatitis. J. Immunol. https:// doi.org/10.4049/jimmunol.1300300.
- Recio, M.C., Giner, R.M., Uriburu, L., Máñez, S., Cerdá, M., De la Fuente, J.R., Ríos, J.L., 2000. In vivo activity of pseudoguaianolide sesquiterpene lactones in acute and chronic inflammation. Life Sci. 66 (26), 2509–2518.
- Sándor, K., Helyes, Z., Elekes, K., Szolcsányi, J., 2009. Involvement of capsaicin-sensitive afferents and the Transient Receptor Potential Vanilloid 1 Receptor in xylene-induced nocifensive behaviour and inflammation in the mouse. Neurosci. Lett. https://doi. org/10.1016/j.neulet.2009.01.016.
- Sasseville, D., 2008. Occupational contact dermatitis. Allergy Asthma Clin. Immunol. https://doi.org/10.1186/1710-1492-4-2-59.
- Silva, C.R., Oliveira, S.M., Rossato, M.F., Dalmolin, G.D., Guerra, G.P., Prudente, A.D.S., Cabrini, D.A., Otuki, M.F., André, E., Ferreira, J., 2011. The involvement of TRPA1 channel activation in the inflammatory response evoked by topical application of cinnamaldehyde to mice. Life Sci. https://doi.org/10.1016/j.lfs.2011.03.017.
- Staruschenko, A., Jeske, N.A., Akopian, A.N., 2010. Contribution of TRPV1-TRPA1 interaction to the single channel properties of the TRPA1 channel. J. Biol. Chem. https://doi.org/10.1074/jbc.M110.106153.
- Taylor-Clark, T.E., Kiros, F., Carr, M.J., McAlexander, M.A., 2009. Transient receptor potential ankyrin 1 mediates toluene diisocyanate-evoked respiratory irritation. Am. J. Respir. Cell Mol. Biol. https://doi.org/10.1165/rcmb.2008-02920C.
- Tian, L.J., DU, Y.R., Xiao, Y., Lv, Z.M., Yu, Y.Q., Cui, X.Y., Chen, J., 2009. Mediating roles of the vanilloid receptor TRPV1 in activation of rat primary afferent nociceptive neurons by formaldehyde. Sheng Li Xue Bao 61 (5), 404–416.
- Wilson, S.R., Nelson, A.M., Batia, L., Morita, T., Estandian, D., Owens, D.M., Lumpkin, E.A., Bautista, D.M., 2013. The ion channel TRPA1 is required for chronic itch. J. Neurosci. https://doi.org/10.1523/jneurosci.5318-12.2013.