



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Assessing the Burden of Nondeferrable Major Uro-oncologic Surgery to Guide Prioritisation Strategies During the COVID-19 Pandemic:

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Assessing the Burden of Nondeferrable Major Uro-oncologic Surgery to Guide Prioritisation Strategies During the COVID-19 Pandemic: Insights from Three Italian High-volume Referral Centres / Campi R., Amparore D., Capitanio U., Checcucci E., Salonia A., Fiori C., Minervini A., Briganti A., Carini M., Montorsi F., Serni S., Porpiglia F.. - In: EUROPEAN UROLOGY. - ISSN 0302-2838. - ELETTRONICO. - (2020),

Availability:

The webpage <https://hdl.handle.net/2158/1193724> of the repository was last updated on 2020-05-20T17:39:01Z

Published version:

DOI: 10.1016/j.eururo.2020.03.054

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proof

Assessing the Burden of Nondeferrable Major Uro-oncologic Surgery to Guide Prioritisation Strategies During the COVID-19 Pandemic: Insights from Three Italian High-volume Referral Centres

Riccardo Campi, Daniele Amparore, Umberto Capitanio, Enrico Checcucci, Andrea Salonia, Cristian Fiori, Andrea Minervini, Alberto Briganti, Marco Carini, Francesco Montorsi, Sergio Serni, Francesco Porpiglia



PII: S0302-2838(20)30229-3

DOI: <https://doi.org/10.1016/j.eururo.2020.03.054>

Reference: EURURO 8796

To appear in: *European Urology*

Accepted Date: 31 March 2020

Please cite this article as: { doi: <https://doi.org/>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

Assessing the Burden of Nondeferrable Major Uro-oncologic Surgery to Guide Prioritisation Strategies During the COVID-19 Pandemic: Insights from Three Italian High-volume Referral Centres

Riccardo Campi ^{a,b,c,†,*}, Daniele Amparore ^{d,†}, Umberto Capitanio ^e, Enrico Checcucci ^d, Andrea Salonia ^{e,f}, Cristian Fiori ^d, Andrea Minervini ^{b,c}, Alberto Briganti ^{e,f}, Marco Carini ^{b,c,‡}, Francesco Montorsi ^{e,f,‡}, Sergio Serni ^{a,c,‡}, Francesco Porpiglia ^{d,‡}

^a Department of Urological Robotic Surgery and Renal Transplantation, Careggi Hospital, University of Florence, Florence, Italy

^b Department of Experimental and Clinical Medicine, University of Florence, Florence, Italy

^c Department of Urological Oncologic Minimally Invasive Robotic Surgery and Andrology, Careggi Hospital, , University of Florence, Florence, Italy

^d Division of Urology, Department of Oncology, School of Medicine, San Luigi Hospital, University of Turin, Orbassano, Turin, Italy

^e Division of Experimental Oncology/Unit of Urology, Urological Research Institute, IRCCS Ospedale San Raffaele, Milan, Italy

^f University Vita-Salute San Raffaele, Milan, Italy

[†] These authors contributed equally to this work.

[‡] These authors contributed equally to senior authorship.

* Corresponding author. Chirurgia Urologica Robotica Mini-Invasiva e dei Trapianti Renali, Azienda Ospedaliera Careggi, Università di Firenze, Università degli Studi di Firenze, Viale San Luca, 50134 Firenze, Italy. Tel. +39 05 52758011; Fax: +39 05 52758014.
E-mail address: riccardo.campi@gmail.com (R. Campi).

Abstract

The coronavirus 2019 (COVID-19) pandemic has led to an unprecedented emergency scenario for all aspects of health care, including urology. At the time of writing, Italy was the country with the highest rates of both infection and mortality. A panel of experts recently released recommendations for prioritising urologic surgeries in a low-resource setting. Of note, major cancer surgery represents a compelling challenge. However, the burden of these procedures and the impact of such recommendations on urologic practice are currently unknown. To fill this gap, we assessed the yearly proportion of high-priority major uro-oncologic surgeries at three Italian high-volume academic centres. Of 2387 major cancer surgeries, 32.3% were classified as high priority (12.6% of radical nephroureterectomy, 17.3% of nephrectomy, 33.9% of radical prostatectomy, and 36.2% of radical cystectomy cases). Moreover, 26.4% of high-priority major cancer surgeries were performed in patients at higher perioperative risk (American Society of Anesthesiologists score ≥ 3), with radical cystectomy contributing the most to this cohort (50%). Our real-life data contextualise ongoing recommendations on prioritisation strategies during the current COVID-19 pandemic, highlighting the need for better patient selection for surgery. We found that approximately two-thirds of elective major uro-oncologic surgeries can be safely postponed or changed to another treatment modality when the availability of health care resources is reduced.

Patient summary: We used data from three high-volume Italian academic urology centres to evaluate how many surgeries performed for prostate, bladder, kidney, and upper tract urothelial cancer can be postponed in times of emergency. We found that approximately two-thirds of patients with these cancers do not require high-priority surgery. Conversely, of patients requiring high-priority surgery, approximately one in four is considered at high perioperative risk. These patients may pose challenges in allocation of resources in critical scenarios such as the current COVID-19 pandemic.

Keywords: Coronavirus; COVID-19; High priority; Italy; Major cancer surgery; Pandemic

The coronavirus 2019 (COVID-19) pandemic has led to an unprecedented emergency scenario for all aspects of health care, including urology [1–4]. At the time of writing, Italy was the country with the highest rates of infection and mortality from COVID-19. According to the updated Italian statistics released on March 25, 2020, 74 386 people had been infected, with 7503 deaths [5]. The dramatic spread of this epidemic is leading to major changes in the management of patients with cancer, including those affected by genitourinary malignancies [6]. In this context, virtually all urology centres have been forced to prioritise surgical interventions for cancer patients by applying a series of restrictions on elective approaches to optimise health care resources and minimise the risk of hospital-acquired infection [4,7–9]. This prioritisation process started early in Italy [3,4].

By looking at the ongoing Italian situation, many opinion leaders worldwide have proposed recommendations for the triage of elective urologic surgeries during the COVID-19 pandemic. A panel of experts from Europe and the USA recently proposed a list of urologic procedures that should be prioritised, taking into account factors that include the aggressiveness of each disease (and its stage), the impact of short-term delays to care, and the availability of alternative treatment modalities [7]. Of note, the focus should be on major cancer surgery, which represents a large and demanding proportion of practice in most referral centres.

However, the exact burden of major urologic cancer procedures is unknown and the impact of such recommendations on urologic practice still needs to be investigated. The importance of such an analysis is even more important given the heterogeneous outcomes for uro-oncologic patients and the possible need to shift surgical candidates to other forms of treatment as an alternative to surgery.

To fill these gaps, we provide a real-life overview of the yearly proportion of high-priority major uro-oncologic surgeries at three academic referral centres in northern and central Italy (San Luigi Hospital in Turin, San Raffaele Hospital in Milan, and Careggi Hospital in Florence, all performing >500 major cancer procedures annually), where the COVID-19 pandemic has put a strain on health care resources.

We queried our respective institutional prospectively collected databases for the most common urologic cancers (prostate, urothelial, and kidney) to select patients undergoing elective major surgery during a 12-mo period (2018 or 2019). Among the surgeries, Stensland et al [7] included radical cystectomy (RC), radical nephroureterectomy (RNU), nephrectomy (NEP) for cT3+ disease, and radical prostatectomy (RP) for high-risk patients if not eligible for radiation. From an analytic standpoint, besides all RC, RNU, and NEP procedures for cT3+ tumours, we also considered NEP for cT2 masses as high-priority surgery, as the decision-making is highly nuanced for several patient-specific considerations and the tumour growth rate [7]. In addition, we included RPs for patients with locally

advanced prostate cancer (defined according to the European Association of Urology guidelines [9]) because even though alternative treatment strategies may be offered, their availability for all eligible patients could be potentially limited by logistic issues in times of emergency.

Overall, 2387 patients were included. Of these, 771 (32.3%) were classified as high priority (Fig. 1). Stratifying these surgeries by cancer type, RNU, NEP, RP, and RC accounted for 12.6%, 17.3%, 33.9%, and 36.2%, respectively, of high-priority procedures. High-priority RPs represented 11.0% of all major cancer surgeries performed in our centres, with NEPs accounting for 5.6%, RCs for 11.7%, and RNUs for 4.1%.

Minimally invasive surgery (MIS) was used for >90% of high-priority RPs, and ranged between 40% for RC and 61% for high-priority NEP among the other surgeries. The average length of hospitalisation (LOH) for high-priority major oncologic surgeries at our centres is higher than the data reported by Stensland et al [7]; this is potentially explained by differences in health care systems across countries. Of note, MIS at our centres appears to have an impact on LOH for all procedures except for RC.

It is notable that a non-negligible proportion of patients undergoing high-priority major cancer surgeries are at higher perioperative risk (with potentially longer LOH and a greater need for postoperative monitoring in intensive care units), posing challenges in resource allocation during the current COVID-19 pandemic [2–4].

Using an American Society of Anesthesiologists (ASA) score of ≥ 3 as a surrogate metric, Figure 2 provides an overview of the burden of these patients. Overall, 26.4% of high-priority major cancer surgeries were performed in patients at higher perioperative risk. Patients undergoing RC contributed the most to this cohort (50%). Stratifying ASA ≥ 3 cases by procedure, high-priority RP accounted for 14% NEP for 26%, RNU for 32.0%, and RC for 36%.

Our study provides data that may contextualise the ongoing recommendations [7] on selection of priority major cancer surgeries in the increasingly challenging health care situation caused by the COVID-19 pandemic.

The key finding is that in times of emergency that necessarily lead to a dramatic reduction in operating room availability for urologic procedures, 67.8% of elective major uro-oncologic surgeries can be postponed. Furthermore, of the remaining 32.2% of patients requiring high-priority surgery, a non-negligible proportion might be shifted to alternative treatment strategies if available, after shared decision-making. Finally, careful selection of surgical candidates is key during the COVID-19 pandemic considering that 26.2% of high-priority major surgeries were performed in patients with high preoperative risk. For these patients, use of alternative treatment modalities may be even more important to further reduce the potential need for intensive care units given the limited resources.

Overall, this study may have practical implications considering that in our three centres a progressive significant reduction in or even complete interruption of all elective procedures was mandated, starting 2 wk after the first certified COVID-19 case in Italy.

Our findings should be contextualised considering that many other factors related to patients, resources, and health care scenarios (eg, high- vs low-volume centre) impact on surgical indications and decision-making when prioritising and scheduling surgeries, especially in times of emergency such as the COVID-19 pandemic.

In conclusion, our real-life data from three Italian academic referral centres offer the urology community and policymakers insights into the burden of major cancer procedures that can be postponed in times of emergency. Our study can help other urology centres worldwide in adapting their management strategies for the inflow of uro-oncologic patients in light of the expected reduction in resources caused by the COVID-19 pandemic.

Author contributions: Riccardo Campi had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Campi, Amparore, Serni, Porpiglia.

Acquisition of data: Campi, Amparore, Checcucci, Capitanio.

Analysis and interpretation of data: Campi, Amparore.

Drafting of the manuscript: Campi, Amparore.

Critical revision of the manuscript for important intellectual content: Checcucci, Fiori, Porpiglia, Capitanio, Salonia, Briganti, Montorsi, Carini, Serni, Minervini.

Statistical analysis: Campi, Amparore.

Obtaining funding: None.

Administrative, technical, or material support: None.

Supervision: Serni, Carini, Montorsi, Porpiglia.

Other: None.

Financial disclosures: Riccardo Campi certifies that all conflicts of interest, including specific financial interests and relationships and affiliations relevant to the subject matter or materials discussed in the manuscript (eg, employment/affiliation, grants or funding, consultancies, honoraria, stock ownership or options, expert testimony, royalties, or patents filed, received, or pending), are the following: None.

Funding/Support and role of the sponsor: None.

References

- 1 World Health Organization. Report of the WHO-China Joint Mission on coronavirus disease 2019 (COVID-19). [www.who.int/publications-detail/report-of-the-who-china-joint-mission-on-coronavirus-disease-2019-\(covid-19\)](http://www.who.int/publications-detail/report-of-the-who-china-joint-mission-on-coronavirus-disease-2019-(covid-19))
- 2 Remuzzi A, Remuzzi G. COVID-19 and Italy: what next? *Lancet*. In press. [https://doi.org/10.1016/S0140-6736\(20\)30627-9](https://doi.org/10.1016/S0140-6736(20)30627-9)
- 3 Naspro R, Da Pozzo LF. Urology in the time of corona. *Nat Rev Urol*. In press. <https://doi.org/10.1038/s41585-020-0312-1>
- 4 Ficarra V, Novara G, Abrate A, et al. Urology practice during COVID-19 pandemic. *Minerva Urol Nefrol*. In press. <https://doi.org/10.23736/S0393-2249.20.03846-1>
- 5 Italian Ministry of Health. Nuovo coronavirus. www.salute.gov.it/nuovocoronavirus
- 6 Gillesen Sommer S, Powles T. Advice for medical oncology care of urological cancer patients during the COVID-19 pandemic. *Eur Urol*. In press.
- 7 Stensland KD, Morgan TM, Moinzadeh A, et al. Considerations in the triage of urologic surgeries during the COVID-19 pandemic. *Eur Urol*. In press.
- 8 American College of Surgeons. COVID-19: guidance for triage of non-emergent surgical procedures. www.facs.org/about-acs/covid-19/information-for-surgeons/triage
- 9 Mottet N, Cornford P, van den Bergh RCN, et al. EAU-EANM-ESTRO-ESUR-SIOG guidelines on prostate cancer, version 2020. <https://uroweb.org/guideline/prostate-cancer/>

Fig. 1 – Cumulative proportion of patients undergoing high-priority major cancer surgeries, stratified by cancer type, procedure, and surgical approach. (A) Overall proportion of patients undergoing high-priority major cancer surgeries and relative contribution of each cancer to the overall number of high-priority surgeries. (B) Cumulative proportion of patients undergoing high-priority radical prostatectomy, nephrectomy, radical cystectomy, and radical nephroureterectomy among all major cancer surgeries ($n = 2387$). (C) Cumulative proportion of high-priority major cancer surgeries via minimally invasive (dark grey) or open surgery (light grey) and average length of hospitalisation (LOH; range in days) for each procedure. All data are for either 2018 (from January 1 to December 31) or 2019 (from January 1 to December 31). BCa = bladder cancer; MIS = minimally invasive surgery; PCa = prostate cancer; RCC = renal cell carcinoma; UTUC = upper tract urothelial carcinoma.




Fig. 2 – Proportion of patients with higher perioperative risk, defined as ASA score ≥ 3 , undergoing high-priority major cancer surgeries. (A) Overall proportion of patients with ASA score ≥ 3 undergoing high-priority major cancer surgeries and relative contribution of each cancer to the overall number of high-priority patients with ASA score ≥ 3 ($n = 204$). (B) Percentage of patients with ASA score ≥ 3 undergoing high-priority radical prostatectomy (RP), nephrectomy (NEP), radical cystectomy (RC), or radical nephroureterectomy (RNU) as a proportion of all high-priority cases for each procedure. All data are for either 2018 (from January 1 to December 31) or 2019 (from January 1 to December 31).

ASA = American Society of Anesthesiologists; BCa = bladder cancer; PCa = prostate cancer; RCC = renal cell carcinoma; UTUC = upper tract urothelial carcinoma.

