



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Rearrangement and Polarization

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Rearrangement and Polarization / GABRIELE BIANCHI, RICHARD J. GARDNER, PAOLO GRONCHI, MARKUS KIDERLEN. - In: ADVANCES IN MATHEMATICS. - ISSN 0001-8708. - ELETTRONICO. - 374:(2020), pp. 0-0. [10.1016/j.aim.2020.107380]

Availability:

The webpage <https://hdl.handle.net/2158/1201955> of the repository was last updated on 2024-12-14T16:20:55Z

Published version:

DOI: 10.1016/j.aim.2020.107380

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

REARRANGEMENT AND POLARIZATION

GABRIELE BIANCHI, RICHARD J. GARDNER, PAOLO GRONCHI, AND MARKUS KIDERLEN

ABSTRACT. The paper has two main goals. The first is to take a new approach to rearrangements on certain classes of measurable real-valued functions on \mathbb{R}^n . Rearrangements are maps that are monotonic (up to sets of measure zero) and equimeasurable, i.e., they preserve the measure of super-level sets of functions. All the principal known symmetrization processes for functions, such as Steiner and Schwarz symmetrization, are rearrangements, and these have a multitude of applications in diverse areas of the mathematical sciences. The second goal is to understand which properties of rearrangements characterize polarization, a special rearrangement that has proved particularly useful in a number of contexts. In order to achieve this, new results are obtained on the structure of measure-preserving maps on convex bodies and of rearrangements generally.

1. INTRODUCTION

The idea of replacing an object by one that retains some of its features but is in some sense more symmetrical has been extremely fruitful. The object may be a set or a function, for example, and the process is then often called symmetrization or rearrangement, respectively. Steiner symmetrization, introduced by Jakob Steiner around 1836 in his attempt to prove the isoperimetric inequality, is still today a potent tool for establishing crucial inequalities in geometry; see, for example, [27, 36, 37, 38]. The influence of such inequalities, which often have analytical versions, extends far beyond geometry to other areas such as analysis and PDEs, and even outside mathematics, to economics and finance. The books [9], [23, Chapter 9], [26, Chapter 9], [43, Chapter 10], and survey [22] should serve as gateways to the literature.

The topic received a huge boost in 1951 from the classic text of Pólya and Szegő [40]. By this time, many other types of symmetrization had been introduced, with similar applications. The general idea is to find a symmetrization that preserves one physical quantity, while not increasing (or sometimes not reducing) another. As well as volume, surface area, and mean width, [40] considers electrostatic capacity, principal frequency (the first eigenvalue of the Laplacian), and torsional rigidity, thereby extending the scope to mathematical physics. In

2010 *Mathematics Subject Classification.* Primary: 28A20, 52A20; secondary: 46E30, 52A38.

Key words and phrases. Convex body, Steiner symmetrization, Schwarz symmetrization, Minkowski symmetrization, two-point symmetrization, rearrangement, polarization.

First and third author supported in part by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). Second author supported in part by U.S. National Science Foundation Grant DMS-1402929. Fourth author supported by the Centre for Stochastic Geometry and Advanced Bioimaging, funded by a grant from the Villum Foundation.

fact, much of this work was motivated by conjectures of the mathematician-engineer de Saint-Venant (1856) and physicist Strutt (a.k.a. Lord Rayleigh) (1877), and subsequent work of Hadamard, Poincaré, and others. The latter included results on rearrangement of functions, already used to great effect in the 1920s by Faber and Krahn. It turns out that rearranging a function is a notion so fertile that applications arise in areas too diverse for a single text to cover them all in detail: Classical analysis, calculus of variations, complex analysis, convex geometry, geometric measure theory, Banach spaces, potential theory, PDEs, fluid dynamics, mechanics, and meteorology, for example. Luckily, a beautiful and quite recent survey by Talenti [47] contains a comprehensive bibliography, conveniently divided between the main periods of development, from which [5, 10, 12, 13, 16, 21, 29, 30, 31, 32, 33, 35, 46], together with the recent book [3], illustrate the list of areas just mentioned.

The Steiner rearrangement of a function f with respect to an $(n - 1)$ -dimensional subspace H in \mathbb{R}^n , and its natural generalization, the Schwarz rearrangement of f with respect to a k -dimensional subspace H , are defined via the corresponding symmetrals of its super-level sets; see, for example, [26, p. 178]. For convenience, we shall denote either rearrangement by $S_H f$. Symmetrals of sets can be identified with rearrangements of their characteristic functions. The related notion of polarization is more recent but has already stimulated much interest. According to Solynin [45, p. 123], it was first considered for planar sets by Wolontis [52] in 1952, and for functions by Ahlfors [1, p. 34] and Baernstein and Taylor [4] in the 1970s. The term itself is due to Dubinin [17]. The standard *polarization* process, sometimes called *two-point symmetrization*, with respect to an *oriented* $(n - 1)$ -dimensional (linear) subspace H , takes a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and replaces it by

$$(1) \quad P_H f(x) = \begin{cases} \max\{f(x), f(x^\dagger)\}, & \text{if } x \in H^+, \\ \min\{f(x), f(x^\dagger)\}, & \text{if } x \in H^-, \end{cases}$$

where \dagger denotes the reflection in H and where H^+ , H^- , are the two closed half-spaces bounded by H and determined by its orientation. If $A \subset \mathbb{R}^n$, then $P_H A$ is the set satisfying $1_{P_H A} = P_H 1_A$, where 1_A denotes the characteristic function of A . The process has several useful properties: It is equimeasurable, monotonic, L^p -contracting, and reduces the modulus of continuity (see Section 3 for the definitions of these terms and references).

The article [4] demonstrated that polarizations can be more efficient than rearrangements in establishing inequalities involving integrals, and was followed by a number of papers applying polarization to inequalities in the theory of capacities. For example, Dubinin [18] generalized the result of Wolontis [52] by showing that the generalized capacity of a condenser in \mathbb{R}^n does not increase under polarization. Then, in 2000, a landmark study by Brock and Solynin [8] gave further significance to polarization by showing that the Steiner or Schwarz rearrangements of a function (or symmetrals of a compact set) with respect to a subspace H can be approximated in $L^p(\mathbb{R}^n)$ (or in the Hausdorff metric, respectively) via successive polarizations with respect to a sequence (H_k) of oriented subspaces. In [8], the sequence (H_k) may depend on the function or set, but this dependence was removed by Van Schaftingen [49, 50]. Indeed,

by [50, Theorem 1 and Section 4.3], the desired approximation of the Steiner or Schwarz rearrangement $S_H f$ of a suitable function f may be obtained by taking any sequence (H_k) dense in the set of oriented subspaces J such that J^+ contains H in its interior and defining $f_1 = f$ and

$$f_{k+1} = (P_{H_k} \circ P_{H_{k-1}} \circ \cdots \circ P_{H_1})f_k$$

for $k \in \mathbb{N}$; then $f_k \rightarrow S_H f$ as $k \rightarrow \infty$. Moreover, polarization is flexible enough to approximate other processes, such as spherical rearrangement and spherical cap symmetrization; see [50]. We refer to [3, Sections 1.7 and 2.3] for a general introduction to polarization. In addition to the references given there and those provided above, more recent work includes [11, 15, 19, 28].

In [7], an investigation was initiated into symmetrization processes defined, like Steiner symmetrization, with respect to a subspace in \mathbb{R}^n . Characterizations of Steiner symmetrization and others such as Minkowski symmetrization were proved, in terms of basic properties they possess. It is natural, then, to undertake a similar study with a view to obtaining characterizations of polarization. Here we consider general maps $T : X \rightarrow X$, where X is $\mathcal{M}(\mathbb{R}^n)$ (or $\mathcal{M}_+(\mathbb{R}^n)$), the space of real-valued (or nonnegative, respectively) measurable functions on \mathbb{R}^n , the space $\mathcal{S}(\mathbb{R}^n)$ of symmetrizable functions in $\mathcal{M}(\mathbb{R}^n)$, or the space $\mathcal{V}(\mathbb{R}^n)$ of functions in $\mathcal{M}_+(\mathbb{R}^n)$ vanishing at infinity. (See Sections 2 and 3 for definitions and terminology.) For any $T : X \rightarrow X$ we can consider the induced map $\diamond_T : \mathcal{L}^n \rightarrow \mathcal{L}^n$, where \mathcal{L}^n is the class of measurable sets in \mathbb{R}^n of finite measure: If $A \in \mathcal{L}^n$, we let $\diamond_T A$ be the set of all $x \in \mathbb{R}^n$ with $T1_A(x) = 1$.

Our new results begin in Section 4 on equimeasurable maps from a subset X of $\mathcal{M}(\mathbb{R}^n)$ to itself, i.e., those that preserve the measure of super-level sets, and rearrangements, equimeasurable maps that are also monotonic. This second and different usage of the term rearrangement—it is now a transformation on a class of functions—is appropriate, since Steiner, Schwarz, and other special rearrangements all have these two properties; see, for example, [31, Section II.2]. Note, however, that the present paper differs in that monotonic really means essentially monotonic, i.e., up to sets of \mathcal{H}^n -measure zero. The first main result is Lemma 4.1(iii), which states that a rearrangement $T : \mathcal{M}(\mathbb{R}^n) \rightarrow \mathcal{M}(\mathbb{R}^n)$ essentially acts as the identity on constant functions. Even the special case $T0 = 0$, essentially, of this natural result seems not to be obvious. This is applied to prove that a rearrangement $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$ essentially satisfies the weak linearity property (18) in Lemma 4.7, which is in turn a crucial ingredient in the second main result, Theorem 4.8. The latter provides the explicit formula (22) for Tf , where $f \in \mathcal{S}(\mathbb{R}^n)$, in terms of the induced map \diamond_T defined above. Theorem 4.8 is also used in proving Theorem 4.9, which establishes the fundamental formula (see (28) below) $\varphi(Tf) = T(\varphi \circ f)$, for $f \in \mathcal{S}(\mathbb{R}^n)$ and right-continuous increasing (i.e., non-decreasing) functions $\varphi : \mathbb{R} \rightarrow \mathbb{R}$. The formula (18), and versions of (22) and Theorem 4.9, appear elsewhere in the literature (compare, for example, [51, p. 138], [8, Equations (3.1) and (3.6), p. 1762], and [51, Definition 4 and Proposition 3(d)]), so we must stress that our approach is quite different and more logical and general. Earlier works such as [8] and [51] begin with set transformations and use them to define special maps on classes of functions, whereas we start with general maps T on classes of functions, define the induced set transformation \diamond_T , and show that in the main situations

of interest, \diamond_T determines T . In particular, [8, Equation (3.1), p. 1762], like [51, Definition 4], is a definition, not a result, and [8, Equation (3.6), p. 1762] and [51, Proposition 3(d)] are deduced from these definitions. See the Appendix for a more detailed comparison of the two approaches.

Polarization has another basic property in addition to those listed above, namely, it is defined *pointwise*, as is clear from (1). General pointwise maps $T : X \rightarrow X$ with respect to an oriented subspace H , defined by (4) below, are the focus of Section 5. Theorem 5.1 gives an explicit formula for maps $T : X \rightarrow X$ that are both pointwise with respect to H and equimeasurable, where $X = \mathcal{M}(\mathbb{R}^n)$, $\mathcal{M}_+(\mathbb{R}^n)$, $\mathcal{S}(\mathbb{R}^n)$, or $\mathcal{V}(\mathbb{R}^n)$. The other main result in this section, Theorem 5.8, shows that once $T : X \rightarrow X$ has these two properties, the others listed above—monotonicity, L^p -contracting for $p \geq 1$, and modulus of continuity reducing—are all equivalent and characterize T as being one of four maps: Id , \dagger , P_H , or $P_H^\dagger = \dagger \circ P_H$, where Id and \dagger denote the identity map and reflection in H , respectively.

The pointwise property is a strong one and for the rest of the paper it is discarded. In Section 4 we examine general maps $T : X \rightarrow X$. Our approach is to gain knowledge first about the induced maps $\diamond_T : \mathcal{L}^n \rightarrow \mathcal{L}^n$. With this aim, in Section 6, we study general maps $\diamond : \mathcal{E} \subset \mathcal{L}^n \rightarrow \mathcal{L}^n$ between sets in terms of various properties, defined in Section 3, but for the most part self-explanatory. The first main result is Theorem 6.6, which gives a formula for maps $\diamond : \mathcal{K}_n^n \rightarrow \mathcal{L}^n$ that are monotonic, measure preserving, respect H -cylinders, and map balls to balls, where \mathcal{K}_n^n is the set of convex bodies in \mathbb{R}^n and H is an $(n-1)$ -dimensional subspace. The formula (see (63)) shows that for such maps there is a contraction $\varphi_\diamond : \mathbb{R} \rightarrow \mathbb{R}$ such that if $K \in \mathcal{K}_n^n$, almost all chords of K orthogonal to H are moved orthogonally to H by a distance determined by φ_\diamond and the position of the chord relative to H . With this in hand, Corollary 6.7 states that $\varphi_\diamond(t) = t, -t, |t|$, or $-|t|$, if and only if \diamond essentially (i.e., up to sets of the appropriate measure zero) equals Id , \dagger , \diamond_{P_H} , or $\diamond_{P_H}^\dagger = \diamond_{P_H^\dagger}$, respectively. The goal then is to find additional or stronger properties that will force φ_\diamond to be one of these four functions. A first attempt replaces the ball-preserving property by one that for measure-preserving maps is stronger and also enjoyed by polarization, namely, that \diamond preserves perimeter on convex bodies. However, Theorem 6.10 shows that maps $\diamond : \mathcal{K}_n^n \rightarrow \mathcal{L}^n$ that are monotonic, measure preserving, respect H -cylinders, and preserve perimeter on convex bodies, are precisely those for which the contraction φ_\diamond satisfies the eikonal equation $|\varphi_\diamond'(t)| = 1$ for almost all $t \in \mathbb{R}$. There are clearly solutions to the latter equation other than $\varphi_\diamond(t) = \pm t$ or $\pm|t|$. To achieve our goal, we find it necessary to focus on maps $\diamond : \mathcal{E} \rightarrow \mathcal{L}^n$, where \mathcal{E} is either the class \mathcal{C}^n of compact sets in \mathbb{R}^n or \mathcal{L}^n , and replace the property that \diamond respects H -cylinders by the stronger one of invariance on H -symmetric unions of two disjoint balls. (Note that while this condition may seem peculiar, it is much weaker than the natural assumption that \diamond is invariant on all H -symmetric sets.) Thus in Theorem 6.16, we prove that if $\mathcal{E} = \mathcal{C}^n$ or \mathcal{L}^n and $\diamond : \mathcal{E} \rightarrow \mathcal{L}^n$ is monotonic, measure preserving, perimeter preserving on convex bodies, and invariant on H -symmetric unions of two disjoint balls, then \diamond essentially equals Id , \dagger , \diamond_{P_H} , or $\diamond_{P_H}^\dagger$.

Since the maps $\diamond : \mathcal{E} \subset \mathcal{L}^n \rightarrow \mathcal{L}^n$ we study in Section 6 include but are not necessarily symmetrizations, this part of our paper may be viewed as a widening of the scope of [7].

Armed with Theorem 6.16, we prove in Theorem 6.21 that if $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$ or $T : \mathcal{V}(\mathbb{R}^n) \rightarrow \mathcal{V}(\mathbb{R}^n)$ is a rearrangement, and the induced map \diamond_T is perimeter preserving on convex bodies and invariant on H -symmetric unions of two disjoint balls, then T essentially equals Id , \dagger , P_H , or P_H^\dagger . For maps $T : X \rightarrow X$, where $X = \mathcal{M}(\mathbb{R}^n)$ or $\mathcal{M}_+(\mathbb{R}^n)$, the same properties allow the same conclusion for the restriction of T to $\mathcal{S}(\mathbb{R}^n)$ or $\mathcal{V}(\mathbb{R}^n)$, respectively, though not for the unrestricted map. This is shown in Theorem 6.22. Both Theorems 6.21 and 6.22 depend on the main results from Section 4.

As was mentioned earlier, polarization is L^p -contracting and reduces the modulus of continuity, but since compositions of polarizations with respect to different oriented subspaces retain these two properties, they do not seem so useful in classifying polarization among rearrangements.

Throughout the paper we provide examples showing that our main results are best possible in the sense that none of the assumed properties can be omitted or significantly weakened.

We are grateful to David Preiss for communicating the construction in Remark 6.12.

2. PRELIMINARIES

As usual, S^{n-1} denotes the unit sphere and o the origin in Euclidean n -space \mathbb{R}^n . Unless stated otherwise, we assume throughout that $n \geq 2$. The standard orthonormal basis for \mathbb{R}^n is $\{e_1, \dots, e_n\}$ and the Euclidean norm is denoted by $\|\cdot\|$. The term *ball* in \mathbb{R}^n will always mean a *closed* n -dimensional ball unless otherwise stated. The unit ball in \mathbb{R}^n will be denoted by B^n and $B(x, r)$ is the ball with center x and radius r . If $x, y \in \mathbb{R}^n$ we write $x \cdot y$ for the inner product and $[x, y]$ for the line segment with endpoints x and y . If $x \in \mathbb{R}^n \setminus \{o\}$, then x^\perp is the $(n-1)$ -dimensional subspace orthogonal to x . Throughout the paper, the term *subspace* means a linear subspace.

If A is a set, we denote by $\text{lin } A$, $\text{conv } A$, $\text{cl } A$, $\text{int } A$, $\text{relint } A$, and $\text{dim } A$ the *linear hull*, *convex hull*, *closure*, *interior*, *relative interior*, and *dimension* (that is, the dimension of the affine hull) of A , respectively. If H is a subspace of \mathbb{R}^n , then $A|H$ is the (orthogonal) projection of A on H and $x|H$ is the projection of a vector $x \in \mathbb{R}^n$ on H .

If A and B are sets in \mathbb{R}^n and $t \geq 0$, then $tA = \{tx : x \in A\}$ and

$$A + B = \{x + y : x \in A, y \in B\}$$

denotes the *Minkowski sum* of A and B .

When H is a fixed subspace of \mathbb{R}^n , we use A^\dagger for the *reflection* of A in H , i.e., the image of A under the map that takes $x \in \mathbb{R}^n$ to $2(x|H) - x$. If $A^\dagger = A$, we say A is *H -symmetric*. If $H = \{o\}$, we instead write $-A = (-1)A$ for the reflection of A in the origin and *o -symmetric* for $\{o\}$ -symmetric. A set A is called *rotationally symmetric* with respect to H if for $x \in H$, $A \cap (H^\perp + x) = r_x(B^n \cap H^\perp) + x$ for some $r_x \geq 0$. If $\text{dim } H = n - 1$, then a compact convex set is rotationally symmetric with respect to H if and only if it is *H -symmetric*. The term

H -symmetric spherical cylinder will always mean a set of the form

$$(B(x, r) \cap H) + s(B^n \cap H^\perp) = (B(x, r) \cap H) \times s(B^n \cap H^\perp),$$

where $r, s > 0$. Of course, H -symmetric spherical cylinders are rotationally symmetric with respect to both H and H^\perp .

The phrase *translate orthogonal to H* means translate by a vector in H^\perp .

We write \mathcal{H}^k for k -dimensional Hausdorff measure in \mathbb{R}^n , where $k \in \{1, \dots, n\}$. When dealing with relationships between sets in \mathbb{R}^n or functions on \mathbb{R}^n , the term *essentially* means up to a set of \mathcal{H}^n -measure zero.

The Grassmannian of k -dimensional subspaces in \mathbb{R}^n is denoted by $\mathcal{G}(n, k)$.

We denote by \mathcal{C}^n , \mathcal{G}^n , \mathcal{B}^n , \mathcal{M}^n , and \mathcal{L}^n the class of nonempty compact sets, open sets, bounded Borel sets, \mathcal{H}^n -measurable sets, and \mathcal{H}^n -measurable sets of finite \mathcal{H}^n -measure, respectively, in \mathbb{R}^n . Let \mathcal{K}^n be the class of nonempty compact convex subsets of \mathbb{R}^n and let \mathcal{K}_n^n be the class of *convex bodies*, i.e., members of \mathcal{K}^n with interior points. If $K \in \mathcal{K}^n$, then

$$h_K(x) = \sup\{x \cdot y : y \in K\},$$

for $x \in \mathbb{R}^n$, defines the *support function* h_K of K . The texts by Gruber [26] and Schneider [43] contain a wealth of useful information about convex sets and related concepts such as the *intrinsic volumes* V_j , $j \in \{1, \dots, n\}$ (see also [23, Appendix A]). In particular, if $K \in \mathcal{K}^n$ and $\dim K = n$ then $2V_{n-1}(K)$ is the *surface area* of K . If $\dim K = k$, then $V_k(K) = \mathcal{H}^k(K)$ is the *volume* of K . By κ_n we denote the volume $\mathcal{H}^n(B^n)$ of the unit ball in \mathbb{R}^n .

Let $\mathcal{M}(\mathbb{R}^n)$ (or $\mathcal{M}_+(\mathbb{R}^n)$) denote the set of real-valued (or nonnegative, respectively) measurable functions on \mathbb{R}^n and let $\mathcal{S}(\mathbb{R}^n)$ denote the set of functions f in $\mathcal{M}(\mathbb{R}^n)$ such that $\mathcal{H}^n(\{x : f(x) > t\}) < \infty$ for $t > \text{ess inf } f$. By $\mathcal{V}(\mathbb{R}^n)$, we denote the set of functions f in $\mathcal{M}_+(\mathbb{R}^n)$ such that $\mathcal{H}^n(\{x : f(x) > t\}) < \infty$ for $t > 0$. The four classes of functions satisfy $\mathcal{V}(\mathbb{R}^n) \subset \mathcal{S}(\mathbb{R}^n) \subset \mathcal{M}(\mathbb{R}^n)$ and $\mathcal{V}(\mathbb{R}^n) \subset \mathcal{M}_+(\mathbb{R}^n) \subset \mathcal{M}(\mathbb{R}^n)$. Members of $\mathcal{S}(\mathbb{R}^n)$ have been called *symmetrizable* (see, e.g., [8]) and those of $\mathcal{V}(\mathbb{R}^n)$ are often said to *vanish at infinity*. Note that the constant functions are symmetrizable but do not vanish at infinity unless they are essentially zero.

If $T : X \rightarrow X$, we shall usually write Tf instead of $T(f)$. If $T_0, T_1 : X \rightarrow X$ are maps, we say that T_0 is *essentially equal* to T_1 if for $f \in X$, $T_0f(x) = T_1f(x)$ for \mathcal{H}^n -almost all $x \in \mathbb{R}^n$, where the exceptional set may depend on f .

3. PROPERTIES OF MAPS

Let $i \in \{1, \dots, n-1\}$, let $H \in \mathcal{G}(n, i)$ be fixed, and recall that K^\dagger is the reflection of K in H . We consider the following properties of a map $\diamond : \mathcal{E} \subset \mathcal{L}^n \rightarrow \mathcal{L}^n$, where it is assumed (here and throughout the paper) that they hold for all $K, L \in \mathcal{E}$, that the class \mathcal{E} is appropriate for the property concerned, and that sets of \mathcal{H}^n -measure zero are ignored.

1. (*Monotonic or strictly monotonic*) $K \subset L \Rightarrow \diamond K \subset \diamond L$ (or $\diamond K \subset \diamond L$ and $K \neq L \Rightarrow \diamond K \neq \diamond L$, respectively).
2. (*Measure preserving*) $\mathcal{H}^n(\diamond K) = \mathcal{H}^n(K)$.
3. (*Invariant on H -symmetric sets*) $K^\dagger = K \Rightarrow \diamond K = K$.

4. (*Invariant on H -symmetric spherical cylinders*) If $K = (B(x, r) \cap H) + s(B^n \cap H^\perp)$, where $r, s > 0$ and $x \in \mathbb{R}^n$, then $\diamond K = K$.

5. (*Maps balls to balls*) If $K = B(x, r)$, then $\diamond K = B(x', r')$.

6. (*Respects H -cylinders*) If $K \subset (B(x, r) \cap H) + H^\perp$, then $\diamond K \subset (B(x, r) \cap H) + H^\perp$.

Clearly invariance on H -symmetric sets implies invariance on H -symmetric spherical cylinders. If $\mathcal{E} \subset \mathcal{B}^n$ and \diamond is monotonic and invariant on H -symmetric spherical cylinders, then \diamond respects H -cylinders. The assumption $\mathcal{E} \subset \mathcal{B}^n$ cannot be omitted here, as the following example for $n = 2$ shows. Let H be a one-dimensional subspace in \mathbb{R}^2 , and let $R_{H^\perp}E$ denote the reflection of $E \in \mathcal{L}^2$ in H^\perp . Define $\diamond : \mathcal{L}^2 \rightarrow \mathcal{L}^2$ by

$$\diamond E = \begin{cases} E, & \text{if } E \text{ is essentially bounded,} \\ E \cup R_{H^\perp}E, & \text{otherwise.} \end{cases}$$

Then \diamond is invariant on all essentially bounded sets, and in particular on all spherical cylinders. The mapping \diamond is also monotonic, but does not respect H -cylinders. In Lemma 6.13, we show that this conclusion can be drawn if additional conditions are imposed.

We need one further property.

7. (*Perimeter preserving on convex bodies*) For each $K \in \mathcal{K}_n^n$, $\diamond K$ is a set of finite perimeter such that $S(\diamond K) = S(K) = 2V_{n-1}(K)$, where S denotes perimeter (see, for example, [20, p. 170]).

Let $X \subset \mathcal{M}(\mathbb{R}^n)$. We consider the following properties of a map $T : X \rightarrow X$, where the properties are assumed to hold for all $f, g \in X$:

1. (*Equimeasurable*)

$$(2) \quad \mathcal{H}^n(\{x : Tf(x) > t\}) = \mathcal{H}^n(\{x : f(x) > t\})$$

for $t \in \mathbb{R}$.

2. (*Monotonic*) $f \leq g$, essentially, implies $Tf \leq Tg$, essentially.

3. (*L^p -contracting*) $\|Tf - Tg\|_p \leq \|f - g\|_p$ when $f - g \in L^p(\mathbb{R}^n)$.

4. (*Modulus of continuity reducing*) $\omega_d(Tf) \leq \omega_d(f)$ for $d > 0$, where

$$(3) \quad \omega_d(f) = \operatorname{ess\,sup}_{\|x-y\| \leq d} |f(x) - f(y)|$$

is the *modulus of continuity* of $f \in X$.

The map T is called a *rearrangement* if it is equimeasurable and monotonic.

If X contains the characteristic functions of sets in \mathcal{L}^n , $T : X \rightarrow X$, and $A \in \mathcal{L}^n$, let $\diamond_T A = \{x : T1_A(x) = 1\}$. In Lemma 4.5(i) below, it is shown that the induced map $\diamond_T : \mathcal{L}^n \rightarrow \mathcal{L}^n$ is well defined when T is equimeasurable and $X = \mathcal{M}(\mathbb{R}^n)$, $\mathcal{M}_+(\mathbb{R}^n)$, $\mathcal{S}(\mathbb{R}^n)$, or $\mathcal{V}(\mathbb{R}^n)$.

Two further properties of T depend on some $H \in \mathcal{G}(n, n-1)$ which in the second case is oriented and bounds closed half-spaces H^+ and H^- .

5. (*Invariant on H -symmetric spherical cylinders*) If $K = (B(x, r) \cap H) + s(B^n \cap H^\perp)$, where $r, s > 0$, then $\diamond_T K$ is well defined and $\diamond_T K = K$, essentially.

6. (*Pointwise with respect to H*)

$$(4) \quad Tf(x) = \begin{cases} F^+(f(x), f^\dagger(x)), & \text{if } x \in H^+, \\ F^-(f(x), f^\dagger(x)), & \text{if } x \in H^-, \end{cases}$$

where $f^\dagger(x) = f(x^\dagger)$ is the reflection of f in H and where $F^+, F^- : D^2 \rightarrow D$ coincide on the diagonal $\{(s, s) : s \in D\}$. Here

$$D = \{f(x) : x \in \mathbb{R}^n, f \in X\}.$$

Thus $D = \mathbb{R}$ if $X = \mathcal{M}(\mathbb{R}^n)$ or $\mathcal{S}(\mathbb{R}^n)$ and $D = [0, \infty)$ if $X = \mathcal{M}_+(\mathbb{R}^n)$ or $\mathcal{V}(\mathbb{R}^n)$, and in each case D^2 is the common domain of F^+ and F^- .

The pointwise property is inspired by the pointwise operations defined in [24]. The functions F^+ and F^- are said to be *associated with T* . One can consider special associated functions such as the *p th means* $M_p(s, t) = (|s|^p + |t|^p)^{1/p}$ for $p > 0$, $M_\infty(s, t) = \max\{s, t\}$, and $M_{-\infty}(s, t) = \min\{s, t\}$. For $p < 0$, one can define $M_p(s, t) = (|s|^p + |t|^p)^{1/p}$, if $st \neq 0$, and $M_p(s, t) = 0$, otherwise. Then polarization (1) corresponds to taking $F^+ = M_\infty$ and $F^- = M_{-\infty}$ in (4).

Again taking a cue from [24], one might consider the following more general version of the pointwise property:

$$(5) \quad Tf(x) = \begin{cases} (f *^+ f^\dagger)(x), & \text{if } x \in H^+, \\ (f *^- f^\dagger)(x), & \text{if } x \in H^-, \end{cases}$$

where $*^+$ and $*^-$ are two operations between functions on \mathbb{R}^n . To assure that (5) is well defined, one would require that $f *^+ f^\dagger = f *^- f^\dagger$ on H . Then (1) corresponds to taking $f *^+ g = \max\{f, g\}$ and $f *^- g = \min\{f, g\}$. However, the apparent restriction in the definition (5) is an illusion, since $f *^+ f^\dagger$ and $f *^- f^\dagger$ may as well be replaced by arbitrary functions of f .

Polarization, defined by (1), has all the properties 1–6. For properties 1 and 4, see [3, Propositions 1.35 and 1.37], respectively, noting that the general assumption in [3, p. 1] that $f \in \mathcal{S}(\mathbb{R}^n)$ is not necessary. Properties 2 and 5 are clear and 6 was discussed above. Property 3 seems only to have been stated with unnecessary extra assumptions, so we provide a proof in Theorem 5.8 below.

4. EQUIMEASURABLE MAPS AND REARRANGEMENTS

If $f \in \mathcal{V}(\mathbb{R}^n)$, then $\text{ess inf } f = 0$ and it follows that if $T : \mathcal{V}(\mathbb{R}^n) \rightarrow \mathcal{V}(\mathbb{R}^n)$, then $\text{ess inf } Tf = \text{ess inf } f$. We now examine the situation for the other classes of functions.

Lemma 4.1. (i) *If $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$ is equimeasurable, then $\text{ess inf } Tf = \text{ess inf } f$ for $f \in \mathcal{S}(\mathbb{R}^n)$.*

(ii) *If $T : \mathcal{M}(\mathbb{R}^n) \rightarrow \mathcal{M}(\mathbb{R}^n)$ is a rearrangement, then $\text{ess inf } Tf \geq \text{ess inf } f$ for $f \in \mathcal{M}(\mathbb{R}^n)$. Hence, $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$.*

(iii) *In either case, $T : \mathcal{V}(\mathbb{R}^n) \rightarrow \mathcal{V}(\mathbb{R}^n)$ and T is essentially the identity on constant functions.*

Proof. (i) Suppose that $\text{ess inf } Tf \neq \text{ess inf } f$ for some $f \in \mathcal{S}(\mathbb{R}^n)$. Choose $t \in \mathbb{R}$ strictly between $\text{ess inf } Tf$ and $\text{ess inf } f$ and note that (2) is violated, since one side is finite and the other infinite.

(ii) Throughout the proof, we shall ignore sets of \mathcal{H}^n -measure zero. Let $f \in \mathcal{M}(\mathbb{R}^n)$. Suppose that $\text{ess inf } f = a > \text{ess inf } Tf$. Then there is a $t > 0$ such that

$$E = \{x : Tf(x) \leq a - t\}$$

has positive \mathcal{H}^n -measure.

Let $f_0(x) = a - \|x\|$ for $x \in \mathbb{R}^n$. Then $f_0 \in \mathcal{M}(\mathbb{R}^n)$ and $f_0 \leq f$, so the monotonicity of T implies that $Tf_0 \leq Tf$. Consequently, we may choose $t_0 > 0$ such that

$$(6) \quad \mathcal{H}^n(\{x : Tf_0(x) > a - t_0\} \cap E) > 0.$$

Note that $t_0 \geq t$ by the definition of E . Define

$$g(x) = \begin{cases} \max\{f_0(x), a - t/2\}, & \text{if } x \in t_0 B^n, \\ f_0(x), & \text{if } x \notin t_0 B^n. \end{cases}$$

Clearly $g \in \mathcal{M}(\mathbb{R}^n)$, $f_0 \leq g \leq f$, and

$$(7) \quad \{x : f_0(x) > a - t_0\} = \{x : g(x) > a - t_0\} = t_0 B^n.$$

We have

$$(8) \quad \{x : Tf_0(x) > a - t_0\} = \{x : Tg(x) > a - t_0\},$$

because the monotonicity of T implies that the set on the right contains the set on the left, and the two sets have the same \mathcal{H}^n -measure, by (7) and the equimeasurability of T .

The monotonicity of T and $g \leq f$ imply that $Tg \leq Tf$. In particular, $Tg(x) \leq a - t$ when $x \in E$, so

$$(9) \quad \{x : Tg(x) > a - 3t/4\} \cap E = \emptyset.$$

Since $t_0 \geq t$, we have $\{x : Tg(x) > a - 3t/4\} \subset \{x : Tg(x) > a - t_0\}$, so (8) yields

$$(10) \quad \{x : Tg(x) > a - 3t/4\} \subset \{x : Tf_0(x) > a - t_0\}.$$

Moreover,

$$(11) \quad \begin{aligned} \mathcal{H}^n(\{x : Tg(x) > a - 3t/4\}) &= \mathcal{H}^n(\{x : g(x) > a - 3t/4\}) \\ &= \mathcal{H}^n(t_0 B^n) = \mathcal{H}^n(\{x : f_0(x) > a - t_0\}) \\ &= \mathcal{H}^n(\{x : Tf_0(x) > a - t_0\}). \end{aligned}$$

Formulas (10) and (11) imply that

$$\{x : Tg(x) > a - 3t/4\} = \{x : Tf_0(x) > a - t_0\}.$$

But this contradicts (6) and (9) and proves that $\text{ess inf } Tf \geq \text{ess inf } f$.

Let $f \in \mathcal{S}(\mathbb{R}^n)$. If $t > \text{ess inf } Tf$, then $t > \text{ess inf } f$, so by (2) and the definition of $\mathcal{S}(\mathbb{R}^n)$, we have

$$\mathcal{H}^n(\{x : Tf(x) > t\}) = \mathcal{H}^n(\{x : f(x) > t\}) < \infty.$$

Therefore $Tf \in \mathcal{S}(\mathbb{R}^n)$, as required.

(iii) If $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$ is equimeasurable and $f \in \mathcal{V}(\mathbb{R}^n)$, then $\text{ess inf } Tf = \text{ess inf } f = 0$ by (i), so $Tf \in \mathcal{V}(\mathbb{R}^n)$. If $T : \mathcal{M}(\mathbb{R}^n) \rightarrow \mathcal{M}(\mathbb{R}^n)$ is a rearrangement and $f \in \mathcal{V}(\mathbb{R}^n)$, then $\text{ess inf } Tf \geq \text{ess inf } f = 0$ by (ii). This and (2) imply that $Tf \in \mathcal{V}(\mathbb{R}^n)$. That T is essentially the identity on constant functions is immediate in case (i). In case (ii), if $f \equiv c$ is constant, we obtain $Tc \geq c$. This and the equimeasurability of T yield $Tc = c$. \square

Example 4.2. Lemma 4.1(ii) and (iii) do not hold in general if $T : \mathcal{M}(\mathbb{R}^n) \rightarrow \mathcal{M}(\mathbb{R}^n)$ is only assumed to be equimeasurable. To see this, let

$$(12) \quad g_1(x) = \begin{cases} -1/\|x\|, & x \neq o, \\ 0, & x = o. \end{cases}$$

If $A \in \mathcal{L}^n$, define

$$T_0 1_A(x) = \begin{cases} 1, & \text{if } x \in A, \\ g_1(x), & \text{if } x \notin A. \end{cases}$$

Then (2) holds with $f = 1_A$ and we may extend T_0 arbitrarily to an equimeasurable map from $\mathcal{M}(\mathbb{R}^n)$ to itself. Since $T_0 1_A \notin \mathcal{S}(\mathbb{R}^n)$, it is not even true that $T_0 : \mathcal{V}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$. For the remaining statements in Lemma 4.1(ii) and (iii), define $T_1 0 = g_1$ and extend the definition of T_1 arbitrarily to an equimeasurable map from $\mathcal{M}(\mathbb{R}^n)$ to itself. Then $\text{ess inf } T_1 0 = -\infty < 0 = \text{ess inf } 0$ and $T_1 0$ is not a constant function. \square

If $T : \mathcal{M}_+(\mathbb{R}^n) \rightarrow \mathcal{M}_+(\mathbb{R}^n)$ is equimeasurable, then $T : \mathcal{V}(\mathbb{R}^n) \rightarrow \mathcal{V}(\mathbb{R}^n)$ follows immediately from (2). However, the following example shows that the other statements in Lemma 4.1(ii) and (iii) do not hold in general if $T : \mathcal{M}_+(\mathbb{R}^n) \rightarrow \mathcal{M}_+(\mathbb{R}^n)$ is a rearrangement.

Example 4.3. There is a rearrangement $T : \mathcal{M}_+(\mathbb{R}^n) \rightarrow \mathcal{M}_+(\mathbb{R}^n)$ such that $\text{ess inf } Tf < \text{ess inf } f$ for some $f \in \mathcal{M}_+(\mathbb{R}^n)$. Indeed, define T by letting

$$Tf(x) = \begin{cases} f(x), & \text{if } x_1 < 0, \\ 0, & \text{if } 0 \leq x_1 \leq 1, \\ f(x - e_1), & \text{if } x_1 > 1, \end{cases}$$

for $f \in \mathcal{M}_+(\mathbb{R}^n)$ and $x = (x_1, \dots, x_n) \in \mathbb{R}^n$. It is easy to check that T is a rearrangement. If $f \equiv c$ is a constant function and $c > 0$, then $\text{ess inf } Tf = 0 < c = \text{ess inf } f$. It is also not true that T essentially maps constant functions to constant functions. \square

Example 4.4. If $X = \mathcal{M}(\mathbb{R}^n)$ or $\mathcal{M}_+(\mathbb{R}^n)$, there are rearrangements $T : X \rightarrow X$ such that $\text{ess inf } Tf > \text{ess inf } f$ for some $f \in X$. To see this, call $f \in X$ of *type I* if $\mathcal{H}^n(\{x : f(x) > t\}) = \infty$ for $t \geq \text{ess inf } f$ and of *type II* otherwise, i.e., if there is a $t_0 \geq \text{ess inf } f$ such that $\mathcal{H}^n(\{x : f(x) > t\}) < \infty$ for $t > t_0$. Then define $Tf = f + 1$ if f is of type I and $Tf = f$ if f is of type II. Clearly, $T : X \rightarrow X$ is equimeasurable. If $f \leq g$, then either f and g are of the same type, or f is of type II and g is of type I. It follows that $Tf \leq Tg$ and hence that T is a rearrangement. If $f_0(x) = \|x\|$ for $x \in \mathbb{R}^n$, then f_0 is of type I, so $\text{ess inf } Tf_0 = \text{ess inf}(f_0 + 1) = 1 > 0 = \text{ess inf } f_0$. \square

Lemma 4.5. *Let $X = \mathcal{M}(\mathbb{R}^n)$, $\mathcal{M}_+(\mathbb{R}^n)$, $\mathcal{S}(\mathbb{R}^n)$, or $\mathcal{V}(\mathbb{R}^n)$, and let $T : X \rightarrow X$ be equimeasurable.*

(i) *The induced map $\diamond_T : \mathcal{L}^n \rightarrow \mathcal{L}^n$ given by*

$$(13) \quad \diamond_T A = \{x : T1_A(x) = 1\}$$

for $A \in \mathcal{L}^n$ is well defined and measure preserving.

(ii) *If $X = \mathcal{M}_+(\mathbb{R}^n)$, $\mathcal{S}(\mathbb{R}^n)$, or $\mathcal{V}(\mathbb{R}^n)$, then T essentially maps characteristic functions of sets in \mathcal{L}^n to characteristic functions of sets in \mathcal{L}^n , in the sense that for each $A \in \mathcal{L}^n$,*

$$(14) \quad T1_A = 1_{\diamond_T A},$$

essentially.

Proof. (i) If $\alpha > 0$ and $A \in \mathcal{L}^n$, the equimeasurability of T yields

$$(15) \quad \mathcal{H}^n(\{x : T(\alpha 1_A)(x) > t\}) = \mathcal{H}^n(\{x : (\alpha 1_A)(x) > t\}) = \begin{cases} 0, & t \geq \alpha, \\ \mathcal{H}^n(A), & 0 \leq t < \alpha, \\ \infty, & t < 0. \end{cases}$$

Hence $T(\alpha 1_A)(x) \in (-\infty, 0] \cup \{\alpha\}$ for \mathcal{H}^n -almost all $x \in \mathbb{R}^n$, and the measurable set

$$(16) \quad A_\alpha = \{x : T(\alpha 1_A)(x) = \alpha\}$$

satisfies $\mathcal{H}^n(A_\alpha) = \mathcal{H}^n(A) < \infty$. This shows that $A_\alpha \in \mathcal{L}^n$. Applying this with $\alpha = 1$ and setting $\diamond_T A = A_1$, we obtain (13) and the measure-preserving property of \diamond_T .

(ii) It follows from (13) that

$$T1_A(x) = 1 \Leftrightarrow x \in \diamond_T A \Leftrightarrow 1_{\diamond_T A}(x) = 1.$$

It now suffices to observe that if $X = \mathcal{M}_+(\mathbb{R}^n)$, $\mathcal{S}(\mathbb{R}^n)$, or $\mathcal{V}(\mathbb{R}^n)$, then $T1_A(x) \in \{0, 1\}$ for \mathcal{H}^n -almost all $x \in \mathbb{R}^n$, where for $X = \mathcal{S}(\mathbb{R}^n)$ we used Lemma 4.1(i). \square

Example 4.6. If in Lemma 4.5(i) we have $X = \mathcal{M}(\mathbb{R}^n)$ and extend the definition (13) to a map $\diamond_T : \mathcal{M}^n \rightarrow \mathcal{M}^n$, then it need not be measure preserving. To see this, let $g_2 : \mathbb{R}^n \rightarrow \mathbb{R}$ be defined by

$$(17) \quad g_2(x) = \begin{cases} \frac{\|x\|-1}{\|x\|}, & x \neq o, \\ 0, & x = o, \end{cases}$$

let $A \in \mathcal{M}^n$ be such that $\mathcal{H}^n(A) = \infty$, and let

$$T1_A(x) = \begin{cases} g_2(x), & \text{if } x \in A, \\ 0, & \text{if } x \notin A. \end{cases}$$

Note that

$$\mathcal{H}^n(\{x : T1_A(x) > t\}) = \mathcal{H}^n(\{x : 1_A(x) > t\}) = \begin{cases} 0, & t \geq 1, \\ \infty, & t < 1, \end{cases}$$

so (15) holds with $\alpha = 1$. Extend the definition of T to an equimeasurable map from $\mathcal{M}(\mathbb{R}^n)$ to $\mathcal{M}(\mathbb{R}^n)$ arbitrarily. Since

$$\diamond_T A = \{x : T1_A(x) = 1\} = \{x \in A : g_2(x) = 1\} = \emptyset,$$

\diamond_T is not measure preserving. □

The equimeasurable map T_0 from Example 4.2 shows that Lemma 4.5(ii) does not hold when $X = \mathcal{M}(\mathbb{R}^n)$.

Lemma 4.7. *Let $X = \mathcal{S}(\mathbb{R}^n)$ or $\mathcal{V}(\mathbb{R}^n)$ and let $T : X \rightarrow X$ be a rearrangement. For $X = \mathcal{S}(\mathbb{R}^n)$, $A \in \mathcal{L}^n$, and $\alpha, \beta \in \mathbb{R}$ with $\alpha \geq 0$, we have*

$$(18) \quad T(\alpha 1_A + \beta) = \alpha T1_A + \beta,$$

essentially. When $X = \mathcal{V}(\mathbb{R}^n)$, (18) holds, essentially, if $\beta = 0$.

Proof. The case when $\alpha = 0$ follows from Lemma 4.1(iii), so henceforth we assume that $\alpha > 0$.

We shall ignore sets of \mathcal{H}^n -measure zero for the remainder of the proof. We first assume that $\beta = 0$. If $0 < \alpha' \leq \alpha$, then $\alpha' 1_A \leq \alpha 1_A$, so $T(\alpha' 1_A) \leq T(\alpha 1_A)$. Now $T(\alpha' 1_A)(x) = \alpha'$ if and only if $x \in A_{\alpha'}$, where $A_{\alpha'}$ is defined by (16) with α replaced by α' , so $T(\alpha 1_A)(x) \geq \alpha'$ for $x \in A_{\alpha'}$. From the proof of Lemma 4.5(i), we see that $T(\alpha 1_A)(x) = \alpha$ for $x \in A_{\alpha'}$ and hence $A_{\alpha'} \subset A_{\alpha}$. By (15), we have $\mathcal{H}^n(A_{\alpha'}) = \mathcal{H}^n(A) = \mathcal{H}^n(A_{\alpha})$, so $A_{\alpha'} = A_{\alpha}$. Consequently, for each $\alpha > 0$ we have $A_{\alpha} = A_1$. Now

$$T(\alpha 1_A)(x) = \alpha \Leftrightarrow x \in A_{\alpha} \Leftrightarrow x \in A_1 \quad \text{and} \quad \alpha 1_{\diamond_T A}(x) = \alpha \Leftrightarrow x \in A_1,$$

so, using (14), we obtain

$$(19) \quad T(\alpha 1_A)(x) = \alpha 1_{\diamond_T A}(x) = \alpha T1_A,$$

as required. This proves (18) when $\beta = 0$ and the second statement in the lemma.

Suppose that $\beta \neq 0$ and for convenience let $h = \alpha 1_A + \beta$. Then $h \in \{\beta, \alpha + \beta\}$. Arguing as in the proof of Lemma 4.5(i), we see that $Th \in (-\infty, \beta] \cup \{\alpha + \beta\}$, and hence, by Lemma 4.1(i), $Th \in \{\beta, \alpha + \beta\}$. It follows that $Th = \alpha 1_B + \beta$ for some \mathcal{H}^n -measurable set B . For $t \in (\beta, \alpha + \beta)$,

$$\mathcal{H}^n(B) = \mathcal{H}^n\{x : Th(x) > t\} = \mathcal{H}^n\{x : h(x) > t\} = \mathcal{H}^n(A).$$

In view of (14), it will suffice to show that $B = \diamond_T A$.

Assume that $\beta > 0$. Then $h \geq (\alpha + \beta) 1_A$, so using the monotonicity of T , and (19) with α replaced by $\alpha + \beta$, we get

$$Th = \alpha 1_B + \beta \geq T((\alpha + \beta) 1_A) = (\alpha + \beta) T1_A = (\alpha + \beta) 1_{\diamond_T A}.$$

Since $\alpha 1_B(x) + \beta = \alpha + \beta \Leftrightarrow x \in B$ and $(\alpha + \beta) 1_{\diamond_T A}(x) = \alpha + \beta \Leftrightarrow x \in \diamond_T A$, we must have $B = \diamond_T A$.

Finally, suppose that $\beta < 0$. Let $\gamma = \max\{\alpha + \beta, 1\}$. Then

$$h = \alpha 1_A + \beta \leq (\gamma - \beta) 1_A + \beta \leq \gamma 1_A,$$

so arguing as above, we find that

$$(20) \quad \alpha 1_B + \beta = Th \leq T((\gamma - \beta) 1_A + \beta) \leq (\gamma - \beta) 1_C + \beta \leq \gamma T1_A = \gamma 1_{\diamond_T A},$$

for some C with $\mathcal{H}^n(C) = \mathcal{H}^n(B) = \mathcal{H}^n(A)$. Since $(\gamma - \beta) 1_C(x) + \beta = \gamma \Leftrightarrow x \in C$ and $\gamma 1_{\diamond_T A}(x) = \gamma \Leftrightarrow x \in \diamond_T A$, the right-hand inequality in (20) yields $C = \diamond_T A$. If $\alpha + \beta \geq 1$, then $\gamma = \alpha + \beta$ and the left-hand inequality in (20) similarly yields $B = C$. If $\alpha + \beta < 1$, then

$\gamma = 1$ and the left-hand inequality in (20) becomes $\alpha 1_B \leq (1 - \beta)1_C$. Now $\alpha 1_B(x) = \alpha > 0 \Leftrightarrow x \in B$ and $(1 - \beta)1_C(x) = 0 \Leftrightarrow x \notin C$, so $\mathcal{H}^n(B \setminus C) = 0$. From this and $\mathcal{H}^n(B) = \mathcal{H}^n(C)$, we conclude that $B = C$. Therefore $B = \diamond_T A$, as required. \square

Theorem 4.8. *Let $X = \mathcal{M}(\mathbb{R}^n)$, $\mathcal{M}_+(\mathbb{R}^n)$, $\mathcal{S}(\mathbb{R}^n)$, or $\mathcal{V}(\mathbb{R}^n)$ and let $T : X \rightarrow X$ be a rearrangement.*

(i) *The map $\diamond_T : \mathcal{L}^n \rightarrow \mathcal{L}^n$ defined by (13) is monotonic.*

(ii) *If $X = \mathcal{S}(\mathbb{R}^n)$ or $\mathcal{V}(\mathbb{R}^n)$ and $f \in X$, then*

$$(21) \quad \{x : Tf(x) \geq t\} = \diamond_T \{x : f(x) \geq t\},$$

essentially, for $t > \text{ess inf } f$. Moreover, T is essentially determined by \diamond_T , since

$$(22) \quad Tf(x) = \max \{ \sup \{ t \in \mathbb{Q}, t > \text{ess inf } f : x \in \diamond_T \{z : f(z) \geq t\} \}, \text{ess inf } f \},$$

essentially.

Proof. (i) If $A \subset B$, then $1_A \leq 1_B$ and hence $1 = T1_A(x) \leq T1_B(x)$ for \mathcal{H}^n -almost all $x \in \diamond_T A = \{x : T1_A(x) = 1\}$. Since $T1_B(x) \in (-\infty, 0] \cup \{1\}$ for \mathcal{H}^n -almost all $x \in \mathbb{R}^n$, it is clear that $\diamond_T A \subset \{x : T1_B(x) = 1\} = \diamond_T B$, essentially. Therefore \diamond_T is monotonic.

(ii) Since $\mathcal{V}(\mathbb{R}^n) \subset \mathcal{S}(\mathbb{R}^n)$, it suffices to consider the case when $X = \mathcal{S}(\mathbb{R}^n)$. Let $f \in \mathcal{S}(\mathbb{R}^n)$ and let $t > \text{ess inf } f$. If (t_m) is an increasing sequence with $\text{ess inf } f < t_m < t$ converging to t , the fact that $\mathcal{H}^n(\{x : f(x) > t_1\}) < \infty$ implies that

$$\mathcal{H}^n(\{x : f(x) \geq t\}) = \mathcal{H}^n(\cap_{m=1}^{\infty} \{x : f(x) > t_m\}) = \lim_{m \rightarrow \infty} \mathcal{H}^n(\{x : f(x) > t_m\}).$$

The same statement holds when f is replaced by Tf . The equimeasurability of T then yields

$$(23) \quad \mathcal{H}^n(\{x : f(x) \geq t\}) = \mathcal{H}^n(\{x : Tf(x) \geq t\}).$$

Assume that $\text{ess inf } f > -\infty$. Let $C = \{x : f(x) \geq t\}$. It is easy to check that

$$f \geq (t - \text{ess inf } f)1_C + \text{ess inf } f.$$

By Lemma 4.7 with $\alpha = t - \text{ess inf } f$ and $\beta = \text{ess inf } f$, and the monotonicity of T , we obtain

$$Tf \geq T((t - \text{ess inf } f)1_C + \text{ess inf } f) = (t - \text{ess inf } f)T1_C + \text{ess inf } f,$$

essentially. This inequality and (13) give

$$\diamond_T \{x : f(x) \geq t\} = \diamond_T C = \{x : T1_C(x) = 1\} \subset \{x : Tf(x) \geq t\},$$

essentially. The left- and right-hand sides are of equal \mathcal{H}^n -measure by (23) and the measure-preserving property of \diamond_T , and are therefore essentially equal. Hence (21) holds when $\text{ess inf } f > -\infty$.

Now suppose that $\text{ess inf } f = -\infty$. Let $s < t$ and define $f_s(x) = \max\{f(x), s\}$ for $x \in \mathbb{R}^n$. Then $\text{ess inf } f_s = s$ and

$$(24) \quad \{x : f(x) \geq t\} = \{x : f_s(x) \geq t\}.$$

Since $f \leq f_s$, the monotonicity of T implies that $\{x : Tf_s(x) \geq t\}$ is essentially contained in $\{x : Tf(x) \geq t\}$. This, (24), and the equimeasurability of T yield $\{x : Tf(x) \geq t\} = \{x : Tf_s(x) \geq t\}$, essentially. By (21) with f replaced by f_s , and (24) again, we obtain

$$\{x : Tf(x) \geq t\} = \{x : Tf_s(x) \geq t\} = \diamond_T \{x : f_s(x) \geq t\} = \diamond_T \{x : f(x) \geq t\},$$

proving that (21) holds generally.

Consequently, for each $t > \text{ess inf } f$, the symmetric difference

$$(25) \quad N_t = \{x : Tf(x) \geq t\} \triangle \diamond_T \{x : f(x) \geq t\}$$

satisfies $\mathcal{H}^n(N_t) = 0$. According to Lemma 4.1(i), there is a set N such that $\mathcal{H}^n(N) = 0$ and $Tf(x) \geq \text{ess inf } f$ when $x \notin N$. If $g : \mathbb{R}^n \rightarrow \mathbb{R}$, then $g(x) = \sup\{t \in \mathbb{Q} : g(x) \geq t\}$ for $x \in \mathbb{R}^n$. Using this with $g = Tf$ and taking (25) into account, we obtain (22) for $x \in \mathbb{R}^n \setminus (\cup\{N_t : t \in \mathbb{Q}, t > \text{ess inf } f\} \cup N)$ and hence for \mathcal{H}^n -almost all $x \in \mathbb{R}^n$. \square

Under the assumptions in Lemma 4.8(ii), it is also true that

$$(26) \quad \{x : Tf(x) > t\} = \diamond_T \{x : f(x) > t\},$$

essentially, for $t > \text{ess inf } f$. Indeed, using (21), we have

$$\begin{aligned} \{x : Tf(x) > t\} &= \cup_{n \in \mathbb{N}} \{x : Tf(x) \geq t + 1/n\} = \cup_{n \in \mathbb{N}} \diamond_T \{x : f(x) \geq t + 1/n\} \\ &= \diamond_T \cup_{n \in \mathbb{N}} \{x : f(x) \geq t + 1/n\} = \diamond_T \{x : f(x) > t\}, \end{aligned}$$

essentially, for $t > \text{ess inf } f$, where the third equality follows easily from the fact that \diamond_T is measure preserving and monotonic. It follows that

$$(27) \quad Tf(x) = \max \{ \sup\{t \in \mathbb{Q}, t > \text{ess inf } f : x \in \diamond_T \{z : f(z) > t\}\}, \text{ess inf } f \},$$

essentially, an alternative formula to (22).

We stress that the exceptional set in Theorem 4.8(ii) cannot be avoided and may depend on f . For example, define $T : \mathcal{V}(\mathbb{R}^n) \rightarrow \mathcal{V}(\mathbb{R}^n)$ by

$$Tf(x) = \begin{cases} 0, & \text{if } f(x) < 1 \text{ and } x \in \mathbb{Q}^n, \\ f(x), & \text{otherwise.} \end{cases}$$

Then T is a rearrangement, but it does not coincide with the identity, although $\diamond_T = \diamond_{\text{id}}$. Note also that the supremum over \mathbb{Q} in (22) cannot be replaced by the supremum over \mathbb{R} , and it is consistent with ZFC that it cannot be replaced by the essential supremum over \mathbb{R} ; see the remarks after Example 7.2.

With obvious modifications, the following result also applies to rearrangements $T : \mathcal{V}(\mathbb{R}^n) \rightarrow \mathcal{V}(\mathbb{R}^n)$. The equality (28) also appears in [51, Proposition 3(d)], where the notation and framework is substantially different, as we explain in the Appendix. Note that [51, Proposition 3(d)] assumes (in our notation) that φ is left-continuous and increasing, but this is not valid in our context, as we show in Example 4.10.

Theorem 4.9. *Let $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$ be a rearrangement and let $f \in \mathcal{S}(\mathbb{R}^n)$. If $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is right-continuous and increasing (i.e., non-decreasing), then $\varphi \circ f \in \mathcal{S}(\mathbb{R}^n)$ and*

$$(28) \quad \varphi(Tf) = T(\varphi \circ f),$$

essentially. It follows that $T(\alpha f + \beta) = \alpha Tf + \beta$, essentially, for $\alpha, \beta \in \mathbb{R}$ with $\alpha \geq 0$.

Proof. We do not require the right-continuity of φ everywhere, but only at $\text{ess inf } f$ when $\text{ess inf } f > -\infty$.

We first claim that for any $g \in \mathcal{S}(\mathbb{R}^n)$ such that φ is right-continuous at $\text{ess inf } g$ when $\text{ess inf } g > -\infty$, we have

$$(29) \quad \text{ess inf } \varphi \circ g = \begin{cases} \varphi(\text{ess inf } g), & \text{if } \text{ess inf } g > -\infty, \\ \inf \varphi, & \text{if } \text{ess inf } g = -\infty. \end{cases}$$

Indeed, if $\text{ess inf } g = -\infty$, then $\text{ess inf } \varphi \circ g \geq \inf \varphi$ is obvious, while

$$(30) \quad \mathcal{H}^n(\{x : \varphi(g(x)) \leq \varphi(t)\}) \geq \mathcal{H}^n(\{x : g(x) < t\}) > 0$$

for $t \in \mathbb{R}$ and hence $\inf \varphi = \lim_{t \rightarrow -\infty} \varphi(t) \geq \text{ess inf } \varphi \circ g$. If $\text{ess inf } g > -\infty$, we have $g \geq \text{ess inf } g$, essentially, so $\varphi \circ g \geq \varphi(\text{ess inf } g)$, essentially, and hence $\text{ess inf } \varphi \circ g \geq \varphi(\text{ess inf } g)$. On the other hand, (30) holds for $t > \text{ess inf } g$, so for such t , $\varphi(t) \geq \text{ess inf } \varphi \circ g$. Then $\varphi(\text{ess inf } g) \geq \text{ess inf } \varphi \circ g$ follows from the right-continuity of φ at $\text{ess inf } g$. This proves (29).

Let $f \in \mathcal{S}(\mathbb{R}^n)$. For $t \in \mathbb{R}$, let $s_t = \inf\{s : \varphi(s) \geq t\}$. Since φ is increasing, we have

$$(31) \quad \{s : \varphi(s) \geq t\} = \begin{cases} [s_t, \infty), & \text{if } s_t \in \mathbb{R} \text{ and } \varphi(s_t) \geq t, \\ (s_t, \infty), & \text{otherwise.} \end{cases}$$

As φ is also right-continuous at $\text{ess inf } f$ when $\text{ess inf } f > -\infty$, then under the latter assumption,

$$(32) \quad t > \varphi(\text{ess inf } f) \Rightarrow s_t > \text{ess inf } f.$$

Suppose that $\text{ess inf } f = -\infty$. If $t > \inf \varphi$, let $t > t' > \inf \varphi$. Then $s_{t'} > -\infty$, so (31) with t replaced by t' implies that

$$(33) \quad \mathcal{H}^n(\{x : \varphi(f(x)) > t\}) \leq \mathcal{H}^n(\{x : \varphi(f(x)) \geq t'\}) \leq \mathcal{H}^n(\{x : f(x) \geq s_{t'}\}) < \infty,$$

since this holds trivially when $s_{t'} = \infty$ and in view of $f \in \mathcal{S}(\mathbb{R}^n)$ otherwise. Now suppose that $\text{ess inf } f > -\infty$. If $t > \varphi(\text{ess inf } f)$, let $t > t' > \varphi(\text{ess inf } f)$. By (32) with t replaced by t' , we conclude that $s_{t'} > \text{ess inf } f$, so (33) holds again since $f \in \mathcal{S}(\mathbb{R}^n)$. Since $\text{ess inf } \varphi \circ f = \varphi(\text{ess inf } f)$, by (29) with $g = f$, this proves that $\varphi \circ f \in \mathcal{S}(\mathbb{R}^n)$.

We claim that $\text{ess inf } \varphi(Tf) = \text{ess inf } T(\varphi \circ f)$. To this end, note that for $g \in \mathcal{S}(\mathbb{R}^n)$, we have $\text{ess inf } Tg = \text{ess inf } g$, by Lemma 4.1(i). We apply this with $g = f$ and $g = \varphi \circ f$ and (29) with $g = Tf$ and $g = f$. If $\text{ess inf } f > -\infty$, we get

$$\text{ess inf } \varphi(Tf) = \varphi(\text{ess inf } Tf) = \varphi(\text{ess inf } f) = \text{ess inf }(\varphi \circ f) = \text{ess inf } T(\varphi \circ f),$$

while if $\text{ess inf } f = -\infty$, then $\text{ess inf } Tf = -\infty$ and we obtain

$$\text{ess inf } \varphi(Tf) = \inf \varphi = \text{ess inf } \varphi \circ f = \text{ess inf } T(\varphi \circ f).$$

This proves the claim.

The next step is to prove that

$$(34) \quad \{x : \varphi(Tf(x)) \geq t\} = \{x : T(\varphi \circ f)(x) \geq t\},$$

essentially, for $t > \text{ess inf } \varphi(Tf) = \text{ess inf } T(\varphi \circ f)$. In fact, the latter inequality and (29) with $g = Tf$ imply that $s_t > \text{ess inf } Tf = \text{ess inf } f$, where (32) was used when $s = \text{ess inf } f > -\infty$. If $\varphi(s_t) \geq t$, we use (21) twice to obtain

$$\begin{aligned} \{x : \varphi(Tf(x)) \geq t\} &= \{x : Tf(x) \geq s_t\} = \diamond_T \{x : f(x) \geq s_t\} \\ &= \diamond_T \{x : \varphi(f(x)) \geq t\} = \{x : T(\varphi \circ f)(x) \geq t\}, \end{aligned}$$

essentially. A similar argument, using (26) instead of (21), yields (34) when $\varphi(s_t) < t$ or $s_t \notin \mathbb{R}$.

The proof of the first statement in the corollary is concluded by noting that if $g, h \in \mathcal{S}(\mathbb{R}^n)$ satisfy $\text{ess inf } g = \text{ess inf } h$ and $\{x : g(x) \geq t\} = \{x : h(x) \geq t\}$, essentially, for all $t > \text{ess inf } g$, then $g = h$, essentially. Indeed, following the proof of [51, Lemma 1], we may otherwise assume that there is an $\varepsilon > 0$ such that $\mathcal{H}^n(\{x : h(x) > g(x) + \varepsilon\}) > 0$. But

$$\{x : h(x) > g(x) + \varepsilon\} \subset \bigcup_{n \in \mathbb{Z}, n\varepsilon \geq \text{ess inf } g} (\{x : h(x) \geq n\varepsilon\} \setminus \{x : g(x) \geq n\varepsilon\}),$$

essentially, and the right-hand side has \mathcal{H}^n -measure zero, a contradiction.

The second statement in the corollary follows immediately from the first on setting $\varphi(t) = \alpha t + \beta$. \square

The proof of the previous theorem, as was mentioned at the beginning of it, actually only requires the right-continuity of φ at $\text{ess inf } f$ when $\text{ess inf } f > -\infty$. The following example shows that this is the weakest possible continuity condition on φ for which the theorem holds.

Example 4.10. If $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is left-continuous and increasing, it is possible that $f \in \mathcal{S}(\mathbb{R}^n)$ but $\varphi \circ f \notin \mathcal{S}(\mathbb{R}^n)$. Indeed, taking $n = 1$ for simplicity, let $\varphi(t) = t$ for $t > 0$ and $\varphi(t) = -1$ for $t \leq 0$. Let $f \in \mathcal{S}(\mathbb{R})$ be any function such that $f(x) > 0$ for $x > 0$ and $f(x) = 0$ for $x \leq 0$. Then $\varphi(f(x)) \geq 0$ for $x > 0$ and $\varphi(f(x)) = -1$ for $x \leq 0$, so $\varphi \circ f \notin \mathcal{S}(\mathbb{R}^n)$. Note that φ is continuous everywhere except at $\text{ess inf } f = 0$, where it is only left-continuous. \square

Equimeasurable maps satisfying (28) for all right-continuous, increasing φ must actually be rearrangements, as we now show. The first part of the proof uses ideas of Van Schaftingen [48, Proposition 2.4.1].

Lemma 4.11. *Let $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$ be equimeasurable. Suppose that $\varphi(Tf) = T(\varphi \circ f)$, essentially, whenever $f \in \mathcal{S}(\mathbb{R}^n)$ and $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is right-continuous and increasing. Then T is monotonic and hence a rearrangement.*

Proof. We shall ignore sets of \mathcal{H}^n -measure zero in this proof. Let $f \in \mathcal{S}(\mathbb{R}^n)$ and for $c \in \mathbb{R}$, define $\varphi_c(t) = 1$ if $t \geq c$ and $\varphi_c(t) = 0$ if $t < c$. Note that φ_c is right-continuous and increasing, and for $g \in \mathcal{S}(\mathbb{R}^n)$, we have $1_{\{x: g(x) \geq c\}} = \varphi_c \circ g \in \mathcal{S}(\mathbb{R}^n)$, by the first part of

the proof of Theorem 4.9 (which does not involve T), with φ and f replaced by φ_c and g , respectively. Using this with $g = Tf$ and $g = f$ and our assumption on T , we obtain

$$1_{\{x: Tf(x) \geq c\}} = \varphi_c(Tf) = T(\varphi_c \circ f) = T1_{\{x: f(x) \geq c\}}.$$

Hence,

$$\{x : Tf(x) \geq c\} = \begin{cases} \diamond_T \{x : f(x) \geq c\}, & \text{if } c > \text{ess inf } f, \\ \mathbb{R}^n, & \text{otherwise,} \end{cases}$$

where (14) was used in the first case and Lemma 4.1(i) in the second case. As $Tf(x) = \sup\{c \in \mathbb{Q} : Tf(x) \geq c\}$, the map T satisfies (22).

Suppose that $A \subset B \subset \mathbb{R}^n$ and let $h = 1_A + 1_B$. It follows easily from (22) that $Th = 1_{(\diamond_TA) \cup \diamond_TB} + 1_{\diamond_TA}$. Since T is equimeasurable, we have $\mathcal{H}^n((\diamond_TA) \cup \diamond_TB) = \mathcal{H}^n(\diamond_TB)$, so $\diamond_TA \subset \diamond_TB$. Thus \diamond_T is monotonic. This implies that if $f, g \in \mathcal{S}(\mathbb{R}^n)$ and $f \leq g$, then $\diamond_T\{z : f(z) \geq t\} \subset \diamond_T\{z : g(z) \geq t\}$, and then $Tf \leq Tg$ is a consequence of (22). \square

Example 4.12. Let $X = \mathcal{M}(\mathbb{R}^n)$ or $\mathcal{M}_+(\mathbb{R}^n)$. There is a rearrangement $T : X \rightarrow X$ such that $T \neq \text{Id}$ but $T = \text{Id}$ on $\mathcal{V}(\mathbb{R}^n)$. In particular, Theorem 4.8(ii) does not hold. Indeed, for $f \in X$, let

$$t_f = \inf\{t \geq 0 : \mathcal{H}^n(\{x : f(x) > t\}) < \infty\}$$

and let $A_f = \{x : f(x) \geq t_f\}$. Define

$$Tf(x) = \begin{cases} f(x), & \text{if } x \in A_f, \\ \min\{f(x) + 1, t_f\}, & \text{if } x \notin A_f. \end{cases}$$

Note that if $f \in \mathcal{V}(\mathbb{R}^n)$, then $t_f = 0$ and $A_f = \mathbb{R}^n$, so $T = \text{Id}$ on $\mathcal{V}(\mathbb{R}^n)$. Let

$$f(x) = \begin{cases} 0, & \text{if } x \in B^n, \\ \frac{\|x\|}{\|x\|-1}, & \text{if } x \notin B^n. \end{cases}$$

Then $f \in \mathcal{M}_+(\mathbb{R}^n)$, $t_f = 1$, and $A_f = \mathbb{R}^n \setminus B^n$, so $Tf = f + 1_{B^n} \neq f$.

We claim that T is a rearrangement. Note first that $Tf \geq f$. Let $f \in X$ and suppose that $t \geq t_f$. If $f(x) > t$, then $x \in A_f$, so $Tf(x) = f(x) > t$. Conversely, if $Tf(x) > t$, then $Tf(x) > t_f$, so $x \in A_f$, implying that $Tf(x) = f(x)$ and thus $f(x) > t$. Hence

$$\{x : Tf(x) > t\} = \{x : f(x) > t\}.$$

Now suppose that $t < t_f$. Then $\mathcal{H}^n(\{x : f(x) > t\}) = \infty$ by the definition of t_f . But if $f(x) > t$, then $Tf(x) \geq f(x) > t$, so $\mathcal{H}^n(\{x : Tf(x) > t\}) = \infty$. This proves that T is equimeasurable.

Let $f, g \in X$ satisfy $f \leq g$. Then $t_f \leq t_g$. If $x \in A_f$, then $Tf(x) = f(x) \leq g(x) \leq Tg(x)$. If $x \in A_g \setminus A_f$, then

$$Tg(x) = g(x) \geq t_g \geq \min\{f(x) + 1, t_g\} \geq \min\{f(x) + 1, t_f\} = Tf(x).$$

Finally, if $x \notin A_f \cup A_g$, then

$$Tg(x) = \min\{g(x) + 1, t_g\} \geq \min\{f(x) + 1, t_f\} = Tf(x).$$

This proves that T is monotonic. \square

5. POINTWISE MAPS BETWEEN FUNCTIONS

Theorem 5.1. *Let $H \in \mathcal{G}(n, n-1)$ be oriented, let $X = \mathcal{M}(\mathbb{R}^n)$, $\mathcal{M}_+(\mathbb{R}^n)$, $\mathcal{S}(\mathbb{R}^n)$, or $\mathcal{V}(\mathbb{R}^n)$, and suppose that $T : X \rightarrow X$ is pointwise with respect to H . Then T is equimeasurable if and only if its associated functions F^+ and F^- satisfy*

$$(35) \quad \{F^+(r, s), F^-(s, r)\} = \{r, s\}$$

for $(r, s) \in D^2$, the common domain of F^+ and F^- .

Proof. We first consider the case when $X = \mathcal{M}(\mathbb{R}^n)$ or $\mathcal{S}(\mathbb{R}^n)$, so that $D = \mathbb{R}$. Assume that T is equimeasurable. We claim that

$$(36) \quad F^+(r, r) = F^-(r, r) = r$$

for $r \in \mathbb{R}$. To see this, let $f \equiv r$ be constant on \mathbb{R}^n . From (4) and the fact that F^+ and F^- coincide on the diagonal of \mathbb{R}^2 , we see that $Tf \equiv F^+(r, r) = F^-(r, r)$ is also constant. This and (2) yield (36) (when $X = \mathcal{S}(\mathbb{R}^n)$, this is also a consequence of Lemma 4.1(iii)).

Now fix $r, s \in \mathbb{R}$. Let $A \subset \text{int } H^+$ be compact with $\mathcal{H}^n(A) > 0$, let $c < \min\{r, s\}$, and let

$$f(x) = r1_A(x) + s1_{A^\dagger}(x) + c1_{\mathbb{R}^n \setminus (A \cup A^\dagger)}(x).$$

Note that $f \in \mathcal{S}(\mathbb{R}^n)$. From (4) and (36), we have

$$(37) \quad Tf(x) = \begin{cases} F^+(r, s), & \text{if } x \in A, \\ F^-(s, r), & \text{if } x \in A^\dagger, \\ c, & \text{otherwise.} \end{cases}$$

If $c < t < \min\{r, s\}$, both sides of (2) equal $2\mathcal{H}^n(A)$, while if $t = \max\{r, s\}$, both sides are zero. Thus

$$(38) \quad \min\{r, s\} \leq F^+(r, s), F^-(s, r) \leq \max\{r, s\}.$$

If $r \neq s$, we can choose t in (2) with $\min\{r, s\} < t < \max\{r, s\}$. Then both sides of (2) equal $\mathcal{H}^n(A)$, so $F^+(r, s) \leq t$ and $F^-(s, r) > t$ or vice versa. As $\min\{r, s\} < t < \max\{r, s\}$ was arbitrary, $F^+(r, s) = \min\{r, s\}$ and $F^-(s, r) = \max\{r, s\}$ or vice versa. This proves (35) when $r \neq s$, and (35) holds trivially when $r = s$ due to (38).

Now assume that (35) holds and let $f \in X$. Define

$$M = \{x \in \mathbb{R}^n : f(x) = f^\dagger(x)\},$$

$$M^\pm = \{x \in H^\pm \setminus M : F^\pm(f(x), f^\dagger(x)) = f(x)\},$$

and

$$M_\dagger^\pm = \{x \in H^\pm \setminus M : F^\pm(f(x), f^\dagger(x)) = f^\dagger(x)\}.$$

By (35), these five sets form a partition of \mathbb{R}^n . Note that if $x \notin M$, then $Tf(x) = f(x)$ if and only if $x \in M^+ \cup M^-$ and $Tf(x) = f^\dagger(x)$ if and only if $x \in M_\dagger^+ \cup M_\dagger^-$. This and the fact that by (35) we have $Tf = f$ on M yield

$$(39) \quad \{x : Tf(x) > t\} = (\{x : f(x) > t\} \cap (M \cup M^+ \cup M^-)) \cup (\{x : f^\dagger(x) > t\} \cap (M_\dagger^+ \cup M_\dagger^-))$$

for $t \in \mathbb{R}$. Using the definitions of M_\dagger^- and M , together with (35), we obtain

$$\begin{aligned} x \in (M_\dagger^-)^\dagger \Leftrightarrow x^\dagger \in M_\dagger^- &\Leftrightarrow x^\dagger \in H^- \setminus M \text{ and } F^-(f(x^\dagger), f^\dagger(x^\dagger)) = f^\dagger(x^\dagger) \\ &\Leftrightarrow x \in H^+ \setminus M \text{ and } F^-(f^\dagger(x), f(x)) = f(x) \\ &\Leftrightarrow x \in H^+ \setminus M \text{ and } F^+(f(x), f^\dagger(x)) = f^\dagger(x) \Leftrightarrow x \in M_\dagger^+. \end{aligned}$$

Consequently, $(M_\dagger^-)^\dagger = M_\dagger^+$ and $(M_\dagger^+)^\dagger = M_\dagger^-$. Therefore

$$(\{x : f^\dagger(x) > t\} \cap (M_\dagger^+ \cup M_\dagger^-))^\dagger = \{x : f(x) > t\} \cap (M_\dagger^- \cup M_\dagger^+).$$

In particular,

$$\mathcal{H}^n(\{x : f^\dagger(x) > t\} \cap (M_\dagger^+ \cup M_\dagger^-)) = \mathcal{H}^n(\{x : f(x) > t\} \cap (M_\dagger^- \cup M_\dagger^+)).$$

It follows from (39) that $\mathcal{H}^n(\{x : Tf(x) > t\}) = \mathcal{H}^n(\{x : f(x) > t\})$, so T is equimeasurable. This completes the proof when $X = \mathcal{M}(\mathbb{R}^n)$ or $\mathcal{S}(\mathbb{R}^n)$.

Now suppose that $X = \mathcal{M}_+(\mathbb{R}^n)$ or $\mathcal{V}(\mathbb{R}^n)$, so that $D = [0, \infty)$. The second part of the above proof can be applied without change. If $X = \mathcal{M}_+(\mathbb{R}^n)$, the first part of the above proof also still applies, but when $X = \mathcal{V}(\mathbb{R}^n)$, we cannot use constant functions other than $f \equiv 0$ and thus can only obtain the weaker version

$$(40) \quad F^+(0, 0) = F^-(0, 0) = 0$$

of (36). Nevertheless, we can follow the argument in the second paragraph when $\min\{r, s\} > 0$ and $c = 0$, and this yields (38) when $r, s > 0$. Setting $r = s > 0$ in (38) and using (40), we retrieve (36) for $r \geq 0$. With (36) in hand, we may assume that $r = 0$ and $s > 0$. Then by using (2) with $t = s$ and with $0 < t \leq s$, one obtains (38) for $r, s \geq 0$ and the conclusion follows easily as before. \square

Corollary 5.2. *Let $H \in \mathcal{G}(n, n-1)$ be oriented, let $X = \mathcal{M}(\mathbb{R}^n)$, $\mathcal{M}_+(\mathbb{R}^n)$, $\mathcal{S}(\mathbb{R}^n)$, or $\mathcal{V}(\mathbb{R}^n)$, and suppose that $T : X \rightarrow X$ is pointwise with respect to H . If T is equimeasurable, then it maps characteristic functions of sets in \mathcal{L}^n to characteristic functions of sets in \mathcal{L}^n and $\diamond_T = \text{Id}$, $\diamond_T = \dagger$, $\diamond_T = \diamond_{P_H}$, or $\diamond_T = \diamond_{P_H}^\dagger$, where Id is the identity map and \dagger is reflection in H .*

Proof. By Theorem 5.1, (35) holds, implying that $F^\pm(0, 0) = 0$, $F^\pm(1, 1) = 1$,

$$\text{either } F^+(1, 0) = 1 \text{ and } F^-(0, 1) = 0, \text{ or } F^+(1, 0) = 0 \text{ and } F^-(0, 1) = 1,$$

and

$$\text{either } F^+(0, 1) = 0 \text{ and } F^-(1, 0) = 1 \text{ or } F^+(0, 1) = 1 \text{ and } F^-(1, 0) = 0.$$

Therefore we can have the following four combinations: (i) $F^+(r, s) = F^-(r, s) = r$ for $r, s \in \{0, 1\}$, (ii) $F^+(r, s) = \max\{r, s\}$ and $F^-(r, s) = \min\{r, s\}$ for $r, s \in \{0, 1\}$, (iii) $F^+(r, s) =$

$\min\{r, s\}$ and $F^-(r, s) = \max\{r, s\}$ for $r, s \in \{0, 1\}$, or (iv) $F^+(r, s) = F^-(r, s) = s$ for $r, s \in \{0, 1\}$. These correspond to $T1_A = 1_A$, $T1_A = P_H 1_A$, $T1_A = (P_H 1_A)^\dagger$, and $T_A = (1_A)^\dagger$, each for all $A \in \mathcal{L}^n$, respectively. In particular, T maps characteristic functions of sets in \mathcal{L}^n to characteristic functions of sets in \mathcal{L}^n , and \diamond_T is Id , \diamond_{P_H} , $\diamond_{P_H^\dagger}$, or \dagger . \square

Despite the previous result, maps T that are both pointwise and equimeasurable need not be one of the four special maps, $T = \text{Id}$, $T = \dagger$, $T = P_H$, or $T = P_H^\dagger$. Indeed, by Theorem 5.1, it is enough to define T via associated functions F^+ and F^- that satisfy (35). For example, one can take

$$F^+(r, s) = \begin{cases} r, & \text{if } r \in \mathbb{Q} \cap D, \\ s, & \text{if } r \in D \setminus \mathbb{Q} \end{cases} \quad \text{and} \quad F^-(s, r) = \begin{cases} s, & \text{if } r \in \mathbb{Q} \cap D, \\ r, & \text{if } r \in D \setminus \mathbb{Q}. \end{cases}$$

The next few results supply further conditions that eliminate such exotic examples.

Lemma 5.3. *Let $H \in \mathcal{G}(n, n-1)$ be oriented, let $X = \mathcal{M}(\mathbb{R}^n)$, $\mathcal{M}_+(\mathbb{R}^n)$, $\mathcal{S}(\mathbb{R}^n)$, or $\mathcal{V}(\mathbb{R}^n)$, and suppose that $T : X \rightarrow X$ is pointwise with respect to H and equimeasurable. If the functions F^+ and F^- associated with T are continuous on D^2 , then $T = \text{Id}$, $T = \dagger$, $T = P_H$, or $T = P_H^\dagger$.*

Proof. As F^+ is continuous on D^2 , (35) implies that either $F^+(r, s) = r$ for $(r, s) \in D^2$ or $F^+(r, s) = s$ for $(r, s) \in D^2$. Let

$$E_1 = \{(r, s) \in D^2 : r \geq s\} \quad \text{and} \quad E_2 = \{(r, s) \in D^2 : r \leq s\}.$$

As E_1 is connected, $\{(r, s) \in E_1 : F^+(r, s) = r\}$ is either empty or E_1 . If it is empty, then $F^+(r, s) = s$ on E_1 . It follows that either $F^+(r, s) = r = \max\{r, s\}$ on E_1 or $F^+(r, s) = s = \min\{r, s\}$ on E_1 . In the same way, either $F^+(r, s) = \max\{r, s\}$ on E_2 or $F^+(r, s) = \min\{r, s\}$ on E_2 . Similar arguments show that the same possibilities hold when F^+ is replaced by F^- . Taking (35) into account, we arrive at four possibilities for F^+ and F^- on D^2 , corresponding to those for T in the statement of the corollary. \square

Motivated by the previous lemma, we now seek conditions ensuring that the associated functions F^+ and F^- are continuous on D^2 .

Lemma 5.4. *Let $H \in \mathcal{G}(n, n-1)$ be oriented, let $X = \mathcal{M}(\mathbb{R}^n)$, $\mathcal{M}_+(\mathbb{R}^n)$, $\mathcal{S}(\mathbb{R}^n)$, or $\mathcal{V}(\mathbb{R}^n)$, and suppose that $T : X \rightarrow X$ is pointwise with respect to H with associated functions F^+ and F^- . Then*

- (i) *T is monotonic if and only if F^+ and F^- are increasing in each variable, and*
- (ii) *if T is a rearrangement, then F^+ and F^- are continuous on D^2 .*

Proof. (i) It follows from the definition of a pointwise map that T is monotonic if F^+ and F^- are increasing in each variable. For the other implication, suppose that T is monotonic. Let $r_1, s_1, r_2, s_2 \in D$ satisfy $r_1 \leq r_2$ and $s_1 \leq s_2$. For $i = 1, 2$, define $f_i \in X$ by

$$(41) \quad f_i(x) = r_i 1_{B^n \cap H^+}(x) + s_i 1_{B^n \cap H^-}(x) + \min\{0, r_i, s_i\} 1_{\mathbb{R}^n \setminus B^n}(x)$$

for $x \in \mathbb{R}^n$. (The last term in (41) is to ensure that $f_i \in X$ when $X = \mathcal{S}(\mathbb{R}^n)$ or $\mathcal{V}(\mathbb{R}^n)$.) For $x \in B^n \cap H^+$ we have

$$Tf_i(x) = F^+(f_i(x), f_i^\dagger(x)) = F^+(r_i, s_i)$$

for $i = 1, 2$. Since $f_1 \leq f_2$ and T is monotonic, $Tf_1(x) \leq Tf_2(x)$ for almost all $x \in B^n \cap H^+$ and hence $F^+(r_1, s_1) \leq F^+(r_2, s_2)$. A similar argument holds for F^- . It follows that F^+ and F^- are increasing in each variable.

(ii) Suppose that T is a rearrangement, i.e., equimeasurable and monotonic. Then (35) holds. Let (r_k, s_k) , $k \in \mathbb{N}$, be a sequence in D^2 converging to (r, s) . We may assume that $r \neq s$, as otherwise (35) implies $F^\pm(r_k, s_k) \rightarrow F^\pm(r, s)$ as $k \rightarrow \infty$. Without loss of generality, suppose that $r > s$. By considering subsequences, we may also assume that $\{(r_k, s_k) : k \in \mathbb{N}\}$ is contained in one of the four sets

$$\begin{aligned} D_{++} &= \{(r', s') \in D^2 : r' \geq r, s' \geq s\}, & D_{+-} &= \{(r', s') \in D^2 : r' \geq r, s' \leq s\}, \\ D_{-+} &= \{(r', s') \in D^2 : r' \leq r, s' \geq s\}, & D_{--} &= \{(r', s') \in D^2 : r' \leq r, s' \leq s\}. \end{aligned}$$

If $\{(r_k, s_k) : k \in \mathbb{N}\} \subset D_{++}$ and $F^+(r, s) = r$, then (i) implies that for sufficiently large k , we have

$$r = F^+(r, s) \leq F^+(r_k, s_k) \leq F^+(r_k, r_k) = r_k.$$

Since $r_k \rightarrow r$, this shows that $F^+(r_k, s_k) \rightarrow F^+(r, s)$, and then $F^-(r_k, s_k) \rightarrow F^-(r, s)$ as $k \rightarrow \infty$ by (35). If $\{(r_k, s_k) : k \in \mathbb{N}\} \subset D_{+-}$ and $F^+(r, s) = s$, we have $F^-(r, s) = r$ by (35), and the same arguments can be applied to F^- .

Suppose that $\{(r_k, s_k) : k \in \mathbb{N}\} \subset D_{-+}$. Then $(r, s_k) \in D_{--}$ and $(r_k, s) \in D_{++}$, so both sides of

$$F^+(r, s_k) \leq F^+(r_k, s_k) \leq F^+(r_k, s),$$

converge to $F^+(r, s)$ and it follows that $F^+(r_k, s_k) \rightarrow F^+(r, s)$ as $k \rightarrow \infty$. Equation (35) now yields $F^-(r_k, s_k) \rightarrow F^-(r, s)$ as $k \rightarrow \infty$. The remaining case $\{(r_k, s_k) : k \in \mathbb{N}\} \subset D_{--}$ is treated in a similar way. \square

Lemma 5.5. *Let $H \in \mathcal{G}(n, n-1)$ be oriented, let $X = \mathcal{M}(\mathbb{R}^n)$ (or $\mathcal{M}_+(\mathbb{R}^n)$, $\mathcal{S}(\mathbb{R}^n)$, or $\mathcal{V}(\mathbb{R}^n)$), and suppose that $T : X \rightarrow X$ is pointwise with respect to H , equimeasurable, and maps linear functions (or piecewise linear continuous functions, respectively) to continuous functions. Then the functions F^+ and F^- associated with T are continuous on D^2 .*

Proof. Assume first that $X = \mathcal{M}(\mathbb{R}^n)$. Since T is equimeasurable, (35) holds, implying that $F^\pm(r, r) = r$ for $r \in D$. Suppose that T maps linear functions to continuous functions. We may assume that $H = e_n^\perp$ and define $f(x) = x \cdot (e_1 + e_n)$ for $x \in \mathbb{R}^n$. Then f is linear and if $x = (x_1, \dots, x_n)$, then $f(x) = x_1 + x_n$ and $f^\dagger(x) = x_1 - x_n$. Thus for $x \in H^+$, we have $Tf(x) = F^+(f(x), f^\dagger(x)) = F^+(x_1 + x_n, x_1 - x_n)$. Let $x_{rs} = ((r+s)/2, 0, \dots, 0, (r-s)/2)$ and note that $x_{rs} \in H^+$ if and only if $(r, s) \in E_1 = \{(r, s) \in D^2 : r \geq s\}$. Consequently,

$$(42) \quad (r, s) \mapsto Tf(x_{rs}) = F^+(r, s)$$

is continuous on E_1 . A similar argument using the linear function $f(x) = x \cdot (e_1 - e_n)$ shows that F^+ is continuous on $E_2 = \{(r, s) \in D^2 : r \leq s\}$. It follows that F^+ is continuous on D^2 , and we arrive at the same conclusion for F^- similarly.

Suppose that $X = \mathcal{M}_+(\mathbb{R}^n)$, $\mathcal{S}(\mathbb{R}^n)$, or $X = \mathcal{V}(\mathbb{R}^n)$. We adopt the notation of the first part of this proof. For $(r_0, s_0) \in E_1$ we can choose a nonnegative piecewise linear continuous function f coinciding with $x \mapsto |x \cdot (e_1 + e_n)| \geq 0$ on the ball tB^n with $t > 2^{-1/2}\|(r_0, s_0)\|$ and vanishing outside an even larger ball. Clearly $f \in X$. Following the arguments above, it can be seen that the restriction of $(r, s) \mapsto F^+(r, s)$ to E_1 is continuous at (r_0, s_0) . Similar arguments for $(r_0, s_0) \in E_2$ and for F^- lead to the desired conclusion. \square

Let $1 \leq p \leq \infty$. A function $F : D^2 \rightarrow \mathbb{R}^2$ is l_2^p -contracting if

$$\|F(r, s) - F(r', s')\|_p \leq \|(r, s) - (r', s')\|_p$$

for $(r, s), (r', s') \in D^2$, where $\|\cdot\|_p$ is the norm in l_2^p . For instance, the function

$$(43) \quad F(r, s) = (M_\infty(r, s), M_{-\infty}(s, r)),$$

with the associated functions of the polarization operation as components, is l_2^p -contracting. In fact, this can be checked directly for $p = \infty$. For $1 \leq p < \infty$, it follows from the inequality

$$|r - s'|^p + |s - r'|^p \leq |r - r'|^p + |s - s'|^p,$$

where $r \leq s$ and $s' \leq r'$, which is in turn a consequence of the convexity of the function $|t|^p$, $p \geq 1$. Indeed, as in [48, p. 43], the latter implies that if $a \in \mathbb{R}$ and $b, c \geq 0$, then

$$|a|^p - |a - b|^p \leq |a + c|^p - |a - b + c|^p,$$

so the required inequality results from setting $a = s - r'$, $b = s - r$, and $c = r' - s'$.

Lemma 5.6. *Let $H \in \mathcal{G}(n, n-1)$ be oriented, let $X = \mathcal{M}(\mathbb{R}^n)$, $\mathcal{M}_+(\mathbb{R}^n)$, $\mathcal{S}(\mathbb{R}^n)$, or $\mathcal{V}(\mathbb{R}^n)$, and let $1 \leq p \leq \infty$. Suppose that $T : X \rightarrow X$ is pointwise with respect to H with associated functions F^+ and F^- . Then*

(i) *T is L^p -contracting if and only if*

$$(44) \quad F(s, t) = (F^+(s, t), F^-(t, s)), \quad (s, t) \in D^2,$$

is l_2^p -contracting, and

(ii) *if T is L^p -contracting, then F^+ and F^- are Lipschitz on D^2 with Lipschitz constant $\sqrt{2}$.*

Proof. (i) We first show that if F is l_2^p -contracting, then T is L^p -contracting. Let $f_1, f_2 \in X$. Suppose that $p < \infty$. As T is pointwise, we have

$$\begin{aligned} \|Tf_1 - Tf_2\|_p^p &= \int_{\mathbb{R}^n} |Tf_1(x) - Tf_2(x)|^p dx \\ &= \int_{H^+} |F^+(f_1(x), f_1^\dagger(x)) - F^+(f_2(x), f_2^\dagger(x))|^p dx \\ &\quad + \int_{H^-} |F^-(f_1(x), f_1^\dagger(x)) - F^-(f_2(x), f_2^\dagger(x))|^p dx. \end{aligned}$$

Substituting x by x^\dagger in the second integral and using the definition of F in (44), we obtain

$$(45) \quad \|Tf_1 - Tf_2\|_p^p = \int_{H^+} \|F(f_1(x), f_1^\dagger(x)) - F(f_2(x), f_2^\dagger(x))\|_p^p dx.$$

In a similar fashion, it is easily seen that

$$(46) \quad \|f_1 - f_2\|_p^p = \int_{H^+} \|(f_1(x), f_1^\dagger(x)) - (f_2(x), f_2^\dagger(x))\|_p^p dx.$$

Now if F is l_2^p -contracting, (45) is bounded from above by (46), so T is L^p -contracting.

The case when $p = \infty$ follows similarly from the equations

$$\|Tf_1 - Tf_2\|_\infty = \operatorname{ess\,sup}_{x \in H^+} \|F(f_1(x), f_1^\dagger(x)) - F(f_2(x), f_2^\dagger(x))\|_\infty$$

and

$$\|f_1 - f_2\|_\infty = \operatorname{ess\,sup}_{x \in H^+} \|(f_1(x), f_1^\dagger(x)) - (f_2(x), f_2^\dagger(x))\|_\infty.$$

To show the other direction, let $r_1, s_1, r_2, s_2 \in D$ and define $f_i \in X$, $i = 1, 2$, by (41) with the term $\min\{0, r_i, s_i\}$ replaced by $\min\{0, r_1, r_2, s_1, s_2\}$. As T is L^p -contracting, we get

$$\frac{\kappa_n}{2} \|F(r_1, s_1) - F(r_2, s_2)\|_p^p \leq \|Tf_1 - Tf_2\|_p^p \leq \|f_1 - f_2\|_p^p = \frac{\kappa_n}{2} \|(r_1, s_1) - (r_2, s_2)\|_p^p,$$

so F is l_2^p -contracting.

(ii) Suppose that T is L^p -contracting. Then F is l_2^p -contracting by (i), so

$$\begin{aligned} |F^+(r_1, s_1) - F^+(r_2, s_2)| &\leq \|F(r_1, s_1) - F(r_2, s_2)\|_p \leq \|(r_1, s_1) - (r_2, s_2)\|_p \\ &\leq \sqrt{2} \|(r_1, s_1) - (r_2, s_2)\|, \end{aligned}$$

as $\|\cdot\|_p \leq \|\cdot\|_1 \leq \sqrt{2}\|\cdot\|$. Hence F^+ is Lipschitz on D^2 with Lipschitz constant $\sqrt{2}$, and the same argument can be applied to F^- . \square

Lemma 5.7. *Let $H \in \mathcal{G}(n, n-1)$ be oriented, let $X = \mathcal{M}(\mathbb{R}^n)$, $\mathcal{M}_+(\mathbb{R}^n)$, $\mathcal{S}(\mathbb{R}^n)$, or $\mathcal{V}(\mathbb{R}^n)$, and suppose that $T : X \rightarrow X$ is pointwise with respect to H with associated functions F^+ and F^- . If T reduces the modulus of continuity, then T is L^∞ -contracting.*

Proof. Suppose that T reduces the modulus of continuity and let $r, s, r', s' \in D$. Choose $f \in X$ and $x, y \in H^+$ such that $f(x) = r$, $f(y) = r'$, $f(x^\dagger) = s$, $f(y^\dagger) = s'$, and

$$\omega_d(f) = \|(f(x), f(x^\dagger)) - (f(y), f(y^\dagger))\|_\infty = \|(r, s) - (r', s')\|_\infty,$$

where $d = \|x - y\|$. As T reduces the modulus of continuity,

$$(47) \quad |F^+(r, s) - F^+(r', s')| = |Tf(x) - Tf(y)| \leq \omega_d(Tf) \leq \omega_d(f) = \|(r, s) - (r', s')\|_\infty.$$

A similar relation for $|Tf(x^\dagger) - Tf(y^\dagger)|$ yields

$$(48) \quad |F^-(s, r) - F^-(s', r')| \leq \|(r, s) - (r', s')\|_\infty.$$

From (47) and (48), we conclude that F (defined by (44)) is l_2^∞ -contracting and the result follows from Lemma 5.6(i) with $p = \infty$. \square

Summarizing, we have the following set of characterizations.

Theorem 5.8. *Let $H \in \mathcal{G}(n, n-1)$ be oriented, let $X = \mathcal{M}(\mathbb{R}^n)$ (or $\mathcal{M}_+(\mathbb{R}^n)$, $\mathcal{S}(\mathbb{R}^n)$, or $\mathcal{V}(\mathbb{R}^n)$), and suppose that $T : X \rightarrow X$ is pointwise with respect to H and equimeasurable. The following statements are equivalent.*

- (i) *The associated functions F^+ and F^- are continuous on D^2 .*
- (ii) *T is monotonic.*
- (iii) *T is a rearrangement.*
- (iv) *T maps linear functions (or piecewise linear continuous functions, respectively) to continuous functions.*
- (v) *T is L^p -contracting for some (or, equivalently, for all) $1 \leq p \leq \infty$.*
- (vi) *T reduces the modulus of continuity.*
- (vii) *$T = \text{Id}$, $T = \dagger$, $T = P_H$, or $T = P_H^\dagger$.*

Proof. It is easy to check that (vii) implies (i)–(iv) and (vi). To see that (vii) \Rightarrow (v), it is enough to apply Lemma 5.6 with $T = P_H$, since the function F in (43) is l_2^p -contracting. Lemma 5.3 gives (i) \Rightarrow (vii) and Lemmas 5.4(ii), 5.5, and 5.6(ii) show that (iii) \Rightarrow (i), (iv) \Rightarrow (i), and (v) \Rightarrow (i), respectively. That (ii) \Rightarrow (iii) follows from the definition of a rearrangement. Finally, the implication (vi) \Rightarrow (v) follows from Lemma 5.7 and the implications already established. \square

6. GENERAL MAPS BETWEEN SETS AND BETWEEN FUNCTIONS

The proof of the following result is essentially the same as that of [7, Theorem 10.1(i)]. The statement is more general, since the map \diamond need not be an i -symmetrization in the sense of [7], the sets concerned need not be compact, and invariance on H -symmetric cylinders is replaced by the weaker condition that \diamond respects H -cylinders.

Lemma 6.1. *Let $i \in \{1, \dots, n-1\}$ and let $H \in \mathcal{G}(n, i)$. Suppose that $\diamond : \mathcal{E} \subset \mathcal{L}^n \rightarrow \mathcal{L}^n$ is monotonic, measure preserving, and respects H -cylinders. Then*

$$(49) \quad \mathcal{H}^{n-i}((\diamond K) \cap (H^\perp + x)) = \mathcal{H}^{n-i}(K \cap (H^\perp + x))$$

for $K \in \mathcal{E}$ and \mathcal{H}^i -almost all $x \in H$.

In the following results, we always assume for convenience that $\mathcal{K}_n^n \subset \mathcal{E}$, even though this assumption can sometimes be weakened.

Lemma 6.2. *Let $H = u^\perp$, $u \in S^{n-1}$, let $\mathcal{K}_n^n \subset \mathcal{E} \subset \mathcal{L}^n$, and suppose that $\diamond : \mathcal{E} \rightarrow \mathcal{L}^n$ is monotonic, respects H -cylinders, and maps balls to balls. Then there is a contraction (i.e., a Lipschitz function with Lipschitz constant 1) $\varphi_\diamond : \mathbb{R} \rightarrow \mathbb{R}$ such that*

$$(50) \quad \diamond B(x + tu, r) = B(x + \varphi_\diamond(t)u, r),$$

essentially, for $r > 0$ and $x \in H$.

Proof. In the proof we ignore sets of \mathcal{H}^n -measure zero. For $r > 0$ and $x \in H$, let $C(x, r)$ be the infinite spherical cylinder of radius r and axis $H^\perp + x$. Let $t \in \mathbb{R}$. By our assumptions, $\diamond B(x + tu, r)$ is a ball contained in $C(x, r)$. We claim that the radius of $\diamond B(x + tu, r)$

is r . Indeed, if it is less than r , there is a ball $B_0 \subset B(x + tu, r)$ whose projection on H is disjoint from that of $\diamond B(x + tu, r)$. However, by the monotonicity of \diamond , $\diamond B_0 \subset \diamond B(x + tu, r)$, contradicting the fact that \diamond respects H -cylinders. It follows that $\diamond B(x + tu, r) = B(x + t'u, r)$ for some $t' = t'(r, t, x) \in \mathbb{R}$.

Fix $t \in \mathbb{R}$ and suppose that neither of the balls $B(x_j + tu, r_j)$, $j = 1, 2$, contains the other. Let $z \in H$ and $s \in \mathbb{R}$ be such that $B(z + tu, s) \supset B(x_1 + tu, r_1) \cup B(x_2 + tu, r_2)$ is tangent to both $B(x_1 + tu, r_1)$ and $B(x_2 + tu, r_2)$ at points in $\partial C(z, s) \cap (H + tu)$. Then there are $t' = t'(s, t, z)$ and $t'_j = t'(r_j, t, x_j)$, $j = 1, 2$, with

$$\begin{aligned} B(z + t'u, s) &= \diamond B(z + tu, s) \supset \diamond B(x_1 + tu, r_1) \cup \diamond B(x_2 + tu, r_2) \\ &= B(x_1 + t'_1 u, r_1) \cup B(x_2 + t'_2 u, r_2), \end{aligned}$$

where we used the monotonicity of \diamond . It follows that $B(z + t'u, s)$ contains $B(x_1 + t'_1 u, r_1)$ and $B(x_2 + t'_2 u, r_2)$ and is tangent to both of them at points in $\partial C(z, s) \cap (H + t'u)$. This forces $t'_1 = t' = t'_2$, so

$$(51) \quad t'(r_1, t, x_1) = t'(r_2, t, x_2).$$

If one of $B(x_j + tu, r_j)$, $j = 1, 2$, is contained in the other, say $B(x_1 + tu, r_1) \subset B(x_2 + tu, r_2)$, choose $B(z + tu, s)$ disjoint from $B(x_2 + tu, r_2)$. Then (51), applied first to the disjoint balls $B(x_1 + tu, r_1)$ and $B(z + tu, s)$, and then to the disjoint balls $B(x_2 + tu, r_2)$ and $B(z + tu, s)$, yields

$$t'(r_1, t, x_1) = t'(s, t, z) = t'(r_2, t, x_2).$$

This shows that (51) holds generally, so $t'(r, t, x)$ is independent of r and x . Then $\varphi_\diamond(t) = t'$ is the required function.

Suppose that there are $s, t \in \mathbb{R}$ and $\varepsilon > 0$ such that

$$|\varphi_\diamond(s) - \varphi_\diamond(t)| = |s - t| + \varepsilon.$$

Let $|s - t|/2 < r < (|s - t| + \varepsilon)/2$. If $K = B(su, r) \cap B(tu, r)$, then there is a ball $B_1 \subset K$ and hence $\emptyset \neq \diamond B_1 \subset \diamond K$. On the other hand, since $K \in \mathcal{E}$, the monotonicity of \diamond also implies that

$$\diamond K \subset \diamond B(su, r) \cap \diamond B(tu, r) = B(\varphi_\diamond(s)u, r) \cap B(\varphi_\diamond(t)u, r) = \emptyset.$$

This contradiction shows that φ_\diamond is Lipschitz with Lipschitz constant 1. \square

Lemma 6.3. *Let $H = u^\perp$, $u \in S^{n-1}$, let $\mathcal{K}_n^n \subset \mathcal{E} \subset \mathcal{L}^n$, and suppose that $\diamond : \mathcal{E} \rightarrow \mathcal{L}^n$ is monotonic, measure preserving, respects H -cylinders, and maps balls to balls. Let $\varphi_\diamond : \mathbb{R} \rightarrow \mathbb{R}$ be the function from Lemma 6.2. Then for each H -symmetric $K \in \mathcal{K}_n^n$ and $t \in \mathbb{R}$, we have*

$$(52) \quad \diamond(K + tu) = K + \varphi_\diamond(t)u,$$

essentially.

Proof. In the proof we ignore sets of \mathcal{H}^n -measure zero. If $r > 0$ and $x \in H$, let $C(x, r)$ be the infinite spherical cylinder of radius r and axis $H^\perp + x$. Suppose first that $K =$

$(B(x, r) \cap H) + [-su, su]$ is an H -symmetric spherical cylinder, where $r, s > 0$ and $x \in H$. Let $t \in \mathbb{R}$ and let $S = \cup_{m \in \mathbb{N}} B(z_m + tu, s)$, where $\{z_m : m \in \mathbb{N}\}$ is dense in $B(x, r) \cap H$. Then

$$\text{int}(K + tu) \subset S \cap C(x, r) \subset K + tu$$

and we can write $K + tu = S \cap C(x, r)$ since we are ignoring sets of \mathcal{H}^n -measure zero. By the monotonicity of \diamond ,

$$(53) \quad S' = \cup_{m \in \mathbb{N}} \diamond B(z_m + tu, s) \subset \diamond S,$$

as there are countably many sets in the union. By Lemma 6.2,

$$S' = \cup_{m \in \mathbb{N}} B(z_m + \varphi_\diamond(t)u, s) = S + (\varphi_\diamond(t) - t)u.$$

This and the measure-preserving property of \diamond yield $\mathcal{H}^n(S') = \mathcal{H}^n(S) = \mathcal{H}^n(\diamond S)$ and therefore (53) implies that $\diamond S = S'$. Since \diamond respects H -cylinders, we have $\diamond(K + tu) \subset C(x, r)$ and hence

$$\diamond(K + tu) \subset \diamond S \cap C(x, r) = (S + (\varphi_\diamond(t) - t)u) \cap C(x, r) = K + \varphi_\diamond(t)u.$$

Now let $K \in \mathcal{K}_n^n$ be an arbitrary H -symmetric set. Let $\{x_m : m \in \mathbb{N}\}$ be a dense set in $K|H$. Since $K + tu$ is symmetric with respect to $H + tu$, it is clear that we can find H -symmetric spherical cylinders $C_k = (B(x_{m_k}, r_k) \cap H) + [-s_k u, s_k u]$, $k \in \mathbb{N}$, where r_k, s_k are positive rationals, such that

$$(54) \quad \text{int}(K + tu) \subset \cup_k (C_k + tu) \subset K + tu.$$

By the previous paragraph, $\diamond(C_k + tu) = C_k + \varphi_\diamond(t)u$, so (54) and the monotonicity of \diamond yields

$$(55) \quad E = \cup_k \diamond(C_k + tu) = \cup_k (C_k + \varphi_\diamond(t)u) = (\cup_k C_k) + \varphi_\diamond(t)u \subset \diamond(K + tu).$$

Since E is a translate of $\cup_k C_k$, (54) and the measure-preserving property of \diamond imply that $\mathcal{H}^n(E) = \mathcal{H}^n(K + tu) = \mathcal{H}^n(\diamond(K + tu))$. Hence, by (54) and (55), we have

$$\diamond(K + tu) = E = (\cup_k C_k) + \varphi_\diamond(t)u = K + \varphi_\diamond(t)u,$$

essentially. □

Since φ_\diamond is a contraction, we have

$$(56) \quad \varphi_\diamond(0) = 0 \quad \Leftrightarrow \quad |\varphi_\diamond(t)| \leq |t|, \text{ for } t \in \mathbb{R}.$$

Indeed, the right-hand side follows from the left-hand side on setting $s = 0$ in $|\varphi_\diamond(s) - \varphi_\diamond(t)| \leq |s - t|$, and the converse is trivial.

Corollary 6.4. *Let $H = u^\perp$, $u \in S^{n-1}$, and suppose that $\diamond : \mathcal{K}_n^n \rightarrow \mathcal{L}^n$ is monotonic, measure preserving, respects H -cylinders, and maps balls to balls. Let $\varphi_\diamond : \mathbb{R} \rightarrow \mathbb{R}$ be the function from Lemma 6.2. Then \diamond is invariant on H -symmetric sets if and only if \diamond is invariant on H -symmetric cylinders, and this occurs if and only if either condition in (56) holds.*

Proof. In the proof we ignore sets of \mathcal{H}^n -measure zero. Suppose that \diamond is invariant on H -symmetric cylinders. If $r > 0$ and $t \in \mathbb{R}$, then

$$B(tu, r) \subset (B(tu, r)|H) + [-(|t| + r)u, (|t| + r)u] = C,$$

say, an H -symmetric cylinder. Then

$$B(\varphi_\diamond(t)u, r) = \diamond B(tu, r) \subset \diamond C = C.$$

This yields

$$[\varphi_\diamond(t) - r, \varphi_\diamond(t) + r] = B(\varphi_\diamond(t)u, r)|H^\perp \subset C|H^\perp = [-(|t| + r)u, (|t| + r)u].$$

Therefore $|\varphi_\diamond(t)| \leq |t|$. Then $\varphi_\diamond(0) = 0$, so setting $t = 0$ in (52) implies that \diamond is invariant on H -symmetric sets. It follows directly from the definitions that the latter implies that \diamond is invariant on H -symmetric cylinders. \square

We shall find it convenient to define, for $K \in \mathcal{K}^n$, $t \in \mathbb{R}$, and $u \in S^{n-1}$,

$$(57) \quad K_t = (K - tu) \cap (K^\dagger + tu).$$

Note that if $H = u^\perp$, then K_t is H -symmetric.

Lemma 6.5. *Let $H = u^\perp$, $u \in S^{n-1}$, and let $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ be an arbitrary contraction. Then*

$$(58) \quad (\cup_{t \in \mathbb{R}} (K_t + \varphi(t)u)) \cap (H^\perp + x) = (K \cap (H^\perp + x)) + (\varphi(t_x) - t_x)u$$

for $K \in \mathcal{K}_n^n$ and $x \in H$, where K_t is defined by (57) and $x + t_x u$ is the midpoint of $K \cap (H^\perp + x)$.

Proof. If $M \in \mathcal{K}^n$, $x \in H$, and $s, t \in \mathbb{R}$, then $x + su \in M_t$ if and only if $x + (t \pm s)u \in M$. Applying this first with $M = K$ and then with $M = K \cap (H^\perp + x)$ shows that

$$(59) \quad K_t \cap (H^\perp + x) = (K \cap (H^\perp + x))_t.$$

Note also that

$$(60) \quad (K \cap (H^\perp + x))_{t_x} = (K \cap (H^\perp + x)) - t_x u.$$

Let $x \in H$ be such that the (possibly degenerate) line segment $S = K \cap (H^\perp + x)$ is nonempty and let $2r_x \geq 0$ be its length. Then $S_{t_x+t} = \emptyset$ for $|t| > r_x$. Suppose that $|t| \leq r_x$. Then $S_{t_x+t} = [x - (r_x - |t|)u, x + (r_x - |t|)u]$. As φ is a contraction, $|\varphi(t_x + t) - \varphi(t_x)| \leq |t|$ and hence

$$S_{t_x+t} + (\varphi(t_x + t) - \varphi(t_x))u \subset S_{t_x+t} + [-|t|u, |t|u] = [x - r_x u, x + r_x u] = S_{t_x}.$$

Rearranging and replacing $t_x + t$ by t , we obtain

$$(61) \quad (K \cap (H^\perp + x))_t + \varphi(t)u = S_t + \varphi(t)u \subset S_{t_x} + \varphi(t_x)u = (K \cap (H^\perp + x))_{t_x} + \varphi(t_x)u$$

whenever the left-hand side is nonempty. Hence (61) holds for $x \in H$ and $t \in \mathbb{R}$.

Applying, in turn, (59), (61), and (60), we obtain

$$(\cup_{t \in \mathbb{R}} (K_t + \varphi(t)u)) \cap (H^\perp + x) \subset (K \cap (H^\perp + x)) + (\varphi(t_x) - t_x)u.$$

The reverse inclusion is a consequence of

$$(\cup_{t \in \mathbb{R}} (K_t + \varphi(t)u)) \cap (H^\perp + x) \supset (K_{t_x} \cap (H^\perp + x)) + \varphi(t_x)u$$

together with (59) and (60). This proves (58). \square

Theorem 6.6. *Let $H = u^\perp$, $u \in S^{n-1}$, let $\diamond : \mathcal{K}_n^n \rightarrow \mathcal{L}^n$, and let $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ be a contraction. The following are equivalent.*

(i) \diamond is monotonic, measure preserving, respects H -cylinders, and maps balls to balls. The mapping φ coincides with the function φ_\diamond from Lemma 6.2.

(ii) For each $K \in \mathcal{K}_n^n$,

$$(62) \quad \diamond K = \bigcup_{t \in \mathbb{R}} \text{int}(K_t + \varphi(t)u),$$

essentially, where K_t is defined by (57).

(iii) For each $K \in \mathcal{K}_n^n$ and \mathcal{H}^{n-1} -almost all $x \in H$,

$$(63) \quad (\diamond K) \cap (H^\perp + x) = (K \cap (H^\perp + x)) + (\varphi(t_x) - t_x)u,$$

up to a set of \mathcal{H}^1 -measure zero, where $x + t_x u$ is the midpoint of $K \cap (H^\perp + x)$.

Proof. (i) \Rightarrow (ii). In the proof we ignore sets of \mathcal{H}^n -measure zero. Let $K \in \mathcal{K}_n^n$ and let $t \in \mathbb{R}$. The set K_t defined by (57) is H -symmetric, so $K_t + tu \subset K$ is symmetric with respect to $H + tu$. From the monotonicity of \diamond and (52), we obtain

$$\bigcup_{t \in \mathbb{Q}} (K_t + \varphi_\diamond(t)u) = \bigcup_{t \in \mathbb{Q}} \diamond(K_t + tu) \subset \diamond K,$$

because there are countably many sets in the union. Let $t \in \mathbb{R}$ with $K_t \neq \emptyset$ and choose $t_m \in \mathbb{Q}$ with $K_{t_m} \neq \emptyset$ such that $t_m \rightarrow t$ as $m \rightarrow \infty$. Since φ_\diamond is a Lipschitz map and $t \mapsto K_t$ is continuous on $\{t \in \mathbb{R} : K_t \neq \emptyset\}$, we have $K_{t_m} + \varphi_\diamond(t_m)u \rightarrow K_t + \varphi_\diamond(t)u$ in the Hausdorff metric as $m \rightarrow \infty$ and hence

$$\text{int}(K_t + \varphi_\diamond(t)u) \subset \bigcup_{m \in \mathbb{N}} (K_{t_m} + \varphi_\diamond(t_m)u).$$

It follows that

$$U = \bigcup_{t \in \mathbb{R}} \text{int}(K_t + \varphi_\diamond(t)u) \subset \diamond K.$$

Let $V = \bigcup_{x \in (\text{int } K)|H} \text{int}(K_{t_x} + \varphi_\diamond(t_x)u)$, where $x + t_x u$ is the midpoint of $K \cap (H^\perp + x)$. Then

$$(64) \quad V \subset U \subset \diamond K.$$

If $x \in (\text{int } K)|H$, then $(K_{t_x} + t_x u) \cap (H^\perp + x) = K \cap (H^\perp + x)$, so

$$(65) \quad \begin{aligned} \mathcal{H}^1(V \cap (H^\perp + x)) &\geq \mathcal{H}^1((\text{int}(K_{t_x} + \varphi_\diamond(t_x)u)) \cap (H^\perp + x)) \\ &= \mathcal{H}^1((K_{t_x} + \varphi_\diamond(t_x)u) \cap (H^\perp + x)) \\ &= \mathcal{H}^1((K_{t_x} + t_x u) \cap (H^\perp + x)) = \mathcal{H}^1(K \cap (H^\perp + x)). \end{aligned}$$

From this, an application of Fubini's theorem and the measure-preserving property of \diamond give $\mathcal{H}^n(V) \geq \mathcal{H}^n(K) = \mathcal{H}^n(\diamond K)$. Therefore, by (64), we have $V = U = \diamond K$, essentially.

(ii) \Rightarrow (iii). Let $K \in \mathcal{K}_n^n$ and let $W = \bigcup_{t \in \mathbb{R}} (K_t + \varphi_\diamond(t)u)$. Then

$$U \cap (H^\perp + x) \subset W \cap (H^\perp + x) = (K \cap (H^\perp + x)) + (\varphi(t_x) - t_x)u$$

for $x \in H$, by (58). Moreover, (64) and (65) imply that

$$\mathcal{H}^1(U \cap (H^\perp + x)) \geq \mathcal{H}^1(V \cap (H^\perp + x)) \geq \mathcal{H}^1(K \cap (H^\perp + x)),$$

so

$$(66) \quad U \cap (H^\perp + x) = (K \cap (H^\perp + x)) + (\varphi(t_x) - t_x)u,$$

up to a set of \mathcal{H}^1 -measure zero. By (ii), $\diamond K = U$, essentially, and (iii) follows.

(iii) \Rightarrow (ii). Let $K \in \mathcal{K}_n^n$. If (iii) holds, then (ii) follows from (66) (which did not require (ii) for its proof) and Fubini's theorem.

(ii) \Rightarrow (i). Assume that (ii) holds. Clearly, \diamond is monotonic due to (62). We have already seen that (ii) implies (iii). That \diamond is measure preserving and respects H -cylinders follows directly from (63) and Fubini's theorem. If $z \in H$, $t \in \mathbb{R}$, and $r > 0$, then (62) implies that

$$\diamond B(z + tu, r) \supset B(z + tu, r)_t + \varphi(t)u = B(z + \varphi(t)u, r),$$

essentially. Together with the measure-preserving property, this proves that \diamond maps balls to balls. \square

None of the properties of \diamond listed in Theorem 6.6(i) can be omitted. Indeed, no three of these properties imply the fourth, as is shown by the map \diamond' from Example 6.8 below, the map taking a convex body to the o -symmetric ball of the same volume, Minkowski symmetrization (see [7, Section 3]), and Example 6.17 below.

In view of (63), a map \diamond satisfying Theorem 6.6(i) may be regarded as a “still” in a parallel chord movement in the direction u , in the sense of convex geometry. The concept of a parallel chord movement was, in a more general form, introduced by Rogers and Shephard [41] (see also [43, p. 543]) and is extremely useful in convex geometry, where, however, it is always assumed that the movement preserves convexity. It is easy to see that \diamond preserves convexity if and only if φ_\diamond is affine; cf. the proof of Theorem 6.20.

Corollary 6.7. *Let $H = u^\perp$, $u \in S^{n-1}$, be oriented with $u \in H^+$ and suppose that $\mathcal{E} = \mathcal{K}_n^n$ or \mathcal{K}^n . Suppose that $\diamond : \mathcal{E} \rightarrow \mathcal{L}^n$ is monotonic, measure preserving, respects H -cylinders, and maps balls to balls. Let $\varphi_\diamond : \mathbb{R} \rightarrow \mathbb{R}$ be the function from Lemma 6.2. Then $\varphi_\diamond(t) = t, -t, |t|$, or $-|t|$, if and only if \diamond essentially equals Id , \dagger , \diamond_{P_H} , or $\diamond_{P_H}^\dagger$, respectively.*

Proof. If \diamond is given, it is easy to check that φ_\diamond has the appropriate form, by applying \diamond to balls and using (50). The converse follows directly from (63). Indeed, we need only consider $K \in \mathcal{K}_n^n$, for otherwise $\mathcal{H}^n(K) = 0$ and there is nothing to prove. Consider, for example, the case when $\varphi_\diamond = |t|$; the other cases are similar. By (63), for \mathcal{H}^{n-1} -almost all $x \in H$, we have

$$\begin{aligned} (\diamond K) \cap (H^\perp + x) &= (K \cap (H^\perp + x)) + (|t_x| - t_x)u \\ &= \begin{cases} K \cap (H^\perp + x), & \text{if } t_x \geq 0, \\ K^\dagger \cap (H^\perp + x), & \text{if } t_x < 0, \end{cases} \\ &= (\diamond_{P_H} K) \cap (H^\perp + x), \end{aligned}$$

up to a set of \mathcal{H}^1 -measure zero. This shows that \diamond essentially equals \diamond_{P_H} , as required. \square

Example 6.8. Let $H = u^\perp$, $u \in S^{n-1}$, be oriented with $u \in H^+$. Define $\diamond' : \mathcal{L}^n \rightarrow \mathcal{L}^n$ by $\diamond'E = E^+ \cup E^-$, where $E^+ = E \cap H^+$ and $E^- \subset H^-$ is given by

$$E^- \cap (H^\perp + x) = [x, x - \lambda u],$$

with $\lambda = \mathcal{H}^1(E \cap H^- \cap (H^\perp + x))$ for each $x \in H$. Thus $\diamond' = \text{Id}$ when applied to subsets of H^+ and \diamond' corresponds to Blaschke shaking [23, Note 2.4] with respect to H when applied to subsets of H^- .

Define $\diamond : \mathcal{L}^n \rightarrow \mathcal{L}^n$ by $\diamond = \diamond' \circ \diamond_{P_H}$. It is easy to see that \diamond' is monotonic, measure preserving, and respects H -cylinders, and hence \diamond also has these three properties.

We claim that $\diamond = \diamond_{P_H}$ on \mathcal{K}^n . To see this, let $K \in \mathcal{K}^n$. Then

$$(\diamond_{P_H} K) \cap H^- = (K \cap K^\dagger) \cap H^-.$$

As $K \cap K^\dagger$ is H -symmetric and convex, we have

$$((K \cap K^\dagger) \cap H^-) \cap (H^\perp + x) = [x, x - \lambda u],$$

where $\lambda = \mathcal{H}^1((K \cap K^\dagger) \cap H^- \cap (H^\perp + x))$ for $x \in H$. Thus, $\diamond K = \diamond'(\diamond_{P_H} K) = \diamond_{P_H} K$, proving the claim.

As a consequence, \diamond maps balls to balls and therefore satisfies all the hypotheses of Corollary 6.7 with $\varphi_\diamond(t) = |t|$, yet it is clear that \diamond is essentially different from \diamond_{P_H} . Indeed, Corollary 6.7 is false for maps $\diamond : \mathcal{K}_n^n \subset \mathcal{E} \rightarrow \mathcal{L}^n$ if \mathcal{E} contains an H -symmetric union of two disjoint balls, since if E is such a union, then $\diamond E \neq \diamond_{P_H} E$. \square

Lemma 6.9. *Let $H \in \mathcal{G}(n, n-1)$, let $\mathcal{K}_n^n \subset \mathcal{E} \subset \mathcal{L}^n$, and suppose that $\diamond : \mathcal{E} \rightarrow \mathcal{L}^n$ is measure preserving and perimeter preserving on convex bodies. Then \diamond maps balls to balls.*

Proof. Let $x \in \mathbb{R}^n$ and let $r > 0$. Our assumptions imply that $\diamond B(x, r)$ is a set of finite perimeter that has the same \mathcal{H}^n -measure and perimeter as $B(x, r)$. It follows from the isoperimetric inequality for sets of finite perimeter and its equality condition (see [14] or [39, Theorem 14.1]) that $\diamond B(x, r)$ is a ball, modulo a set of \mathcal{H}^n -measure zero. \square

Theorem 6.10. *Let $H = u^\perp$, $u \in S^{n-1}$, be oriented with $u \in H^+$ and suppose $\diamond : \mathcal{K}_n^n \rightarrow \mathcal{L}^n$ is monotonic, measure preserving, respects H -cylinders, and perimeter preserving on convex bodies. If φ_\diamond is the contraction defined in Lemma 6.2, then $|\varphi'_\diamond(t)| = 1$ for \mathcal{H}^1 -almost all $t \in \mathbb{R}$.*

Conversely, if $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is a contraction satisfying $|\varphi'(t)| = 1$ for \mathcal{H}^1 -almost all $t \in \mathbb{R}$, then (63) defines a map $\diamond : \mathcal{K}^n \rightarrow \mathcal{L}^n$ that is monotonic, measure preserving, respects H -cylinders, and perimeter preserving on convex bodies.

Proof. We may assume that $u = e_n$ and write $H^\perp = \langle e_n \rangle$ for the x_n -axis. If $x \in H = e_n^\perp$, write $x = (x_1, \dots, x_{n-1})$. If $t \in \mathbb{R}$ and $r > 0$, let $D(t, r) = B(te_1, r) \cap e_n^\perp$.

Suppose that $t > 0$. For $0 < r < t$, define

$$K(t, r) = \{x + x_n e_n : x \in D(t, r), 0 \leq x_n \leq 2x_1\}.$$

Then $K(t, r) \in \mathcal{K}_n^n$ and if $x \in D(t, r)$, then $x + x_1 e_n$ is the midpoint of $K(t, r) \cap (\langle e_n \rangle + x)$. Lemma 6.9 implies that \diamond maps balls to balls, so (63) holds for \diamond . Hence

$$(67) \quad \diamond K(t, r) = \{x + x_n e_n : x \in D(t, r), \varphi_\diamond(x_1) - x_1 \leq x_n \leq \varphi_\diamond(x_1) + x_1\},$$

essentially. Since φ_\diamond is Lipschitz by Lemma 6.2, $\diamond K(t, r)$ is a set of finite perimeter; see [2, Proposition 3.62]. From (67) (or see (63)), we have

$$(\diamond K(t, r)) \cap (\langle e_n \rangle + x) = (K(t, r) \cap (\langle e_n \rangle + x)) + (\varphi_\diamond(x_1) - x_1)e_n,$$

up to a set of \mathcal{H}^1 -measure zero, for \mathcal{H}^{n-1} -almost all $x \in D(t, r)$. Also, $\|\nabla(2x_1)\| = 2$, $\|\nabla 0\| = 0$, and $\|\nabla(\varphi_\diamond(x_1) \pm x_1)\| = |\varphi'_\diamond(x_1) \pm 1|$. As \diamond is perimeter preserving on convex bodies, $K(t, r)$ and $\diamond(K(t, r))$ have equal perimeters, so we obtain (see e.g. [20, p. 101])

$$\begin{aligned} & \int_{D(t, r)} \sqrt{1 + 2^2} dx + \int_{D(t, r)} \sqrt{1 + 0^2} dx \\ &= \int_{D(t, r)} \sqrt{1 + (\varphi'_\diamond(x_1) + 1)^2} dx + \int_{D(t, r)} \sqrt{1 + (\varphi'_\diamond(x_1) - 1)^2} dx. \end{aligned}$$

Dividing the previous equation by $\mathcal{H}^{n-1}(D(t, r))$ and taking the limit as $r \rightarrow 0$, Lebesgue's differentiation theorem (see e.g. [42, Theorem 8.8]) yields

$$\sqrt{5} + 1 = \sqrt{1 + (\varphi'_\diamond(t) + 1)^2} + \sqrt{1 + (\varphi'_\diamond(t) - 1)^2}$$

for \mathcal{H}^1 -almost all $t \in (0, \infty)$. It is easy to check that the only solutions of the previous equation are $\varphi'_\diamond(t) = \pm 1$ for \mathcal{H}^1 -almost all $t \in (0, \infty)$.

The above argument can be repeated with $t < 0$, $0 < r < -t$, and

$$K'(t, r) = \{x + t e_n : x \in D(t, r), -2x_1 \leq x_n \leq 0\}.$$

This yields that $\varphi'_\diamond(t) = \pm 1$ for \mathcal{H}^1 -almost all $t \in (-\infty, 0)$, so the first statement in the theorem is proved.

Conversely, suppose that φ is a contraction such that $|\varphi'(t)| = 1$ for $t \in \mathbb{R} \setminus N$, where $\mathcal{H}^1(N) = 0$. Then (63) defines a map $\diamond : \mathcal{K}^n \rightarrow \mathcal{L}^n$. Theorem 6.6 implies that \diamond is monotonic, measure preserving, and respects H -cylinders on \mathcal{K}_n^n , but it is clear that these properties hold on \mathcal{K}^n .

It remains to prove that \diamond preserves perimeter on convex bodies. Let $K \in \mathcal{K}_n^n$. We may assume without loss of generality that $\diamond : \mathcal{K}^n \rightarrow \mathcal{L}^n$ is defined by (63) for all $x \in H$ and without the exceptional sets of \mathcal{H}^1 -measure zero. Indeed, the difference, by Fubini's theorem, is a set of \mathcal{H}^n -measure zero, which does not change perimeter (see [39, Exercise 12.16]). Let

$$f^+(x) = \max_{x + t e_n \in K} t, \quad f^-(x) = \min_{x + t e_n \in K} t, \quad g^+(x) = \max_{x + t e_n \in \diamond K} t, \quad \text{and} \quad g^-(x) = \min_{x + t e_n \in \diamond K} t$$

be the functions from $K|H$ to \mathbb{R} whose graphs are the top and bottom parts of ∂K and $\partial(\diamond K)$. With the already established notation from Lemma 6.5, we have

$$(68) \quad t_x = (f^+(x) + f^-(x))/2$$

and

$$(69) \quad g^\pm(x) = \varphi(t_x) \pm (f^+(x) - f^-(x))/2.$$

Put $\Omega = \text{relint}(K|H)$. For $E \subset \mathbb{R}^n$, let

$$E_1 = (\partial E) \cap (((K|H) \setminus \Omega) + H^\perp) \quad \text{and} \quad E_2 = (\partial E) \cap (\Omega + H^\perp).$$

Clearly ∂K (or $\partial(\diamond K)$) is the disjoint union of K_1 and K_2 (or $(\diamond K)_1$ and $(\diamond K)_2$, respectively). We show below that

$$(70) \quad \mathcal{H}^{n-1}((\diamond K)_i) = \mathcal{H}^{n-1}(K_i)$$

for $i = 1, 2$. Assuming this is true, the proof is completed as follows. Since K is a convex body, it has Lipschitz boundary. The functions $\pm f^\pm$ are convex and hence locally Lipschitz on Ω (see [43, Theorem 1.5.3]), so it follows from (68) and (69) that g^\pm are also locally Lipschitz on Ω . From this and the fact that φ is Lipschitz, it follows that $\diamond K$ also has Lipschitz boundary. Note that $\partial K = \partial(\text{int } K)$, $\partial(\diamond K) = \partial(\text{int } (\diamond K))$, and $\mathcal{H}^{n-1}(\partial(\diamond K)) = \mathcal{H}^{n-1}(\partial K) < \infty$ by (70). Then it follows from [2, Equation (3.63)] and [2, Proposition 3.62)], applied to $\text{int } K$ and $\text{int } (\diamond K)$, and (70), that

$$S(\diamond K) = \mathcal{H}^{n-1}(\partial(\diamond K)) = \mathcal{H}^{n-1}(\partial K) = S(K),$$

as required.

To prove (70) for $i = 1$, note that because $(K|H) \setminus \Omega$ is the boundary of a convex body in H , each $x \in (K|H) \setminus \Omega$ has a relative neighborhood $U \subset ((K|H) \setminus \Omega)$ such that there is a Lipschitz bijection from $U + H^\perp$ to \mathbb{R}^{n-1} , with Lipschitz inverse, that maps vertical lines isometrically to vertical lines. Then the desired result follows directly from the area formula [20, Theorem 1, p. 96], the fact that

$$\mathcal{H}^1((\diamond K) \cap (H^\perp + x)) = \mathcal{H}^1(K \cap (H^\perp + x))$$

for all $x \in (K|H) \setminus \Omega$, and Fubini's theorem in \mathbb{R}^{n-1} .

It therefore remains to prove that $\mathcal{H}^{n-1}((\diamond K)_2) = \mathcal{H}^{n-1}(K_2)$. To this end, let $M = \{x \in \Omega : t_x \in N\}$. Applying the coarea formula [20, p. 112] to the locally Lipschitz function $x \mapsto t_x$ gives

$$\int_M |\nabla t_x| dx = \int_N \mathcal{H}^{n-1}(\{x : t_x = s\}) ds = 0,$$

as $\mathcal{H}^1(N) = 0$. Therefore $\mathcal{H}^{n-1}(\{x \in M : \nabla t_x \neq o\}) = 0$. As all the functions f^\pm and g^\pm are locally Lipschitz on Ω , their gradients exist for \mathcal{H}^{n-1} -almost all $x \in \Omega$. Using (68) and (69), a direct calculation shows that

$$(71) \quad \nabla g^\pm(x) = \begin{cases} \nabla f^\pm(x), & \text{if } \varphi'(t_x) = 1 \text{ and } \nabla t_x \neq o, \\ -\nabla f^\mp(x), & \text{if } \varphi'(t_x) = -1 \text{ and } \nabla t_x \neq o. \end{cases}$$

Using the fact that φ is a contraction, for $x \in \Omega$ with $\nabla t_x = 0$ and $h \in H$ with $\|h\|$ sufficiently small, we have

$$\frac{|\varphi(t_{x+h}) - \varphi(t_x)|}{\|h\|} \leq \frac{|t_{x+h} - t_x|}{\|h\|} = \frac{|t_{x+h} - t_x - \langle \nabla t_x, h \rangle|}{\|h\|} \rightarrow 0$$

as $h \rightarrow 0$, implying that $\nabla \varphi(t_x)$ exists and is zero. Then, taking gradients in (68) and (69), we obtain

$$(72) \quad \nabla g^\pm(x) = \pm \nabla f^+(x) = \mp \nabla f^-(x), \quad \text{if } \nabla t_x = 0,$$

for \mathcal{H}^{n-1} -almost all $x \in \Omega$. Using (71), (72), and [20, p. 101], we get

$$\begin{aligned} \mathcal{H}^{n-1}((\diamond K)_2) &= \int_{\Omega} \left((1 + \|\nabla g^+(x)\|^2)^{1/2} + (1 + \|\nabla g^-(x)\|^2)^{1/2} \right) dx \\ &= \int_{\Omega} \left((1 + \|\nabla f^+(x)\|^2)^{1/2} + (1 + \|\nabla f^-(x)\|^2)^{1/2} \right) dx = \mathcal{H}^{n-1}(K_2), \end{aligned}$$

as required. \square

The equation $|\varphi'(t)| = 1$, on a given domain and usually stated with boundary conditions, is a special case of the *eikonal equation*; see, for example, [6, p. 47]. In addition to the four functions $\varphi(t) = t$, $-t$, $|t|$, and $-|t|$ in Corollary 6.7, there are infinitely many other solutions \mathcal{H}^1 -almost everywhere on \mathbb{R} , including the function in the following example.

Example 6.11. Let $H = u^\perp$, $u \in S^{n-1}$, be oriented with $u \in H^+$. Define the contraction $\varphi(t) = \min_{k \in \mathbb{Z}} |t - k|$ for $t \in \mathbb{R}$ and let $\diamond : \mathcal{K}^n \rightarrow \mathcal{L}^n$ be defined by (63). Then $|\varphi'(t)| = 1$ for \mathcal{H}^1 -almost all $t \in \mathbb{R}$. By Theorem 6.10, \diamond is monotonic, measure preserving, respects H -cylinders, and is perimeter preserving on convex bodies. By Lemma 6.9, \diamond maps balls to balls and hence, by Corollary 6.4, it is invariant on H -symmetric sets. However, \diamond is not essentially equal to Id , \dagger , \diamond_{P_H} , or $\diamond_{P_H}^\dagger$. \square

Remark 6.12. The functions appearing in Theorem 6.10, i.e., Lipschitz functions $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ such that $|\varphi'(t)| = \pm 1$ almost everywhere, may be non-differentiable at an uncountable number of points. In fact, given any set N such that $\mathcal{H}^1(N) = 0$, there is a function φ of this type such that φ' does not exist at any point in N . To see this, first note that by [25, Theorem 1], there is a Borel set E such that at each point in N , the upper Lebesgue density of E is 1 and the lower Lebesgue density of E is 0. (To construct E , let (G_k) be a sequence of open sets such that $N \subset G_{k+1} \subset G_k$, $\mathcal{H}^1(G_1) \leq 1$, and for each component C of G_k , $\mathcal{H}^1(C \cap G_{k+1}) \leq (1/k)\mathcal{H}^1(C)$. Then let $E = \cup_k (G_{2k-1} \setminus G_{2k})$.) Now define

$$\varphi(t) = \int_0^t (2(1_E(s)) - 1) ds$$

for $t \in \mathbb{R}$. Then φ is Lipschitz with $\varphi'(t) = 1$ or -1 at each density point of E or $\mathbb{R} \setminus E$, respectively, and hence almost everywhere by Lebesgue's density theorem, and it follows directly from the just-mentioned properties of E that φ' does not exist at any point in N .

Lemma 6.13. *Let $H = u^\perp$, $u \in S^{n-1}$, and let $\mathcal{K}_n^n \subset \mathcal{E} \subset \mathcal{L}^n$, where \mathcal{E} is closed under intersections with H -symmetric spherical cylinders. If $\diamond : \mathcal{E} \rightarrow \mathcal{L}^n$ is monotonic, measure preserving, and invariant on H -symmetric spherical cylinders, then \diamond respects H -cylinders.*

Proof. In the proof, we ignore sets of \mathcal{H}^n -measure zero. If $r > 0$ and $x \in H$, let $C(x, r)$ be the infinite spherical cylinder of radius r and axis $H^\perp + x$. Suppose that $E \in \mathcal{E}$ and $E \subset C(x, r)$. Since $E \in \mathcal{L}^n$, we can choose $s_m > 0$, $m \in \mathbb{N}$, such that if $C_m = (C|H) + s_m[-u, u]$ and $E_m = E \cap C_m$, then $\mathcal{H}^n(E \setminus E_m) \leq 1/m$. As $E_m \in \mathcal{E}$, \diamond is monotonic, and C_m is an H -symmetric spherical cylinder, we have $\diamond E_m \subset \diamond C_m = C_m$ and hence $F = \cup_m \diamond E_m \subset C(x, r)$. Also, $\diamond E_m \subset \diamond E$ for $m \in \mathbb{N}$, implying that $F \subset \diamond E$. Now

$$\mathcal{H}^n(F) \geq \mathcal{H}^n(\diamond E_m) = \mathcal{H}^n(E_m) \geq \mathcal{H}^n(E) - 1/m = \mathcal{H}^n(\diamond E) - 1/m$$

for each m , so $\mathcal{H}^n(F) \geq \mathcal{H}^n(\diamond E)$. Thus $\diamond E = F \subset C(x, r)$, as required. \square

All seven properties of maps $\diamond : \mathcal{E} \subset \mathcal{L}^n \rightarrow \mathcal{L}^n$ listed in Section 3 are shared by \diamond_{P_H} and the map \diamond of Example 6.11, so a further property is needed to distinguish \diamond_{P_H} . When \mathcal{E} contains H -symmetric unions of two disjoint balls, we can take inspiration from Example 6.8 and use the new property in the next result. Note that the property is much weaker than invariance on all H -symmetric sets.

Lemma 6.14. *Let $H = u^\perp$, $u \in S^{n-1}$, and let $\mathcal{K}_n^n \subset \mathcal{E} \subset \mathcal{L}^n$, where \mathcal{E} is closed under intersections with H -symmetric spherical cylinders. If $\diamond : \mathcal{E} \rightarrow \mathcal{L}^n$ is monotonic, measure preserving, and defined and invariant on H -symmetric unions of two disjoint balls, then \diamond is invariant on H -symmetric spherical cylinders and hence respects H -cylinders.*

Proof. Let $C = (B(x, r) \cap H) + s(B^n \cap H^\perp)$, where $r, s > 0$, be an H -symmetric spherical cylinder. Let $A = \text{int}(C \setminus H)$. Choose H -symmetric unions U_m of two disjoint balls, $m \in \mathbb{N}$, such that $A = \cup_m U_m$. By monotonicity and invariance on H -symmetric unions of two disjoint balls,

$$A = \cup_m U_m = \cup_m \diamond U_m \subset \diamond C.$$

Therefore, $C \subset \diamond C$, essentially. The measure-preserving property of \diamond now implies that $\diamond C = C$, essentially. Thus \diamond is invariant on H -symmetric spherical cylinders and hence respects H -cylinders by Lemma 6.13. \square

Lemma 6.15. *Let $H = u^\perp$, $u \in S^{n-1}$, be oriented with $u \in H^+$ and let $\mathcal{E} = \mathcal{C}^n$ or \mathcal{L}^n . Suppose that $\diamond : \mathcal{E} \rightarrow \mathcal{L}^n$ is monotonic, measure preserving, maps balls to balls, and invariant on H -symmetric unions of two disjoint balls. Let $\varphi_\diamond : \mathbb{R} \rightarrow \mathbb{R}$ be the function from Lemma 6.2. Then $\varphi_\diamond(t) = t, -t, |t|$, or $-|t|$, if and only if \diamond essentially equals Id , \dagger , \diamond_{P_H} , or $\diamond_{P_H}^\dagger$, respectively.*

Proof. By Lemma 6.14, our assumptions on \diamond imply that it also respects H -cylinders, allowing previous results to be applied.

If \diamond is given, it is easy to check that φ_\diamond has the appropriate form, by applying \diamond to balls and using (50).

It will suffice show that if $\varphi_\diamond = |t|$, then $\diamond = \diamond_{P_H}$. Indeed, the case when $\varphi_\diamond = -|t|$ then follows by applying the previous case to \diamond^\dagger . The cases when $\varphi_\diamond = \pm t$ are even simpler, not requiring the assumption that \diamond is invariant on H -symmetric unions of two disjoint balls, but only that \diamond respects H -cylinders.

In the rest of the proof, we shall ignore sets of \mathcal{H}^n -measure zero.

Assume initially that $E \in \mathcal{E}$ is a countable union of balls, $E = \bigcup_{j=1}^\infty B_j$. Let I^- be the set of all indices j for which the center of B_j lies in H^- , and put $I^+ = \mathbb{N} \setminus I^-$. By (50), $\diamond B_j = B_j$ for $j \in I^+$. Since \diamond is monotonic, we obtain

$$(73) \quad \bigcup_{j \in I^+} B_j = \bigcup_{j \in I^+} \diamond B_j \subset \diamond E.$$

For $j \in I^-$ we have $\diamond B_j = B_j^\dagger$ by (50), so a similar argument gives

$$(74) \quad (\bigcup_{j \in I^-} B_j)^\dagger \subset \diamond E.$$

We claim that

$$(75) \quad E \cap E^\dagger \subset \diamond E.$$

To see this, note that

$$E \cap E^\dagger = \bigcup_{j,k=1}^\infty (B_j \cap B_k^\dagger) = \bigcup_{j \leq k} E_{jk},$$

where $E_{jk} = (B_j \cap B_k^\dagger) \cup (B_k \cap B_j^\dagger)$ is an H -symmetric union of (at most) two compact convex sets. It will therefore suffice to show that $\text{int } E_{jk} \subset \diamond E$ for $j, k \in \mathbb{N}$. To this end, let C_1, C_2, \dots be balls such that

$$\text{int } (E_{jk} \cap H^+) = \bigcup_{i=1}^\infty C_i.$$

As E_{jk} is H -symmetric, this implies that

$$C_i \cup C_i^\dagger \subset (\text{int } E_{jk}) \setminus H \subset E \cap E^\dagger \subset E.$$

Hence,

$$\text{int } E_{jk} \subset \bigcup_{i=1}^\infty (C_i \cup C_i^\dagger) = \bigcup_{i=1}^\infty \diamond(C_i \cup C_i^\dagger) \subset \diamond E,$$

where the equality is justified as \diamond is invariant on H -symmetric unions of two disjoint balls, and the final containment follows by the monotonicity of \diamond . This proves (75).

By (73), (74), and (75), if we can prove the first containment in

$$(76) \quad \diamond_{P_H} E \subset (\bigcup_{j \in I^+} B_j) \cup (\bigcup_{j \in I^-} B_j)^\dagger \cup (E \cap E^\dagger) \subset \diamond E,$$

the measure-preserving properties of \diamond and \diamond_{P_H} will show that $\diamond E = \diamond_{P_H} E$. To prove the inclusion in question, fix $x \in (\diamond_{P_H} E) \setminus H^-$. Then $\{x, x^\dagger\} \cap E \neq \emptyset$ and there is a $j \in \mathbb{N}$ with $x \in B_j$ or $x \in B_j^\dagger$. Suppose that $x \in B_j$. If $j \in I^+$, the first inclusion in (76) holds trivially, and if $j \in I^-$ it holds as $x \in B_j \setminus H^- \subset B_j^\dagger$. Similar arguments can be used if $x \in B_j^\dagger$. When $x \in (\diamond_{P_H} E) \cap H^-$, we have $\{x, x^\dagger\} \subset E$, so $x \in E \cap E^\dagger$. Concluding, $\diamond E = \diamond_{P_H} E$ whenever $E \in \mathcal{E}$ is a countable union of balls. In particular, this is true if $E \in \mathcal{E}$ is an open set.

Let (E_m) be a decreasing sequence of sets in \mathcal{E} and let $E = \bigcap_m E_m \in \mathcal{E}$. We claim that if $\diamond E_m = \diamond_{P_H} E_m$ for $m \in \mathbb{N}$, then $\diamond E = \diamond_{P_H} E$. We first show that

$$(77) \quad \bigcap_m (\diamond_{P_H} E_m) = \diamond_{P_H} E.$$

To prove (77), observe that for $x \in H^-$ and $A \in \mathcal{L}^n$, we have $x \in \diamond_{P_H} A$ if and only if $\{x, x^\dagger\} \subset A$. Hence, when $x \in H^-$ we have

$$x \in \bigcap_m (\diamond_{P_H} E_m) \Leftrightarrow \{x, x^\dagger\} \subset E_m, \text{ for } m \in \mathbb{N} \Leftrightarrow \{x, x^\dagger\} \subset E \Leftrightarrow x \in \diamond_{P_H} E.$$

For $x \in H^+$, we have $x \in \diamond_{P_H} A$ if and only if $\{x, x^\dagger\} \cap A \neq \emptyset$. Therefore

$$x \in \bigcap_m (\diamond_{P_H} E_m) \Leftrightarrow \{x, x^\dagger\} \cap E_m \neq \emptyset, \text{ for } m \in \mathbb{N} \Leftrightarrow \{x, x^\dagger\} \cap E \neq \emptyset \Leftrightarrow x \in \diamond_{P_H} E,$$

where for the second equivalence, we used the fact that (E_m) is decreasing, implying that if one of x or x^\dagger is not in E_m for some m , then the other must be in E_m for all m . This proves (77). From the monotonicity of \diamond , the assumption $\diamond E_m = \diamond_{P_H} E_m$ and (77), we obtain

$$\diamond E \subset \bigcap_m \diamond E_m = \bigcap_m \diamond_{P_H} E_m = \diamond_{P_H} E,$$

and the measure-preserving properties of \diamond and \diamond_{P_H} show that $\diamond E = \diamond_{P_H} E$. Concluding, the subclass of \mathcal{E} where \diamond and \diamond_{P_H} essentially coincide is closed under decreasing limits, if the intersection is contained in \mathcal{E} .

Let $\mathcal{E} = \mathcal{L}^n$. If $E \in \mathcal{E}$, there is a decreasing sequence (E_m) of open sets whose intersection is essentially E . By what was proved earlier, this concludes the proof for this case.

Now let $\mathcal{E} = \mathcal{C}^n$ and let $E \in \mathcal{E}$. By compactness, there is a set E_1 that contains E in its interior and is a finite union of balls with radius 1. Using compactness again, we can find a finite union E_2 of balls with radius at most $1/2$ containing E in its interior, such that $E_2 \subset E_1$. Continuing this way, a decreasing sequence (E_m) of finite unions of balls is constructed with $\bigcap_m E_m = E$. The first part of the proof shows that $\diamond E_m = \diamond_{P_H} E_m$, and the second part of the proof gives $\diamond E = \diamond_{P_H} E$. \square

Let \mathcal{U}^n be the family of countable unions of balls in \mathbb{R}^n . Suppose that $\mathcal{E} \subset \mathcal{L}^n$ is a class of measurable sets containing all balls and such that for each $E \in \mathcal{E}$, there is a decreasing sequence (E_m) of sets in \mathcal{U}^n whose intersection is essentially E . Then the proof of the previous lemma shows that it holds for the class \mathcal{E} .

Theorem 6.16. *Let $H = u^\perp$, $u \in S^{n-1}$, be oriented with $u \in H^+$, let $\mathcal{E} = \mathcal{C}^n$ or \mathcal{L}^n , and suppose that $\diamond : \mathcal{E} \rightarrow \mathcal{L}^n$ is monotonic, measure preserving, perimeter preserving on convex bodies, and invariant on H -symmetric unions of two disjoint balls. Then \diamond essentially equals Id , \dagger , \diamond_{P_H} , or $\diamond_{P_H}^\dagger$.*

Proof. By Lemmas 6.9 and 6.14, our assumptions imply that \diamond respects H -cylinders and maps balls to balls, so Theorem 6.6 implies that the restriction of \diamond to \mathcal{K}^n is determined by (63). Let $t_0 \neq 0$. For $0 < r < |t_0|$, the balls $B(\pm t_0 u, r)$ are disjoint. Since \diamond is monotonic and invariant on H -symmetric unions of two disjoint balls, either $\diamond B(t_0 u, r) = B(t_0 u, r)$ or $\diamond B(t_0 u, r) = B(-t_0 u, r)$. It follows from (50) that $\varphi_\diamond(t_0) = \pm t_0$. The continuity of φ_\diamond implies that $\varphi_\diamond(t) = t, -t, |t|$, or $-|t|$, so the desired conclusion is provided by Lemma 6.15. \square

The map $\diamond : \mathcal{L}^n \rightarrow \mathcal{L}^n$ in Example 6.8 can also be considered as a map $\diamond : \mathcal{C}^n \rightarrow \mathcal{C}^n$. We showed in Example 6.8 that $\diamond = \diamond_{P_H}$ on \mathcal{K}^n . This implies that \diamond is perimeter preserving on convex bodies. Consequently, \diamond has all the properties assumed in the previous theorem except that it is not invariant on H -symmetric unions of two disjoint balls, showing that the latter property cannot be replaced by the weaker assumption of respecting H -cylinders.

The following examples deal with the other assumptions in Theorem 6.16, where it is always assumed that $H = u^\perp$, $u \in S^{n-1}$.

Example 6.17. Given $E \in \mathcal{L}^n$ with $\mathcal{H}^n(E) > 0$, let

$$c_E = \frac{1}{\mathcal{H}^n(E)} \int_E x \, dx$$

be the center of gravity of E and let $\diamond E$ be the reflection of E in the hyperplane $H + c_E$. Then $\diamond : \mathcal{E} \rightarrow \mathcal{E}$ for $\mathcal{E} = \mathcal{C}^n$ or \mathcal{L}^n is measure preserving, perimeter preserving on convex bodies, and invariant on H -symmetric unions of two disjoint balls. (Indeed, \diamond is clearly invariant on all H -symmetric sets.) It is not monotonic, as can be seen by considering the double cone $\text{conv}([-u, u] \cup (B^n \cap H))$ and its subset, the cone $\text{conv}(\{u\} \cup (B^n \cap H))$. \square

Example 6.18. Let $\mathcal{E} = \mathcal{C}^n$ or \mathcal{L}^n and define $\diamond : \mathcal{E} \rightarrow \mathcal{E}$ by $\diamond E = \text{cl}(\text{int } E)$. Then \diamond is monotonic, perimeter preserving on convex bodies, and invariant on H -symmetric unions of two disjoint balls, but not measure preserving. \square

Example 6.19. For $x \in \mathbb{R}^n$, let

$$F(x) = \begin{cases} x, & \text{if } d(x, H) > 1, \\ x^\dagger, & \text{if } d(x, H) \leq 1 \end{cases}$$

and define $\diamond : \mathcal{E} \rightarrow \mathcal{L}^n$ for $\mathcal{E} = \mathcal{C}^n$ or \mathcal{L}^n by $\diamond E = F(E)$. Then \diamond is monotonic, measure preserving, and invariant on H -symmetric unions of two disjoint balls (indeed, on all H -symmetric sets), but does not preserve perimeter on convex bodies. \square

For maps $\diamond : \mathcal{K}^n \rightarrow \mathcal{L}^n$, invariance on H -symmetric unions of two disjoint balls is not available. We therefore resort to a different and rather strong condition; we say that \diamond is *convexity preserving away from H* if $\diamond K$ is essentially convex (that is, $\diamond K$ coincides with a convex set up to a set of \mathcal{H}^n -measure zero) for all $K \in \mathcal{K}^n$ with $K \cap H = \emptyset$.

Theorem 6.20. *Let $H = u^\perp$, $u \in S^{n-1}$, be oriented with $u \in H^+$ and let $\diamond : \mathcal{K}_n^n \rightarrow \mathcal{L}^n$ be monotonic, measure preserving, invariant on H -symmetric sets, perimeter preserving on convex bodies, and convexity preserving away from H . Then \diamond essentially equals Id , \dagger , \diamond_{P_H} , or $\diamond_{P_H}^\dagger$.*

Proof. We may assume that $u = e_n$. Let $C = \{0\} \times [0, 1]^{n-2} \times \{0\}$ be the unit cube in $H \cap e_1^\perp$. For $r, t > 0$, define

$$K(t, r) = \{x + x_n e_n : x \in C + [0, te_1], r \leq x_n \leq r + 2x_1\}.$$

Then $K(t, r) \in \mathcal{K}_n^n$ is disjoint from H and if $x \in C + [o, te_1]$, then $x + (r + x_1)e_n$ is the midpoint of $K(t, r) \cap (\langle e_n \rangle + x)$. From Theorem 6.10, we know that (63) holds, where $|\varphi'_\diamond(t)| = 1$ for \mathcal{H}^1 -almost all t . Then

$$(78) \quad \diamond K(t, r) = \{x + x_n e_n : x \in C + [o, te_1], \varphi_\diamond(r + x_1) - x_1 \leq x_n \leq \varphi_\diamond(r + x_1) + x_1\},$$

essentially. Since \diamond is convexity preserving away from H , there is a convex set $L = L(t, r)$ essentially equal to $\diamond K(t, r)$. As the boundary of a convex set has \mathcal{H}^n -measure zero, we may assume that L is closed. The sets $M_1 = C + \varphi_\diamond(r)e_n$ and $M_2 = C + [(\varphi_\diamond(r + t) - t)e_n, (\varphi_\diamond(r + t) + t)e_n] + te_1$ are contained in L , because any point in either set can be approximated by points in the right side of (78) that are also points in L . As L is convex, $M = \text{conv}(M_1 \cup M_2) \subset L$. The wedge M has the same volume as $K(t, r)$, so the measure-preserving property of \diamond yields $\diamond K(t, r) = L = M$, up to sets of \mathcal{H}^n -measure zero. Comparing the set in (78) with M and using the continuity of φ_\diamond , we obtain

$$\varphi_\diamond(r + x_1) = \left(1 - \frac{x_1}{t}\right) \varphi_\diamond(r) + \frac{x_1}{t} \varphi_\diamond(r + t)$$

for $0 \leq x_1 \leq t$. Letting $r \rightarrow 0+$, we conclude that φ_\diamond is affine on $[0, t]$ for $t > 0$. As \diamond is monotonic and invariant on H -symmetric sets, it respects H -cylinders, and our assumptions and Lemma 6.9 ensure that it also maps balls to balls. Therefore, by Corollary 6.4, $\varphi_\diamond(0) = 0$ and so φ_\diamond is linear. As $|\varphi'_\diamond| = 1$ almost everywhere, this implies that $\varphi_\diamond = \pm \text{Id}$ on $[0, \infty)$. Similar arguments applied to

$$K'(t, r) = \{x + x_n e_n : x \in C + [te_1, o], r - 2x_1 \leq x_n \leq r\},$$

where $t, r < 0$, show that $\varphi_\diamond = \pm \text{Id}$ on $(-\infty, 0]$ and hence that $\varphi_\diamond(t) = t, -t, |t|$, or $-|t|$. The proof is completed by Corollary 6.7. \square

Example 6.17 shows that the assumption that \diamond is monotonic cannot be dropped in the previous theorem. If $o \neq x_0 \in H$, the map $\diamond K = K + x_0$ has all the properties in Theorem 6.20 except invariance on H -symmetric sets. If we define \diamond via (63), where $\varphi_\diamond(t) = t/2$, then it is easily checked that \diamond has all the properties except that it is not perimeter preserving. Example 6.11 shows that the assumption that \diamond is convexity preserving away from H cannot be omitted.

We do not have an example showing that the measure-preserving assumption is necessary, and this relates to an open problem stated as a variant of [7, Problem 11.1], namely, does there exist a map from the convex bodies in \mathbb{R}^n to those that are symmetric with respect to a fixed hyperplane H that is monotonic, invariant on H -symmetric sets, and perimeter-preserving? If such a map existed, it would provide the required example.

Theorem 6.21. *Let $H = u^\perp$, $u \in S^{n-1}$, be oriented with $u \in H^+$. Let $X = \mathcal{S}(\mathbb{R}^n)$ or $\mathcal{V}(\mathbb{R}^n)$ and suppose that $T : X \rightarrow X$ is a rearrangement. If the induced map \diamond_T defined by (13) is perimeter preserving on convex bodies and invariant on H -symmetric unions of two disjoint balls, then T essentially equals Id , \dagger , P_H , or P_H^\dagger .*

Proof. By Lemma 4.5 and Theorem 4.8(i), \diamond_T is well defined, monotonic, and measure preserving. Together with our assumptions on \diamond_T , we can apply Theorem 6.16 with $\mathcal{E} = \mathcal{L}^n$ to

conclude that \diamond_T essentially equals Id , \dagger , \diamond_{P_H} , or $\diamond_{P_H}^\dagger$. The proof is completed by Theorem 4.8(ii). \square

We now address the question of finding versions of the previous theorem for maps $T : X \rightarrow X$, where $X = \mathcal{M}(\mathbb{R}^n)$ or $\mathcal{M}_+(\mathbb{R}^n)$.

Theorem 6.22. *Let $H = u^\perp$, $u \in S^{n-1}$, be oriented with $u \in H^+$. Let $X = \mathcal{M}(\mathbb{R}^n)$ (or $X = \mathcal{M}_+(\mathbb{R}^n)$) and let $T : X \rightarrow X$ be a rearrangement. If the induced map \diamond_T defined by (13) is perimeter preserving on convex bodies and invariant on H -symmetric unions of two disjoint balls, then the restriction of T to $\mathcal{S}(\mathbb{R}^n)$ (or $\mathcal{V}(\mathbb{R}^n)$, respectively) essentially equals Id , \dagger , P_H , or P_H^\dagger .*

Proof. If $X = \mathcal{M}(\mathbb{R}^n)$ (or $X = \mathcal{M}_+(\mathbb{R}^n)$) and $T : X \rightarrow X$ is a rearrangement, then $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$ (or $T : \mathcal{V}(\mathbb{R}^n) \rightarrow \mathcal{V}(\mathbb{R}^n)$, respectively). This follows from Lemma 4.1(ii) (or the definitions of $\mathcal{V}(\mathbb{R}^n)$ and equimeasurability, respectively). Since the restricted maps satisfy the hypotheses of Theorem 6.21, the result follows. \square

Example 4.12 shows that the conclusion of Theorem 6.22 cannot be drawn for the unrestricted maps $T : \mathcal{M}(\mathbb{R}^n) \rightarrow \mathcal{M}(\mathbb{R}^n)$ or $T : \mathcal{M}_+(\mathbb{R}^n) \rightarrow \mathcal{M}_+(\mathbb{R}^n)$.

7. APPENDIX

The purpose here is to compare our approach to rearrangements on $\mathcal{S}(\mathbb{R}^n)$ in Section 4 with that of [8] and [51]. Recall that we begin with a rearrangement $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$ and show in Theorem 4.8(ii) that T is essentially determined by the associated measure-preserving and monotonic map $\diamond_T : \mathcal{L}^n \rightarrow \mathcal{L}^n$, defined by (13), via the formula

$$(79) \quad Tf(x) = \max \{ \sup \{ t \in \mathbb{Q}, t > \text{ess inf } f : x \in \diamond_T \{ z : f(z) \geq t \} \}, \text{ess inf } f \},$$

which holds for \mathcal{H}^n -almost all $x \in \mathbb{R}^n$, or alternatively via (27).

Brock and Solynin [8, Section 3] and Van Shaftingen and Willem [51] reverse the procedure, starting with a set transformation \diamond and defining function transformations. The latter paper, particularly, allows other possibilities, but we may focus on the case when $\diamond : \mathcal{L}^n \rightarrow \mathcal{L}^n$ is measure preserving and *pointwise monotonic* (meaning that if $K \subset L$, the containment $\diamond K \subset \diamond L$ must hold everywhere and not just almost everywhere). Consider defining a function T on $\mathcal{S}(\mathbb{R}^n)$ by

$$(80) \quad Tf(x) = \max \{ \sup \{ t > \inf f : x \in \diamond \{ z : f(z) > t \} \}, \inf f \},$$

for $x \in \mathbb{R}^n$. The formula (80) is equivalent to [8, Equation (3.1), p. 1762], where f is assumed continuous and it is shown that $Tf \in \mathcal{S}(\mathbb{R}^n)$. For general $f \in \mathcal{S}(\mathbb{R}^n)$, Brock and Solynin suggest replacing the supremum in (80) by the essential supremum and then claim that $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$ is a rearrangement. In [51], the authors work with *admissible functions*, which in our context is equivalent to demanding that $\inf f = \text{ess inf } f$. In [51, Definition 4], the map T defined by (80) is considered for admissible f and denoted by \check{S} , as well as a map

\overline{S} where in (80), $\{z : f(z) > t\}$ is replaced by $\{z : f(z) \geq t\}$. Among other results, it is stated in [51, Propositions 1–3] that under the above assumptions on \diamond , $T = \overline{S} = \check{S}$ and the formula

$$(81) \quad Tf(x) = \max\{\sup\{t > \text{ess inf } f : x \in \diamond\{z : f(z) \geq t\}\}, \text{ess inf } f\},$$

for $x \in \mathbb{R}^n$, defines a rearrangement $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$.

Since the definition (81) appears to be a thoroughly measure-theoretic one, it is natural to ask whether the pointwise monotonicity of \diamond is really required, or whether it could be replaced by monotonicity in our sense. The following example shows that this is not the case.

Example 7.1. Let $n = 1$ and for $A \in \mathcal{L}^1$, define

$$\diamond A = \begin{cases} A \cup \{\text{ess sup } A + 1\}, & \text{if } \text{ess sup } A < \infty, \\ A, & \text{otherwise.} \end{cases}$$

Then \diamond is essentially the identity on \mathcal{L}^1 and thus measure preserving and monotonic, but it is not pointwise monotonic. For the function $f(x) = -|x|$ in $\mathcal{S}(\mathbb{R}^n)$ and $t \leq 0$,

$$\diamond\{f \geq t\} = \diamond[t, -t] = [t, -t] \cup \{1 - t\},$$

so by (81), we have

$$\begin{aligned} Tf(x) &= \sup\{t \leq 0 : x \in [t, -t] \cup \{1 - t\}\} \\ &= \sup\{t \leq 0 : t \leq -|x| \text{ or } t = 1 - x\} = \begin{cases} 1 - x & \text{if } x \geq 1, \\ -|x|, & \text{otherwise.} \end{cases} \end{aligned}$$

Hence, T is not equimeasurable and therefore not a rearrangement. \square

One may wonder whether an alternative definition, namely,

$$(82) \quad Tf(x) = \max\{\text{ess sup}\{t > \text{ess inf } f : x \in \diamond\{z : f(z) \geq t\}\}, \text{ess inf } f\},$$

for $x \in \mathbb{R}^n$, would allow the pointwise monotonicity assumption on \diamond to be weakened, but the following example shows that it is consistent with ZFC that this is also not true.

Example 7.2. Let $n = 1$. Assuming the continuum hypothesis CH, Sierpiński [44] constructed a set $S \subset [0, 1]^2$ such that for $t \in [0, 1]$, the horizontal section $S_t = \{x \in [0, 1] : (x, t) \in S\}$ is countable and for $x \in [0, 1]$, the vertical section $S_x = \{t \in [0, 1] : (x, t) \in S\}$ is such that $[0, 1] \setminus S_x$ is countable. For $A \in \mathcal{L}^1$, define

$$\diamond A = \begin{cases} A \cup S_t, & \text{if } A = [-t, 0] \text{ for some } t \in [0, 1], \\ A, & \text{otherwise.} \end{cases}$$

Then \diamond is essentially the identity on \mathcal{L}^1 and thus measure preserving and monotonic, but it is not pointwise monotonic. Put

$$f(x) = \begin{cases} 1 + x, & \text{if } x \leq 0, \\ -x, & \text{otherwise.} \end{cases}$$

Then $f \in \mathcal{S}(\mathbb{R}^n)$ and direct calculation using (82) shows that

$$Tf(x) = \begin{cases} 1, & \text{if } x \in [0, 1], \\ f(x), & \text{otherwise,} \end{cases}$$

so T is not equimeasurable and hence not a rearrangement. \square

It would suffice in the previous example if the sections of S satisfy $\mathcal{H}^1(S_t) = 0$ for all $t \in [0, 1]$ and $\mathcal{H}^1(S_x) = 1$ for all $x \in [0, 1]$. The existence of such an S can be proved using Martin's Axiom MA and is therefore consistent with the negation of CH, while it is also consistent with ZFC that no such set exists. See [34, p. 673] for these and other related remarks.

The supremum over \mathbb{Q} in (79) cannot be replaced by the supremum over \mathbb{R} , i.e., the formula in Theorem 4.8(ii) cannot be replaced by (81). Indeed, let $n = 1$ and for $f \in \mathcal{S}(\mathbb{R})$, define

$$Tf = \begin{cases} 1_{A \cup \{\text{ess sup } A + 1\}}, & \text{if } f = 1_A, \text{ where } A \in \mathcal{L}^1 \text{ and } \text{ess sup } A < \infty, \\ f, & \text{otherwise.} \end{cases}$$

Then T is essentially the identity and therefore a rearrangement on $\mathcal{S}(\mathbb{R})$, so by Theorem 4.8(ii), (79) holds. From its definition (13), we see that \diamond_T is the map \diamond from Example 7.1, so it follows from that example that if T satisfied (81), it could not be equimeasurable, a contradiction. In a similar way, using Example 7.2 instead of Example 7.1, we see that it is consistent with ZFC that the supremum over \mathbb{Q} in (79) cannot be replaced by the essential supremum over \mathbb{R} .

To the best of our knowledge, our result in Section 4 that every rearrangement $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$ arises from the map $\diamond_T : \mathcal{L}^n \rightarrow \mathcal{L}^n$ defined by (13) has not been proved before. The question of when a function transformation arises from a set transformation is addressed in [51, Proposition 4]. This result appears to be based on [48, Proposition 2.4.1], where we find the statement and proof clearer. Restricting to our setting, it states that if an otherwise arbitrary $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$ is such that

$$(83) \quad \varphi \circ f \in \mathcal{S}(\mathbb{R}^n) \quad \text{and} \quad \varphi(Tf) = T(\varphi \circ f)$$

for $f \in \mathcal{S}(\mathbb{R}^n)$, whenever $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is right-continuous and increasing, then there is a $\diamond : \mathcal{L}^n \rightarrow \mathcal{L}^n$ such that T arises from \diamond via the formula (81). Note that T need not be a rearrangement for (83) to hold. For example, let $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$ be the pointwise map defined by (4) with $F(s, t) = \max\{s, t\}$, i.e., $Tf(x) = \max\{f(x), f(x^1)\}$ for $x \in \mathbb{R}^n$. Then T is monotonic but not equimeasurable, while it is easy to check that it satisfies (83). (However, Lemma 4.11 implies that an equimeasurable map satisfying (83) must be monotonic and hence a rearrangement.) On the other hand, [48, Proposition 2.4.1] and [51, Proposition 4] say nothing about rearrangements $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$ until it is known that (83) is true. This is just what we show holds, essentially, in Theorem 4.9, with a proof allowing the weakest possible continuity assumption on φ .

REFERENCES

- [1] L. V. Ahlfors, *Conformal Invariants: Topics in Geometric Function Theory*, McGraw-Hill Book Co., New York-Düsseldorf-Johannesburg, 1973.
- [2] L. Ambrosio, N. Fusco, and D. Pallara, *Functions of Bounded Variation and Free Discontinuity Problems*, Oxford University Press, New York, 2000.
- [3] A. Baernstein II, *Symmetrization in Analysis*, with D. Drasin and R. S. Laugesen, Cambridge University Press, Cambridge, 2019.
- [4] A. Baernstein II and B. A. Taylor, Spherical rearrangements, subharmonic functions, and $*$ -functions in n -space, *Duke Math. J.* **43** (1976), 245–268.
- [5] C. Bandle, *Isoperimetric Inequalities and Applications*, Pitman, Boston, MA, 1980.
- [6] M. Bardi and I. Capuzzo-Dolcetta, *Optimal Control and Viscosity Solutions of Hamilton-Jacobi-Bellman equations*, Birkhäuser, Boston, 1997.
- [7] G. Bianchi, R. J. Gardner, and P. Gronchi, Symmetrization in geometry, *Adv. Math.* **306** (2017), 51–88.
- [8] F. Brock and A. Y. Solynin, An approach to symmetrization via polarization, *Trans. Amer. Math. Soc.* **352** (2000), 1759–1796.
- [9] Y. D. Burago and V. A. Zalgaller, *Geometric inequalities*, Springer, New York, 1980.
- [10] A. Burchard, Cases of equality in the Riesz rearrangement inequality, *Ann. of Math. (2)* **143** (1996), 499–527.
- [11] A. Burchard and M. Fortier, Random polarizations, *Adv. Math.* **234** (2013), 550–573.
- [12] G. R. Burton, Rearrangements of functions, saddle points and uncountable families of steady configurations for a vortex, *Acta Math.* **163** (1989), 291–309.
- [13] K. M. Chong and N. M. Rice, *Equimeasurable Rearrangements of Functions*, Queen’s University, Kingston, Ont., 1971.
- [14] E. De Giorgi, Sulla proprietà isoperimetrica dell’ipersfera, nella classe degli insiemi aventi frontiera orientata di misura finita, *Atti Accad. Naz. Lincei. Mem. Cl. Sci. Fis. Mat. Nat. Sez. I* **5** (1958), 33–44.
- [15] J. Dekeyser and J. Van Schaftingen, Approximation of symmetrizations by Markov processes, *Indiana Univ. Math. J.* **66** (2017), 1145–1172.
- [16] R. J. Douglas, Rearrangements of functions with applications to meteorology and ideal fluid flow, in: *Large-Scale Atmosphere-Ocean Dynamics, Vol. I*, pp. 288–341, Cambridge University Press, Cambridge, 2002.
- [17] V. N. Dubinin, Transformation of condensers in space, *Dokl. Akad. Nauk SSSR* **296** (1987), 18–20; translation in *Soviet Math. Dokl.* **36** (1988), 217–219.
- [18] V. N. Dubinin, Capacities and geometric transformations of subsets in n -space, *Geom. Funct. Anal.* **3** (1993), 342–369.
- [19] V. N. Dubinin, *Condenser Capacities and Symmetrization in Geometric Function Theory*, Springer, Basel, 2014.
- [20] L. C. Evans and R. F. Gariepy, *Measure Theory and Fine Properties of Functions*, CRC Press, Boca Raton, FL, 1992.
- [21] L. E. Fraenkel, *An Introduction to Maximum Principles and Symmetry in Elliptic Problems*, Cambridge University Press, Cambridge, 2000.
- [22] R. J. Gardner, The Brunn-Minkowski inequality, *Bull. Amer. Math. Soc.* **39** (2002), 355–405.
- [23] R. J. Gardner, *Geometric Tomography*, second edition, Cambridge University Press, New York, 2006.
- [24] R. J. Gardner and M. Kiderlen, Operations between functions, *Comm. Anal. Geom.* **26** (2018), 787–855.
- [25] C. Goffman, On Lebesgue’s density theorem, *Proc. Amer. Math. Soc.* **1** (1950), 384–388.
- [26] P. M. Gruber, *Convex and Discrete Geometry*, Springer, Berlin, 2007.
- [27] C. Haberl and F. E. Schuster, General L_p affine isoperimetric inequalities, *J. Differential Geom.* **83** (2009), 1–26.
- [28] T. Hack and P. Pivovarov, Randomized Urysohn-type inequalities, [arXiv:1910.11654](https://arxiv.org/abs/1910.11654).

- [29] W. K. Hayman, *Multivalent Functions*, second edition, Cambridge University Press, Cambridge, 1994.
- [30] A. Henrot, *Extremum Problems for Eigenvalues of Elliptic Operators*, Birkhäuser, Basel, 2006.
- [31] B. Kawohl, *Rearrangements and Convexity of Level Sets in PDE*, Lecture Notes in Mathematics, 1150, Springer, Berlin, 1985.
- [32] S. Kesavan, *Symmetrization & Applications*, World Scientific Publishing, Hackensack, NJ, 2006.
- [33] S. G. Krantz and H. R. Parks, *The Geometry of Domains in Space*, Birkhäuser, Boston, MA, 1999.
- [34] M. Laczkovich, Two constructions of Sierpiński and some cardinal invariants of ideals, *Real Anal. Exchange* **24** (1998/99), 663–676.
- [35] E. H. Lieb and M. Loss, *Analysis*, second edition, American Mathematical Society, Providence, RI, 2001.
- [36] E. Lutwak, D. Yang, and G. Zhang, L_p affine isoperimetric inequalities, *J. Differential Geom.* **56** (2000), 111–132.
- [37] E. Lutwak, D. Yang, and G. Zhang, Orlicz centroid bodies, *J. Differential Geom.* **84** (2010), 365–387.
- [38] E. Lutwak and G. Zhang, Blaschke-Santaló inequalities, *J. Differential Geom.* **47** (1997), 1–16.
- [39] F. Maggi, *Sets of Finite Perimeter and Geometric Variational Problems. An Introduction to Geometric Measure Theory*, Cambridge University Press, Cambridge, 2012.
- [40] G. Pólya and G. Szegő, *Isoperimetric Inequalities in Mathematical Physics*, Princeton University Press, Princeton, NJ, 1951.
- [41] C. A. Rogers and G. C. Shephard, Some extremal problems for convex bodies, *Mathematika* **5** (1958), 93–102.
- [42] W. Rudin, *Real and Complex Analysis*, third edition, McGraw-Hill, New York, 1987.
- [43] R. Schneider, *Convex Bodies: The Brunn-Minkowski Theory*, second edition, Cambridge University Press, Cambridge, 2014.
- [44] W. Sierpiński, Sur un théorème équivalent à l’hypothèse du continu, *Bull. Intern. Acad. Sci. Cracovie, A* 1919, 1–3.
- [45] A. Y. Solynin, Continuous symmetrization via polarization, *Algebra i Analiz* **24** (2012), 157–222; translation in *St. Petersburg Math. J.* **24** (2013), 117–166.
- [46] D. W. Stroock, *Essentials of Integration Theory for Analysis*, Springer, New York, 2011.
- [47] G. Talenti, The art of rearranging, *Milan J. Math.* **84** (2016), 105–157.
- [48] J. Van Schaftingen; Réarrangements et problèmes elliptiques non linéaires. Thesis (Ph.D.)-Université Catholique de Louvain. 2002. 94 pp.
- [49] J. Van Schaftingen, Universal approximation of symmetrizations by polarizations, *Proc. Amer. Math. Soc.* **134** (2005), 177–186.
- [50] J. Van Schaftingen, Explicit approximation of the symmetric rearrangement by polarizations, *Arch. Math.* **93** (2009), 181–190.
- [51] J. Van Schaftingen and M. Willem, Set transformations, symmetrizations and isoperimetric inequalities, in: *Nonlinear analysis and applications to physical sciences*, pp. 135–152, Springer Italia, Milan, 2004.
- [52] V. Wolontis, Properties of conformal invariants, *Amer. J. Math.* **74** (1952), 587–606.

DIPARTIMENTO DI MATEMATICA E INFORMATICA “U. DINI”, UNIVERSITÀ DI FIRENZE, VIALE MORGAGNI 67/A, FIRENZE, ITALY I-50134

Email address: gabriele.bianchi@unifi.it

DEPARTMENT OF MATHEMATICS, WESTERN WASHINGTON UNIVERSITY, BELLINGHAM, WA 98225-9063, USA

Email address: Richard.Gardner@wwu.edu

DIPARTIMENTO DI MATEMATICA E INFORMATICA “U. DINI”, UNIVERSITÀ DI FIRENZE, PIAZZA Ghiberti 27, FIRENZE, ITALY I-50122

Email address: paolo.gronchi@unifi.it

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF AARHUS, NY MUNKEGADE, DK-8000
AARHUS C, DENMARK

Email address: `kiderlen@math.au.dk`