



Letter to Editors

Doxycycline or minocycline may be a viable treatment option against SARS-CoV-2



ARTICLE INFO

Keywords:

SARS-CoV-2

COVID-19

Doxycycline

Minocycline

Tetracyclines

As of June 21, coronavirus SARS-CoV-2 has infected almost 9 millions of people and the number of laboratory-confirmed Coronavirus Disease 2019 (COVID-19) deaths are 461,715, worldwide [1]. The virus causes acute respiratory distress syndrome (ARDS) and, eventually, death from multiple organ failure, in a non-neglectable portion of patients [2]. As such, COVID-19 is an international public health emergency. Although some recent reports suggested the efficacy of diverse agents, there is currently no standard therapy for COVID-19.

Several preclinical studies showed that doxycycline (DOX) and minocycline (MIN), semi-synthetic tetracyclines frequently used in clinical practice against a variety of infective agents and well tolerated, are also effective against some RNA viruses [3,4]. It is commonly recognized that tetracyclines act selectively inhibiting microbial protein synthesis by binding to the 16S rRNA [5]. Recent evidence suggested that the precise site of interaction between DOX/MIN and 16S rRNA as well as other cellular RNA molecules could be double-stranded RNAs (dsRNAs) [6]. The dsRNA-binding mechanism could explain the tetracyclines broad antimicrobial and antiviral properties [6]. Indeed, rRNAs form numerous double-stranded structures that are essential to protein synthesis. Furthermore, dsRNAs have been observed as intermediates of the viral replication of positive-stranded viruses, including SARS coronaviruses [7,8]. SARS-CoV-2 is a positive-stranded RNA virus and, interestingly, DOX/MIN antiviral efficacy was hitherto reported mostly against positive-sense RNA viruses [3,4]. Besides, replication of RNA viruses occurs in the host cell cytoplasm, where Mg²⁺ concentration is most favorable to DOX/MIN binding [5]. As such, DOX/MIN may be effective also against SARS-CoV-2 infection. In this regard, accumulating evidence shows that SARS is associated with an aberrant induction of inflammatory cytokines/chemokines, mostly activated by dsRNA intermediates of viral replication [9,10]. In particular, the robust viral replication and delayed IFN- γ signaling accompanying the initial steps of SARS seem to be consequence of the coronavirus ability to initially evade the host dsRNA-sensors [10,11]. Therefore, it is plausible that early administration of dsRNA-binding DOX/MIN may reduce the risk of SARS. This hypothesis is strengthened by the fact that both DOX and MIN are known to have relevant anti-inflammatory properties [12]. In detail, the tetracyclines inhibit pro-inflammatory cytokines and matrix metalloproteinases that play a key role in coronavirus acute infection and are involved in chemokine activation and

in tissue destruction, such as perivascular basement membrane and parenchymal extracellular matrix, thereby facilitating inflammatory cells infiltration [12,13]. Of note, also this immunomodulatory effect seems to be dsRNA-mediated [5]. Hence, DOX/MIN may inhibit both the viral replication and the host exuberant inflammatory response. Taken together, all evidence suggests that DOX/MIN may be active against SARS-CoV-2 and clinical trials investigating these drugs are urgently warranted.

Conflict of Interest

All authors report no conflict of interest.

Role of funding Source

There was no funding source.

References

- [1] World Health Organization: Coronavirus disease 2019 (COVID-19) Situation Report – 153. <https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200621-covid-19-sitrep-153-date-june-21-2020.pdf> Date: June 21, 2020 Date accessed: June 22, 2020.
- [2] Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in china: summary of a report of 72 314 cases from the Chinese center for disease control and prevention. *JAMA* 2020;323(13):1239. <https://doi.org/10.1001/jama.2020.2648>. [Epub ahead of print].
- [3] Michaelis M, Kleinschmidt MC, Doerr HW, Cinatl Jr. J. Minocycline inhibits West Nile virus replication and apoptosis in human neuronal cells. *J Antimicrob Chemother* 2007;2007(60):981–6.
- [4] Rothan Hussin A, Mohamed Zulqarnain, Paydar Mohammadjavad, Rahman Noorsaadah Abd, Yusof Rohana. Inhibitory effect of doxycycline against dengue virus replication in vitro. *Arch Virol* 2014;159:711–8.
- [5] Chukwudi Chinwe U. rRNA binding sites and the molecular mechanism of action of the tetracyclines. *Antimicrob Agents Chemother* 2016;60:4433–41.
- [6] Chinwe U, Chukwudi CU, Good Liam. Interaction of the tetracyclines with double-stranded RNAs of random base sequence: new perspectives on the target and mechanism of action. *J Antibiot (Tokyo)* 2016;69:622–30.
- [7] Weber Friedemann, Wagner Valentina, Rasmussen Simon B, Hartmann Rune, Paludan Søren R. Double-stranded RNA Is produced by positive-strand RNA viruses and DNA viruses but not in detectable amounts by negative-strand RNA viruses. *J Virol* 2006;80:5059–64.
- [8] Knoop Kévin, Kikkert Marjolein, van den Worm Sjoerd HE, Zevenhoven-Dobbe Jessica C, van der Meer Yvonne, Koster Abraham J, Mieke Mommaas A, Snijder Eric J. SARS-coronavirus replication is supported by a reticulovesicular network of modified endoplasmic reticulum. *PLoS Biol* 2008;6:e226.

<https://doi.org/10.1016/j.mehy.2020.110054>

Received 22 June 2020; Accepted 26 June 2020

0306-9877/ © 2020 Elsevier Ltd. All rights reserved.

- [9] Ding Shou-Wei, Olivier Voinnet O. Antiviral immunity directed by small RNAs. *Cell* 2007;130:413–26.
- [10] Cheung Chung Y, Poon Leo LM, Ng Iris HY, Luk Winsie, Sia Sin-Fun, Wu Mavis HS, Chan Kwok-Hung, Yuen Kwok-Yung, Gordon Siamon, Guan Yi, Peiris Joseph SM. Cytokine responses in severe acute respiratory syndrome coronavirus-infected macrophages in vitro: possible relevance to pathogenesis. *J Virol* 2005;79:7819–26.
- [11] Deng Xufang, Hackbart Matthew, Mettelman Robert C, O'Brien Amornrat, Mielech Anna M, Guanghui Yi C, Kao Cheng, Baker Susan C. Coronavirus nonstructural protein 15 mediates evasion of dsRNA sensors and limits apoptosis in macrophages. *PNAS* 2017;114:E4251–60.
- [12] Griffin Michael O, Fricovsky Eduardo, Ceballos Guillermo, Villarreal Francisco. Tetracyclines: a pleiotropic family of compounds with promising therapeutic properties. *Am J Physiol Cell Physiol* 2010;299:C539–48.
- [13] Marten NW, Zhou J. The role of metalloproteinases in corona virus infection. In: Lavi E, Constantinescu CS, editors. *Experimental models of multiple sclerosis*. Boston, MA: Springer; 2005.

Edoardo Francini
*Department of Experimental and Clinical Medicine, University of Florence,
Florence 50134, Italy*
E-mail address: edoardo.francini@unifi.it.

Salvatora T. Miano
Unit of Oncology, Santa Maria alle Scotte Hospital, Siena, Italy

Anna I. Fiaschi
Department of Pharmacology, Santa Maria alle Scotte Hospital, Siena, Italy

Guido Francini
University of Siena, Siena, Italy