



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Unexpected ethyltellurenylation of epoxides with elemental tellurium under lithium triethylborohydride conditions

Questa è la versione Preprint (Submitted version) della seguente pubblicazione:

Original Citation:

Unexpected ethyltellurenylation of epoxides with elemental tellurium under lithium triethylborohydride conditions / Damiano Tanini; Antonella Capperucci. - In: CHEMISTRY. - ISSN 2624-8549. - ELETTRONICO. - 2:(2020), pp. 652-661. [10.3390/chemistry2030041]

Availability:

The webpage <https://hdl.handle.net/2158/1215087> of the repository was last updated on 2020-11-10T15:03:24Z

Published version:

DOI: 10.3390/chemistry2030041

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

Conformità alle politiche dell'editore / Compliance to publisher's policies

Questa versione della pubblicazione è conforme a quanto richiesto dalle politiche dell'editore in materia di copyright.

This version of the publication conforms to the publisher's copyright policies.

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

1 Type of the Paper (Article, Review, Communication, etc.)

2 Unexpected ethyltellurenylation of epoxides with 3 elemental tellurium under lithium 4 triethylborohydride conitions

5 Damiano Tanini^{1,*} and Antonella Capperucci

6 ¹ Department of Chemistry "Ugo Schiff", Università di Firenze, Via della Lastruccia 3-13, 50019 Sesto
7 Fiorentino, Italy; antonella.capperucci@unifi.it (A.C.); damiano.tanini@unifi.it (D.T.).

8 Affiliation 1; e-mail@e-mail.com

9)

10 Received: date; Accepted: date; Published: date

11 **Abstract:** The one-pot multistep ethyltellurenylation reaction of epoxides with elemental tellurium
12 and lithium triethylborohydride has been described. The reaction mechanism has been
13 experimentally investigated. Lithium ditelluride and triethyl borane, formed from elemental
14 tellurium and lithium triethylborohydride, were shown to be the key species involved in the
15 reaction mechanism. Epoxides undergo ring-opening reaction with lithium ditelluride to afford
16 β -hydroxy ditellurides, which are sequentially converted into the corresponding β -hydroxy-alkyl-
17 ethyltellurides by transmetalation with triethyl borane, reasonably proceeding through S_N2
18 mechanism.

19 **Keywords:** tellurium; tellurides; ditellurides; superhydride; boranes; ring-opening-reactions;
20 epoxides; transmetalation; radicals.

21

22 1. Introduction

23 Organoselenium [1,2] and organotellurium [3] compounds continue to find wide application in
24 chemical sciences and biology [4-8]. Tellurium-containing derivatives play an important role in
25 organic synthesis [3,9], materials sciences [10,11], and medicinal chemistry [8,12-15]. The
26 incorporation of tellurium atoms into organic structures is often a rewarding strategy in developing
27 new enzyme modulators [14-17], catalysts [18], smart materials [10,11], and
28 glutathione-peroxidase-like antioxidants [19-25]. Additionally, often undergoing regio- and
29 stereo-selective transformations, organotellurium compounds can be employed in synthetically
30 useful functional group conversion reactions [26,27] and carbon-carbon bond-forming
31 processes.[28-30] Owing to these features, tellurenylation reactions provide an attractive functional
32 handle for further elaboration. The development of new, reliable, and general methodologies
33 towards these chalcogen-containing organic molecules is thus highly sought after in organic
34 synthesis. Particularly, the possibility to access densely functionalised and sp^3 -rich compounds,
35 characterised by high molecular complexity, enables the possibility to define and explore new
36 chemical space and plays a key role in terms of successfully developing new catalysts and drug
37 candidates [31,32]. Furthermore, sp^3 -rich organochalcogens bearing O- and N-containing
38 functionalities have been demonstrated to possess improved catalytic and pharmacological
39 properties [15-17,20,23]. However, although a number of methods towards selenides and tellurides
40 have been reported, a number of limitations remained, including functional-group compatibility
41 and the harsh reaction conditions. Therefore, the development of mild procedures for the synthesis
42 of densely functionalised molecules still remains challenging.

43 Three-membered heterocycles such as epoxides and aziridines, often undergoing regioselective
 44 nucleophilic ring-opening reactions (NRORs), represent convenient starting materials for the
 45 synthesis of functionalised chalcogen-containing systems [33]. A number of ring-opening-based
 46 procedures towards hydroxy- and amino-substituted selenides and tellurides have been developed
 47 over the last decade [34-39]. Such a functionalised chalcogenides have also been employed as
 48 intermediates for the synthesis of valuable compounds [40,41] and as organocatalysts for the
 49 asymmetric addition of diethylzinc to aldehydes [42].


50 In this communication, as a part of our growing interest in the study of the chemistry of
 51 organotellurium compounds, we report a study on the mechanism of an unexpected reaction of
 52 epoxides with elemental tellurium and lithium triethylborohydride, leading to the formation of
 53 β -hydroxy-alkyl-ethyl-tellurides.

54 2. Results

55 During the course of our studies on the reactivity of strained heterocycles with
 56 selenium-centered nucleophiles we developed convenient routes towards a variety of hydroxy-,
 57 amino-, and mercapto-substituted Se-containing systems [43-45]. For example, through the tuning of
 58 the stoichiometry and the conditions of the reaction of $(\text{Me}_3\text{Si})_2\text{Se}$ [(bis(trimethylsilyl)selenide, a
 59 synthetic equivalent of hydrogen selenide] with epoxides, thiiranes, and aziridines, we were able to
 60 successfully achieve a range of functionalised selenols [43], selenides, and diselenides [46].

61 Attracted by the synthetic utility and versatility of organotellurium compounds, we recently
 62 moved to evaluate the chemistry of tellurium-centered nucleophiles with strained heterocycles
 63 [47,48]. The poor stability of $(\text{Me}_3\text{Si})_2\text{Te}$ [49] prompted us to employ lithium telluride and lithium
 64 ditelluride, generated from elemental tellurium and lithium triethylborohydride (superhydride), as
 65 tellurenylation reagents for the NRORs of epoxides and aziridines [48]. However, while ring
 66 opening of epoxides with Li_2Te provided access to symmetrical β -hydroxy-tellurides **1** (Scheme 1,
 67 part a), the reaction with Li_2Te_2 gave almost exclusively ethyl-alkyl-tellurides **2** instead of the
 68 expected β -hydroxy-ditellurides **3**, which were isolated only in traces amount (Scheme 1, part b).
 69 Intrigued by this result, we wished to deeper investigate such a transformation in order to establish
 70 the mechanism involved in the formation of unsymmetrical hydroxy-ethyl-tellurides **2**.

71



72


73

74 **Scheme 1.** Reactivity of epoxides with Li_2Te and Li_2Te_2 , generated from elemental tellurium under
 75 lithium triethylborohydride conditions. **a**, Synthesis of tellurides **1**. **b**, Unexpected formation of
 76 ethyl-alkyl-tellurides **2**.

77 A plausible explanation for the formation of unsymmetrical tellurides **2** involves the
 78 transmetalation of triethylborane with β -hydroxy-ditellurides **3**. However, an alternative path could
 79 proceed through the ring-opening of epoxides with tris(ethyltelluro)borane **4** which, in principle,
 80 could be generated from lithium ditelluride and triethyl borane. A series of control experiments
 81 were therefore undertaken in order to test these hypotheses.

82 We initially evaluated whether tris(ethyltelluro)borane **4** could be generated upon treatment of
 83 elemental tellurium with lithium triethylborohydride. However, the formation of **4** was not
 84 observed under the standard reaction conditions (Scheme 2, reaction *a*). Traces of **4** were detected
 85 neither performing the reaction in a coaxial NMR tube and monitoring its progress over the time.

123 ditelluride **3** [56]. Furthermore, the reaction of **8** with oxygen or borylperoxyl radicals (V) would
 124 afford reactive tellurenyl peroxides which plausibly decompose, thus explaining the rather low yield
 125 of the transmetalation reaction and the absence of ditelluride **3**, or unreacted epoxide in the crude
 126 mixture.
 127



128
 129
 130
 131

Scheme 3. Proposed mechanism for the formation of unsymmetrical hydroxy-ethyltellurides **2**.

132 3. Conclusions

133 In conclusion, we have described a one-pot multistep reaction in which epoxides are converted
 134 into the corresponding unsymmetrical β -hydroxy-ethyltellurides upon treatment with elemental
 135 tellurium under lithium triethylborohydride reducing conditions. The reaction mechanism has been
 136 experimentally investigated; β -hydroxy ditellurides and triethyl borane were demonstrated to be the
 137 key species involved in this one-pot ethyltellurenylation reaction. The transmetalation of triethyl
 138 borane with hydroxy-dialkyl-ditellurides, reasonably occurring through an oxygen-induced S_{H2}
 139 mechanism, represents the key step of the process. The findings here described can be exploited for
 140 the development of novel general methodologies towards the synthesis of synthetically and
 141 biologically valuable complex sp^3 -rich unsymmetrical tellurides. Further studies on the application
 142 of this reaction to functionalised boranes (and boronic esters) for the preparation and the elaboration
 143 of poly-functionalised unsymmetrical tellurides are currently on-going in our laboratories.

144 4. Materials and Methods

145 4.1 Experimental Section

146 All reactions were carried out in an oven-dried glassware. Solvents were dried using a solvent
 147 purification system (Pure-Solv™). All commercial materials were purchased from various
 148 commercial sources and used as received, without further purification. Flash column
 149 chromatography purifications were performed with Silica gel 60 (230-400 mesh). Thin layer
 150 chromatography was performed with TLC plates Silica gel 60 F₂₅₄, which was visualised under UV
 151 light, or by staining with an ethanolic acid solution of *p*-anisaldehyde followed by heating. High
 152 resolution mass spectra (HRMS) were recorded by Electrospray Ionization (ESI).

153 ¹H and ¹³C NMR spectra were recorded in CDCl₃ with Mercury 400, Bruker 400 Ultrashield, and
 154 Varian Gemini 200 spectrometers operating at 400 MHz for ¹H and 100 MHz for ¹³C. NMR signals

155 were referenced to nondeuterated residual solvent signals: 7.26 ppm for ^1H and 77.0 ppm for ^{13}C .
156 ^{125}Te NMR spectra were recorded in CDCl_3 at 126 MHz with a Bruker Ultrashield 400 Plus
157 instrument. $(\text{PhTe})_2$ was used as an external reference ($\delta = 420$ ppm). Chemical shifts (δ) are given in
158 parts per million (ppm), and coupling constants (J) are given in Hertz (Hz), rounded to the nearest
159 0.1 Hz. ^1H NMR data are reported as follows: chemical shift, integration, multiplicity (s = singlet, d =
160 doublet, t = triplet, q = quartet, m = multiplet, dd = doublet of doublet, bs = broad singlet, ecc.),
161 coupling constant (J), and assignment.

162 Ditelluride **3a** [50] and diselenide **6a** [46] were prepared from 2-((benzyloxy)methyl)oxirane
163 according to literature reported procedures.

164 4.2 General Procedure for the synthesis of β -hydroxy-alkyl-ethyl-tellurides **2**

165 Li_2Te_2 was generated according to literature [49] from 1.0 mL of a 1 M THF solution of LiEt_3BH
166 (1.0 mmol, 1.0 eq.) and elemental tellurium powder (128 mg, 1.0 mmol, 1.0 eq.), stirred at ambient
167 temperature under inert atmosphere (N_2) for 6 h. The dark red suspension of Li_2Te_2 in THF was
168 treated with the epoxide (1.0 mmol, 1.0 eq.) and the reaction was stirred for 6 h at ambient
169 temperature. Afterwards, the mixture was diluted with Et_2O (10 mL), filtered through a short pad of
170 celite, washed with sat. aq. NH_4Cl and then with H_2O (2×5 mL). The organic phase was dried over
171 Na_2SO_4 , filtered and evaporated under reduced pressure. The crude residue was then purified by
172 flash chromatography to yield β -hydroxy-alkyl-ethyl-tellurides **2**.

173 4.2.1 Synthesis of 1-(Benzyloxy)-3-(ethyltellanyl)propan-2-ol **2a**

174 According to the general procedure, 2-((benzyloxy)methyl)oxirane (152 μL , 1.0 mmol) and
175 elemental tellurium (128 mg, 1.0 mmol) gave after flash chromatography (Et_2O /petroleum ether 1:1)
176 **2a** as a colourless oil (61 mg, 38%). ^1H NMR (200 MHz, CDCl_3): δ (ppm) 1.60 (3H, t, $J = 7.6$ Hz, CH_3),
177 2.63 (2H, ap q, $J = 7.6$ Hz, $\text{CH}_3\text{CH}_2\text{Te}$), 2.63 (1H, bs, OH), 2.76-2.89 (2H, m, CH_2Te), 3.41-3.48 (1H, m,
178 $\text{CH}_a\text{H}_b\text{O}$), 3.59 (1H, dd, $J = 4.2, 9.6$ Hz, $\text{CH}_a\text{H}_b\text{O}$), 3.45-3.97 (1H, m, CHOH), 4.55 (2H, ap s, CH_2Ph),
179 7.26-7.40 (5H, m). ^{13}C NMR (100 MHz, CDCl_3): δ (ppm) -4.5($\text{CH}_3\text{CH}_2\text{Te}$), 7.7 (CH_2Te), 17.8, 70.6, 73.4,
180 74.3, 127.8, 127.8, 128.4, 137.9. ^{125}Te NMR (126 MHz, CDCl_3): δ (ppm) 213.6.

181

182 **Conflicts of Interest:** The authors declare no conflict of interest.

183 References

- 184 1. Lenardão, E. J.; Santi, C.; Sancineto, L. *New Frontiers in Organoselenium Compounds*, Springer, New
185 York, 2018.
- 186 2. Wirth, T. *Organoselenium Chemistry: Synthesis and Reactions*, Wiley-VCH Verlag GmbH & Co, 2012.
- 187 3. Petragani, N.; Stefani, H. A. *Tellurium in Organic Synthesis*, 2nd ed., Elsevier, Amsterdam, 2007.
- 188 4. [4] Chem biol.....
- 189 5. Gandin, V.; Khalkar, P.; Braude, J.; Fernandes, A. P. Organic selenium compounds as potential
190 chemotherapeutic agents for improved cancer treatment. *Free Radic. Biol. Med.* **2018**, *127*, 80-97.
- 191 6. Kumar, S.; Yan, J.; Poon, J.; Singh, V. P.; Lu, X.; Karlsson Ott, M.; Engman, L.; Kumar, S.
192 Multifunctional Antioxidants: Regenerable Radical-Trapping and Hydroperoxide-Decomposing
193 Ebselenols. *Angew. Chem. Int. Ed.* **2016**, *55*, 3729-3733.
- 194 7. Ba, L. A.; Doring, M.; Jamier, V.; Jacob, C. Tellurium: an element with great biological potency and
195 potential. *Org. Biomol. Chem.* **2010**, *8*, 4203-4216.
- 196 8. For a review see: Nogueira, C. W.; Zeni, G.; Rocha, J. B. T. Organoselenium and organotellurium
197 compounds: toxicology and pharmacology. *Chem. Rev.*, **2004**, *104*, 6255-6286 and references cited
198 therein.
- 199 9. Comasseto, J. V.; Gariani, R. A. Biotransformations on Organic Selenides and Tellurides: Synthetic
200 Applications. *Tetrahedron*, **2009**, *65*, 8447-8459.

- 201 10. Liu, J.; Ma, X.; Tong, Y.; Lang, M. Self-healing polyurethane based on ditelluride bonds. *Appl. Surf. Sci.*
202 2018, 455, 318-325.
- 203 11. Xia, X.; Xiang, X.; Huang, F.; Zhang, Z.; Han, L. A tellurysulfide bond-containing redox-responsive
204 superparamagnetic nanogel with acid-responsiveness for efficient anticancer therapy. *Chem. Commun.*
205 2017, 53, 13141-13144.
- 206 12. Engman, L.; Al-Maharik, N.; McNaughton, M.; Birmingham, A.; Powis, G. Thioredoxin reductase and
207 cancer cell growth inhibition by organotellurium antioxidants. *Anti-Cancer Drugs*, 2003, 14, 153-161.
- 208 13. Rooseboom, M.; Vermeulen, N. P. E.; Durgut, F.; Commandeur, J. N. M. Comparative study on the
209 bioactivation mechanisms and cytotoxicity of Te-phenyl-L-tellurocysteine,
210 Se-Phenyl-L-selenocysteine, and S-Phenyl-L-cysteine. *Chem. Res. Toxicol.* 2002, 15, 1610-1618.
- 211 14. Lin, T.; Ding, Z.; Li, N.; Xu, J.; Luo, G.; Liu, J.; Shen, J. 2-Tellurium-bridged β -cyclodextrin, a
212 thioredoxin reductase inhibitor, sensitizes human breast cancer cells to TRAIL-induced apoptosis
213 through DR5 induction and NF- κ B suppression. *Carcinogenesis*, 2010, 32, 154-167.
- 214 15. Tanini, D.; Ricci, L.; Capperucci, A.; Di Cesare Mannelli, L.; Ghelardini, C.; Peat, T. S.; Carta, F.;
215 Angeli, A.; Supuran, C. T. Synthesis of novel tellurides bearing benzensulfonamide moiety as
216 carbonic anhydrase inhibitors with antitumor activity. *Eur. J. Med. Chem.* 2019, 181, 111586.
- 217 16. Tanini, D.; Capperucci, A.; Supuran, C. T., Angeli, A. Sulfur, selenium and tellurium containing
218 amines act as effective carbonic anhydrase activators. *Bioorg. Chem.* 2019, 87, 516-522.
- 219 17. Angeli, A.; Tanini, D.; Capperucci, A.; Supuran, C.T. First evaluation of organotellurium derivatives
220 as carbonic anhydrase I, II, IV, VII and IX inhibitors. *Bioorg. Chem.* 2018, 76, 268-272.
- 221 18. Rettig, I. D.; Van, J.; Brauer, J. B.; Luo, W.; McCormick, T. M. Tellurorhodamine photocatalyzed
222 aerobic oxidation of organo-silanes and phosphines by visible-light. *Dalton Trans.* 2019, 48, 5665-5673.
- 223 19. Tanini, D.; Lupori, B.; Malevolti, G.; Ambrosi, M.; Lo Nostro, P.; Capperucci, A. Direct biocatalysed
224 synthesis of first sulfur-, selenium- and tellurium- containing L-ascorbyl hybrid derivatives with
225 radical trapping and GPx-like properties. *Chem. Commun.* 2019, 55, 5705-5708.
- 226 20. Tanini, D.; Grechi, A.; Ricci, L.; Dei, S.; Teodori, E.; Capperucci, A. Novel functionalized
227 organotellurides with enhanced thiol peroxidase catalytic activity. *New J. Chem.* 2018, 42, 6077-6083.
- 228 21. Bortoli, M.; Torsello, M.; Bickelhaupt, F. M.; Orian, L. Role of the Chalcogen (S, Se, Te) in the
229 Oxidation Mechanism of the Glutathione Peroxidase Active Site. *ChemPhysChem.* 2017, 18, 2990-2998.
- 230 22. Singh, V. P.; Poon, J. F.; Engman, L. Catalytic Antioxidants: Regenerable Tellurium Analogues of
231 Vitamin E. *Org. Lett.* 2013, 15, 6274-6277.
- 232 23. Braga, A. L.; Alberto, E. E.; Soares, L. C.; Rocha, J. B. T.; Sudati, J. H.; Roos, D. H. Synthesis of
233 telluroamino acid derivatives with remarkable GPx like activity. *Org. Biol. Chem.* 2009, 7, 43-45.
- 234 24. Giles, G. I.; Fry, F. H.; Tasker, K. M.; Holme, A. L.; Peers, C.; Green, K. N.; Klotz, L. O.; Sies, H.; Jacob,
235 C. Evaluation of sulfur, selenium and tellurium catalysts with antioxidant potential. *Org. Biomol.*
236 *Chem.* 2003, 1, 4317-4322.
- 237 25. Tiano, L.; Fedeli, D.; Santroni, A. M.; Villarini, M.; Engman, L.; Falcioni, G. Effect of three diaryl
238 tellurides, and an organoselenium compound in trout erythrocytes exposed to oxidative stress in
239 vitro. *Mutat. Res.* 2000, 464, 269-277.
- 240 26. Petragnani, N.; Stefani, H. A. Advances in organic tellurium chemistry. *Tetrahedron*, 2005, 61,
241 1613-1679.
- 242 27. Comasseto, J. V.; Barrientos-Astigarraga, R. E. Add a Little Tellurium to Your Synthetic Plans!
243 *Aldrichim. Acta*, 2000, 33, 66-78.
- 244 28. Stefani, H. A.; Pena, J. M.; Manarin, F.; Ando, R. A.; Leal, M.; Petragnani, N. Negishi cross-coupling of
245 organotellurium compounds: Synthesis of biaryls, aryl-, and diaryl acetylenes. *Tetrahedron Lett.* 2011,
246 52, 4398-4401.
- 247 29. Berlin, S.; Ericsson, C.; Engman, L. Radical Carbonylation/Reductive Cyclization for the Construction
248 of Tetrahydrofuran-3-ones and Pyrrolidin-3-ones. *J. Org. Chem.* 2003, 68, 8386-8396.
- 249 30. Tucci, F. C.; Chieffi, A.; Comasseto, J. V.; Marino, J. P. Tellurium in Organic Synthesis. Preparation of
250 Z-Vinyl Cuprates from Z-Vinyl Tellurides and Their Reaction with Enones and Epoxides. *J. Org.*
251 *Chem.*, 1996, 61, 4975-4989.
- 252 31. Hirata, K.; Kotoku, M.; Seki, N.; Maeba, T.; Maeda, K.; Hirashima, S.; Sakai, T.; Obika, S.; Hori, A.;
253 Hase, Y.; *et al.* SAR Exploration Guided by LE and Fsp³: Discovery of a Selective and Orally
254 Efficacious ROR γ Inhibitor. *ACS Med. Chem. Lett.*, 2016, 7, 23-27.

- 255 32. Lovering, F.; Bikker, J.; Humblet, C. Escape from Flatland: Increasing Saturation as an Approach to
256 Improving Clinical Success. *J. Med. Chem.* **2009**, *52*, 6752-6756.
- 257 33. Tanini, D.; Capperucci, A. Ring opening reactions of heterocycles with selenium and tellurium
258 nucleophiles. *New J. Chem.* **2019**, *43*, 11451-11468.
- 259 34. Silva, P. C.; Borges, E. L.; Lima, D. B.; Jacob, R. G.; Lenardão, E. J.; Perin G.; Silva, M. S. A simple and
260 non-conventional method for the synthesis of selected β -arylalkylchalcogeno substituted alcohols,
261 amines and carboxylic acids. *ARKIVOC*, **2016**, *5*, 376-389.
- 262 35. Ganesh, V.; Chandrasekaran, S. One-Pot Synthesis of β -Amino/ β -Hydroxy Selenides and Sulfides
263 from Aziridines and Epoxides. *Synthesis*, **2009**, 3267-3278.
- 264 36. Braga, A. L.; Schwab, R. S.; Alberto, E. E.; Salman, S. M.; Vargas J.; Azaredo, J. B. Ring opening of
265 unprotected aziridines by zinc selenolates in a biphasic system. *Tetrahedron Lett.* **2009**, *50*, 2309-2311.
- 266 37. Santi, C.; Santoro, S.; Battistelli, B.; Testaferri, L.; Tiecco, M. Preparation of the First Bench-Stable
267 Phenyl Selenolate: an Interesting "On Water" Nucleophilic Reagent. *Eur. J. Org. Chem.* **2008**,
268 5387-5390.
- 269 38. Vargas, F.; Comasseto, J. V. Practical synthesis of chiral β -telluro amines by ring-opening reaction of
270 aziridines. *J. Organomet. Chem.* **2009**, *694*, 122-126.
- 271 39. Leng, T.; Wu, G.; Zhou, Y.-B.; Gao, W.; Ding, J.; Huang, X.; Liu M.; Wu, H. Silver-Catalyzed One-Pot
272 Three-Component Selective Synthesis of β -Hydroxy Selenides. *Adv. Synth. Catal.* **2018**, *360*, 4336-4340.
- 273 40. Berlin, S.; Ericsson, C.; Engman, L. Radical carbonylation/reductive cyclization for the construction of
274 tetrahydrofuran-3-ones and pyrrolidin-3-ones. *J. Org. Chem.*, **2003**, *68*, 8386-8396.
- 275 41. Liu, L.; Sun, Y.; Wang, J.; Ou, W.; Wang, X.; Huang, S. A New Formal Synthetic Route to Entecavir.
276 *Synlett*, **2019**, 748-752.
- 277 42. Braga, A. L.; Paixão, M. W.; Lüdtke, D. S.; Silveira, C. C.; Rodrigues, O. E. D. Synthesis of new chiral
278 aliphatic amino diselenides and their application as catalysts for the enantioselective addition of
279 diethylzinc to aldehydes. *Org. Lett.*, **2003**, *5*, 2635-2638.
- 280 43. Tanini, D.; Tiberi, C.; Gellini, C.; Salvi, P.R.; Capperucci, A. A Straightforward access to stable
281 β -functionalized alkyl selenols. *Adv. Synth. Catal.* **2018**, *360*, 3367-3375.
- 282 44. Tanini, D.; Scarpelli, S.; Ermini, E.; Capperucci, A. Seleno-Michael reaction of stable functionalised
283 alkyl selenols: A versatile tool for the synthesis of acyclic and cyclic unsymmetrical alkyl and vinyl
284 selenides. *Adv. Synth. Catal.* **2019**, *361*, 2337-2346.
- 285 45. Tanini, D.; D'Esopo, V.; Tatini, D.; Ambrosi, M.; Lo Nostro, P.; Capperucci, A. Selenated and
286 Sulfurated Analogues of triacyl glycerols: selective synthesis and structural characterization. *Chem.*
287 *Eur. J.*, **2020**, *26*, 2719-2725.
- 288 46. Tanini, D.; Capperucci, A.; Degl'Innocenti, A. Bis-(trimethylsilyl)selenide in the Selective Synthesis of
289 β -Hydroxy, β -Mercapto, and β -Amino Diorganyl Diselenides and Selenides Through Ring Opening
290 of Strained Heterocycles. *Eur. J. Org. Chem.* **2015**, 357-369.
- 291 47. Tanini, D.; Borgogni, C.; Capperucci, A. Mild and selective silicon-mediated access to enantioenriched
292 1,2-mercaptoamines and β -amino arylchalcogenides. *New J. Chem.* **2019**, *43*, 6388-6393.
- 293 48. Tanini, D.; Grechi, A.; Dei, S.; Teodori, E.; Capperucci, A. An easy one-step procedure for the
294 synthesis of novel β -functionalised tellurides. *Tetrahedron* **2017**, *73*, 5646-5653.
- 295 49. Detty, M. R.; Seidler, M. D. Bis(trialkylsilyl) chalcogenides. 1. Preparation and reduction of group
296 VIA oxides. *J. Org. Chem.* **1982**, *47*, 1354-1356.
- 297 50. Tanini, D.; Ricci, L.; Capperucci, A. Rongalite-Promoted *on Water* Synthesis of Functionalised
298 Tellurides and Ditellurides. *Adv. Synth. Catal.* **2020**, *362*, 1323-1332.
- 299 51. Ollivier, C.; Renaud, P. Organoboranes as a Source of Radicals. *Chem. Rev.* **2001**, *101*, 3415-3434.
- 300 52. Russell, G. A.; Tashtoush H. Free-radical chain-substitution reactions of alkylmercury halides. *J. Am.*
301 *Chem. Soc.* **1983**, *105*, 1398-1399
- 302 53. Abe, T.; Aso, Y.; Otsuro, T.; Ogura, F. Oxygen-Induced Transmetalation of Organoboranes with
303 Diphenyl Ditelluride. *Chem. Lett.* **1990**, 1671-1674.
- 304 54. Krenske, E. H.; Pryor, W.; Houk, K. N. Mechanism of S_H2 Reactions of Disulfides: Frontside vs
305 Backside, Stepwise vs Concerted. *J. Org. Chem.* **2009**, *74*, 5356-5360.
- 306 55. Wu, Y.-W.; Huang, S.-H.; Tseng, T.-F.; Yang, J.-F. The Mechanism Study of Free Radical S_H2' Reactions
307 by Leaving Group Effect and Secondary α -Deuterium Kinetic Isotope Effect. *J. Chin. Chem. Soc.* **2004**,
308 *51*, 1005-1011.

309 56. If regenerated, ditellurides **3** reasonably undergoes transmetallation with ethyl radicals affording
310 β -hydroxy-alkyl-ethyl-tellurides **2** and radicals **8**, following the proposed mechanism reported in the
311 Scheme 3.



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

312