



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Stable isotopes ($\delta^{13}\text{C}$, $\delta^{18}\text{O}$) and element ratios (Mg/Ca, Sr/Ca) of Jurassic belemnites, bivalves and brachiopods from the Neuquén

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Stable isotopes ($\delta^{13}\text{C}$, $\delta^{18}\text{O}$) and element ratios (Mg/Ca, Sr/Ca) of Jurassic belemnites, bivalves and brachiopods from the Neuquén Basin (Argentina): challenges and opportunities for palaeoenvironmental reconstructions / Matthias Alberti; Horacio Parent; Alberto C. Garrido; Nils Andersen; Dieter Garbe-Schönberg; Silvia Danise. - In: JOURNAL OF THE GEOLOGICAL SOCIETY. - ISSN 0016-7649. - ELETTRONICO.

Availability:

The webpage <https://hdl.handle.net/2158/1216979> of the repository was last updated on 2020-12-18T15:01:39Z

Published version:

DOI: 10.1144/jgs2020-163

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

Stable isotopes ($\delta^{13}\text{C}$, $\delta^{18}\text{O}$) and element ratios (Mg/Ca, Sr/Ca) of Jurassic belemnites, bivalves, and brachiopods from the Neuquén Basin (Argentina): challenges and opportunities for palaeoenvironmental reconstructions

Matthias Alberti^{1*}, Horacio Parent², Alberto C. Garrido^{3,4}, Nils Andersen⁵, Dieter Garbe-Schönberg¹, Silvia Danise⁶

¹*Institut für Geowissenschaften, Christian-Albrechts-Universität zu Kiel, Ludwig-Meyn-Straße 10, 24118 Kiel, Germany*

²*Laboratorio de Paleontología, IFG-FCEIA, Universidad Nacional de Rosario, Pellegrini 250, 2000 Rosario, Argentina*

³*Museo Provincial de Ciencias Naturales “Prof. Dr. Juan Olsacher”, Dirección Provincial de Minería, Elena de Vega 472, 8340 Zapala, Argentina*

⁴*Centro de Investigación en Geociencias de la Patagonia - CIGPat, Departamento de Geología y Petróleo, Facultad de Ingeniería, Universidad Nacional del Comahue, Buenos Aires 1.400, Neuquén, Argentina*

⁵*Leibniz Laboratory for Radiometric Dating and Stable Isotope Research, Christian-Albrechts-Universität zu Kiel, Max-Eyth-Straße 11, 24118 Kiel, Germany*

⁶*Dipartimento di Scienze della Terra, Università degli Studi di Firenze, Via La Pira 4, 50121, Firenze, Italy*

**Corresponding author (email: matthias.alberti@ifg.uni-kiel.de)*

Abstract: Fossils from the Jurassic succession of the Neuquén Basin (Argentina) were analysed for their stable isotope ($\delta^{13}\text{C}$, $\delta^{18}\text{O}$) and elemental (Mg/Ca, Sr/Ca) composition. Mg/Ca ratios point to comparatively stable temperature conditions from the Bajocian to Early Oxfordian and during the Tithonian, but do not allow a reliable reconstruction of absolute water temperatures. Sr/Ca ratios follow the general global pattern indicating water exchange between the basin and the open ocean. The $\delta^{18}\text{O}$ values can be translated into water temperatures between 20 to 25 °C for most of the studied intervals with possible shorter cold spells in the Late Pliensbachian, Bajocian, and Late Tithonian. However, precise temperature reconstructions are complicated by bivalve shells from the northern/central part of the basin pointing to local fluctuations in the $\delta^{18}\text{O}$ values of seawater. Potential reasons for these variations are discussed, but it seems most likely that they are caused by phases of enhanced

freshwater input leading to meso- to brachyhaline conditions in the northern study areas. The present article therefore exemplifies the particular challenges for temperature reconstructions in marginal seas and highlights the opportunities of combining different geochemical proxies to disentangle the influence of different environmental parameters.

Keywords: Jurassic, Argentina, Neuquén Basin, geochemistry, temperatures, palaeoclimate.

Over the last decades, stable oxygen isotope analyses of fossil hardparts ($\delta^{18}\text{O}_{\text{shell}}$) were used to continuously improve our understanding of the Jurassic world and climate (e.g., Bowen, 1963; Stevens and Clayton, 1971; Ditchfield et al., 1994; Price and Sellwood, 1997; Dromart et al., 2003; Dera et al., 2011; Korte et al., 2015). Nevertheless, such studies focusing mainly on the reconstruction of water temperatures are still dominated by data for European localities with other regions being comparatively understudied. In order to differentiate between global and regional trends, data from other regions are necessary. The present study concentrates on the Jurassic of Argentina and is part of a series of articles aimed at deepening our understanding of environmental conditions of Gondwanan localities (Alberti et al., 2012a, b, 2013, 2017, 2019a, b, 2020).

South America has extensive Jurassic successions, which have been studied by geoscientists for considerable time (e.g., pioneer studies by Steuer, 1897; Burckhardt, 1900, 1903; Haupt, 1907; Weaver, 1931). Their rich fossil content has sparked various palaeontological studies and also allowed the establishment of a biostratigraphic framework based on ammonites with an ever-increasing resolution. Although well-preserved fossils are common in many stratigraphic intervals, attempts at reconstructing environmental conditions during the Jurassic based on geochemical analyses of well-dated shells are still few (e.g., Bowen, 1963; Volkheimer et al., 2008; Gómez-Dacal et al., 2018; Alberti et al., 2019b). Temperature reconstructions based on $\delta^{18}\text{O}_{\text{shell}}$ analyses in particular necessitate assumptions on the depositional setting including water depth or the $\delta^{18}\text{O}$ value of seawater ($\delta^{18}\text{O}_{\text{sea}}$) during the life time of the analysed fossil organisms. Influences such as evaporation, river discharge, or rainfall patterns are particularly important for marginal seas such as the Neuquén Basin, which was separated from the open ocean by a volcanic arc (Fig. 1; compare Lazo et al., 2008). In order to differentiate environmental parameters, the present study combines the analyses of stable isotopes ($\delta^{18}\text{O}$, $\delta^{13}\text{C}$) and element ratios (Mg/Ca, Sr/Ca) of 179 fossil specimens from the Neuquén Basin. Elemental analyses (Fe, Mn) were also used to evaluate the preservational quality of the studied fossils.

Geological overview

The Neuquén Basin is situated between 34-41° S and 66-71° W in southwestern South America covering approximately 120,000 km² of present-day Chile and Argentina (Fig. 2; Howell et al., 2005; Parent et al., 2013). The Triassic-Paleogene succession consists of several thousand metres of sedimentary rocks including an almost complete Jurassic-Early Cretaceous marine record (Howell et al., 2005). During most of its development and until today, the Neuquén Basin was delimited by the Sierra Pintada Massif in the northeast and the North Patagonian Massif in the south, while the Andean volcanic arc separated the basin from the open Pacific Ocean (Fig. 1). It is unknown to which degree water exchange with the open ocean was possible or where exactly seaways for such an exchange existed. Some reconstructions depict the volcanic arc as a loose chain of islands (e.g., Howell et al., 2005; Fig. 1A). In contrast, Vicente (2005) proposed a more restricted situation with a single connection north of the basin (i.e. the Curepto Strait; Fig. 1B). Eventually, the continued Andean orogeny led to the uplift and folding of the Mesozoic strata, which became subsequently exposed in the western part of the Neuquén Basin (Howell et al., 2005).

The material used in the present study has been collected from two groups of outcrops: (1) the northern sections in the vicinity of Chos Malal and (2) the southern sections near Zapala (Fig. 2). The two areas are approximately 180 km apart and characterized by slightly different depositional settings (Fig. 3). The northern sections near Chos Malal include the localities of Vega de la Veranada (VV) and Pampa Tril (PT). Studied outcrops in this area include Bajocian to lower Cretaceous rocks with marine fossils occurring at several levels. In the Bathonian to Oxfordian strata, bivalves occur regularly, while belemnites and ammonites become more common in the Tithonian. Ammonites allow precise age assignments for large parts of the succession (compare Parent and Garrido, 2015; Parent et al., 2015, 2020). In general, siliciclastic rocks (partly cross-bedded) dominate the Middle Jurassic, but the Upper Callovian to Oxfordian La Manga Formation is characterized by carbonates interpreted as a distal platform (Gulisano, 1992). The Middle Oxfordian to Kimmeridgian strata consist of massive evaporites and continental rocks. The Tithonian to lowermost Cretaceous units are again dominated by fine-grained siliciclastics which were interpreted as basinal to outer ramp deposits (e.g., Spalletti et al., 1999). Finally, the Lower Valanginian Mulichinco Formation consists of calcareous sandstones with oysters (Schwarz and Howell, 2005).

The following sections were sampled near Zapala in the southern Neuquén Basin: Portada Covunco (PC), Cerrito Caracoles (CC), Cerro Granito (CG), Picún Leufú (PL-1&2), Picún Leufú Campamento Vialidad (PL-CV), and Charahuilla (CH). Some of these localities represent classic sections studied since many decades (e.g., Suero, 1951; Westermann and Riccardi, 1979; Leanza, 1990, 1993; Leanza et al., 2013). In general, the depositional setting in the southern Neuquén Basin has been described as more shallow than further north due to the Huincul Arch (compare Parent et al., 2013). Consequently, the fossil fauna is often more diverse including bivalves, ammonites, brachiopods, gastropods, echinoderms, hermatypic corals, and serpulids (e.g. Armella et al., 2007, 2008; Garrido and Parent, 2013; Parent et al., 2013). The succession is dominated by marine siliciclastics with intercalations of continental rocks (e.g., in the Callovian and Kimmeridgian; Fig. 3). A noteworthy exception is the Tithonian Picún Leufú Formation, which consists of bioclastic limestones in addition to calcareous siltstones. The individual localities and sections are described in more detail in the Supplementary Material (including geographic coordinates and information on biostratigraphy).

Material and methods

In total, 119 bivalves, 50 belemnites, seven brachiopods, and three aptychi were analysed in the present study (for photographs of exemplary specimens see the Supplementary Material). Most of these fossils were collected during a field survey in February 2018 at the localities of Vega de la Veranada, Pampa Tril, and Picún Leufú 1. Additional shells were selected from the collections of the Museo Provincial de Ciencias Naturales “Prof. Dr. Juan A. Olsacher” in Zapala, Argentina. The vast majority of the selected shells are oysters belonging to the genus *Gryphaea*. In addition, few shells of *Actinostreon*, *Aetostreon*, *Placunopsis*, and *Trichites* were analysed (Aberhan, 1994; Rubilar, 2005; Lazo, 2007; Rubilar and Lazo, 2009; Bressan and Palma, 2010). Jurassic belemnites of the Neuquén Basin are still relatively poorly known. The majority of the Bajocian rostra have a short conical form without a groove and could be assigned to the genus *Brevibelus*. The Callovian-Oxfordian specimens are characterized by a depressed cross-section with a prominent ventral groove and can be assigned to the genus *Belemnopsis*. The Tithonian belemnites belonging to the genus *Hibolites* show mostly elongate rostra, which are cylindrical in cross-section and have no or only a weak shallow ventral groove (Howlett, 1989; Doyle, 1992; Doyle et al., 1996, 1997). Analysed brachiopods were identified as the rhynchonellid species *Rhynchonelloides lamberti*, *Piarorhynchia*

136 *keideli*, and *Rhynchonella variabilis*. Due to occasional fragmentary preservation, not all
137 collected fossils could be identified to generic level.

138 Most of the collected fossils can be attributed to a particular horizon and ammonite
139 zone in already previously published sections (see Supplementary Material for details). The
140 individual sections were correlated based on regional and Tethyan ammonite biostratigraphy
141 and "theoretical" absolute ages were assigned to each sample based on the Geological Time
142 Scale 2016 (Ogg et al., 2016). This age model is theoretical in the extent that dating and
143 correlating of ammonite zones in South America are on-going processes and continuous
144 changes are expected in the future (e.g., Lena et al., 2019). Even though such changes and
145 improvements in dating might change the assigned absolute ages of the samples, the overall
146 trends discussed in the present study will likely remain the same. The northern sections near
147 Chos Malal are represented with samples from the Lower-Middle Bathonian to Lower
148 Oxfordian and the Tithonian. Fossils of the southern sections near Zapala were collected from
149 the Pliensbachian, Lower Bajocian, Bathonian, and Tithonian. The sampling focus lay on
150 Middle and Upper Jurassic sections, but no fossils could be collected from the Middle
151 Oxfordian to Kimmeridgian interval, as the corresponding strata consist of continental to
152 evaporitic deposits, which do not contain marine fossils (Fig. 3).

153 The collected fossils were examined macroscopically and the seemingly best
154 preserved specimens were subsequently cleaned and sampled with a hand-held dental drill.
155 For bivalves and brachiopods, the areas close to the umbo/hinge and near muscle scars were
156 avoided, since these are often influenced by vital effects during shell formation (e.g.,
157 Carpenter and Lohmann, 1995). Similarly, belemnite rostra were not sampled close to the
158 outer rim or near the alveole, since these are commonly prone to alteration. Large oyster
159 shells were rare, but one specimen of *Gryphaea* sp. from the Lower Oxfordian could be cut
160 longitudinally and sampled at high-resolution with 27 samples taken along a transect
161 perpendicular to the growth lines. In addition, five samples of sediment and one sample of
162 cement filling the alveole of a belemnite rostrum were collected. All specimens are
163 permanently stored at the Museo Provincial de Ciencias Naturales "Prof. Dr. Juan A.
164 Olsacher" in Zapala, Argentina.

165 The collected carbonate powder was analysed using a carbonate preparation device
166 (Kiel IV) connected to a ThermoScientific MAT 253 mass spectrometer at the Leibniz
167 Laboratory for Radiometric Dating and Stable Isotope Research at the Christian-Albrechts-
168 Universität zu Kiel, Germany. The carbonate samples were reacted within the preparation
169 device with 100 % orthophosphoric acid at 75 °C and the evolved CO₂ gas was then analysed

using the mass spectrometer. On daily routine, different laboratory internal carbonate standards and two international carbonate standards (NBS-19; IAEA-603) were analysed to control the precision of measured $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values. All values are reported in per mil relative to the Vienna Pee Dee Belemnite (VPDB) scale using NBS-19. Analytical precision of stable isotope analysis was better than $\pm 0.08\text{‰}$ ($\pm 1\text{SD}$) for $\delta^{18}\text{O}$ and better than $\pm 0.05\text{‰}$ ($\pm 1\text{SD}$) for $\delta^{13}\text{C}$.

In most cases, the sample size was large enough for elemental analyses in addition to stable isotope analyses. Samples were dissolved in dilute nitric acid and analysed for their Mg/Ca and Sr/Ca ratios and Fe, Mn mass fractions using an ICP-OES instrument (Spectro Ciros SOP) at the Institut für Geowissenschaften, Christian-Albrechts-Universität zu Kiel, Germany. Average uncertainty for Sr/Ca was around 0.9‰ and for Mg/Ca around 1.2‰. Reference materials Coral JCp-1, Tridacna JCt-1, and carbonate ECRM 752 were used as secondary standards.

Within the present study, potential correlations between different analytical results in the acquired datasets were evaluated with the help of the Spearman correlation coefficient (r_s). Illustrated linear trend lines are based on reduced major axis (RMA) regression.

Results

Detailed results of the geochemical analyses are listed in the Supplementary Material.

Preservation of the fossil material

A thorough check of the preservational quality is very important in any study using fossil shells for geochemical analyses. Cathodoluminescence microscopy has become a standard procedure to examine whether diagenetic alteration changed the chemical composition of fossil shells (e.g., Wierzbowski, 2002, 2004; Wierzbowski and Joachimski, 2007; Ullmann and Korte, 2015; Arabas, 2016; Arabas et al., 2017), but unfortunately such an easy check was not possible in the current study since the fossil material had to remain in Argentina. Instead, iron and manganese contents were analysed for most of the used specimens, which also allow an evaluation of preservational quality (e.g., Brand and Veizer, 1980, 1981; Price and Sellwood, 1997; Wierzbowski and Joachimski, 2007; Wierzbowski et al., 2009; Fujioka et al., 2019). Since concentrations of both elements are relatively low in pristine shells, cut-off values can be defined which may be used to separate potentially altered samples from the

database. Results from sediment and cement samples show high iron and manganese contents, thereby attesting to the availability of both elements in pore waters. Consequently, all fossils with an iron content above 300 $\mu\text{g/g}$ and a manganese content above 100 $\mu\text{g/g}$ were removed from environmental reconstructions (compare similar cut-off values used by Wierzbowski and Joachimski, 2007; Wierzbowski et al., 2009; Nunn and Price, 2010; Alberti et al., 2012b; Arabas et al., 2017). Furthermore, all specimens for which elemental analysis could not be performed were considered unreliable and excluded from further interpretations. Of the 119 sampled bivalves, twelve did not yield enough carbonate powder for elemental analysis and 47 were deemed unreliable because of exceeding the cut-off grades in their iron and/or manganese content. This left 60 presumably well-preserved bivalve shells. Of the 50 sampled belemnite rostra, 48 could be analysed for their element content. Five belemnite rostra were removed from further interpretations due to high iron and/or manganese contents. This left 43 seemingly well-preserved belemnites. Most of the collected brachiopods showed some abrasion on the outer shell surface. Due to their thin inner shell layer, only two of the seven brachiopod shells yielded enough sample material for elemental analyses and both were considered well-preserved. Finally, all three aptychi were deemed unreliable since their data reflect either diagenetically altered carbonate or contamination by carbonate cements filling the abundant pores in the shell plates. After elemental analyses, the collection was separated into seemingly well-preserved specimens and possibly unreliable fossils. While both datasets are included in the Supplementary Material, only results of the well-preserved fossils were used for interpretations.

Another potential indicator for the alteration of the stable isotope composition of fossil shells is a correlation of their $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values (e.g., Hodgson, 1966; Ullmann and Korte, 2015). Even though such a correlation can occasionally be primary in origin (e.g., due to a stronger primary productivity at higher temperatures), diagenetic alteration commonly leads to a decrease in $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values (e.g., Hodgson, 1966; Hudson, 1977; Nelson and Smith, 1996). This can be demonstrated by the results from the sediment samples, which show much lower $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values than well-preserved fossils. Similarly, several fossil shells, which were considered unreliable and potentially diagenetically altered because of their high iron and/or manganese contents, exhibit lowered $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values (Fig. 4). Consequently, an increasing alteration of shells should lead to a positive correlation between the stable isotope values. In contrast, the 60 well-preserved bivalve shells show a negative correlation ($r_s = -0.60$, $p < 0.05$) which is believed to be caused by the strong variation of analytical results between the different study areas instead of diagenetic alteration. If only the bivalves of the

northern sections are considered, the negative correlation is weaker ($r_s = -0.44$, $p < 0.05$; Fig. 4). The oysters of the southern sections show no significant correlation ($r_s = -0.16$, $p = 0.58$; Fig. 4). Similarly, the $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values of the 43 well-preserved belemnites show no significant correlation ($r_s = 0.00$, $p = 0.97$; Fig. 4), also if separated by study area.

Results of stable isotope and trace element analyses

Since the results are markedly different between the northern and the southern sections, both will be described separately in the following paragraphs.

Northern sections near Chos Malal. The $\delta^{13}\text{C}$ values of well-preserved oyster shells near Chos Malal (Fig. 5A) show an increase from the Early-Middle Bathonian (average: 2.09 ‰) to the Late Callovian (Dimorphosus Zone, average: 3.83 ‰) and then a slight decrease into the Early Oxfordian (Pressulus Zone, average: 3.31 ‰). No samples are available from the Kimmeridgian, but in the Early Tithonian $\delta^{13}\text{C}$ values are low (Picunleufuense Zone, average: 0.91 ‰). Values then increase slightly towards the Middle Tithonian (Zitteli/Mendozanus Zone, average: 1.74 ‰) and then decrease again until the Late Tithonian (Alternans Zone, minimum: -0.33 ‰). The two oyster shells with a Valanginian age have again slightly higher $\delta^{13}\text{C}$ values (Riveroi Zone, average: 1.10 ‰). The number of belemnites from the northern sections is comparatively limited, but the results show a similar pattern to the oysters with higher $\delta^{13}\text{C}$ values in the Early Oxfordian (Pressulus Zone, average: 0.57 ‰) and lower values in the Tithonian (average: -1.11 ‰).

The $\delta^{18}\text{O}$ values of well-preserved Jurassic oyster shells of the northern sections (Fig. 5B) are all strikingly low (below -5 ‰). The data show a gradual increase from the Early-Middle Bathonian (average: -8.59 ‰) to the Early Oxfordian (Pressulus Zone, average: -7.54 ‰). After a gap in data, this trend can be followed further from the Early Tithonian (Picunleufuense Zone, average: -6.45 ‰) to the Late Tithonian (Alternans Zone, maximum: -5.11 ‰). The two Valanginian oysters show much higher $\delta^{18}\text{O}$ values (Riveroi Zone, average: -2.66 ‰). In contrast to the oysters, the belemnites of the northern sections show no characteristic trend through time, but much higher absolute values. In the Early Oxfordian (Pressulus Zone), $\delta^{18}\text{O}$ values of belemnites vary around an average of -0.67 ‰. In the Tithonian, values are again a bit lower (average: -1.59 ‰) but also quite scattered (between -4.20 and -0.40 ‰).

The Mg/Ca ratios of well-preserved oysters of the northern sections remain largely stable through time (Fig. 5C). The record begins in the Early-Middle Bathonian (average: 2.42 mmol/mol) and changes little until the Early Oxfordian (Pressulus Zone, average: 2.36 mmol/mol). Values are similar in the Early (Picunleufuense Zone, average: 2.85 mmol/mol) and Middle Tithonian (Zitteli/Mendozanus Zone, average: 2.13 mmol/mol; Internispinosum Zone, average: 2.90 mmol/mol). Mg/Ca-ratios in the Late Tithonian (Alternans Zone) fluctuate mostly around an average of 1.63 mmol/mol, except for one outlier with a Mg/Ca ratio of 8.16 mmol/mol (sample MOZ-PI 11834/1). The two oysters from the Valanginian (Riveroi Zone) have Mg/Ca ratios around an average of 2.51 mmol/mol. In comparison, the belemnites have much higher Mg/Ca ratios. The two specimens of the Early Oxfordian (Pressulus Zone) have values around an average of 8.52 mmol/mol. In the Tithonian, the ratios fluctuate strongly around an average of 11.35 mmol/mol (between 7.92 and 14.42 mmol/mol).

The Sr/Ca ratios of oyster shells (Fig. 5D) of the northern sections are relatively stable from the Early-Middle Bathonian (average: 0.59 mmol/mol) to the early Late Callovian (Primus Zone, average: 0.60 mmol/mol). Following this, the Sr/Ca ratio decreases and reaches lowest values in the Early Oxfordian (Pressulus Zone, average: 0.46 mmol/mol). In the Tithonian, Sr/Ca ratios are generally higher and show an increase from the Early Tithonian (Picunleufuense Zone, average: 0.75 mmol/mol) to the Middle Tithonian (Zitteli/Mendozanus Zone, average: 1.07 mmol/mol) and a subsequent decrease until the Late Tithonian (Alternans Zone, minimum: 0.73 mmol/mol). The two Valanginian (Riveroi Zone) oysters show Sr/Ca ratios around an average of 0.86 mmol/mol. The Sr/Ca ratios of belemnites of the Early Oxfordian (Pressulus Zone) vary around an average of 1.39 mmol/mol. In the Tithonian, the values fluctuate more strongly around an average of 1.78 mmol/mol (between 1.33 and 2.17 mmol/mol).

Southern sections near Zapala. Two well-preserved rhynchonellid brachiopod shells with a Pliensbachian age show $\delta^{13}\text{C}$ values around an average of 3.96 ‰ (Fig. 6A). A well-preserved Early Bajocian (Giebeli Zone) oyster shell has a $\delta^{13}\text{C}$ value of 3.70 ‰ (sample MOZ-PI 11830/1) and another oyster shell from the Bathonian has a $\delta^{13}\text{C}$ value of 3.35 ‰ (sample MOZ-PI 11254). Considerably more data is available from well-preserved oyster shells with a Late Tithonian (Alternans Zone) age, which show $\delta^{13}\text{C}$ values around an average of 0.61 ‰. Even though the dataset is limited (especially for the Bajocian and Bathonian), the values seem to show an overall decrease in $\delta^{13}\text{C}$ values from the Middle to Late Jurassic, similar to

the northern sections (Fig. 5A). In addition, the recorded absolute $\delta^{13}\text{C}$ values of oyster shells are similar between the northern and southern sections (particularly for the Late Tithonian where more data is available for both areas). The 34 well-preserved belemnites with an Early Bajocian age (Giebeli Zone) show $\delta^{13}\text{C}$ values between 0.55 and 3.04 ‰ (average: 1.69 ‰).

The $\delta^{18}\text{O}$ record starts with two rhynchonellid brachiopods of Pliensbachian age with values ranging around an average of -3.24 ‰ (Fig. 6B). Two well-preserved oysters from the Middle Jurassic give $\delta^{18}\text{O}$ values of -1.44 ‰ for the Early Bajocian (Giebeli Zone) and -3.08 ‰ for the Bathonian. The $\delta^{18}\text{O}$ values of oysters from the Late Tithonian (Alternans Zone) fluctuate around an average of -1.47 ‰. These values are much higher than those recorded from oysters of the northern sections. Furthermore, they do not show a very prominent trend through time. Belemnites of the Early Bajocian (Giebeli Zone) recorded $\delta^{18}\text{O}$ values between -1.52 and -0.16 ‰, except for one outlier with -3.19 ‰. Their overall average is -0.81 ‰, which is largely comparable to values of belemnites from the northern sections, even though the latter have different ages.

The two brachiopods with a Pliensbachian age recorded Mg/Ca ratios with an average of 4.83 mmol/mol (Fig. 6C). The Mg/Ca ratios of well-preserved oysters are slightly lower but relatively stable through time. The oyster shells from the Early Bajocian (Giebeli Zone) and the Bathonian show Mg/Ca ratios of 2.10 and 2.41 mmol/mol respectively. In the Late Tithonian (Alternans Zone), Mg/Ca ratios of well-preserved oysters vary around an average of 2.91 mmol/mol. In general, the results are similar to those of oysters from the northern sections in absolute values and by being relatively stable through time. The belemnites with an Early Bajocian age (Giebeli Zone) show much higher Mg/Ca ratios around an average of 12.80 mmol/mol (ranging between 7.80 and 17.10 mmol/mol).

The two Pliensbachian brachiopod shells show Sr/Ca ratios around an average of 1.08 mmol/mol (Fig. 6D). The Sr/Ca ratios of well-preserved oyster shells seem relatively constant through time. Oysters from the Early Bajocian (Giebeli Zone) and the Bathonian have Sr/Ca ratios of around 0.72 mmol/mol. The Sr/Ca ratios of well-preserved bivalve shells from the Late Tithonian (Alternans Zone) fluctuate around an average of 0.71 mmol/mol. The belemnites with an Early Bajocian age (Giebeli Zone) have Sr/Ca ratios around an average of 1.58 mmol/mol (ranging between 1.38 and 1.80 mmol/mol).

Results of the high-resolution stable isotope analysis

One oyster shell (*Gryphaea* sp.) from the Lower Oxfordian (Pressulus Zone) of the Vega de la Veranada has been sampled at high-resolution across growth layers approximately 1 cm below the umbo (Fig. 7; Supplementary Material). A total of 27 samples was taken with an average resolution of three samples per millimeter. These samples were analysed for their stable isotope composition. The $\delta^{18}\text{O}$ values show a cyclic nature around an average value of -7.06‰ with a maximum at -5.77‰ and a minimum at -8.33‰ (Fig. 7A). There are a total of three (possibly four) cycles visible in the $\delta^{18}\text{O}$ values with the amplitude becoming increasingly weaker towards the younger side of the shell. The $\delta^{13}\text{C}$ values fluctuate less strongly around an average of 3.45‰ (maximum: 3.79‰ ; minimum: 3.09‰ ; Fig. 7B). They show a broad positive excursion in the older half of the shell and a broad negative excursion in the younger half. Overall, the $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values show a weak correlation ($r_s = 0.45$; $p = 0.02$; Fig. 7C).

Discussion

Differences between water temperature proxies

Several geochemical methods have been developed and applied to reconstruct absolute water temperatures in Earth's history including the Jurassic (e.g., Urey et al., 1951; Epstein et al., 1951; McArthur et al., 2007; Jenkyns et al., 2012; Li et al., 2013; Wierzbowski et al., 2018; Vickers et al., 2019). Among these, stable isotope ($\delta^{18}\text{O}_{\text{shell}}$) analysis has certainly become the most commonly used procedure leading to fundamental improvements of our understanding of the climate development in the Jurassic (e.g., Dera et al., 2011; Martinez and Dera, 2015; Korte et al., 2015). While results for different benthic taxa (e.g., bivalves and rhynchonellid brachiopods) are generally similar (e.g., Alberti et al., 2012a), belemnite rostra have often been found to record higher $\delta^{18}\text{O}_{\text{shell}}$ values than co-occurring benthic organisms (e.g., Prokoph et al., 2008; Mutterlose et al., 2010; Alberti et al., 2012a, 2019a). When using the same method for temperature reconstructions, this difference leads to the reconstruction of water temperatures commonly $4\text{--}5\text{ °C}$ lower for belemnites compared to shells of co-occurring bivalves or rhynchonellid brachiopods (also compare Dera et al., 2011). A series of reasons has been suggested for this seemingly systematic difference, including different life habits (e.g., a migratory behavior of belemnites) or vital effects during the formation of the belemnite rostrum (e.g., Mutterlose et al., 2010; Hoffmann and Stevens, 2019). Since belemnites are an extinct faunal group, deciphering all processes affecting the stable isotope

composition of their hardparts is difficult. In any case, it is clear that results from belemnites should be interpreted cautiously and separately from those of benthic organisms. For Jurassic calcitic shells, absolute palaeotemperatures are most commonly calculated by the equation given by Anderson and Arthur (1983) with a $\delta^{18}\text{O}_{\text{sea}}$ value of -1‰ during shell formation (as suggested for an ice-free world; Shackleton and Kennett, 1975). Following this approach, stable isotope analyses for belemnite rostra of the northern and southern sections as well as for the oysters and brachiopods of the southern sections lead to reasonable water temperatures (Figs 5, 6). Oxfordian and Tithonian belemnites from the northern sections show a slightly higher variation indicating water temperatures between 13.1 and 20.5 °C with an outlier at 30.6 °C (overall average: 17.8 °C). The Bajocian belemnites of the southern sections indicate water temperatures between 12.6 and 18.2 °C with one outlier at 25.7 °C (overall average: 15.2 °C). As expected, the $\delta^{18}\text{O}_{\text{shell}}$ values of oysters and brachiopods of the southern sections would translate into slightly higher temperatures between 16.6 and 27.0 °C (average: 19.4 °C). In contrast, the oysters of the northern sections near Chos Malal recorded $\delta^{18}\text{O}_{\text{shell}}$ values below -5‰ , which would correspond to unrealistic temperatures of 35 to 60 °C .

Reconstructing absolute water temperatures based on several different element ratios has become another standard method (particularly using foraminifers and corals; e.g., Eggins et al., 2003; Corrège, 2006; McArthur et al., 2007; Cléroux et al., 2008; Hetzinger et al., 2016; Pfeiffer et al., 2017). Experiments on recent bivalve shells have shown that Mg/Ca ratios indeed reflect water temperatures and might be more independent of freshwater influence or enhanced evaporation compared to $\delta^{18}\text{O}_{\text{shell}}$ values (e.g., Klein et al., 1996; Bougeois et al., 2016). However, these studies also revealed that the exact relationship between water temperature and Mg/Ca ratio in the shell depends strongly on the examined species. Consequently, different equations for temperature reconstructions have been developed for different taxa (compare Surge and Lohmann, 2008; Nunn and Price, 2010; Mouchi et al., 2013; Tynan et al., 2017). Since the species used in the current study are long extinct, their Mg/Ca-temperature relationship cannot be measured anymore. Consequently, all temperature reconstructions based on Mg/Ca ratios of extinct organisms should be treated cautiously. Nevertheless, even though absolute temperature estimates based on Mg/Ca ratios might be unreliable, temperature trends through time might be captured with this method.

Belemnites, which do not have closely related, living relatives are especially difficult to interpret. Nunn and Price (2010) proposed an equation to translate Mg/Ca ratios of belemnite rostra into water temperatures. Using their equation, the Oxfordian and Tithonian specimens of the northern sections would indicate water temperatures between 17.2 and 23.3

°C (average: 19.6 °C; Fig. 5C), values that are in the same range of those estimated from $\delta^{18}\text{O}_{\text{shell}}$ values. Similarly, the Bajocian specimens of the southern sections would translate into temperatures between 17.0 and 24.2 °C (average: 21.4 °C; Fig. 6C). However, whether the Mg/Ca ratios of belemnite rostra can be used for temperature reconstructions is still debated. While authors such as Nunn and Price (2010) have found a negative correlation between Mg/Ca ratios and $\delta^{18}\text{O}_{\text{shell}}$ values in belemnites and assumed that temperatures determine the oxygen isotope and Mg/Ca ratio in the rostra (sometimes also Sr/Ca; e.g., McArthur et al., 2007), Li et al. (2013) have found no such correlation in their analyses of Jurassic and Cretaceous belemnites and concluded that this ratio is unreliable as a palaeo-temperature proxy. In the current dataset, there is a weak negative correlation between $\delta^{18}\text{O}_{\text{shell}}$ values and Mg/Ca ratios of belemnites, whether examined for the entire collection ($r_s = -0.35$, $p = 0.02$) or separately for specimens of the north ($r_s = -0.68$, $p = 0.05$) and south ($r_s = -0.32$, $p = 0.07$). At the same time, there is no correlation between $\delta^{18}\text{O}_{\text{shell}}$ values and Sr/Ca ratios of belemnites ($r_s = -0.04$, $p = 0.82$; compare Supplementary Material).

If the equation developed by Mouchi et al. (2013) for modern, juvenile *Crassostrea gigas* is used for the Jurassic oysters of the Neuquén Basin, their Mg/Ca ratios translate into relatively cool water temperatures. The Mg/Ca ratios of the oyster shells of the northern sections do not vary much from the Bathonian up until the Tithonian (Fig. 5C) and would indicate temperatures between 7.6 and 15.1 °C with one outlier of 32.6 °C in the Late Tithonian (overall average: 11.5 °C). Similarly, the oysters of the southern sections show relatively little variation in their Mg/Ca ratios, which would translate into temperatures between 6.4 and 18.0 °C (average: 12.5 °C). However, these absolute temperature values depend very strongly on the used equation and there is no reason to assume that the equation of Mouchi et al. (2013) applies to the presently used Jurassic species. Nevertheless, another equation developed by Surge and Lohmann (2008) for the modern *Crassostrea virginica* in an estuarine setting leads to even colder temperatures, if applied to the present dataset. In any case, it should be noted that the Mg/Ca ratios are more or less the same for material from the northern and southern sections.

In summary, it seems that $\delta^{18}\text{O}_{\text{shell}}$ values of rhynchonellid brachiopods and oysters allow the most reliable and realistic absolute water temperature reconstructions, if the $\delta^{18}\text{O}_{\text{sea}}$ value can be approximated properly. In contrast, the $\delta^{18}\text{O}_{\text{shell}}$ values of belemnite rostra lead to an underestimation of water temperatures if following the traditional approach by using the equation of Anderson and Arthur (1983) or require the usage of a separate equation difficult

to establish for this extinct group. Finally, Mg/Ca ratios are strongly dependent on species-specific fractionation factors.

The Sr/Ca ratio of seawater

Previous authors proposed that Sr/Ca ratios of fossil hardparts reflect ancient water temperatures based on a negative correlation with $\delta^{18}\text{O}_{\text{shell}}$ values (compare McArthur et al., 2007; Sosdian et al., 2012). However, subsequent research failed to confirm a strong link between water temperatures and Sr/Ca ratios (Korte and Hesselbo, 2011). Similarly, the present dataset from Argentina shows no correlation between Sr/Ca ratios and $\delta^{18}\text{O}_{\text{shell}}$ values. Instead it seems that the ratio is largely a function of species-specific factors (e.g., fractionation factor, metabolism) and the original Sr/Ca ratio of the surrounding water body (compare Steuber and Veizer, 2002; Ullmann et al., 2013). Ullmann et al. (2013) reconstructed Sr/Ca ratios of Jurassic seawater by using a Sr distribution coefficient of 0.10 for bivalve shells and a coefficient of 0.32 for belemnite rostra. If these coefficients are applied on the current dataset of the Neuquén Basin, the results of the northern and southern sections are similar (Fig. 8) and compare well to the proposed global seawater Sr/Ca curve compiled from data of Ullmann et al. (2013, 2016). The results of the two Pliensbachian brachiopod shells are also comparable with the global curve if a Sr distribution coefficient of 0.32 is used. The most striking features of the curve of Ullmann et al. (2013, 2016) are a decrease in seawater Sr/Ca ratios throughout the Middle Jurassic, a minimum in the Oxfordian, a subsequent increase in values towards the mid-Tithonian, and a slight decrease again towards the Jurassic-Cretaceous boundary. This trend is attributed to global tectonic events (Ullmann et al., 2013) and is also similar to the global $^{87}\text{Sr}/^{86}\text{Sr}$ curve as compiled by McArthur et al. (2012) and Wierzbowski et al. (2017). While both proxies do not have to be directly related, Ullmann et al. (2013) suggested a common cause for the parallel fluctuations in an interplay between continental input and mid-ocean ridge activity. In any case, the current data suggest that water exchange between the Neuquén Basin and the open ocean occurred during the sampled time intervals, even though both were separated by a volcanic arc (Fig. 1). The slightly higher absolute values could be explained by uncertainties in the Sr distribution coefficients for the used fossil taxa. Alternatively, an influx of water masses with higher Sr/Ca ratios into the basin has to be postulated. In general, modern rivers have Sr/Ca ratios lower than those of modern oceans (e.g., Sosdian et al., 2012) except for some arid

regions in which riverine Sr/Ca ratios can be up to 16.0 mmol/mol (Holmden and Hudson, 2003).

The $\delta^{18}\text{O}$ value of seawater

Reconstructions of absolute water temperatures based on $\delta^{18}\text{O}_{\text{shell}}$ values of fossil hardparts require knowledge of the $\delta^{18}\text{O}_{\text{sea}}$ value. Most studies focusing on the Jurassic time interval use a $\delta^{18}\text{O}_{\text{sea}}$ value of -1‰ to acknowledge the lack of polar ice shields (Shackleton and Kennett, 1975). In general, this is an oversimplification which neglects a likely latitudinal gradient in $\delta^{18}\text{O}_{\text{sea}}$ values, with considerably higher values in the tropics and lower values at high latitudes, caused by the hydrological cycle (compare Zachos et al., 1994; Roche et al., 2006; LeGrande and Schmidt, 2006). However, the Neuquén Basin was situated at a palaeolatitude of ca. 40°S throughout most of the Jurassic (Besse and Courtillot, 2002; Torsvik et al., 2012; van Hinsbergen et al., 2015), which corresponds approximately to the point where the latitudinal $\delta^{18}\text{O}_{\text{sea}}$ gradient could have reached the value -1‰ (compare Alberti et al., 2020). Thus, using this value in the equation of Anderson and Arthur (1983) leads to the reconstruction of reasonable water temperatures for the studied well-preserved belemnites, as well as the oyster and brachiopod shells of the southern sections. In contrast, the $\delta^{18}\text{O}_{\text{shell}}$ values of the bivalves from the northern sections are very negative and would correspond to very high, unrealistic water temperatures. Several options are theoretically possible to explain such negative $\delta^{18}\text{O}_{\text{shell}}$ values, which will be discussed in the following (compare Fig. 9).

(1) Possibly the simplest explanation for very negative $\delta^{18}\text{O}_{\text{shell}}$ values in any fossil record is a poor preservation of the analysed specimens, since diagenetic alteration generally leads to a shift of $\delta^{18}\text{O}_{\text{shell}}$ values to more negative values. However, as described above, no signs of pronounced alteration are present in the used oyster shells. Furthermore, the presence of cyclic changes in $\delta^{18}\text{O}_{\text{shell}}$ values in the oyster used for high-resolution stable isotope analysis (Fig. 7) does not support strong diagenetic alteration, which would have most likely led to a more uniform stable isotope composition throughout the shell. Consequently, alteration seems to be an unlikely cause for the documented negative $\delta^{18}\text{O}_{\text{shell}}$ values of bivalves from the northern sections.

(2) If diagenetic alteration is ruled out as a possible factor and average water temperatures between 20 to 25°C are assumed for the Middle to Late Jurassic, the very negative $\delta^{18}\text{O}_{\text{shell}}$ values would correspond to $\delta^{18}\text{O}_{\text{sea}}$ values at the northern study areas (= northern/central Neuquén Basin) of -7 to -6‰ in the Bathonian to Early Oxfordian and -5 to

506 -4‰ in the Tithonian. Such low $\delta^{18}\text{O}_{\text{sea}}$ values characterize polar waters, which are
 507 commonly enriched in ^{16}O (e.g., $\delta^{18}\text{O}_{\text{sea}}$ values of -5‰ and below have been recorded in
 508 present-day high latitudes and it can be assumed that similar and lower values were possible
 509 in the Jurassic; Schmidt et al., 1999; Thomas and Mol, 2018). Theoretically, north-bound
 510 currents along the South American west coast could have transported polar ocean waters
 511 northwards during the Jurassic (similar to today). However, the Neuquén Basin was situated
 512 at a palaeolatitude of 40°S during the Middle and Late Jurassic and it seems unlikely that
 513 ocean currents could transport a negative $\delta^{18}\text{O}_{\text{sea}}$ signal so far to the north. In fact, models for
 514 absolute $\delta^{18}\text{O}_{\text{sea}}$ values in the Cretaceous do not predict particularly low values at comparable
 515 latitudes in the southern hemisphere (Zhou et al., 2008). Danise et al. (2020) described the
 516 Jurassic temperature development of the Sundance Seaway of northwestern North America.
 517 In this region (at a palaeolatitude of ca. 40°N), Middle Jurassic oysters with very negative
 518 $\delta^{18}\text{O}_{\text{shell}}$ values were explained by an influx of Arctic waters into the epicontinental basin. The
 519 palaeogeography of the Sundance Seaway with only one connection to the open ocean at high
 520 latitudes in northern North America is relatively well known. In contrast, it is still debated to
 521 which extent the volcanic arc separated the Neuquén Basin from the open ocean (Fig. 1).
 522 Some palaeogeographic reconstructions indicate a rather loose chain of islands, which would
 523 allow water exchange through a series of channels west of the actual basin (e.g., Spalletti et
 524 al., 2000: fig. 8; Howell et al., 2005: fig. 4). Similarly, the Sr/Ca ratios measured for the
 525 current study point to water exchange with the open ocean during the studied time intervals
 526 (see above). In contrast, Scherer and Goldberg (2007: fig. 1) seem to imply that the main
 527 connection to the open ocean was situated at high latitudes in southern South America.
 528 However, if an influx of polar waters from the south into an otherwise restricted Neuquén
 529 Basin is used to explain the very negative $\delta^{18}\text{O}_{\text{shell}}$ values of the oysters from the northern
 530 sections, it seems unclear, why the remaining fossils do not show similar values. Furthermore,
 531 Vicente (2005: fig. 13; 2006) proposed that the major connection of the Neuquén Basin with
 532 the open ocean was actually situated in the north (i.e. the Curepto Strait; Fig. 2B) and the
 533 basin might have been closed towards the south (also compare Howell et al., 2005: fig. 4;
 534 Parent, 2006; Kietzmann et al., 2014: fig. 1; Godoy, 2015: fig. 2). It might be speculated that
 535 upwelling along the South American west coast could bring polar water masses with low
 536 $\delta^{18}\text{O}_{\text{sea}}$ values into the Neuquén Basin. While seasonal upwelling along western South
 537 America has been predicted by some Jurassic climate models (Price et al., 1995), no evidence
 538 for this process has been found yet in sediments, fossil faunas, or geochemistry (e.g., Li
 539 enrichments in shells; Sadatzki et al., 2019).

(3) The formation of sea ice leads to water masses with higher salinities and lower $\delta^{18}\text{O}_{\text{sea}}$ values that sink to the sea floor (compare Barrera et al., 1987; Ravelo and Hillaire-Marcel, 2007). However, at a palaeolatitude of 40° S for the Neuquén Basin in the Middle and Late Jurassic, the formation of extensive sea ice is very unlikely (even though some Jurassic climate models suggest sub-zero temperatures at comparable palaeolatitudes in India; Sellwood et al., 2000). Furthermore, changes in $\delta^{18}\text{O}_{\text{sea}}$ values via sea ice formation would only be seasonal.

(4) The breakdown of volcanoclastics into smectite and mixed layer clays at the sediment-water interface and/or within the sediment can lower $\delta^{18}\text{O}$ values by several per mil in bottom and pore waters (Lawrence et al., 1979; Price and Sellwood, 1997). The existence of an active volcanic arc allowed a continuous supply of volcanoclastics into the Neuquén Basin. However, it is not clear why this process should only affect the northern study areas. Furthermore, this geochemical process alone is not strong enough to cause the proposed negative $\delta^{18}\text{O}_{\text{sea}}$ values.

(5) Freshwater is generally enriched in ^{16}O and commonly used to explain negative $\delta^{18}\text{O}_{\text{sea}}$ values. River discharge or strong rainfalls modify $\delta^{18}\text{O}_{\text{sea}}$ values particularly in surface waters, because freshwater forms lenses on top of the heavier saline water. Such freshwater influence restricted to the northern study areas could explain the very negative $\delta^{18}\text{O}_{\text{shell}}$ values of the bivalves. Meso- to brachyhaline conditions (salinities 16-20) in the Bathonian, Late Callovian, and Early Oxfordian would correspond to $\delta^{18}\text{O}_{\text{sea}}$ values of -7 to -6 ‰ at average water temperatures of 20 °C (based on the method of Lazo et al., 2008: fig. 3). The proposed $\delta^{18}\text{O}_{\text{sea}}$ values of -5 to -4 ‰ for the Tithonian would correspond to brachyhaline conditions (salinities 23-27). While oysters are generally tolerant towards fluctuations in salinities and live in marine as well as brackish habitats, ammonites and belemnites are considered stenohaline. However, since these cephalopods are active swimmers, they might have migrated throughout the basin and did not necessarily live within the presumably brackish waters in the northern study areas. Separate habitats would explain the higher $\delta^{18}\text{O}_{\text{shell}}$ values of the belemnites from the northern sections. Post-mortem drift of cephalopod shells over wide distances is also not unlikely.

Climate models for the Jurassic of South America predict the position of the Intertropical Convergence Zone (ITCZ) towards the north of the Neuquén Basin (Scherer and Goldberg, 2007). While the area of the basin itself was situated in a dry region (e.g., Volkheimer et al., 2008), rainfalls towards the north of the basin might have occurred regularly and fueled rivers draining into the northern Neuquén Basin. Since palaeolatitudes

did not change markedly during the Middle and Late Jurassic, such a situation might have
 been stable for long time intervals. Because very low $\delta^{18}\text{O}$ values are limited to the northern
 sections, this area might have been more restricted than the southern part. This is supported by
 the presence of two thick evaporitic units in the northern/central Neuquén Basin (Fig. 3).
 During these phases, the Curepto Strait must have been closed (as suggested by Vicente,
 2005) and the northern basin was thus separated from the open ocean (due to sea-level
 changes and local tectonic movements; Hallam, 2001). Lazo et al. (2008; see also Aguirre-
 Urreta et al., 2008) studied the stable oxygen isotope composition of Early Cretaceous oysters
 in the northern/central Neuquén Basin north of Zapala and near Chos Malal. Similar to the
 Jurassic data from this study, their results include very negative $\delta^{18}\text{O}_{\text{shell}}$ values. The authors
 explained these values by freshwater influence in the basin during certain intervals in the
 Cretaceous. It seems therefore reasonable to propose salinity fluctuations in the sampled
 northern/central Neuquén Basin throughout the Jurassic and Cretaceous as a result of an
 interplay between changing river influx and sea water exchange leading either to brackish
 conditions or the formation of evaporites. Such a scenario changes the general understanding
 of the northern/central Neuquén Basin somewhat as this area was originally considered to
 represent a more distal and deeper area. Even though the Tithonian is dominated by fine-
 grained sediments possibly deposited below wave base, the Bathonian to Oxfordian strata
 sampled here commonly show cross-bedded horizons, which cannot be deposited at high
 water depths. Interestingly, the $\delta^{18}\text{O}_{\text{shell}}$ values of bivalves are less negative in the Tithonian
 compared to the Bathonian to Oxfordian, possibly indicating a weaker freshwater influence
 towards the end of the Jurassic, when sea level was generally higher.

Seasonal temperature changes

The Lower Oxfordian oyster used for high-resolution analysis shows a cyclic signal in its
 $\delta^{18}\text{O}_{\text{shell}}$ values interpreted to reflect seasonal patterns. However, as in other shells from the
 northern sections, the $\delta^{18}\text{O}_{\text{shell}}$ values are very low. Using a $\delta^{18}\text{O}_{\text{sea}}$ value of -1‰ , the $\delta^{18}\text{O}_{\text{shell}}$
 data would translate into unrealistic temperatures between 39 and 53 °C (Fig. 7A). On the
 other hand, the shell's Mg/Ca ratio would indicate an unrealistically low temperature of 12.5
 °C by using the equation of Mouchi et al. (2013). If a $\delta^{18}\text{O}_{\text{sea}}$ value of -6.5‰ is applied
 accounting for the proposed freshwater influence, a seasonality of about 11 °C is
 reconstructed (minimum: 13.0 °C, maximum: 24.0 °C; Fig. 7A). Due to the problems in
 reconstructing precise $\delta^{18}\text{O}_{\text{sea}}$ values, these absolute temperatures are less reliable. However,

the temperature amplitude (= seasonality) might be in fact a reasonable estimate for the Neuquén Basin in the Oxfordian at a palaeolatitude of 40° S. In addition, fossils analysed in the present study and collected from one stratigraphic interval show a comparable variation, possibly reflecting a similarly strong seasonality (also compare ranges in results of Bowen, 1963). Nevertheless, additional seasonal fluctuations in the $\delta^{18}\text{O}_{\text{sea}}$ values caused, for example, by seasonal rainfall cannot be excluded completely. In addition, the cyclic nature of the $\delta^{18}\text{O}_{\text{shell}}$ data of the oyster from Vega de la Veranada shows that the northern study areas were at least occasionally shallow enough to experience seasonal environmental changes (i.e., above the thermocline).

A synopsis on Jurassic water temperatures of South America

So far, reconstructions of absolute water temperatures for the southern hemisphere in the Jurassic are comparatively few in number. Figure 10 combines Jurassic temperature data from South America including the present dataset from the Neuquén Basin as well as results of Bowen (1963), Gómez-Dacal et al. (2018), and Alberti et al. (2019b).

Mg/Ca ratios of oysters and belemnites of the Neuquén Basin do not allow the reliable reconstruction of absolute water temperatures, but point to more or less stable temperature conditions throughout the studied Middle to Late Jurassic intervals. Pliensbachian brachiopods show higher Mg/Ca ratios than Bajocian and Bathonian oysters, but since the relationship between the Mg/Ca ratio and temperature differs strongly between species, it is not clear whether this decrease in values reflects a temperature decrease (Fig. 10A).

The Jurassic stable oxygen isotope record of South America (Fig. 10B) starts with data from Chile published recently by Alberti et al. (2019b). While fossils near Potrerillos in northern Chile show possible freshwater influence, the specimens analysed from sections around El Transito likely recorded water temperatures (compare Alberti et al., 2019b). These shells indicate temperatures around an average of 25.9 °C in the Late Sinemurian, identical to the average of 25.9 °C recorded by the two Pliensbachian brachiopods of the Neuquén Basin. One brachiopod from the latest Pliensbachian of Chile might reflect the likely global Late Pliensbachian Cooling Event with a comparatively low temperature of 19.6 °C (compare Alberti et al., 2019b). Late Toarcian temperatures of Chile are again relatively high around an average of 24.4 °C based on brachiopod and bivalve shells. Bowen (1963) analysed seven seemingly well-preserved, but poorly dated belemnite rostra of the late Early Jurassic. A total of thirty $\delta^{18}\text{O}_{\text{shell}}$ values from the seven fossils translate into water temperatures between 12.4

and 25.3 °C. The poor stratigraphic resolution somehow diminishes the value of these temperature reconstructions. The Middle Jurassic record starts with one Bajocian oyster of the Neuquén Basin, which recorded a relatively low temperature of 17.8 °C. Another oyster from the Bathonian shows again a higher temperature of 25.2 °C. Middle to Late Jurassic belemnites of the Neuquén Basin analysed in the present study indicate relatively constant water temperatures around averages of 15.2 °C for the Early Bajocian, 14.6 °C for the Early Oxfordian, and 18.7 °C for the Tithonian. These values are largely comparable to results of Bowen (1963), who analysed eight samples of two belemnite rostra from the Middle Bajocian with reconstructed temperatures between 14.4 and 23.4 °C. As discussed above, $\delta^{18}\text{O}_{\text{shell}}$ values of the oysters from the northern sections point to a change in $\delta^{18}\text{O}_{\text{sea}}$ values in this area, possibly caused by enhanced freshwater influence. If water temperatures between 20 and 25 °C are assumed, then a $\delta^{18}\text{O}_{\text{sea}}$ value of -6.5 ‰ for the Bathonian to Early Oxfordian and -4.5 ‰ for the Tithonian can be proposed (Fig. 10B). While $\delta^{18}\text{O}_{\text{shell}}$ values of these oysters can therefore not be used to reconstruct reliable water temperatures, the stability of Mg/Ca ratios throughout this time interval suggests the absence of major temperature changes. Oyster shells from the southern sections recorded water temperatures around 18.0 °C in the Late Tithonian, before temperatures increase again into the Valanginian (Early Cretaceous) with two oysters pointing to temperatures around 23.2 °C. Gómez-Dacal et al. (2018) analysed Tithonian to Valanginian oyster shells from three sections in the northern/central Neuquén Basin with results matching the present data comparatively well (Fig. 10B). Tithonian oysters of Gómez-Dacal et al. (2018) show a wide variability in $\delta^{18}\text{O}_{\text{shell}}$ values translating into water temperatures between 20.0 and 33.4 °C (for a $\delta^{18}\text{O}_{\text{sea}}$ value of -1 ‰), possibly reflecting freshwater influence on some of the shells. During the Berriasian, reconstructed water temperatures are more confined around an average of 23.2 °C, but in the Early Valanginian values are again quite scattered (between 21.5 and 31.7 °C). These scattered values might indicate that freshwater influx in the northern/central Neuquén Basin occurred also in the Early Cretaceous. Such a scenario is also supported by Lazo et al. (2008) who recorded several intervals with very negative $\delta^{18}\text{O}_{\text{shell}}$ values of oysters in the Valanginian to Barremian of the northern/central Neuquén Basin and postulated lowered salinities in this area.

Other Jurassic water temperature records of the southern hemisphere

Apart from South America, research on Jurassic water temperatures at comparatively high southern latitudes has focused on James Ross Island in Antarctica, New Zealand, and the Malvinas (Falkland) Plateau.

Ditchfield et al. (1994) analysed Jurassic and Cretaceous macrofossils from James Ross Island in Antarctica. Their collection included ammonites, belemnites, and bivalves with a Tithonian age. Of these, eight belemnite rostra were considered to be well-preserved and yielded $\delta^{18}\text{O}_{\text{shell}}$ values between -1.20 and -0.26 ‰, corresponding to water temperatures of 16.8 to 13.0 °C. Ditchfield et al. (1994) mentioned a palaeolatitude of around 60° S for the James Ross Island, but newer reconstructions point to a palaeolatitude of 45 - 50° S (van Hinsbergen et al., 2015).

Stevens and Clayton (1971) analysed belemnites from New Zealand with Bajocian to Tithonian ages at a palaeolatitude of around 80° S (van Hinsbergen et al., 2015). Their specimens recorded $\delta^{18}\text{O}_{\text{shell}}$ values up to 0.43 ‰, but also considerably lower values around -4 ‰ (with outliers as low as -8.5 ‰), some of which might be diagenetically altered. Podlaha et al. (1998) also analysed $\delta^{18}\text{O}_{\text{shell}}$ values of Late Jurassic belemnites of New Zealand and recorded a large variability in the results (-4.40 to 1.86 ‰; with one outlier at -10.99 ‰). Similarly, Gröcke et al. (2003) analysed Late Jurassic belemnites from New Zealand, noted a high variability in their oxygen ratios, and connected those to changes in the $\delta^{18}\text{O}_{\text{sea}}$ values (e.g., via the formation of ice sheets or snow) instead of strongly fluctuating water temperatures. Ullmann et al. (2013, 2016) reconstructed water temperatures based on a large number of stable isotope analyses including Late Jurassic belemnites from New Zealand. Their specimens recorded quite variable $\delta^{18}\text{O}_{\text{shell}}$ values ranging between -4.1 to 0.8 ‰ for the Oxfordian, -3.0 to 0.5 ‰ for the Kimmeridgian, and -1.3 to 0.8 ‰ for the Early Tithonian. The highest values (up to 1.6 ‰) were reached in the Late Tithonian (Ullmann et al., 2016). Translating these values into absolute water temperatures might be difficult due to uncertainties regarding $\delta^{18}\text{O}_{\text{sea}}$ values at very high latitudes.

Price and Sellwood (1997) analysed 26 belemnite rostra and three inoceramid bivalves with a Late Jurassic age from sites of the Deep Sea Drilling Project on the Malvinas (Falkland) Plateau. The authors mentioned a palaeolatitude of 55 - 60° S for the study area in the Late Jurassic, but more recent reconstructions point to a slightly more northern location (53 - 40° S; van Hinsbergen et al., 2015). The studied taxa recorded surprisingly negative $\delta^{18}\text{O}_{\text{shell}}$ values corresponding to very warm temperatures. While Price and Sellwood (1997) argued that the inoceramids were poorly preserved ($\delta^{18}\text{O}_{\text{shell}}$ values between -2.8 and -4.2 ‰), they considered most of the belemnites to have a pristine composition. The authors

explain the relatively high reconstructed water temperatures (averages for the two study areas of 17.2 and 17.9 °C) by freshwater influx in the semi-enclosed basin. Price and Gröcke (2002) later analysed more Late Jurassic belemnites of the same study area with $\delta^{18}\text{O}_{\text{shell}}$ values ranging between -2.22 and -0.04 ‰ (translating into water temperatures of 12.1 to 21.2 °C). Jenkyns et al. (2012) published TEX_{86} sea-surface temperature reconstructions ranging between 26 to 30 °C for the Malvinas (Falkland) Plateau for the Middle to Late Jurassic. In order to explain these much warmer temperatures, the authors proposed that the analysed belemnites from the same locality lived in colder waters below the thermocline. Most recently, Vickers et al. (2019) used clumped isotope analyses on Late Jurassic to Early Cretaceous belemnites of the Malvinas (Falkland) Plateau and reconstructed warm temperatures between 21 to 28 °C (average: 25 °C). In combination with $\delta^{18}\text{O}_{\text{shell}}$ values of the belemnite rostra, the authors reconstructed surprisingly high $\delta^{18}\text{O}_{\text{sea}}$ values of around $+1$ ‰. They explained this surprisingly high value with increased evaporation in a semi-enclosed basin.

In summary, previous stable isotope analyses of Jurassic fossils from high-latitude locations in the southern hemisphere show relatively scattered and occasionally surprisingly negative $\delta^{18}\text{O}_{\text{shell}}$ values (conventionally indicating very warm water temperatures). Similar to interpretations for the Neuquén Basin, most previous authors have explained this with factors affecting the $\delta^{18}\text{O}_{\text{sea}}$ values (such as freshwater influence). In this regard, the present South American data matches the previous records of other restricted basins very well. At the same time, some previous authors discarded fossils with particularly negative $\delta^{18}\text{O}_{\text{shell}}$ values as poorly preserved, instead of considering other alternative explanations such as freshwater influence or lowered $\delta^{18}\text{O}_{\text{sea}}$ values of polar waters. The validity of very high temperatures reconstructed via the TEX_{86} -palaeothermometer has been questioned by previous authors (compare Vickers et al., 2019). Similarly, the concept of very warm temperatures reconstructed by clumped isotope analyses faces challenges. A $\delta^{18}\text{O}_{\text{sea}}$ value of $+1$ ‰ at high latitudes during the Late Jurassic should be indeed only local/regional in extent (such as in a restricted basin as proposed by Vickers et al., 2019). Other Jurassic temperature reconstructions for the southern hemisphere exist for the Tethys Ocean (i.e. in India and Madagascar; Fürsich et al., 2005; Alberti et al., 2012a, b, 2019a). These records reflect plate tectonic movements during the rifting between western and eastern Gondwana and do not contribute to the discussion in the present study.

Conclusions

105 well-preserved belemnites, bivalves, and brachiopods from two main study areas within the Neuquén Basin were analysed for their stable isotope ($\delta^{13}\text{C}$, $\delta^{18}\text{O}$) and elemental (Mg/Ca, Sr/Ca) composition. The combination of the different geochemical proxies allowed the disentanglement of different environmental parameters influencing the study area. Very negative $\delta^{18}\text{O}_{\text{shell}}$ values of oysters in the northern/central part of the basin likely reflect a variable freshwater influence (meso- to brachyhaline conditions) in this region during the Bathonian to Early Oxfordian and Tithonian. Mg/Ca and $\delta^{18}\text{O}_{\text{shell}}$ data from the remaining localities point to rather stable temperature conditions through the studied time intervals. After considering these limitations, it seems likely that water temperatures in the Neuquén Basin stayed between 20 and 25 °C for most of the studied Jurassic time intervals, possibly interrupted by short colder spells in the Late Pliensbachian, Bajocian, and Late Tithonian. High-resolution $\delta^{18}\text{O}_{\text{shell}}$ analysis of an oyster from the Lower Oxfordian points to a seasonality of around 11 °C, if the $\delta^{18}\text{O}_{\text{shell}}$ fluctuations are explained only by temperature.

Acknowledgements

The authors thank Karen Bremer and Verena Heinath for support with geochemical analyses and two anonymous reviewers for their constructive criticism. The study was financially supported by the German Research Foundation (DFG; project AL 1740/3-1).

References

- Aberhan, M. 1994. Early Jurassic Bivalvia of northern Chile. Part I. Subclasses Palaeotaxodonta, Pteriomorpha, and Isofilibranchia. *Beringeria*, **13**, 1–115.
- Aguirre-Urreta, M.B., Price, G.D., Ruffell, A.H., Lazo, D.G., Kalin, R.M., Ogle, N. & Rawson, P.F. 2008. Southern Hemisphere Early Cretaceous (Valanginian–Early Barremian) carbon and oxygen isotope curves from the Neuquén Basin, Argentina. *Cretaceous Research*, **29**, 87–99.
- Alberti, M., Fürsich, F.T. & Pandey, D.K. 2012a. The Oxfordian stable isotope record ($\delta^{18}\text{O}$, $\delta^{13}\text{C}$) of belemnites, brachiopods, and oysters from the Kachchh Basin (western India) and its potential for palaeoecologic, palaeoclimatic, and palaeogeographic reconstructions. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **344–345**, 49–68.

775 Alberti, M., Fürsich, F.T., Pandey, D.K. & Ramkumar, M. 2012b. Stable isotope analyses of
776 belemnites from the Kachchh Basin, western India: paleoclimatic implications for the
777 Middle to Late Jurassic transition. *Facies*, **58**, 261–278.

778 Alberti, M., Fürsich, F.T. & Pandey, D.K. 2013. Seasonality in low latitudes during the
779 Oxfordian (Late Jurassic) reconstructed via high-resolution stable isotope analysis of the
780 oyster *Actinostreon marshi* (J. Sowerby, 1814) from the Kachchh Basin, western India.
781 *International Journal of Earth Sciences*, **102**, 1321–1336.

782 Alberti, M., Fürsich, F.T., Abdelhady, A.A. & Andersen, N. 2017. Middle to Late Jurassic
783 equatorial seawater temperatures and latitudinal temperature gradients based on stable
784 isotopes of brachiopods and oysters from Gebel Maghara, Egypt. *Palaeogeography*,
785 *Palaeoclimatology*, *Palaeoecology*, **468**, 301–313.

786 Alberti, M., Arabas, A., Fürsich, F.T., Andersen, N. & Ziólkowski, P. 2019a. The Middle to
787 Upper Jurassic stable isotope record of Madagascar: Linking temperature changes with
788 plate tectonics during the break-up of Gondwana. *Gondwana Research*, **73**, 1–15.

789 Alberti, M., Fürsich, F.T. & Andersen, N. 2019b. First steps in reconstructing Early Jurassic
790 sea water temperatures in the Andean Basin of northern Chile based on stable isotope
791 analyses of oyster and brachiopod shells. *Journal of Palaeogeography*, **8**, 33.

792 Alberti, M., Leshno, Y., Fürsich, F.T., Edelman-Furstenberg, Y., Andersen, N. & Garbe-
793 Schönberg, D. 2020. Stress in the tropics? Impact of a latitudinal seawater $\delta^{18}\text{O}$ gradient
794 on Middle Jurassic temperature reconstructions at low latitudes. *Geology*, **48**.

795 Anderson, T.F. & Arthur, M.A. 1983. Stable isotopes of oxygen and carbon and their
796 application to sedimentological and paleoenvironmental problems. In: Arthur, M.A.,
797 Anderson, T.F., Kaplan, I.R., Veizer, J. & Land, L. (eds.) *Stable Isotopes in Sedimentary*
798 *Geology*. SEPM Short Course, 1–151.

799 Arabas, A. 2016. Middle-Upper Jurassic stable isotope records and seawater temperature
800 variations: New palaeoclimate data from marine carbonate and belemnite rostra (Pieniny
801 Klippen Belt, Carpathians). *Palaeogeography*, *Palaeoclimatology*, *Palaeoecology*, **446**,
802 284–294.

803 Arabas, A., Schlögl, J. & Meister, C. 2017. Early Jurassic carbon and oxygen isotope records
804 and seawater temperature variations: Insights from marine carbonate and belemnite rostra
805 (Pieniny Klippen Belt, Carpathians). *Palaeogeography*, *Palaeoclimatology*,
806 *Palaeoecology*, **485**, 119–135.

807 Armella, C., Cabaleri, N. & Leanza, H.A. 2007. Tidally dominated, rimmed-shelf facies of the
808 Picún Leufú Formation (Jurassic/Cretaceous boundary) in southwest Gondwana,
809 Neuquén Basin, Argentina. *Cretaceous Research*, **28**, 961–979.

810 Armella, C., Cabaleri, N. & Leanza, H.A. 2008. Facies de patch reefs en la Formación Picún
811 Leufú (límite Jurásico/Cretácico) en la region de Zapala, Cuenca Neuquina. *Revista del*
812 *Museo Argentino de Ciencias Naturales, N.S.*, **10**, 63–70.

813 Barrera, E., Huber, B.T., Savin, S.M. & Webb, P.-N. 1987. Antarctic marine temperatures:
814 Late Campanian through Early Paleocene. *Paleoceanography*, **2**, 21–47.

815 Besse, J. & Courtillot, V. 2002. Apparent and true polar wander and the geometry of the
816 geomagnetic field over the last 200 Myr. *Journal of Geophysical Research*, **107**, 2300.

817 Bougeois, L., de Rafélis, M., Reichart, G.-J., de Nooijer, L.J. & Dupont-Nivet, G. 2016.
818 Mg/Ca in fossil oyster shells as palaeotemperature proxy, an example from the
819 Palaeogene of Central Asia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **441**,
820 611–626.

821 Bowen, R. 1963. O¹⁸/O¹⁶ paleotemperature measurements on Mesozoic Belemnoida from
822 Neuquen and Santa Cruz provinces, Argentina. *Journal of Paleontology*, **37**, 714–718.

823 Brand, U. & Veizer, J. 1980. Chemical diagenesis of a multicomponent carbonate system - 1:
824 trace elements. *Journal of Sedimentary Petrology*, **50**, 1219–1236.

825 Brand, U. & Veizer, J. 1981. Chemical diagenesis of a multicomponent carbonate system - 2:
826 stable isotopes. *Journal of Sedimentary Petrology*, **51**, 987–997.

827 Bressan, G.S. & Palma, R.M. 2010. Taphonomic analysis of fossil concentrations from La
828 Manga Formation (Oxfordian), Neuquén Basin, Mendoza Province, Argentina. *Journal of*
829 *Iberian Geology*, **36**, 55–71.

830 Burckhardt, C. 1900. Profils géologiques transversaux de la Cordillère Argentino-Chilienne.
831 Stratigraphie et Tectonique. *Annales del Museo de la Plata, Sección Geología y Minería*,
832 **2**, 1–136.

833 Burckhardt, C. 1903. Beiträge zur Kenntnis der Jura- und Kreideformation der Cordillere.
834 *Palaeontographica*, **50**, 1–144.

835 Carpenter, S.J. & Lohmann, K.C. 1995. $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values of modern brachiopod shells.
836 *Geochimica et Cosmochimica Acta*, **59**, 3749–3764.

837 Cléroux, C., Cortijo, E., Anand, P., Labeyrie, L., Bassinot, F., Caillon, N. & Duplessy, J.-C.
838 2008. Mg/Ca and Sr/Ca ratios in planktonic foraminifera: Proxies for upper water column
839 temperature reconstruction. *Paleoceanography*, **23**, PA3214.

840 Corrège, T. 2006. Sea surface temperature and salinity reconstruction from coral geochemical
841 tracers. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **232**, 408–428.

842 Danise, S., Price, G.D., Alberti, M. & Holland, S.M. 2020. Isotopic evidence for partial
843 geochemical decoupling between a Jurassic epicontinental sea and the open ocean.
844 *Gondwana Research*, **82**, 97–107.

845 Dera, G., Brigaud, B., Monna, F., Laffont, R., Pucéat, E., Deconinck, J.-F., Pellenard, P.,
846 Joachimski, M.M. & Durlet, C. 2011. Climatic ups and downs in a disturbed Jurassic
847 world. *Geology*, **39**, 215–218.

848 Ditchfield, P.W., Marshall, J.D. & Pirrie, D. 1994. High latitude palaeotemperature variation:
849 New data from the Tithonian to Eocene of James Ross Island, Antarctica.
850 *Palaeogeography, Palaeoclimatology, Palaeoecology*, **107**, 79–101.

851 Doyle, P. 1992. The British Toarcian (Lower Jurassic) belemnites. Part 2. *Monograph of the*
852 *Palaeontographical Society*, 50–79.

853 Doyle, P., Kelly, S.R.A., Pirrie, D., Riccardi, A.C. & Olivero, E. 1996. Jurassic belemnite
854 biostratigraphy of the Southern Hemisphere: a comparative study of Antarctica and
855 Argentina. *Revista de la Asociación Geológica Argentina*, **51**, 331–338.

856 Doyle, P., Kelly, S.R.A., Pirrie, D. & Riccardi, A.C. 1997. Jurassic belemnite distribution
857 patterns: implications of new data from Antarctica and Argentina. *Alcheringa*, **21**, 219–
858 228.

859 Dromart, G., Garcia, J.-P., Picard, S., Atrops, F., Lécuyer, C. & Sheppard, S.M.F. 2003. Ice
860 age at the Middle-Late Jurassic transition? *Earth and Planetary Science Letters*, **213**,
861 205–220.

862 Eggins, S., De Deckker, P. & Marshall, J. 2003. Mg/Ca variation in planktonic foraminifera
863 tests: implications for reconstructing palaeo-seawater temperature and habitat migration.
864 *Earth and Planetary Science Letters*, **212**, 291–306.

865 Epstein, S., Buchsbaum, R., Lowenstam, H.A. & Urey, H.C. 1951. Carbonate-water isotopic
866 temperature scale. *Geological Society of America Bulletin*, **62**, 417–426.

867 Fujioka, H., Takayanagi, H., Yamamoto, K. & Iryu, Y. 2019. The effects of meteoric
868 diagenesis on the geochemical composition and microstructure of Pliocene fossil
869 *Terebratalia coreanica* and *Laqueus rubellus* brachiopod shells from northeastern Japan.
870 *Progress in Earth and Planetary Science*, **6**, 45.

871 Fürsich, F.T., Singh, I.B., Joachimski, M., Krumm, S., Schlirf, M. & Schlirf, S. 2005.
872 Palaeoclimatic reconstructions of the Middle Jurassic of Kachchh (western India): An

873 integrated approach based on palaeoecological, oxygen isotopic, and clay mineralogical
874 data. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **217**, 289–309.

875 Garrido, A.C. & Parent, H. 2013. Estratigrafía y fauna de amonites de los depósitos
876 “Lotenianos” (Caloviano Medio-Oxfordiano Inferior?) del anticlinal de Picún Leufú,
877 Cuenca Neuquina - Subcuenca de Picún Leufú, Argentina. *Boletín del Instituto de*
878 *Fisiografía y Geología*, **83**, 35–68.

879 Godoy, E. 2015. The north-western margin of the Neuquén Basin in the headwater region of
880 the Maipo drainage, Chile. *Geological Society, London, Special Publications*, 399, 155–
881 165.

882 Gómez-Dacal, A.R., Gómez-Peral, L.E., Spalletti, L.A., Sial, A.N., Siccardi, A. & Poiré, D.G.
883 2018. First record of the Valanginian positive carbon isotope anomaly in the Mendoza
884 shelf, Neuquén Basin, Argentina: palaeoclimatic implications. *Andean Geology*, 45, 111–
885 129.

886 Gröcke, D.R., Price, G.D., Ruffell, A.H., Mutterlose, J. & Baraboshkin, E. 2003. Isotopic
887 evidence for Late Jurassic-Early Cretaceous climate change. *Palaeogeography,*
888 *Palaeoclimatology, Palaeoecology*, **202**, 97–118.

889 Gulisano, C.A. 1992. Palaeogeographic evolution of west-central Argentina. In: Westermann,
890 G.E.G. (ed.) *The Jurassic of the Circum-Pacific*. Cambridge University Press, New York,
891 150–151.

892 Hallam, A. 2001. A review of the broad pattern of Jurassic sea-level changes and their
893 possible causes in the light of current knowledge. *Palaeogeography, Palaeoclimatology,*
894 *Palaeoecology*, **167**, 23–37.

895 Haupt, O. 1907. Beiträge zur Fauna des oberen Malm und der unteren Kreide in der
896 argentinischen Cordillere. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie,*
897 *Beilage-Bände*, **21**, 187–236.

898 Hetzinger, S., Pfeiffer, M., Dullo, W.-C., Zinke, J. & Garbe-Schönberg, D. 2016. A change in
899 coral extension rates and stable isotopes after El Niño-induced coral bleaching and
900 regional stress events. *Scientific Reports*, **6**, 32879.

901 Hodgson, W.A. 1966. Carbon and oxygen isotope ratios in diagenetic carbonates from marine
902 sediments. *Geochimica et Cosmochimica Acta*, **30**, 1223–1233.

903 Hoffmann, R. & Stevens, K. 2019. The palaeobiology of belemnites - foundation for the
904 interpretation of rostrum geochemistry. *Biological Reviews*.

- 905 Holmden, C. & Hudson, J.D. 2003. $^{87}\text{Sr}/^{86}\text{Sr}$ and Sr/Ca investigation of Jurassic mollusks
906 from Scotland: implications for paleosalinities and the Sr/Ca ratio of seawater. *GSA*
907 *Bulletin*, **115**, 1249–1264.
- 908 Howell, J.A., Schwarz, E., Spalletti, L.A. & Veiga, G.D. 2005. The Neuquén Basin: an
909 overview. In: Veiga, G.D., Spalletti, L.A., Howell, J.A. & Schwarz, E. (eds.) *The*
910 *Neuquén Basin, Argentina: A case study in sequence stratigraphy and basin dynamics*.
911 Geological Society, London, Special Publications, **252**, 1–14.
- 912 Howlett, P.J. 1989. Late Jurassic-Early Cretaceous cephalopods of eastern Alexander Island,
913 Antarctica. *Special Papers in Palaeontology*, 41, 1–72.
- 914 Hudson, J.D. 1977. Stable isotopes and limestone lithification. *Journal of the Geological*
915 *Society, London*, **133**, 637–660.
- 916 Jenkyns, H.C., Schouten-Huibers, L., Schouten, S. & Sinninghe Damsté, J.S. 2012. Warm
917 Middle Jurassic-Early Cretaceous high-latitude sea-surface temperatures from the
918 Southern Ocean. *Climate of the Past*, **8**, 215–226.
- 919 Kietzmann, D.A., Palma, R.M., Riccardi, A.C., Martín-Chivelet, J. & López-Gómez, J. 2014.
920 Sedimentology and sequence stratigraphy of a Tithonian-Valanginian carbonate ramp
921 (Vaca Muerta Formation): A misunderstood exceptional source rock in the southern
922 Mendoza area of the Neuquén Basin, Argentina. *Sedimentary Geology*, **302**, 64–86.
- 923 Klein, R.T., Lohmann, K.C. & Thayer, C.W. 1996. Bivalve skeletons record sea-surface
924 temperature and $\delta^{18}\text{O}$ via Mg/Ca and $^{18}\text{O}/^{16}\text{O}$ ratios. *Geology*, **24**, 415–418.
- 925 Korte, C. & Hesselbo, S.P. 2011. Shallow marine carbon and oxygen isotope and elemental
926 records indicate icehouse-greenhouse cycles during the Early Jurassic.
927 *Paleoceanography*, **26**, PA4219.
- 928 Korte, C., Hesselbo, S.P., Ullmann, C.V., Dietl, G., Ruhl, M., Schweigert, G. & Thibault, N.
929 2015. Jurassic climate mode governed by ocean gateway. *Nature Communications*, **6**,
930 10015.
- 931 Lawrence, J.R., Drever, J.I., Anderson, T.F. & Brueckner, H.K. 1979. Importance of
932 alteration of volcanic material in the sediments of Deep Sea Drilling Site 323: chemistry,
933 $^{18}\text{O}/^{16}\text{O}$ and $^{87}\text{Sr}/^{86}\text{Sr}$. *Geochimica et Cosmochimica Acta*, **43**, 573–588.
- 934 Lazo, D.G. 2007. Early Cretaceous bivalves of the Neuquén Basin, west-central Argentina:
935 notes on taxonomy, palaeobiogeography and palaeoecology. *Geological Journal*, **42**,
936 127–142.
- 937 Lazo, D.G., Aguirre-Urreta, M.B., Price, G.D., Rawson, P.F., Ruffell, A.H. & Ogle, N. 2008.
938 Palaeosalinity variations in the Early Cretaceous of the Neuquén Basin, Argentina:

939 Evidence from oxygen isotopes and palaeoecological analysis. *Palaeogeography*,
940 *Palaeoclimatology, Palaeoecology*, **260**, 477–493.

941 Leanza, H.A. 1990. Estratigrafía del Paleozoico y Mesozoico anterior a los movimientos
942 intermálmicos en la comarca del Cerro Chachil, Provincia del Neuquén. *Revista de la*
943 *Asociación Geológica Argentina*, **45**, 272–299.

944 Leanza, H.A. 1993. Estratigrafía del Mesozoico posterior a los movimientos intermálmicos en
945 la comarca del Cerro Chachil, Provincia del Neuquén. *Revista de la Asociación*
946 *Geológica Argentina*, **48**, 71–84.

947 Leanza, H.A., Mazzini, A., Corfu, F., Llambías, E.J., Svensen, H., Planke, S. & Galland, O.
948 2013. The Chachil Limestone (Pliensbachian-earliest Toarcian) Neuquén Basin,
949 Argentina: U-Pb age calibration and its significance on the Early Jurassic evolution of
950 southwestern Gondwana. *Journal of South American Earth Sciences*, **42**, 171–185.

951 LeGrande, A.N. & Schmidt, G.A. 2006. Global gridded data set of the oxygen isotopic
952 composition in seawater. *Geophysical Research Letters*, **33**, L12604.

953 Lena, L., López-Martínez, R., Lescano, M., Aguirre-Urreta, B., Concheyro, A., Vennari, V.,
954 Naipauer, M., Samankassou, E., Pimentel, M., Ramos, V.A. & Schaltegger, U. 2019.
955 High-precision U-Pb ages in the early Tithonian to early Berriasian and implications for
956 the numerical age of the Jurassic-Cretaceous boundary. *Solid Earth*, **10**, 1–14.

957 Li, Q., McArthur, J.M., Doyle, P., Janssen, N., Leng, M.J., Müller, W. & Reboulet, S. 2013.
958 Evaluating Mg/Ca in belemnite calcite as a palaeo-proxy. *Palaeogeography*,
959 *Palaeoclimatology, Palaeoecology*, **388**, 98–108.

960 Martinez, M. & Dera, G. 2015. Orbital pacing of carbon fluxes by a ~9-my eccentricity cycle
961 during the Mesozoic. *Proceedings of the National Academy of Sciences*, **112**, 12604–
962 12609.

963 McArthur, J.M., Doyle, P., Leng, M.J., Reeves, K., Williams, C.T., Garcia-Sanchez, R. &
964 Howarth, R.J. 2007. Testing palaeo-environmental proxies in Jurassic belemnites:
965 Mg/Ca, Sr/Ca, Na/Ca, $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$. *Palaeogeography, Palaeoclimatology*,
966 *Palaeoecology*, **252**, 464–480.

967 McArthur, J.M., Howarth, R.J. & Shields, G.A. 2012. Strontium isotope stratigraphy. In:
968 Gradstein, F.M., Ogg, J.G., Schmitz, M. & Ogg, G. (eds.) *The Geological Time Scale*
969 2012. Elsevier, Amsterdam, 127–144.

970 Mouchi, V., de Rafélis, M., Lartaud, F., Fialin, M. & Verrecchia, E. 2013. Chemical labelling
971 of oyster shells used for time-calibrated high-resolution Mg/Ca ratios: A tool for

972 estimation of past seasonal temperature variations. *Palaeogeography, Palaeoclimatology,*
973 *Palaeoecology*, **373**, 66–74.

974 Mutterlose, J., Malkoc, M., Schouten, S., Sinninghe Damsté, J.S. & Forster, A. 2010. TEX₈₆
975 and stable $\delta^{18}\text{O}$ paleothermometry of early Cretaceous sediments: implications for
976 belemnite ecology and paleotemperature proxy applications. *Earth and Planetary Science*
977 *Letters*, **298**, 286–298.

978 Nelson, C.S. & Smith, A.M. 1996. Stable oxygen and carbon isotope compositional fields for
979 skeletal and diagenetic components in New Zealand Cenozoic nontropical carbonate
980 sediments and limestones: A synthesis and review. *New Zealand Journal of Geology and*
981 *Geophysics*, **39**, 93–107.

982 Nunn, E.V. & Price, G.D. 2010. Late Jurassic (Kimmeridgian-Tithonian) stable isotopes
983 ($\delta^{18}\text{O}$, $\delta^{13}\text{C}$) and Mg/Ca ratios: New palaeoclimate data from Helmsdale, northeast
984 Scotland. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **292**, 325–335.

985 Ogg, J.G., Ogg, G.M., Gradstein, F.M. 2016. *A concise geologic time scale*. Elsevier, 240 p.

986 Parent, H. 2006. Oxfordian and Late Callovian ammonite faunas and biostratigraphy of the
987 Neuquén-Mendoza and Tarapacá basins (Jurassic, Ammonoidea, western South-
988 America). *Boletín del Instituto de Fisiografía y Geología*, **76**, 1–70.

989 Parent, H. & Garrido, A.C. 2015. The ammonite fauna of the La Manga Formation (Late
990 Callovian-Early Oxfordian) of Vega de la Veranada, Neuquén Basin, Argentina. *Neues*
991 *Jahrbuch für Geologie und Paläontologie*, **275**, 163–217.

992 Parent, H., Garrido, A.C., Schweigert, G. & Scherzinger, A. 2013. The Tithonian stratigraphy
993 and ammonite fauna of the transect Portada Covunco-Cerrito Caracoles (Neuquén Basin,
994 Argentina). *Neues Jahrbuch für Geologie und Paläontologie*, **269**, 1–50.

995 Parent, H., Garrido, A.C., Scherzinger, A., Schweigert, G. & Fözy, I. 2015. The Tithonian-
996 Lower Valanginian stratigraphy and ammonite fauna of the Vaca Muerta Formation in
997 Pampa Tril, Neuquén Basin, Argentina. *Boletín del Instituto de Fisiografía y Geología*,
998 **86**, 1–96.

999 Parent, H., Garrido, A.C., Brambilla, L. & Alberti, M. 2020. Upper Bathonian ammonites
1000 from Chacay Melehué (Neuquén Basin, Argentina) and the chronostratigraphy of the
1001 Steinmanni Zone. *Boletín del Instituto de Fisiografía y Geología*, **90**, 1–37.

1002 Pfeiffer, M., Zinke, J., Dullo, W.C., Garbe-Schönberg, D., Latif, M. & Weber, M. 2017.
1003 Indian Ocean corals reveal crucial role of World War II bias for twentieth century
1004 warming. *Scientific Reports*, **7**, 14434.

- Podlaha, O.G., Mutterlose, J. & Veizer, J. 1998. Preservation of $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ in belemnite rostra from the Jurassic/Early Cretaceous successions. *American Journal of Science*, **298**, 324–347.
- Price, G.D. 1999. The evidence and implications of polar ice during the Mesozoic. *Earth-Science Reviews*, **48**, 183–210.
- Price, G.D. & Gröcke, D.R. 2002. Strontium-isotope stratigraphy and oxygen- and carbon-isotope variation during the Middle Jurassic-Early Cretaceous of the Falkland Plateau, South Atlantic. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **183**, 209–222.
- Price, G.D. & Sellwood, B.W. 1997. “Warm” palaeotemperatures from high Late Jurassic palaeolatitudes (Falkland Plateau): Ecological, environmental or diagenetic controls? *Palaeogeography, Palaeoclimatology, Palaeoecology*, **129**, 315–327.
- Price, G.D., Sellwood, B.W. & Valdes, P.J. 1995. Sedimentological evaluation of general circulation model simulations for the “greenhouse” Earth: Cretaceous and Jurassic case studies. *Sedimentary Geology*, **100**, 159–180.
- Price, G.D., Baker, S.J., VanDeVelde, J. & Clémence, M.-E. 2016. High-resolution carbon cycle and seawater temperature evolution during the Early Jurassic (Sinemurian-Early Pliensbachian). *Geochemistry, Geophysics, Geosystems*, **17**, 3917–3928.
- Prokoph, A., Shields, G.A. & Veizer, J. 2008. Compilation and time-series analysis of a marine carbonate $\delta^{18}\text{O}$, $\delta^{13}\text{C}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{34}\text{S}$ database through Earth history. *Earth-Science Reviews*, **87**, 113–133.
- Ravelo, A.C. & Hillaire-Marcel, C. 2007. The use of oxygen and carbon isotopes of Foraminifera in Paleoceanography. In: Hillaire-Marcel, C. & de Vernal, A. (eds.) *Proxies in Late Cenozoic Paleoceanography*. Developments in Marine Geology, **1**, 735–764.
- Roche, D.M., Donnadieu, Y., Pucéat, E. & Paillard, D. 2006. Effect of changes in $\delta^{18}\text{O}$ content of the surface ocean on estimated sea surface temperatures in past warm climate. *Paleoceanography*, **21**, PA2023.
- Rubilar, A. 2005. Heterochrony in Middle Jurassic species of *Gryphaea* (Ostreoidea, Gryphaeidae) from southern South America. – *Geologica Acta* 3: 185-203.
- Rubilar, A.E. & Lazo, D.G. 2009. Description of *Aetostreon pilmatuegrossum* sp. nov. from the Lower Cretaceous of Argentina (Neuquén Basin), and significance of the conservative left valve morphology in oysters of the genus *Aetostreon* Bayle. *Cretaceous Research*, **30**, 727–748.

1037 Sadatzki, H., Alberti, M., Garbe-Schönberg, D., Andersen, N., Strey, P., Fortunato, H.,
1038 Andersson, C. & Schäfer, P. 2019. Paired Li/Ca and $\delta^{18}\text{O}$ peaks in bivalve shells from the
1039 Gulf of Panama mark seasonal coastal upwelling. *Chemical Geology*, **529**, 119295.

1040 Scherer, C.M.S. & Goldberg, K. 2007. Palaeowind patterns during the latest Jurassic-earliest
1041 Cretaceous in Gondwana: Evidence from aeolian cross-strata of the Botucatu Formation,
1042 Brazil. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **250**, 89–100.

1043 Schmidt, G.A., Bigg, G.R. & Rohling, E.J. 1999. Global Seawater Oxygen-18 Database -
1044 v1.22. <https://data.giss.nasa.gov/o18data/>

1045 Schwarz, E. & Howell, J.A. 2005. Sedimentary evolution and depositional architecture of a
1046 lowstand sequence set: the Lower Cretaceous Mulichinco Formation, Neuquén Basin,
1047 Argentina. In: Veiga, G.D., Spalletti, L.A., Howell, J.A. & Schwarz, E. (eds.) *The*
1048 *Neuquén Basin, Argentina: A case study in sequence stratigraphy and basin dynamics*.
1049 Geological Society, London, Special Publications, **252**, 109–138.

1050 Sellwood, B.W. & Valdes, P.J. 2006. Mesozoic climates: general circulation models and the
1051 rock record. *Sedimentary Geology*, **190**, 269–287.

1052 Sellwood, B.W., Valdes, P.J. & Price, G.D. 2000. Geological evaluation of multiple general
1053 circulation model simulations of Late Jurassic palaeoclimate. *Palaeogeography,*
1054 *Palaeoclimatology, Palaeoecology*, **156**, 147–160.

1055 Shackleton, N.J. & Kennett, J.P. 1975. Paleotemperature history of the Cenozoic and the
1056 initiation of Antarctic glaciation: oxygen and carbon isotope analyses in DSDP sites 277,
1057 279, 281. *Initial Reports of the Deep Sea Drilling Project*, **29**, 743–755.

1058 Sosdian, S.M., Lear, C.H., Tao, K., Grossman, E.L., O'Dea, A. & Rosenthal, Y. 2012.
1059 Cenozoic seawater Sr/Ca evolution. *Geochemistry, Geophysics, Geosystems*, **13**, Q10014.

1060 Spalletti, L.A., Gasparini, Z., Veiga, G., Schwarz, E., Fernández, M. & Matheos, S. 1999.
1061 Facies anóxicas, procesos deposicionales y herpetofauna de la rampa marina tithoniana-
1062 berriasiana en la Cuenca Neuquina (Yesera del Tromen), Neuquén, Argentina. *Revista*
1063 *Geológica de Chile*, **26**, 109–123.

1064 Spalletti, L.A., Franzese, J.R., Matheos, S.D. & Schwarz, E. 2000. Sequence stratigraphy of a
1065 tidally dominated carbonate-siliciclastic ramp; the Tithonian-Early Berriasian of the
1066 southern Neuquén Basin, Argentina. *Journal of the Geological Society, London*, **157**,
1067 433–446.

1068 Steuber, T. & Veizer, J. 2002. Phanerozoic record of plate tectonic control of seawater
1069 chemistry and carbonate sedimentation. *Geology*, **30**, 1123–1126.

- 1070 Steuer, A. 1897. Argentinische Jura-Ablagerungen. Ein Beitrag zur Kenntnis der Geologie
1071 und Paläontologie der argentinischen Anden. *Palaeontologische Abhandlungen*, **7**, 129–
1072 222.
- 1073 Stevens, G.R. & Clayton, R.N. 1971. Oxygen isotope studies on Jurassic and Cretaceous
1074 belemnites from New Zealand and their biogeographic significance. *New Zealand*
1075 *Journal of Geology and Geophysics*, **14**, 829–897.
- 1076 Suan, G., Mattioli, E., Pittet, B., Mailliot, S. & Lécuyer, C. 2008. Evidence for major
1077 environmental perturbation prior to and during the Toarcian (Early Jurassic) oceanic
1078 anoxic event from the Lusitanian Basin, Portugal. *Paleoceanography*, **23**, PA1202.
- 1079 Suan, G., Mattioli, E., Pittet, B., Lécuyer, C., Suchéras-Marx, B., Duarte, L.V., Philippe, M.,
1080 Reggiani, L. & Martineau, F. 2010. Secular environmental precursors to Early Toarcian
1081 (Jurassic) extreme climate changes. *Earth and Planetary Science Letters*, **290**, 448–458.
- 1082 Suero, T. 1951. Descripción geológica de la Hoja 36c, Cerro Lotena (Neuquén). *Boletín de la*
1083 *Dirección Nacional de Geología y Minería*, **76**, 1–67.
- 1084 Surge, D. & Lohmann, K. 2008. Evaluating Mg/Ca ratios as a temperature proxy in the
1085 estuarine oyster, *Crassostrea virginica*. *Journal of Geophysical Research*, **113**, G02001.
- 1086 Thomas, H. & Mol, J. 2018. Dissolved inorganic carbon, total alkalinity, $\delta^{18}\text{O}\text{-H}_2\text{O}$, $\delta^{13}\text{C}$ -
1087 DIC, temperature, and salinity collected from discrete bottle samples from CCGS
1088 Amundsen in the Beaufort Sea during August and September 2014, PANGAEA.
1089 <https://doi.pangaea.de/10.1594/PANGAEA.886238>, 2018.
- 1090 Torsvik, T.H., Van der Voo, R., Preeden, U., MacNiocaill, C., Steinberger, B., Doubrovine,
1091 P.V., van Hinsbergen, D.J.J., Domeier, M., Gaina, C., Tohver, E., Meert, J.G.,
1092 McCausland, P.J.A. & Cocks, L.R.M. 2012. Phanerozoic polar wander, palaeogeography
1093 and dynamics. *Earth-Science Reviews*, **114**, 325–368.
- 1094 Tynan, S., Opdyke, B.N., Walczak, M., Eggins, S. & Dutton, A. 2017. Assessment of Mg/Ca
1095 in *Saccostrea glomerata* (the Sydney rock oyster) shell as a potential temperature record.
1096 *Palaeogeography, Palaeoclimatology, Palaeoecology*, **484**, 79–88.
- 1097 Ullmann, C.V. & Korte, C. 2015. Diagenetic alteration in low-Mg calcite from macrofossils: a
1098 review. *Geological Quarterly*, **59**, 3–20.
- 1099 Ullmann, C.V., Hesselbo, S.P. & Korte, C. 2013. Tectonic forcing of Early to Middle Jurassic
1100 seawater Sr/Ca. *Geology*, **41**, 1211–1214.
- 1101 Ullmann, C.V., Campbell, H.J., Frei, R. & Korte, C. 2016. Oxygen and carbon isotope and
1102 Sr/Ca signature of high-latitude Permian to Jurassic calcite fossils from New Zealand and
1103 New Caledonia. *Gondwana Research*, **38**, 60–73.

1104 Urey, H.C., Lowenstam, H.A., Epstein, S. & McKinney, C.R. 1951. Measurement of
 1105 paleotemperatures and temperatures of the Upper Cretaceous of England, Denmark, and
 1106 the southeastern United States. *Geological Society of America Bulletin*, **62**, 399–416.

1107 Valdes, P.J. & Sellwood, B.W. 1992. A palaeoclimate model for the Kimmeridgian.
 1108 *Palaeogeography, Palaeoclimatology, Palaeoecology*, **95**, 47–72.

1109 van Hinsbergen, D.J.J., de Groot, L.V., van Schalk, S.J., Spakman, W., Bijl, P.K., Sluijs, A.,
 1110 Langereis, C.G. & Brinkhuis, H. 2015. A paleolatitude calculator for paleoclimate
 1111 studies. *PLoS One*, **10**, e0126946.

1112 Vicente, J.C. 2005. Dynamic paleogeography of the Jurassic Andean Basin: pattern of
 1113 transgression and localisation of main straits through the magmatic arc. *Revista de la*
 1114 *Asociación Geológica Argentina*, **60**, 221–250.

1115 Vicente, J.-C. 2006. Dynamic paleogeography of the Jurassic Andean Basin: pattern of
 1116 regression and general considerations on main features. *Revista de la Asociación*
 1117 *Geológica Argentina*, **61**, 408–437.

1118 Vickers, M.L., Bajnai, D., Price, G.D., Linckens, J. & Fiebig, J. 2019. Southern high-latitude
 1119 warmth during Jurassic-Cretaceous: New evidence from clumped isotope thermometry.
 1120 *Geology*, **47**, 724–728.

1121 Volkheimer, W., Rauhut, O.W.M., Quattrocchio, M.E. & Martinez, M.A. 2008. Jurassic
 1122 paleoclimates in Argentina, a review. *Revista de la Asociación Geológica Argentina*, **63**,
 1123 549–556.

1124 Weaver, A. 1931. Paleontology of the Jurassic and Cretaceous of West Central Argentina.
 1125 *Memoirs of the University of Washington*, **1**, 1–496.

1126 Westermann, G.E.G. & Riccardi, A.C. 1979. Middle Jurassic ammonoid fauna and
 1127 biochronology of the Argentine-Chilean Andes. Part II: Bajocian Stephanocerataceae.
 1128 *Palaeontographica Abt. A*, **164**, 85–188.

1129 Wierzbowski, H. 2002. Detailed oxygen and carbon isotope stratigraphy of the Oxfordian in
 1130 Central Poland. *International Journal of Earth Sciences*, **91**, 304–314.

1131 Wierzbowski, H. 2004. Carbon and oxygen isotope composition of Oxfordian-Early
 1132 Kimmeridgian belemnite rostra: palaeoenvironmental implications for Late Jurassic seas.
 1133 *Palaeogeography, Palaeoclimatology, Palaeoecology*, **203**, 153–168.

1134 Wierzbowski, H. & Joachimski, M. 2007. Reconstruction of late Bajocian-Bathonian marine
 1135 palaeoenvironments using carbon and oxygen isotope ratios of calcareous fossils from the
 1136 Polish Jura Chain (central Poland). *Palaeogeography, Palaeoclimatology,*
 1137 *Palaeoecology*, **254**, 523–540.

- Wierzbowski, H., Dembicz, K. & Praszker, T. 2009. Oxygen and carbon isotope composition of Callovian-Lower Oxfordian (Middle-Upper Jurassic) belemnite rostra from Central Poland: A record of a Late Callovian global sea-level rise? *Palaeogeography, Palaeoclimatology, Palaeoecology*, **283**, 182–194.
- Wierzbowski, H., Anczkiewicz, R., Pawlak, J., Rogov, M.A. & Kuznetsov, A.B. 2017. Revised Middle-Upper Jurassic strontium isotope stratigraphy. *Chemical Geology*, **466**, 239–255.
- Wierzbowski, H., Bajnai, D., Wacker, U., Rogov, M.A., Fiebig, J. & Tesakova, E.M. 2018. Clumped isotope record of salinity variations in the Subboreal Province at the Middle-Late Jurassic transition. *Global and Planetary Change*, **167**, 172–189.
- Zachos, J.C., Stott, L.D. & Lohmann, K.C. 1994. Evolution of early Cenozoic marine temperatures. *Palaeoceanography*, **9**, 353–387.
- Zhou, J., Poulsen, C.J., Pollard, D. & White, T.S. 2008. Simulation of modern and middle Cretaceous marine $\delta^{18}\text{O}$ with an ocean-atmosphere general circulation model. *Paleoceanography*, **23**, PA3223.

Figure captions

Fig. 1. Palaeogeographic reconstructions for the Neuquén Basin during the Jurassic (modified after Howell et al., 2005). It is still debated whether the volcanic arc separating the basin from the open ocean consisted of a chain of individual islands (A; Howell et al., 2005) or constituted a continuous landmass pierced by only one seaway (B; Vicente, 2005). The study areas are around Zapala (southern sections) and Chos Malal (northern sections).

Fig. 2. Schematic map of the study area in the Neuquén Basin (shaded in grey) in southwestern South America showing the location of the studied sections near Chos Malal in the north and Zapala in the south.

Fig. 3. General lithostratigraphic framework and major facies types for the Jurassic and Lower Cretaceous of the Neuquén Basin (modified after Howell et al., 2005).

Fig. 4. $\delta^{18}\text{O}$ versus $\delta^{13}\text{C}$ values of fossils and sediment samples of the northern sections near Chos Malal and the southern sections near Zapala.

Fig. 5. Results of stable isotope ($\delta^{18}\text{O}$, $\delta^{13}\text{C}$) and element (Mg/Ca, Sr/Ca) analyses of well-preserved fossils of the northern sections near Chos Malal. Temperatures for $\delta^{18}\text{O}_{\text{shell}}$ values were reconstructed with the equation of Anderson and Arthur (1983) and a $\delta^{18}\text{O}_{\text{sea}}$ value of -1 ‰ for an ice-free Jurassic world as suggested by Shackleton and Kennett (1975). Temperatures for Mg/Ca ratios were calculated with the equations of Mouchi et al. (2013) for oysters and Nunn and Price (2010) for belemnites. Main facies types are based on Howell et al. (2005) and the age model is based on Ogg et al. (2016).

Fig. 6. Results of stable isotope ($\delta^{18}\text{O}$, $\delta^{13}\text{C}$) and element (Mg/Ca, Sr/Ca) analyses of well-preserved fossils of the southern sections near Zapala. Temperatures for $\delta^{18}\text{O}_{\text{shell}}$ values were reconstructed with the equation of Anderson and Arthur (1983) and a $\delta^{18}\text{O}_{\text{sea}}$ value of -1 ‰ for an ice-free Jurassic world as suggested by Shackleton and Kennett (1975). Temperatures for Mg/Ca ratios were calculated with the equations of Mouchi et al. (2013) for oysters and Nunn and Price (2010) for belemnites. Main facies types are based on Howell et al. (2005) and the age model is based on Ogg et al. (2016).

Fig. 7. A, B. Results of high-resolution stable isotope ($\delta^{18}\text{O}$, $\delta^{13}\text{C}$) analysis of a specimen of *Gryphaea* sp. (MOZ-PI 11847/13) from the Lower Oxfordian of the Vega de la Veranada near Chos Malal. Temperatures were calculated with the equation of Anderson and Arthur (1983) and a $\delta^{18}\text{O}_{\text{sea}}$ value of -1 ‰ (Shackleton and Kennett, 1975) and alternatively -6.5 ‰. **C.** $\delta^{18}\text{O}$ versus $\delta^{13}\text{C}$ values of the oyster used for the high-resolution stable isotope analysis shows a weak positive correlation.

Fig. 8. Sr/Ca_{sea} ratios reconstructed from fossils of the Neuquén Basin compared to data of Ullmann et al. (2013, 2016) and the $^{87}\text{Sr}/^{86}\text{Sr}$ curve of Wierzbowski et al. (2017).

Fig. 9. Possible processes affecting the $\delta^{18}\text{O}_{\text{sea}}$ and $\delta^{18}\text{O}_{\text{shell}}$ values in marginal seas. Influx of marine waters from the open ocean can change the $\delta^{18}\text{O}_{\text{sea}}$ values positively or negatively depending on their origin. Diagenetic alteration generally leads to a decrease in $\delta^{18}\text{O}_{\text{shell}}$ values. Weathering of volcanoclastics on the sea floor, sea-ice formation, and freshwater influx lower $\delta^{18}\text{O}_{\text{sea}}$ values. In contrast, evaporation leads to an increase in $\delta^{18}\text{O}_{\text{sea}}$ values.

Fig. 10. Compilation of available temperature reconstructions for the Jurassic and Early Cretaceous of South America based on Mg/Ca ratios and $\delta^{18}\text{O}_{\text{shell}}$ values of bivalves,

1206 belemnites, and brachiopods of Argentina and Chile (data combined from the present study
1207 and Bowen, 1963; Gómez-Dacal et al., 2018; Alberti et al., 2019b). Temperatures for Mg/Ca
1208 ratios were calculated with the equations of Mouchi et al. (2013) for oysters and brachiopods
1209 and Nunn and Price (2010) for belemnites. Temperatures for $\delta^{18}\text{O}_{\text{shell}}$ values were
1210 reconstructed with the equation of Anderson and Arthur (1983) and a $\delta^{18}\text{O}_{\text{sea}}$ value of -1‰
1211 for an ice-free Jurassic world as suggested by Shackleton and Kennett (1975). Temperatures
1212 calculated for bivalves of the northern sections were tentatively corrected by using more
1213 negative $\delta^{18}\text{O}_{\text{sea}}$ values to acknowledge a likely freshwater influence in this region. The trend
1214 lines for benthic taxa and belemnites are based on average values and only serve as
1215 orientation and broad indicators for potential long-term trends as some time intervals are not
1216 covered by data. The age model is based on Ogg et al. (2016).