

Crops and Soils Research Paper

Cite this article: Salinger MJ *et al.* (2020). Linking crop yields in Tuscany, Italy, to large-scale atmospheric variability, circulation regimes and weather types. *The Journal of Agricultural Science* **158**, 606–623. <https://doi.org/10.1017/S0021859620001021>

Received: 1 April 2020
Revised: 9 November 2020
Accepted: 18 November 2020
First published online: 7 January 2021

Key words:

Climate change; climate variability; dry conditions; teleconnections; warm season


Author for correspondence:

Alessandro Messeri,
E-mail: alessandro.messeri@unifi.it

© The Author(s), 2021. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

CAMBRIDGE
UNIVERSITY PRESS

Linking crop yields in Tuscany, Italy, to large-scale atmospheric variability, circulation regimes and weather types

M. James Salinger¹, Anna Dalla Marta¹, Giovannagelo Dalu²,
Alessandro Messeri^{3,4} , Marina Baldi², Gianni Messeri^{3,5}, Roberto Vallorani^{3,5},
Marco Morabito^{3,4}, Simone Orlandini^{1,4}, Filiberto Altobelli⁶ and Leonardo Verdi¹

¹Department of Agriculture, Food, Environment and Forestry (DAGRI), University of Florence, 50144 Florence, Italy; ²CNR Institute of BioEconomy, CNR-IBE, 00185 Rome, Italy; ³CNR Institute of BioEconomy, CNR-IBE, 50019 Florence, Italy; ⁴Centre of Bioclimatology (CIBIC), University of Florence, 50144 Florence, Italy; ⁵LaMMA Consortium, Consortium CNR Institute of BioEconomy (CNR-IBE) and Tuscany Region, 50019 Sesto Fiorentino, Italy and ⁶CREA Research Centre for Agricultural Policies and Bioeconomy, Via Po 14, 00198 Roma, Italy

Abstract

The paper presents results from a study examining the relationship between large-scale modes of climate variability with the fluctuations in the yield of barley, durum wheat, olives and sunflower crops in Tuscany, Italy. In particular, the blocking circulation over the growing season, with associated hot and dry conditions, decreased yield for olive crops, barley and durum wheat. The teleconnections analysed in this study are the winter North Atlantic Oscillation (NAO) and the Summer North Atlantic Oscillation (SNAO); the West African Monsoon (WAM) and the Intertropical Front (ITF); and although NAO, SNAO, ITF and WAM are not strictly related to each other, the values of these indices are strongly related to the atmospheric circulation regimes and related weather types. Thus, they have an impact on precipitation and temperature patterns in Italy and on yields of important crops in Tuscany. Results show that the large-scale temperate and tropical variability directly influences the crop yield through three main circulation regimes. These patterns illustrate the importance of the large-scale modes, which, together with the associated weather types, have an impact directly on Tuscan crop yields; both barley and olive yields decline significantly when the ITF is further north with warmer and drier conditions in Italy.

Introduction

Several studies have shown that the expected growth of the world's population to almost 9 billion by 2050, dietary changes, and increase in biofuel consumption will increase the agricultural demand by some 50% compared to 2013 (FAO, 2017; Hoegh-Guldberg *et al.*, 2018; Searchinger *et al.*, 2019). At the same time, satisfying increased agricultural demand is a challenge due to several factors, such as political instabilities (Macours and Swinnen, 2000), pest and disease outbreaks (Oerke, 2006), decrease of available arable land, soil salinization, water scarcity, deforestation, land degradation and changes of climate conditions (Ciais *et al.*, 2005; Herbert *et al.*, 2015).

Different authors (Chen *et al.*, 2004; Osborne and Wheeler, 2013; Ray *et al.*, 2015; Najafi *et al.*, 2019) have demonstrated that climate variability and their teleconnections have led to changes in maize, rice, wheat and soybean crop yields at different scales. Their results show that 60% of the crop yield variability is related to regional climate, whilst global climate variability accounts for roughly a third (32–39%) of the total crop yield variability.

Specifically, the agriculture in Europe is characterized by high-quality production and by a variety of locally adapted farming systems; therefore, it is highly vulnerable to regional climate variability and localized extreme meteorological events.

Recently, Nobre *et al.* (2019) have shown the importance of climate variability in driving the interannual changes of meteorological variables. These authors have shown that teleconnection indices (ENSO, North Atlantic Oscillation (NAO), Scandinavian Pattern, East Atlantic Pattern, and East Atlantic/West Russian Pattern) have an influence on the regional precipitation patterns and on the regional agricultural production. Considering NAO, for example, although its influence is more evident for north European areas, some studies demonstrate that NAO also impacts on weather variables in the Mediterranean at a local scale (Del Pozo *et al.*, 2019), which are correlated with grapevine yield and quality (Dalla Marta *et al.*, 2010), wheat yield (Dalla Marta *et al.*, 2011) and olive pollen production (Avolio *et al.*, 2008).

The climatic trend has also a sensitive effect on Mediterranean tree crops, and winter weather plays a key role. When temperature drops below -10°C , olive trees are stressed and start

showing physiological changes that, in extreme cases, can lead to plant death (Barranco and Ruiz, 2005). Nevertheless, the most vulnerable phase of olives is flowering. Bonofiglio *et al.* (2008) confirmed that 30°C is a threshold above which the development of the ovary pollen tube can be inhibited. In Italy, flowering occurs in late spring when temperatures rarely reach those extreme values. Haworth *et al.* (2018) observed that when temperatures exceed 40°C, plant photosynthetic and vegetative activity stops.

On the other hand, during recent decades, with the aim of responding to an increasing demand for quality and quantity of crop yields, farmers have made great progress in agricultural techniques. In particular, the agricultural technology, the introduction of new cultivars and the dynamic changes in fertilization practices can affect the most important crop yields (Fontana *et al.*, 2015; Agnolucci and De Lipsis, 2019). Parent *et al.* (2018) have shown how farmer's adaptation measures and strategies can reduce the maize yield gradient among European sub-regions. An increase in maize production can be obtained by using genetic variation of the crop cycle duration. Sunflower, instead, despite its high tolerance to water stress can increase yields by 53% when fully irrigated (Demir *et al.*, 2006). In particular, the most sensitive phase to water scarcity is flowering, followed by seed formation (Yawson *et al.*, 2011; Verdi *et al.*, 2019). However, due to different factors such as climatic areas, crop variety and agricultural management strategies, and irrigation strategies, sunflower has a wider water consumption from 200 to 900 mm (Erdem *et al.*, 2001) that require specific consideration to ensure high yields. Thus, these adaptation measures strongly depend on a good knowledge of the regional climate variability. Numerous studies have shown, for example, that an appropriate irrigation of olive crops can improve yields and could be considered as a mitigation strategy for climate change (Iniesta *et al.*, 2009). Goldhamer *et al.* (1994) observed an increase in fruit yields of roughly 50% in irrigated system with respect to rainfed. For that reason, many studies show that much of yield variation between years occurs because of differences in the agricultural technology used, the effect of climate is difficult to detect from the raw yield data (Lu *et al.*, 2017) and detrending of crop yield data is required. Several approaches are available to detrend a time series of data such as linear, non-linear, first-differencing. In our case, the linear approach has been used on barley, durum wheat, olives and sunflower, but no significant trend was identified. In light of this result, it is assumed that there have been no significant changes in these crops yield attributable to change in agriculture technology. In this sense, several authors observed how the climatic trend represents a factor that can significantly modify the agricultural productions (Dalla Marta *et al.*, 2010; Dalla Marta *et al.*, 2011; Fontana *et al.*, 2015). This is confirmed by Teasdale and Cavigelli (2017) reporting as yield trends between conventional and organic agriculture, although with higher values in the conventional, follow the same trend driven by rainfall and temperature conditions. In a recent study investigating the links between the large-scale atmospheric circulation and European crop yields, Ceglar *et al.* (2017) affirmed that technological improvements can only mask the effect of interannual variability on the trend of crop yields. Similarly, Agnolucci and De Lipsis (2019) stated that climatic trend represents the main affecting factor on crop yields, significantly exceeding other factors such as the use of fertilizers, genetic evolution, irrigation, etc.

The strong climate variability in the European region also results in a different vulnerability to climate change (Orlandini

et al., 2008, 2011). Bartolini *et al.* (2013), Vergni and Todisco (2011) and Orlandini *et al.* (2008) affirm that there is an increasing trend of precipitation in North Europe, while southern Europe is currently affected by a decrease in the annual precipitation and by a change in seasonal rainfall patterns. Brunetti *et al.* (2004) observed that in Italy, as for other Mediterranean areas (Alpert *et al.*, 2002), there is a decrease in the number of days with precipitations, despite an increase in rainfall intensity. This trend was also confirmed by studies carried out in Tuscany region by Bartolini *et al.* (2013, 2018), who analysed temporal and spatial changes in precipitation from 1955 to 2013. The study showed how the effects of climate change are evident and have led to a variation in the rainfall regime in recent decades. In particular, a significant decrease in annual, winter and spring rainfall was observed in north-western areas and north-eastern Apennine Mountains but also a significant decrease in annual, winter and spring wet days was observed in all Tuscany areas. Crop phenology and yields are strictly related to climatic conditions and therefore they are sensitive indicators of climate change.

These variations in climate have an influence on the cropping systems by increasing nutrient leaching, runoff and soil erosion and by altering crop-specific water consumption, which are increasingly difficult to satisfy, especially for rainfed cultivations. Hence, agriculture must adopt new irrigation strategies to guarantee high and constant yields and in order to mitigate deleterious effects on the environment (i.e. soil degradation).

Concerning temperatures, a higher and faster increase of daily minimum than of daily maximum has been observed (IPCC, 2007). This situation results in a reduction of the mean diurnal temperature range (ΔTR) with potentially detrimental effects on evaporation and transpiration processes (Vergni *et al.*, 2020). Nevertheless, this tendency is not homogeneous, and in Italy, ΔTR shows a large spatial variability and a positive tendency in the period 1865–1996 (Brunetti *et al.*, 2004). This latter finding is related to the increasing occurrence of subtropical anticyclones over the western Mediterranean region producing clearer skies. On the contrary, Todisco and Vergni (2008) found a significant decrease in ΔTR in Central Italy, approximately from the half of the last century. However, there is a general agreement on the severe effects on Italian agriculture caused by the combined effect of global warming and precipitation.

Different studies have shown that average crop yields from Italian agriculture decreased by 5% over the last 30 years (Moore and Lobell, 2015). In extreme cases, as for barley and winter wheat, yield losses were up to 10%. Moreover, Ray *et al.* (2015) affirmed that from 31 to 51% of wheat yield variability in Italy was caused by climate change. Orlandini *et al.* (2008) and Kukul and Irmak (2018) observed that climate change and variability are affecting crop yields in different ways based on the environmental zones. In particular, the main negative effects in Mediterranean countries are related to a number of factors, namely the reduction of growing season, seasonal climatic variability, drought, heat stresses, hailstorm, pest and disease attacks, weed development, soil erosion and nitrogen losses.

In this study, we analyse some teleconnection indices (Table 1), in particular: the NAO, Summer North Atlantic Oscillation (SNAO), the Intertropical Front (ITF) and the West African Monsoon (WAM) on interannual to decadal time scales, in order to relate their variability to the summer crop yields in Italy. The choice of these teleconnection indices is motivated by their impact on the central Mediterranean climate. When the winter NAO is in its negative phase, there is a precipitation increase

Table 1. Teleconnection indices mentioned in the paper

| Abbreviation | Full name | Description |
|--------------|-----------------------------------|---|
| NAO | North Atlantic Oscillation | It is based on the surface sea-level pressure difference between the subtropical (Azores) high and the subpolar low |
| SNAO | Summer North Atlantic Oscillation | It is NAO during the summer period |
| ITF | Intertropical Front | It separates the wedge of warm moist southwesterly monsoon flow off the tropical Atlantic from much hotter and very dry northeasterly wind from the Sahara Desert |
| WAM | West African Monsoon | |
| ITCZ | Intertropical Convergence Zone | It is a belt of low pressure which circles the Earth generally near the equator where the trade winds of the Northern and Southern Hemispheres come together |
| SAM | South Asian Monsoon | It is among several geographically distributed global monsoons and it affects the Indian subcontinent |
| SHL | Saharan Heat Low | It is an important driver of the West African Monsoon across a range of timescales of variability |

over Italy, with the opposite occurring during its positive phase (Hurrell, 1995; Hurrell *et al.*, 2003). While, when the summer SNAO is in its positive phase, there is a precipitation increase over Italy, with the opposite occurring during the negative phase (Folland *et al.*, 2009; Mariotti and Dell'Aquila, 2012). The WAM and its ITF have an influence on the southern Mediterranean climate patterns (Chen, 2005; Alpert *et al.*, 2006; Gaetani and Fontaine, 2013).

Rodwell and Hoskins (1996) have examined the subsidence of the South Asian Monsoon (SAM) over the eastern Mediterranean Sea and over North Africa; in periods of SAM's strong activity, this subsidence can reach the central Mediterranean region. Whilst Chen (2005) has examined the formation of the mid-tropospheric high over Libya; this high, in periods of WAM's strong activity, can invade the central Mediterranean region. The WAM index (Li and Zeng, 2005) is strongly related to the summer precipitation in the region 10°N–20°N and 30°W–20°E, with teleconnections with Mediterranean region (Gaetani and Fontaine, 2013). WAM's dynamics are rather complex, although African Outgoing Longwave Radiation (OLR) relates well with the 500 hPa heights over the central Mediterranean. The variability of the ITF ranges between 17°N and 21°N, and it is linked to the latitudinal position of WAM (Lélé and Lamb, 2010). WAM originates in the Gulf of Guinea when the Intertropical Convergence Zone (ITCZ) makes its arrival, to establish itself at 10°N–12°N in its mature phase (Thornicroft *et al.*, 2011). In this phase, the southerly desert winds driven by the Saharan Heat Low (SHL) encounter the moist northerly monsoonal winds (Mathbout *et al.*, 2019). Thus, the precipitation over the Sudan-Sahel region is positively related to the ITF's latitude (Lélé and Lamb, 2010). When the ITF is in its northernmost position, the Libyan high resides over the southern-central Mediterranean Sea, bringing hot and dry conditions over Italy.

Italian climate variability is related to these large-scale features, which have, in turn, an impact on the weather types over the Italian peninsula (Hess and Brezowsky, 1952; Lamb, 1972). In this work, we adopt a software package developed in the framework of COST 733 action 'Harmonization and Applications of Weather Type classification for Europe Regions', which offers several classification methods (Philipp *et al.*, 2014; Huth *et al.*, 2015). The weather type classification used in this study is based on two methods for temperature and precipitation: a method based on principal component analysis (the T-mode PCA obliquely rotated

(PCT)), and an algorithm for optimizing partitions (simulated annealing), successfully used in previous works (Messeri *et al.*, 2015, 2018; Salinger *et al.*, 2015; Hidalgo and Jouglá, 2018; Manzanás *et al.*, 2018; Vallorani *et al.*, 2018).

The aim of the study is to investigate the relationship between climate variability and the fluctuations in the yield of barley, durum wheat, olives and sunflower crops in Tuscany, Italy.

Materials and methods

Indices of large-scale climate variability (NAO, SNAO, WAM and ITF)

SNAO values were obtained from the Hadley Centre, United Kingdom Meteorological Office, whilst NAO values were taken from the NOAA Climate Prediction Center (<https://www.cpc.ncep.noaa.gov/data/teledoc/nao.shtml>). WAM index values were obtained from the NOAA Climate Prediction Center (<http://ljp.gcess.cn/dct/page/65579>) (Li and Zeng, 2005). ITF index values were taken from <https://www.cpc.ncep.noaa.gov/products/international/itf/itcz.shtml>. Then, these large-scale modes were correlated with both weather regimes and crop yields in Tuscany.

Weather regimes: Italian weather types

The Tuscany climate has different characteristics from separate areas, being influenced both by the sea that bathes the region to the west, and by the morphology of the Apennines that delimits the territory both to the north and to the east. These differences are reflected in the spatial and temporal distribution of total precipitation (Appendix 1). Plain and hill areas, where barley, durum wheat and olive are concentrated, have annual cumulative precipitation between 1000 mm Versilia Coast – 1300 mm in Lucca and Garfagnana areas – and 400 mm southern coast – 600 mm in Siena and Arezzo areas. As regards the precipitation temporal distribution, they are concentrated in October–December as well as during the spring (especially in April). The highest total rainfall is recorded in November with values between 150 mm (Versilia hill areas) and 80 mm (Grosseto and Siena plane areas). The circulation regimes and weather types can be extremely useful in understanding the spatial and temporal distribution of precipitation over a heterogeneous territory such as Tuscany.

The weather types used in this work are constructed following Vallorani *et al.* (2018), by using mean sea-level pressure from the




Fig. 1. Colour online. Circulation type classification for Italy based on PCT methods.

NCEP-DOE Reanalysis 2 (<https://www.esrl.noaa.gov/psd/>) with a resolution of $2.5^\circ \times 2.5^\circ$ for the period 1979–2015. The nine weather types (WT classes), calculated with PCT method, were then combined in three classes of circulation regimes: cyclonic, anti-cyclonic (blocking) and zonal (westerly) (Fig. 1).

Crop data

Crop data, cultivated areas and production, were obtained from the database of the Italian Institute of Statistics (<http://dati.istat.it/>). We use data for the period 1981–2018 (barley and durum wheat) and for the period 2006–2018 (olives and sunflower) for the Tuscany region, and yields per hectare for each crop were calculated. ISTAT data are based on direct checks of local experts on the territory, as well as on indications from external sources (i.e. professional bodies and producer associations, administrative sources). The local trends of the main crops (arable, agricultural and fodder groups) are recorded annually with insights on vines and olives for wine and oil production, respectively.

Statistical analysis

To investigate associations between crop yields with climate indices and teleconnections and weather types, Pearson's correlation analysis has been performed between the crop and climate/weather data sets for various relevant months and seasons. Significance at the 5 and 1%, and sometimes 10% levels of confidence was identified for further exploration.

Results

Relationships between weather types and precipitation in Tuscany

The 'blocking' weather types are characterized by precipitation lower than the climatological average throughout the March–

October period. The 'zonal' regime has generally low rainfall with WT8 drier than WT6, particularly in September (Fig. 2). In most months, WT6 shows precipitation higher than the average limited to north-west areas; on the contrary, WT8 has negative anomalies. With cyclonic weather types (WT2, WT4, WT7 and WT9), positive anomalies prevail with few exceptions. During the sowing period or in the first phenological phase of barley, durum wheat and sunflower (March and April), the precipitation ranges between less than 20 mm up to 50 mm in the blocking regimes, from 20 mm up to 80 mm in the zonal, and from 30–40 mm up to 100 mm for the trough regime weather types (Figs 3 and 4). During the subsequent phenological phases and the harvest period (June, July, August, September and October), the precipitation ranges between 0 and 40 mm for blocking, 0 and 100 mm with the maximum values in October and September occurring with WT6. For trough types, the minimum values occurred in July with a few mm in WT4 (Fig. 5); in contrast, the maximum values occurred in October for WT4 and WT7 with more than 200 mm (Fig. 6). Additional maps for all months of the cropping year are available in Appendix 2.

Teleconnections between large-scale patterns of climate variability, circulation regimes and weather types

Large scale

For temperate variability, the NAO and SNAO were naturally correlated significantly (Table 2) for July, and July/August. In the tropics, both WAM and SHL contribute to the positioning of the ITF, with a major role played by SHL during the period 1979–2010. This is when the SHL intensity increased the temperature gradient between the Gulf of Guinea and the Sahara (Lavaysse *et al.*, 2016). For temperate/tropical interactions, there is a very strong relationship between the July ITF position and the SNAO (1% significance). With a positive SNAO, the location of higher sea-level pressures over northern Europe allows a more




Fig. 2. Colour online. Precipitation anomalies of September for Italy produced by PCT circulation types with a resolution of 25 km. The anomalies refer to the 1981–2010 base period.




Fig. 3. Colour online. Total precipitation of March for Italy produced by PCT circulation types with a resolution of 25 km. The anomalies refer to the 1981–2010 base period.

northern location of the ITF, while there is almost no correlation between NAO and ITF.

Weather regimes and types

Circulation regimes and weather types reveal significant linkages (Table 3). Those with temperate modes of variability were strong. The links of NAO with blocking, especially over central Europe

(WT5 and WT3) in June, July and August (JJA), were significantly high. Anticyclones over central Europe were prominent during positive NAO events, but not further west (WT1). Zonal westerlies (WT6) tended to be absent especially in the late summer of July, August and September (JAS). Similarly, with positive SNAO episodes, although lack of zonal flow (WT6) occurred with easterly circulation over the western central Mediterranean.




Fig. 4. Colour online. Total precipitation of April for Italy produced by PCT circulation types with a resolution of 25 km. The anomalies refer to the 1981–2010 base period.




Fig. 5. Colour online. Total precipitation of July for Italy produced by PCT circulation types with a resolution of 25 km. The anomalies refer to the 1981–2010 base period.

The tropical ITF teleconnections occurred only in late summer JAS. This was amazingly strong in August ($r = 0.89$) with WT7, when the ITF was at its northernmost position, indicating more cyclonic circulation regimes (troughing) over southern central Europe when the ITF was further south than normal. In this same month, WT1 was more prevalent when the ITF

was displaced northward. For JJA, troughing had a tendency towards high frequencies when the ITF was displaced south. When the WAM was strong, blocking was prevalent over western central Europe (WT1) compared with central Europe (WT5), and sometimes more zonal flow in early summer (WT6).




Fig. 6. Colour online. Total precipitation of October for Italy produced by PCT circulation types with a resolution of 25 km. The anomalies refer to the 1981–2010 base period.

Table 2. Teleconnections between the North Atlantic Oscillation (NAO), Summer North Atlantic Oscillation (SNAO), Intertropical Front (ITF) and the West African Monsoon (WAM)

| ITF | NAO | WAM | SNAO |
|-----|---------|-------|---------|
| a | −0.02 | 0.06 | |
| b | 0.12 | 0.00 | 0.60*** |
| c | −0.03 | 0.19 | −0.04 |
| d | −0.29 | 0.03 | |
| e | −0.31 | 0.07 | −0.13 |
| f | −0.32 | −0.01 | |
| g | −0.12 | −0.05 | |
| h | −0.24 | −0.01 | |
| NAO | SNAO | WAM | |
| a | | −0.06 | |
| b | 0.46** | −0.09 | |
| c | 0.39 | −0.19 | |
| d | | −0.09 | |
| e | 0.54** | −0.08 | |
| f | | −0.18 | |
| g | | 0.21 | |
| h | | −0.23 | |
| WAM | SNAO | | |
| b | −0.36** | | |
| c | −0.34** | | |
| e | −0.43* | | |

Period: a June; b July; c August; d September; e July–August; f June–July–August; g July–August–September; h June–July–August–September.
P levels (**P* < 0.05, ***P* < 0.01, ****P* < 0.001).

Links between climate variability and crop yields

All the crops examined indicated a significant relationship between yield and large-scale circulation (Table 4). Barley yields are clearly linked to the latitude of the ITF. During summer, the further north (south) the ITF was, the lower (higher) the yield. In contrast, durum wheat yields were weakly associated with the latitude of the ITF.

Olive crops had highly significant correlations (1%) with the latitude of the ITF – higher yields occurred when the ITF was further south (north). Sunflower yields displayed some teleconnections with the WAM. When this feature was more vigorous, yields were lower.

Effects of circulation regimes and weather types on crop yields

Depending on the time of year, the circulation regimes and weather types had impacts on winter and spring crop yields (Table 5). For barley yields, WT6 and WT9 during the crop maturation phase (May–June), were the most important types. WT6, zonal flow over Europe, produces lower precipitation over Tuscany (Fig. 2), whereas WT9 with summer troughs in the central Mediterranean give higher precipitation, increasing barley yields. For durum wheat, troughing during the ripening period was especially important (WT2) increasing moisture whilst blocking (WT5) led to lower yields. Troughing regime prevalence increased yields. For warm season crops (olives and sunflowers), the important regimes and types differed. The presence or absence throughout the season of WT2 was important – north easterlies overall, bringing cooler drier conditions in Tuscany favoured increased sunflower yields. Conversely, lack of WT8 and the zonal regime was detrimental between April and June, where there are very significant correlations (>1%). Easterly circulation tends to be warmer and drier.

Table 3. Relationships between the NAO, SNAO, WAM and the ITF

| WT | a | | | b | | | | c | | | | d | | |
|----|---------|----------|----------|---------|----------|---------|---------|---------|----------|---------|---------|----------|---------|----------|
| | NAO | WAM | ITF | NAO | SNAO | WAM | ITF | NAO | SNAO | WAM | ITF | NAO | WAM | ITF |
| 1 | -0.34** | 0.23 | 0.09 | -0.13 | -0.29 | 0.43*** | -0.11 | -0.23 | -0.36** | 0.36** | 0.40** | -0.10 | 0.34** | 0.01 |
| 2 | -0.30* | 0.02 | -0.23 | -0.39** | -0.08 | 0.17 | -0.02 | -0.34** | -0.05 | 0.28* | 0.07 | -0.23 | 0.15 | -0.12 |
| 3 | 0.47** | -0.21 | 0.25* | 0.23 | 0.14 | -0.30* | 0.22 | 0.35** | 0.16 | -0.20 | -0.20 | 0.20 | -0.12 | -0.44*** |
| 4 | -0.18 | 0.00 | 0.10 | -0.01 | -0.08 | -0.03 | -0.26* | 0.23 | 0.22 | -0.19 | -0.01 | -0.44*** | -0.29* | 0.19 |
| 5 | 0.61*** | -0.04 | -0.17 | 0.43** | 0.38** | -0.33** | 0.19 | 0.41** | 0.38** | -0.42** | -0.16 | 0.08 | 0.13 | -0.07 |
| 6 | -0.29* | 0.28* | 0.03 | -0.20 | -0.45*** | 0.16 | 0.10 | 0.01 | -0.28* | 0.37** | 0.22 | -0.20 | -0.54** | 0.37** |
| 7 | 0.09 | -0.08 | -0.07 | -0.32** | -0.30* | 0.04 | -0.04 | -0.05 | -0.09 | 0.18 | -0.89** | -0.40** | -0.10 | 0.19 |
| 8 | -0.30 | -0.06 | 0.14 | -0.03 | 0.02 | 0.18 | 0.10 | -0.32** | -0.20 | -0.08 | -0.02 | 0.07 | 0.10 | -0.02 |
| 9 | 0.18 | -0.27 | 0.00 | 0.13 | 0.22 | -0.22 | -0.15 | 0.34** | 0.17 | -0.16 | 0.07 | 0.28* | 0.24 | 0.00 |
| Tr | -0.31* | -0.22 | -0.14 | -0.34** | -0.02 | 0.03 | -0.34** | -0.06 | 0.13 | 0.09 | -0.05 | -0.18 | 0.07 | 0.23 |
| Bl | 0.55*** | 0.00 | -0.07 | 0.48** | 0.23 | -0.19 | 0.18 | 0.38** | 0.15 | -0.21 | 0.03 | 0.21 | 0.24 | -0.46*** |
| Zo | -0.40 | 0.17 | 0.21 | -0.31* | -0.29* | 0.25 | 0.06 | 0.26 | -0.31** | 0.14 | 0.01 | -0.05 | -0.43 | 0.23 |
| | e | | | | f | | | g | | | h | | | |
| | NAO | SNAO | WAM | ITF | NAO | WAM | ITF | NAO | WAM | ITF | NAO | WAM | ITF | |
| 1 | -0.35** | -0.43 | 0.49*** | 0.09 | -0.36** | 0.62*** | 0.00 | -0.41** | 0.60*** | 0.15 | -0.49** | 0.64*** | 0.09 | |
| 2 | -0.31 | -0.08** | 0.20 | -0.17 | -0.31 | 0.31 | -0.24** | -0.12 | 0.32** | -0.07 | -0.25 | 0.40** | -0.06 | |
| 3 | 0.28* | 0.22 | -0.24 | 0.08 | 0.43*** | -0.41** | 0.08 | 0.27 | -0.22 | 0.03 | 0.25 | -0.44*** | 0.02 | |
| 4 | 0.06 | 0.14 | -0.22 | -0.47** | -0.18 | -0.12 | -0.32** | -0.21 | -0.11 | -0.10 | -0.28 | -0.20 | -0.05 | |
| 5 | 0.55*** | 0.48*** | -0.45*** | 0.06 | 0.57*** | -0.26 | 0.12 | 0.41** | -0.56*** | -0.10 | 0.46 | -0.37** | -0.04 | |
| 6 | -0.36** | -0.53*** | 0.34*** | 0.20 | -0.30* | 0.16 | 0.12 | -0.33** | 0.08 | 0.25 | -0.33** | 0.10 | 0.25 | |
| 7 | -0.16 | -0.25 | 0.09 | -0.20 | 0.03 | 0.15 | -0.22 | 0.02 | 0.26 | -0.30* | 0.20 | 0.23 | -0.29* | |
| 8 | -0.02 | -0.15 | -0.03 | -0.01 | 0.21 | 0.08 | -0.02 | 0.00 | -0.20 | -0.08 | -0.02 | -0.14 | 0.00 | |
| 9 | 0.03 | 0.29 | -0.12 | 0.08 | 0.21 | 0.10 | 0.02 | -0.43** | 0.05 | 0.03 | -0.35** | 0.05 | -0.05 | |
| Tr | -0.26 | 0.08 | 0.06 | -0.27* | -0.16 | 0.18 | -0.36** | -0.45** | 0.28* | -0.14 | -0.46** | 0.29* | -0.19 | |
| Bl | 0.39** | 0.24 | -0.18 | 0.15 | 0.45** | -0.04 | 0.12 | 0.23 | -0.22 | 0.06 | 0.27 | 0.16 | 0.05 | |
| Zo | -0.25 | -0.45*** | 0.20 | 0.11 | -0.07 | 0.15 | 0.05 | -0.16 | -0.09 | 0.10 | -0.19 | 0.12 | 0.17 | |

Circulation regime: TR, trough; BL, block; Zo, zonal.

Period: a June; b July; c August; d September; e July–August; f June–July–August; g July–August–September; h June–July–August–September.

P levels (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$).

Table 4. Relationships between crop yields and the North Atlantic Oscillation (NAO), West African Monsoon (WAM) and the Intertropical Front (ITF)

| | Barley | Durum wheat | Olive oil | Sunflower |
|------|----------|-------------|-----------|-----------|
| NAOb | -0.2 | -0.29 | 0.22 | -0.41* |
| NAOd | -0.03 | -0.04 | -0.06 | 0.37* |
| NAOg | -0.08 | -0.16 | 0.28* | -0.19 |
| WAMa | -0.11 | -0.09 | -0.25 | -0.42* |
| WAMc | 0.22 | 0.04 | -0.10 | -0.28* |
| WAMf | 0.11 | -0.04 | -0.16 | -0.44** |
| WAMh | 0.18 | 0.06 | -0.17 | -0.43** |
| ITFa | -0.36** | -0.32 | -0.48** | 0.05 |
| ITFb | -0.34* | -0.21 | -0.10 | -0.07 |
| ITFc | -0.44** | -0.31* | -0.43** | 0.23 |
| ITFd | -0.29 | -0.10 | -0.62*** | 0.21 |
| ITFe | -0.50*** | -0.34* | -0.35 | 0.08 |
| ITFf | -0.49*** | -0.36** | -0.42* | 0.09 |
| ITFg | -0.49*** | -0.28 | -0.51*** | 0.17 |
| ITFh | -0.53*** | -0.36** | -0.55*** | 0.12 |

Period: a June; b July; c August; d September; e July–August; f June–July–August; g July–August–September; h June–July–August–September.
P levels (**P* < 0.05, ***P* < 0.01, ****P* < 0.001).

Olive yields responded to more troughs, especially WT9 which produces above-normal precipitation between April and June. WT8 was important later during July and August. For the entire season, blocking circulation which produces warmer and drier conditions was detrimental to yield.

Discussion

Climate variability in the temperate and the subtropical parts of Europe and North Africa has significant effects on crop yields in Tuscany. Positive NAO leads to drier than normal winters in Tuscany, with the reverse when it is negative. Conversely, a positive (negative) SNAO produces above-normal JA precipitation. When ITF moves north, hot air masses are displaced from the Libyan Desert over Italy, bringing hot and dry conditions in Tuscany. In short, when the WAM is strong, the ITF reaches its northernmost position, and blocking is prevalent over north west (WT1) in contrast to central Europe (WT5), with a lack of troughing over Italy.

Both barley and olive oil yields decline significantly when the ITF is further north suggesting that this feature nudges the Libyan high over the central Mediterranean, producing warmer and drier conditions. Durum wheat is affected by the ITF to a lesser extent, probably because it is more drought-resistant. Sunflower production showed associations with the WAM, but the correlation is not statistically significant.

During JJA, blocking regimes WT5 and WT3 active over central Europe were highly significant. Anticyclones over central Europe were prominent during positive NAO events, but not further west (WT1). Zonal westerlies (WT6) tended to be absent, especially in JAS. Similar conditions prevail with negative SNAO episodes, with a lack of zonal flow (WT6), when easterly circulation occurs over the western central Mediterranean.

Table 5. Relationships between crop yields and weather regimes and types

| WT | Barley | | | | Durum wheat | | | | Sunflower | | | | Olives | | | | | | |
|----|--------|---------|--------|---------|-------------|----------|---------|---------|-----------|---------|---------|-------|---------|---------|--------|---------|-------|---------|----------|
| | i | l | m | n | o | p | q | r | e | q | r | e | q | r | e | q | r | s | t |
| 1 | 0.18 | 0.24 | 0.27* | 0.24 | 0.29* | 0.30* | 0.32 | 0.41*** | 0.27 | 0.32 | 0.41*** | 0.27 | 0.32 | 0.41*** | -0.34 | 0.03 | 0.03 | 0.03 | -0.38*** |
| 2 | -0.02 | 0.11 | 0.06 | 0.24 | 0.09 | 0.37*** | 0.41* | 0.32** | -0.21 | 0.41* | 0.32** | -0.21 | 0.41* | 0.32** | -0.01 | -0.06** | 0.10 | -0.06** | 0.10 |
| 3 | 0.03 | 0.00 | 0.02 | 0.04 | -0.06 | -0.04 | 0.13 | -0.01 | -0.24 | 0.13 | -0.01 | -0.24 | 0.13 | -0.01 | 0.21 | -0.13 | -0.13 | -0.13 | -0.05 |
| 4 | 0.05 | 0.06 | 0.08 | 0.14 | 0.08 | 0.07 | 0.25 | 0.16 | 0.03 | 0.25 | 0.16 | 0.03 | 0.25 | 0.16 | 0.00 | -0.10 | -0.10 | -0.10 | -0.02 |
| 5 | 0.02 | -0.16 | -0.14 | -0.34** | 0.19 | -0.45*** | -0.30 | -0.27* | 0.21 | -0.30 | -0.27* | 0.21 | -0.30 | -0.27* | -0.01 | -0.16 | -0.16 | -0.16 | -0.20 |
| 6 | -0.09 | -0.38** | -0.28* | -0.40** | -0.22 | -0.21 | -0.37* | -0.32** | -0.01 | -0.37* | -0.32** | -0.01 | -0.37* | -0.32** | -0.17 | 0.34 | 0.34 | 0.34 | 0.22 |
| 7 | -0.04 | -0.09 | -0.09 | -0.05 | -0.03 | -0.17 | -0.10 | -0.05 | 0.34 | -0.10 | -0.05 | 0.34 | -0.10 | -0.05 | 0.17 | 0.26 | 0.26 | 0.26 | 0.28 |
| 8 | 0.20 | 0.00 | 0.15 | 0.02 | -0.15 | 0.05 | -0.47** | -0.07 | -0.29 | -0.47** | -0.07 | -0.29 | -0.47** | -0.07 | 0.51** | -0.20 | -0.20 | -0.20 | 0.19 |
| 9 | -0.20 | 0.38** | 0.03 | 0.33** | -0.03 | 0.24 | -0.13 | 0.16 | -0.01 | -0.13 | 0.16 | -0.01 | -0.13 | 0.16 | -0.06 | 0.03 | 0.03 | 0.03 | 0.16 |
| Tr | -0.16 | 0.28* | 0.05 | 0.39** | 0.05 | 0.40*** | 0.28 | 0.32** | -0.09 | 0.28 | 0.32** | -0.09 | 0.28 | 0.32** | -0.01 | -0.03 | -0.03 | -0.03 | 0.28 |
| Bl | 0.12 | 0.00 | 0.11 | -0.12 | 0.23 | -0.24 | 0.13 | 0.09 | 0.21 | 0.13 | 0.09 | 0.21 | 0.13 | 0.09 | -0.15 | -0.18 | -0.18 | -0.18 | -0.45** |
| Zo | 0.02 | -0.29* | -0.16 | -0.28* | -0.27 | -0.11 | -0.55** | -0.31 | 0.29 | -0.55** | -0.31 | 0.29 | -0.55** | -0.31 | -0.05 | 0.23 | 0.23 | 0.23 | 0.26 |

Circulation regime: TR, trough; BL, block; Zo, zonal.
 Period: e July–August; i March and April; l May and June; m from March to June; n from March to July; o from March to May; p June and July; q from April to August; r from April to November; s from September to November; t from April to November.
P levels (**P* < 0.05, ***P* < 0.01, ****P* < 0.001).

For both barley and durum wheat, warm dry conditions during the grain filling and ripening period decreased, whilst wetter conditions increased yields in Tuscany (see WT5 and through circulation precipitation anomaly in Appendix 2). According to Porter and Gawith (1999), barley and winter wheat seed germination occurs under a wide range of temperatures (3.5–32.7°C), as well as flowering and grain filling stages (9.5–31 and 9.2–35.4°C, respectively), and consequently, an increase in global temperatures and changes in precipitation patterns can contribute to an increase of crop evapotranspiration, making water a potentially limiting factor for crop production in many areas, especially in the Mediterranean. Moore and Lobell (2015) confirm that water is among the most important inputs in agriculture for enhancing and stabilizing crop production and is a limiting factor in warm areas such as the Mediterranean region where water resources are limited.

Anticyclones over central Europe induce a decrease in the yield of both crops, as a zonal flow does for barley. Conversely, the presence of troughs over southern Italy increases barley yields, as do, generally, the presence of a trough circulation for both crops. These results are confirmed by Dolferus *et al.* (2011) for barley and wheat which, like other cereal crops, are vulnerable to water stress depending on which phenological stages the stress occurs (Dolferus *et al.*, 2011). In particular, when water stress occurs before or at anthesis, it mainly affects the grain number by inducing pollen sterility, while stress at grain filling and ripening may reduce kernel weight and size (Labuschagne *et al.*, 2009).

The effects of climate variability on sunflower crops were random, probably because of being a more drought-tolerant crop suitable for regions affected with water scarcity, and because of crop management and cultural practices (irrigation and pesticide application which is a common practice) (Debaeke *et al.*, 2017). Many studies confirm that vulnerability to climate variability can significantly vary from crop to crop and, for example, summer crops have a physiological higher tolerance to warm climates (Gay *et al.*, 1991; Najeeb *et al.*, 2019).

For olive crops, the blocking circulation over the entire season, with associated hot and dry conditions, decreased yields, thus confirming results from other studies showing that olive crops are affected by adverse environmental factors (i.e. water scarcity, heat and high irradiance). For example, despite olives having a strong tolerance to drought (Baldi *et al.*, 2019), some experiments performed in Spain observed as crop water demand for olive range between 485 and 600 mm per growing season (Palomo *et al.*, 2002). Consequently, the cultivation of olive tree crops in the Mediterranean area can be challenging during the extremely dry summers expected to be more frequent due to climate change (Brito *et al.*, 2019). Troughing circulation, especially in southern Italy (WT9), was most important in late spring, likely due to the positive anomaly precipitation in April and June as well as the zonal flow (WT8) in early summer (Appendix 2).

Conclusion

The weather types and circulation regimes approach leads to good results in explaining the alternation of crop yields in Tuscany both for winter crops (barley, durum wheat) and for warm season crops (olives and sunflower), and in particular, the effects proved to be different according to the crop phenological phases. The study highlights that blocking circulation during the growing season, with associated hot and dry conditions, decreases yield for olive crops, barley and durum wheat. For this reason, the extension of this study to other crops and other geographical areas

could allow the creation of a useful database to predict the crop yields according to the prevailing forecast weather types (i.e. by using seasonal forecasts). Finally, this study focused on the impact of climate on crops but other factors, such as diseases and pests, variability of cultivated areas, soil variability, variability of agronomic practices and crop management, are also important. Their impact can be investigated in future work.

Acknowledgements. A particular thanks to Nicola Gambaro who provided support in the creation of the circulation regime datasets.

Financial support. This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Conflict of interest. None.

Ethical standards. Not applicable.

References


- Agnolucci P and De Lipsis V (2019) Long-run trend in agricultural yield and climatic factors in Europe. *Climatic Change* **159**, 385–405.
- Alpert P, Ben-Gai T, Baharad A, Benjamini Y, Yekutieli D, Colacino M, Diodato L, Ramis C, Homar V, Romero R, Michaelides S and Manes A (2002) The paradoxical increase of Mediterranean extreme daily rainfall in spite of decrease in total values. *Geophysical Research Letters* **29**, 311–314.
- Alpert P, Baldi M, Ilani R, Krichak S, Price C, Rodo X, Saaroni H, Ziv B, Kishcha P, Karkan J, Mariotti A and Xoplaki E (2006) Relations between climate variability in the Mediterranean region and the tropics: ENSO, South Asian and African monsoons, hurricanes and Saharan dust. In Lionello P, Malanotte-Rizzoli P and Boscolo R (eds.), *Mediterranean Climate Variability*. Amsterdam: Elsevier, pp. 149–177.
- Avolio E, Pasqualoni L, Federico S, Fornaciari M, Bonfiglio T, Orlandi F, Bellecci C and Romano B (2008) Correlation between large-scale atmospheric fields and the olive pollen season in Central Italy. *International Journal of Biometeorology* **52**, 787–796.
- Baldi A, Brandani G, Petralli M, Messeri A, Cecchi S, Vivoli R and Mancini M (2019) Thermo-pluviometric variability of Val d'Orcia Olive Orchards area (Italy). *Italian Journal of Agrometeorology* **2**, 11–20.
- Barranco D and Ruiz N (2005) Frost tolerance of eight olive cultivars. *Hortscience* **40**, 558–560.
- Bartolini G, Messeri A, Grifoni D, Mannini D and Orlandini S (2013) Recent trends in seasonal and annual precipitation indices in Tuscany (Italy). *Theoretical and Applied Climatology* **118**, 147–157.
- Bartolini G, Grifoni D, Magno R, Torrigiani T and Gozzini B (2018) Changes in temporal distribution of precipitation in a Mediterranean area (Tuscany, Italy) 1955–2013. *International Journal of Climatology* **38**, 1366–1374.
- Bonfiglio T, Orlandi F, Sgromo C, Romano B and Fornaciari M (2008) Influence of temperature and rainfall on timing of olive (*Olea europaea*) flowering in southern Italy. *New Zealand Journal of Crop and Horticultural Science* **36**, 59–69.
- Brito C, Dinis LT, Moutinho-Pereira J and Correia CM (2019) Drought stress effects and olive tree acclimation under a changing climate. *Plants* **8**, 232.
- Brunetti M, Buffoni L, Mangianti F, Maugeri M and Nanni T (2004) Temperature, precipitation and extreme events during the last century in Italy. *Global and Planetary Change* **40**, 141–149.
- Ceglar A, Turco M, Toreti A and Doblas-Reyes FJ (2017) Linking crop yield anomalies to large-scale atmospheric circulation in Europe. *Agricultural and Forest Meteorology* **240–241**, 35–45.
- Chen TC (2005) Maintenance of the Midtropospheric North African summer circulation: Saharan High and African Easterly Jet. *Journal of Climate* **18**, 2943–2962.
- Chen CC, McCarl BA and Schimmelpfennig DE (2004) Yield variability as influenced by climate: a statistical investigation. *Climatic Change* **66**, 239–261.
- Ciais P, Reichstein M, Viovy N, Granier A, Ogee J, Allard V, Aubinet M, Buchmann N, Bernhofer C, Carrara A, Chevallier F, De Noblet N, Friedl AD, Friedlingstein P, Grünwald T, Heinesch B, Keronen P, Knohl A, Krinner G, Loustau D, Manca G, Matteucci G, Miglietta F,

- Orcival JM, Papale D, Pilegaard K, Rambal S, Seufert G, Soussana JF, Sanz MJ, Schulze ED, Vesala T and Valentini R (2005) Europe-wide reduction in primary productivity caused by the heat and drought in 2003. *Nature* **437**, 529–533.
- Cornes R, Van der Schrier G, Van den Besselaar EJM and Jones PD (2018) An ensemble version of the E-OBS temperature and precipitation datasets. *Journal of Geophysical Research: Atmospheres* **123**, 9391–9409. doi: 10.1029/2017JD028200.
- Dalla Marta A, Grifoni D, Mancini M, Storchi P, Zipoli G and Orlandini S (2010) Analysis of the relationships between climate variability and grapevine phenology in the Nobile di Montepulciano wine production area. *The Journal of Agricultural Science* **18**, 657–666.
- Dalla Marta A, Grifoni D, Mancini M, Zipoli G and Orlandini S (2011) The influence of climate on durum wheat quality in Tuscany, Central Italy. *International Journal of Biometeorology* **55**, 87–96.
- Debaeke P, Casadebaig P, Flenet F and Langlade N (2017) Sunflower crop and climate change: vulnerability, adaptation, and mitigation potential from case-studies in Europe. *OCL* **24**, D102.
- Del Pozo A, Brunel-Saldias N, Engler A, Ortega-Farias S, Acevedo-Opazo C, Lobos GA, Jara-Rojas R and Molina-Montenegro MA (2019) Climate change impacts and adaptation strategies of agriculture in Mediterranean-Climatic Regions (MCRs). *Sustainability* **11**, 2769.
- Demir AO, Goksoy AT, Buyukcangaz H, Turan ZN and Koksals ES (2006) Deficit irrigation of sunflower (*Helianthus annuus* L.) in a sub-humid climate. *Irrigation Science* **24**, 279–289.
- Dolferus R, Ji X and Richards RA (2011) Abiotic stress and control of grain number in cereals. *Plant Science* **181**, 331–341.
- Erdem T, Delibas L and Orta AH (2001) Water-use characteristics of sunflower (*Helianthus annuus* L.) under deficit irrigation. *Pakistan Journal of Biological Sciences* **4**, 766–769.
- FAO (2017) *The Future of Food and Agriculture – Trends and Challenges*. Rome: FAO. Available at <http://www.fao.org/3/a-i6583e.pdf>.
- Folland CK, Knight J, Linderholm HW, Fereday D, Ineson S and Hurrell JW (2009) The summer North Atlantic Oscillation: past, present and future. *Journal of Climate* **22**, 1082–1103.
- Fontana G, Toreti A, Ceglar A and De Sanctis G (2015) Early heat waves over Italy and their impacts on durum wheat yields. *Natural Hazards and Earth System Sciences* **15**, 1631–1637.
- Gaetani M and Fontaine B (2013) Interaction between the West African Monsoon and the summer Mediterranean climate: an overview. *Fisica de la Tierra* **25**, 41–55.
- Gay C, Corbinau F and Come D (1991) Effects of temperature and oxygen on seed germination and seedling growth in sunflower (*Helianthus annuus* L.). *Environmental and Experimental Botany* **31**, 193–200.
- Goldhamer DA, Dunai J and Ferguson LF (1994) Irrigation requirements of olive trees and responses to sustained deficit irrigation. *Acta Horticulturae* **356**, 172–175.
- Haworth M, Marino G, Brunetti C, Killi D, De Carlo A and Centritto M (2018) The impact of heat stress and water deficit on the photosynthetic and stomatal physiology of olive (*Olea europaea* L.): a case study of the 2017 HeatWave. *Plants* **7**, 76.
- Herbert ER, Boon P, Burgin AI, Neubauer S, Franklin RB, Ardon M, Hopfensperger KN, Lamers LPM and Gell P (2015) A global perspective on wetland salinization: ecological consequences of a growing threat to freshwater wetlands. *Ecosphere (Washington, DC)* **6**, 1–43.
- Hess P and Brezowsky H (1952) Katalog der Groß wetterlagen Europas (catalog of the European large-scale weather types). *Berichte des Deutschen Wetterdienstes* **33**, 39.
- Hidalgo J and Jouglar R (2018) On the use of local weather types classification to improve climate understanding: an application on the urban climate of Toulouse. *PLoS ONE* **13** (12), doi: 10.1371/journal.pone.0208138.
- Hoegh-Guldberg O, Jacob D, Taylor M, Bindi M, Brown S, Camilloni I, Diedhiou A, Djalante R, Ebi KL, Engelbrecht F, Guiot J, Hijioka Y, Mehrotra S, Payne A, Seneviratne IA, Thomas A, Warren R and Zhou G (2018) Impacts of 1.5°C Global Warming on Natural and Human Systems. In: *Global Warming of 1.5°C – IPCC Special Report*.
- Hurrell JW (1995) Decadal trends in the North-Atlantic Oscillation – regional temperatures and precipitation. *Science (New York, NY)* **269**, 676–679.
- Hurrell JW, Kushnir Y, Ottersen G and Visbeck M (2003) The North Atlantic oscillation: climate significance and environmental impact. *Geophysical Monograph Series* **134**, 1–35. doi: 10.1029/134GM01.
- Huth R, Beck C and Kucerova M (2015) Synoptic-climatological evaluation of the classifications of atmospheric circulation patterns over Europe. *International Journal of Climatology* **3**, 2710–2726.
- Iniesta F, Testi L, Orgaz F and Villalobos FJ (2009) The effects of regulated and continuous deficit irrigation on the water use, growth and yield of olive trees. *European Journal of Agronomy* **30**, 258–265.
- IPCC (2007) *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, UK, 976pp.
- Kukul MS and Irmak S (2018) Climate-driven crop yield and yield variability and climate change impacts on the U.S. Great Plains agricultural production. *Scientific Reports* **8**, 3450.
- Labuschagne MT, Elago O and Koen EJ (2009) The influence of temperature extremes on some quality and starch characteristics in bread, biscuit and durum wheat. *Journal of Cereal Science* **49**, 184–189.
- Lamb HH (1972) British Isles weather types and a register of daily sequence of circulation patterns, 1861–1971. *Geophysical Memoirs* **116**, 85L.
- Lavaysse C, Flamant C, Evan A, Janicot S and Gaetani M (2016) Recent climatological trend of the Saharan heat low and its impact on the West African climate. *Climate Dynamics* **47**, 3479.
- Lélé IM and Lamb PJ (2010) Variability of the intertropical front (ITF) and rainfall over the West African Sudan–Sahel zone. *Journal of Climate* **23**, 3984–4004.
- Li JP and Zeng QC (2005) A new monsoon index, its interannual variability and relation with monsoon precipitation. *Climatic and Environmental Research* **10**, 351–336.
- Lu J, Carbone GJ and Gao P (2017) Detrending crop yield data for spatial visualization of drought impacts in the United States, 1895–2014. *Agricultural and Forest Meteorology* **237–238**, 196–208. doi: 10.1016/j.agrformet.2017.02.001.
- Macours K and Swinnen JFM (2000) Causes of output decline in economic transition: the case of central and eastern European agriculture. *Journal of Comparative Economics* **28**, 172–206. Available at <https://EconPapers.repec.org/RePEc:eee:jcecon:v:28:y:2000:i:1:p:172-206>.
- Manzanas R, Gutierrez JM, Fernandez J, Van Meijgard E, Calmanti S, Magarino ME, Colfino AS and Herrera S (2018) Dynamical and statistical downscaling of seasonal temperature forecasts in Europe: added value for user applications. *Climate Services* **9**, 44–56.
- Mariotti A and Dell'Aquila A (2012) Decadal climate variability in the Mediterranean region: roles of large-scale forcings and regional processes. *Climate Dynamics* **38**, 1129–1145.
- Mathbout A, Lopez-Bustins JA, Royè D, Martin-Vide J and Benhamrouche A (2019) Spatiotemporal variability of daily precipitation concentration and its relationship to teleconnection patterns over the Mediterranean during 1975–2015. *International Journal of Climatology* **40**, 1435–1455. doi: 10.1002/joc.6278.
- Messori A, Morabito M, Messori G, Brandani G, Petralli M, Natali F, Grifoni D, Crisci A, Gensini G and Orlandini S (2015) Weather-related flood and landslide damage: a risk index for Italian regions. *PLoS ONE* **10**, 12.
- Messori G, Benedetti G, Crisci A, Gozzini B, Rossi M, Vallorani R and Maracchi G (2018) A new framework for probabilistic seasonal forecasts based on circulation type classifications and driven by an ensemble global model. *Advances in Scientific Research* **15**, 183–190.
- Moore FC and Lobell DB (2015) The fingerprint of climate trends on European crop yields. *Proceedings of the National Academy of Sciences* **112**, 2670–2675.
- Najafi E, Pal I and Khanbilvardi R (2019) Climate drives variability and joint variability of global crop yields. *Science of the Total Environment* **662**, 361–372.
- Najeeb U, Tan DKY, Sarwar M and Ali S (2019) Adaptation of crops to warmer climates: morphological and physiological mechanisms. In Sarkar A, Sensarma S and vanLoon G (eds), *Sustainable Solutions for Food Security*, Cham: Springer, pp. 53–76.
- Nobre G, Hunink JE, Aerts JC and Ward PJ (2019) Translating large-scale climate variability into crop production forecast in Europe. *Nature Scientific Reports* **9**, 1277.

- Oerke E-C** (2006) Crop losses to pests. *Journal of Agricultural Science* **144**, 31–43.
- Orlandini S, Nejedlik P, Eitzinger J, Alexandrov V, Toulous L, Calanca P, Trnka M and Olesen JE** (2008) Impacts of climate change and variability on European agriculture. Results of inventory analysis in COST 734 countries. Trends and directions in climate research. *Annals of the New York Academy of Science* **1146**, 338–353.
- Orlandini S, Dalla Marta A, Mancini M, Grifoni D et al.** (2011) Impacts of the NAO on Mediterranean crop production. In Vicente-Serrano SM and Trigo RM (eds), *Hydrological, Socioeconomic and Ecological Impacts of the North Atlantic Oscillation in the Mediterranean Region SE - 8. Advances in Global Change Research*. The Netherlands: Springer, pp. 103–112. doi: 10.1007/978-94-007-1372-7_8.
- Osborne TM and Wheeler TR** (2013) Evidence for a climate signal in trends of global crop yield variability over the past 50 years. *Environmental Research Letters* **8**, 1–9.
- Palomo MJ, Moreno F, Fernández JE, Díaz-Espejo A and Girón IF** (2002) Determining water consumption in olive orchards using the water balance approach. *Agricultural Water Management* **55**, 15–35.
- Parent B, Leclere M, Lacube S, Semenov MA, Welcker C, Martre P and Tardieu F** (2018) Maize yields over Europe may increase in spite of climate change, with an appropriate use of the genetic variability of flowering time. *Proceedings of the National Academy of Sciences* **115**, 10642–10647.
- Philipp A, Beck C, Huth R and Jacobei J** (2014) Development and comparison of circulation type classifications using the COST 733 dataset and software. *International Journal of Climatology* **36**, 2673–2691.
- Porter JR and Gawith M** (1999) Temperatures and the growth and development of wheat: a review. *European Journal of Agronomy* **10**, 23–36.
- Ray D, Gerber J, MacDonald G and West PC** (2015) Climate variation explains a third of global crop yield variability. *Nature Communications* **6**, 5989.
- Rodwell MJ and Hoskins BJ** (1996) Monsoons and the dynamics of deserts. *Quarterly Journal of the Royal Meteorological Society* **122**, 1385–1404.
- Salinger J, Baldi M, Grifoni D, Jones G, Bartolini G, Cecchi S, Messeri G, Dalla Marta A, Orlandini S, Dalu GA and Maracchi G** (2015) Seasonal differences in climate in the Chianti region of Tuscany and the relationship to vintage wine quality. *International Journal of Biometeorology* **59**, 1799–1811.
- Searchinger T, Hanson C, Ranganathan J, Lipinski B, Waite R, Winterbottom R, Dinshaw A and Heimlich R** (2019) Creating a sustainable food future – a menu of solutions to sustainably feed more than 9 billion people by 2050. *Technical Report of the World Resources Institute*, Washington, DC. DOI: Available at https://agritrop.cirad.fr/593176/1/WRR_Food_Full_Report_0.pdf.
- Teasdale JR and Cavigelli MA** (2017) Meteorological fluctuations define long-term crop yield patterns in conventional and organic production systems. *Scientific Reports* **7**, 688.
- Thorncroft CD, Nguyen H, Zhang C and Peyrille P** (2011) Annual cycle of the West African monsoon: regional circulations and associated water vapor transport. *Quarterly Journal of the Royal Meteorological Society* **137**, 129–147.
- Todisco F and Vergni L** (2008) Climatic changes in central Italy and their potential effects on corn water consumption. *Agricultural and Forest Meteorology* **148**, 1–11, doi: 10.1016/j.agrformet.2007.08.014.
- Vallorani R, Bartolini G, Betti G, Crisci A, Gozzini B, Grifoni D, Iannuccilli M, Messeri A, Messeri G, Morabito M and Maracchi G** (2018) Circulation type classifications for temperature and precipitation stratification in Italy. *International Journal of Climatology* **30**, 915–931. doi: 10.1002/joc.5219.
- Verdi L, Napoli M, Santoni M, Dalla Marta A and Ceccherini MT** (2019) Soil carbon dioxide emission flux from organic and conventional farming in a long term experiment in Tuscany. 2019 IEEE International Workshop on Metrology for Agriculture and Forestry, MetroAgriFor 2019 – Proceedings, 2019, pp. 85–89, 8909242.
- Vergni L and Todisco F** (2011) Spatio-temporal variability of precipitation, temperature and agricultural drought indices in Central Italy. *Agricultural and Forest Meteorology* **151**, 301–313.
- Vergni L, Todisco F, Di Lena B and Mannocchi F** (2020) Bivariate analysis of drought duration and severity for irrigation planning. *Agricultural Water Management* **229**, issue C, 301–313, doi: 10.1016/j.agwat.2019.105926.
- Yawson DO, Bonsu M, Arman FA and Afrifa EK** (2011) Water requirement of sunflower (*Helianthus annuus* L.) in a tropical humid-coastal savanna zone. *ARPN Journal of Agricultural and Biological Science* **4**, 1–8.


Appendix 1

Total precipitation for Tuscany during the period 1981–2010 (Cornes et al., 2018). Colour online




Appendix 2


Precipitation anomalies and total precipitation for Italy produced by PCT circulation types (resolution of 25 km) for the months not discussed in the text. The anomalies refer to the 1981–2010 base period.




Precipitation Anomalies-March. Colour online.




Precipitation Anomalies-April. Colour online.




Precipitation Anomalies-May. Colour online.




Precipitation Anomalies-June. Colour online.




Precipitation Anomalies-July. Colour online.




Precipitation Anomalies- August. Colour online.




Precipitation Anomalies- October. Colour online.




Total Precipitation-May. Colour online.



Total Precipitation-June. Colour online.



Total Precipitation-August. Colour online.



Total Precipitation-September. Colour online.