

Malignancies among children and young people with HIV in Western and Eastern Europe and Thailand

The European Pregnancy and Paediatric Infections Cohort Collaboration (EPPICC) study group*

Objectives: Investigate trends over time and predictors of malignancies among children and young people with HIV.

Design: Pooled data from 17 cohorts in 15 countries across Europe and Thailand.

Methods: Individuals diagnosed with HIV and presenting to paediatric care less than 18 years of age were included. Time at risk began at birth for children with documented vertically acquired HIV, and from first HIV-care visit for others. Children were followed until death, loss-to-follow-up, or last visit in paediatric or adult care (where data after transfer to adult care were available). Rates of reported malignancies were calculated overall and for AIDS-defining malignancies (ADM) and non-AIDS-defining malignancies (NADM) separately. Risk factors for any malignancy were explored using Poisson regression, and for mortality following a malignancy diagnosis using Cox regression.

Results: Among 9632 individuals included, 140 (1.5%) were ever diagnosed with a malignancy, of which 112 (80%) were ADM. Overall, the rate of any malignancy was 1.18 per 1000 person-years; the rate of ADM decreased over time whereas the rate of NADM increased. Male sex, being from a European cohort, vertically acquired HIV, current severe immunosuppression, current viral load greater than 400 copies/ml, older age, and, for those not on treatment, earlier calendar year, were risk factors for a malignancy diagnosis. Fifty-eight (41%) individuals with a malignancy died, a median 2.4 months (IQR 0.6–8.8) after malignancy diagnosis.

Conclusion: The rate of ADM has declined since widespread availability of combination ART, although of NADM, there was a small increase. Mortality following a malignancy was high, warranting further investigation.

Copyright © 2021 The Author(s). Published by Wolters Kluwer Health, Inc.

AIDS 2021, **35**:1973–1985

Keywords: adolescents, children, Europe, HIV, malignancies, Thailand

Introduction

Cancer is relatively rare in childhood in the general population, with the incidence rate to 19 years of age across Western and Eastern Europe estimated at 0.14 per 1000 person-years [1]. However, as a result of immunosuppression, systemic inflammation, persistent HIV viremia, and increased susceptibility to oncogenic viruses, such as Epstein–Barr Virus (EBV) and human herpesvirus 8 (HHV-8), people with HIV are at higher risk of some malignancies, including both those considered AIDS-

defining (ADM), such as Kaposi sarcoma and non-Hodgkin lymphoma, and non-AIDS-defining (NADM), such as Hodgkin lymphoma [2,3].

In adults living with HIV, studies have reported declining incidence of ADM but increasing incidence of NADM in recent years [4–6]. However, children who acquire HIV vertically have lifelong exposure to the virus, and their long-term risk of malignancies is unclear, with the only studies that have followed them into young adulthood limited by small sample sizes [7]. Studies in children have

Correspondence to Elizabeth Chappell, MRC Clinical Trials Unit at UCL, Institute of Clinical Trials and Methodology, 90 High Holborn, 2nd Floor, London WC1V 6LJ, UK.

Tel: +44 20 7670 4851; e-mail: e.chappell@ucl.ac.uk

* The names of the authors of the European Pregnancy and Paediatric Infections Cohort Collaboration (EPPICC) study group are listed in the Acknowledgements section.

Received: 11 January 2021; revised: 18 May 2021; accepted: 24 May 2021.

DOI:10.1097/QAD.0000000000002965

ISSN 0269-9370 Copyright © 2021 The Author(s). Published by Wolters Kluwer Health, Inc. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

shown those on antiretroviral therapy (ART) and initiating ART at younger ages are at lower risk [8–10]. In this study, we describe rates, risk factors for malignancy and mortality following a malignancy diagnosis in children and young people in the European Pregnancy and Paediatric Infections Cohort Collaboration (EPPICC).

Methods

Seventeen observational cohorts in EPPICC from 15 countries across Europe and Thailand contributed individual-level demographic, clinical, laboratory and treatment-related data, which were pseudo-anonymized and pooled electronically using a modified HIV Cohorts Data Exchange Protocol (HICDEP, www.hicdep.org). All cohorts received approval from local/national ethics committees.

All children diagnosed with HIV, presenting to paediatric HIV care before 18 years of age, and with more than 1 day of follow-up were included. Time at risk began at birth for those with vertically acquired HIV, or on the date first seen in HIV care (defined as their first CD4⁺ cell count, viral load, use of ART or AIDS-defining event) for those with other or unknown modes of acquisition as date of HIV infection was often unknown. Data after transfer to adult care were available for some individuals but if not they were censored at their last visit in paediatric care. Individuals were followed until death, loss to follow-up, drop out from cohort (for example, if they moved clinic), or their last visit prior to 1 October 2016.

None of the contributing cohorts are linked to cancer registries, and so data on malignancy events were collected from routine HIV healthcare records. Cervical intraepithelial neoplasia, a precancerous condition, was not included in the analysis. Malignancies were categorized as ADM or NADM based on the US Centers for Disease Control and Prevention (CDC) 2014 surveillance criteria [11]. The number and rate (per 1000 person-years) of malignancy events were summarized over calendar time, overall and separately for ADM and NADM. To allow for the possibility of multiple malignancy diagnoses, rates were calculated as the number of malignancy events divided by the total duration of follow-up. Rates were presented among all patients, as well as among patients on 'stable ART', defined as those on ART for at least 6 months, who currently had viral load 400 copies/ml or less and with no HIV-related immunosuppression (defined based on WHO criteria as CD4⁺ >35% for those age <1 year; >30% for 1 to <3 years; >25% for 3–<5 years; >500 cells/μl for ≥5 years [12]). Rates of specific malignancies were presented where more than one event was reported. Characteristics at the time of malignancy

diagnoses were compared among those with an ADM or NADM, using a chi-squared or Fisher's exact test (where numbers were <5) for categorical variables and Wilcoxon's rank-sum test for continuous variables.

Risk factors for any malignancy were explored using Poisson regression. Factors considered included sex, region of cohort (UK/Ireland, Thailand, Russia/Ukraine, rest of Europe), mode of HIV acquisition (vertical vs. other). In addition, the following time-updated characteristics were considered (with measurements carried forward up to 12 months): current ART status (not initiated ART vs. <6 months since ART initiation vs. ≥6 months since ART initiation), current calendar year (<1996 vs. 1996–2003 vs. 2004–2009 vs. ≥2010), current age (<5 vs. 5 to <10 vs. 10 to <15 vs. ≥15 years), current WHO immunological stage (severe vs. not severe, with severe immunosuppression defined based on WHO criteria as CD4⁺ <25% for those age <1 year; <20% for age 1 to <3 years; <15% for 3 to <5 years; <200 cells/μl or <15% for ≥5 years [12]), current viral load (≤400 vs. >400 copies/ml), and current BMI-for-age *z*-score (>2 vs. –2 to 2 vs. <–2, based on WHO reference data [13]). Ethnicity and place of birth could not be included given that one of the areas of focus was comparing by region, and there were few children born abroad or from different ethnic groups in Thailand and Russia/Ukraine; the effects of these variables in the other cohorts were explored in a supplementary analysis. Two-way interactions between current ART status, calendar year and age were considered, using a cut-off of *P*=0.15 for inclusion. Associations were explored using univariable models, and all factors were included in a multivariable model. Missing data on sex (missing for <1% of patients), mode of HIV acquisition (5% of patients) and current immunological stage (48% of patient-time), viral load (48% of patient-time) and BMI-for-age *z*-score (66% of patient-time) were imputed using multiple imputation with chained equations; 20 imputed datasets were created, and coefficients combined using Rubin's rules [14]. Risk factors for ADM and NADM were subsequently explored separately, using the same methods.

The probability of all-cause mortality by 3 years after malignancy diagnosis was estimated using Kaplan–Meier methods, overall and by calendar year. Cox regression was used to explore associations between mortality and the following (at malignancy diagnosis, and categorized as above if not specified): calendar year, malignancy type (ADM vs. NADM), recent presentation to HIV care (<6 months vs. ≥6 months), sex, region, mode of HIV acquisition, age, ART status, WHO immunological stage and BMI-for-age *z*-score.

Events occurring prior to the date first seen in HIV care (*n*=1) or with unknown diagnosis date (*n*=3) in those with other/unknown mode of infection were excluded

Table 1. Patient characteristics.

	UK/Ireland (N = 2069)	Thailand (N = 877)	Russia/Ukraine (N = 2280)	Rest of Europe (N = 4406)	Total (N = 9632)
	<i>n</i> (%) or median (IQR)				
Sex					
Female	1090 (53%)	473 (54%)	1241 (54%)	2245 (51%)	5049 (52%)
Male	978 (47%)	404 (46%)	1036 (45%)	2160 (49%)	4578 (48%)
Unknown	1 (<0.5%)	0	3 (<0.5%)	1 (<0.5%)	5 (<0.5%)
Mode of HIV acquisition					
Vertical	1922 (93%)	874 (100%)	2083 (91%)	3473 (79%)	8352 (87%)
Blood transfusion	40 (2%)	2 (<0.5%)	8 (<0.5%)	557 (13%)	607 (6%)
Sexual contact	0	1 (<0.5%)	37 (2%)	45 (1%)	83 (1%)
Other	1 (<0.5%)	0	16 (1%)	56 (1%)	73 (1%)
Unknown	106 (5%)	0	136 (6%)	275 (6%)	517 (5%)
Ethnicity					
White	179 (9%)	0	264 (99%)	1455 (33%)	3898 (40%)
Black African	1616 (78%)	0	0	453 (10%)	2069 (21%)
Other	227 (11%)	877 (100%)	2 (<0.5%)	162 (4%)	1268 (13%)
Unknown	47 (2%)	0	14 (1%)	2336 (53%)	2397 (25%)
Born abroad					
Yes	1164 (56%)	16 (2%)	6 (<0.5%)	964 (22%)	2150 (22%)
No	876 (42%)	782 (89%)	2262 (99%)	3377 (77%)	7297 (76%)
Unknown	29 (1%)	79 (9%)	12 (1%)	65 (1%)	185 (2%)
Region of country of birth if born abroad					
Europe	22 (2%)	0	0	49 (5%)	71 (3%)
sub-Saharan Africa	654 (56%)	0	0	306 (32%)	960 (45%)
Other	29 (2%)	12 (88%)	0	64 (7%)	107 (5%)
Unknown	459 (39%)	2 (13%)	6 (100%)	545 (57%)	1012 (47%)
Entry to HIV care ^a					
<1996	719 (35%)	319 (36%)	6 (<0.5%)	2192 (50%)	3233 (34%)
1996–2003	954 (46%)	466 (53%)	622 (27%)	1470 (33%)	3515 (36%)
2004–2009	311 (15%)	77 (8%)	1109 (45%)	521 (12%)	1927 (20%)
≥2010	85 (4%)	15 (2%)	633 (28%)	223 (5%)	957 (10%)
Ever initiated ART	1847 (89%)	877 (100%)	2085 (91%)	3970 (90%)	8779 (91%)
Age at ART initiation (years)	6.7 (2.4–11.2)	6.5 (2.3–9.6)	3.0 (1.0–6.5)	3.8 (0.9–8.9)	4.3 (1.2–9.1)
Duration of follow-up (years)	16.1 (11.2–17.9)	15.2 (11.0–18.3)	8.4 (4.7–11.9)	13.9 (7.1–18.3)	12.9 (7.1–17.5)
Current follow-up status					
Still in paediatric care	837 (40%)	225 (26%)	1656 (73%)	1581 (36%)	4299 (45%)
Transferred to adult care	920 (44%)	70 (8%)	100 (5%)	931 (21%)	2021 (21%)
Dropped out of cohort	117 (6%)	338 (39%)	65 (3%)	227 (5%)	747 (8%)
Lost-to-follow-up	80 (4%)	170 (19%)	406 (18%)	994 (23%)	1650 (17%)
Died	115 (6%)	74 (8%)	53 (2%)	673 (15%)	915 (10%)
Data available after transfer to adult care	149 (16%)	70 (100%)	62 (62%)	272 (29%)	553 (27%)
Age at transfer to adult care (years)	17.7 (16.8–18.5)	17.7 (16.8–18.3)	17.9 (17.8–17.9)	18.7 (17.7–20.1)	18.1 (17.2–19.1)
Duration of follow-up in adult care (years)	2.8 (1.2–4.3)	2.8 (1.2–4.5)	1.4 (0.5–2.6)	3.6 (1.7–6.3)	2.8 (1.2–5.0)

ART, antiretroviral therapy; IQR, interquartile range.

from all analyses. Further, children with vertically acquired HIV who had an event with unknown diagnosis date ($n = 4$) were included in the overall malignancy rate but excluded from analyses of the rate over calendar time and of risk factors.

Statistical analyses were conducted using Stata 16.1 (StataCorp, College Station, Texas, USA).

Results

Overall, 9632 patients had more than 1 day of follow-up in paediatric HIV care, and were therefore, included in this analysis (Table 1). Two thousand and sixty-nine (21%) patients were from UK/Ireland, 877 (9%) from Thailand, 2280 (24%) from Russia or Ukraine, and 4406 (46%) from

the rest of Europe. The mode of HIV acquisition was documented as vertically acquired for 8352 (87%) patients, blood transfusion for 607 (6%), sexual contact for 83 (1%), other modes for 73 (1%), and was unknown for 517 (5%). Three thousand eight hundred and ninety-eight (40%) children were white, 2069 (21%) were black African, 1268 (13%) were of other ethnicity, and ethnicity was not reported for 2397 (25%). The median duration of follow-up was 12.9 [interquartile range (IQR) 7.1–17.5] years, with a total of 118 418 person-years of follow-up, of which 87 072 person-years was among those born within the country of their cohort and 28 946 person-years those born outside. Eight thousand, seven hundred and seventy-nine (91%) patients ever initiated ART. Calendar year of entry to HIV care ranged from 1980 to 2016. At last follow-up, 4299 (45%) patients were still in paediatric care, 1650 (17%) had been lost to follow-up, 747 (8%) had dropped out for other reasons, 915 (10%)

Table 2. Malignancy events.

Event	N = 140 [n (%)]
AIDS-defining malignancies	112 (80%)
Non-Hodgkin lymphoma	83
Diffuse large B-cell (immunoblastic or centroblastic)	43
Burkitt (classical or atypical)	11
Primary brain	10
Unspecified	19
Kaposi sarcoma	25
Cervical cancer	1
Unspecified	3
Non-AIDS-defining malignancies	27 (19%)
Hodgkin lymphoma	15
Hepatocellular carcinoma ^a	2
Leiomyosarcoma	1
Other ^b	8
Unspecified	1
Unspecified	1 (1%)

^aOne patient had hepatitis B co-infection, and one had hepatitis C co-infection.

^bDescription of the eight other non-AIDS-defining malignancies: brain ganglioglioma; ganglioneuroblastoma; chest wall tumour (unspecified); disseminated adenocarcinoma; malignant fibrous histiocytoma; neuroendocrine tumour of the pancreas; cystic teratoma; acute T-lymphoblastic leukaemia.

had died in paediatric care, and the remaining 2021 (21%) had transferred to adult care, of whom 553 (27%) had data available after transfer with a median duration of adult care follow-up of 2.8 (1.2, 5.0) years.

One hundred and forty (1.5%) patients had a malignancy event (Table 2). No patient experienced more than one event. Of the 140 events, 112 (80%) were ADM, with the most common being non-Hodgkin lymphoma ($n=83$) and Kaposi sarcoma ($n=25$). A further 27 (19%) were NADM, including Hodgkin lymphoma ($n=15$) and hepatocellular carcinoma ($n=2$, one each in patients co-infected with hepatitis B and C).

Across all events, two (1%) patients were diagnosed with a malignancy prior to entry to HIV care, 8 (6%) were diagnosed at the same time as entry to HIV care, and 19 (14%) patients were diagnosed within 6 months. Among these 29 patients, 22 acquired HIV through vertical transmission [median age at malignancy diagnosis 5.1 (IQR 2.8–9.4) years], two through blood transfusion, three through sexual contact, and the mode of acquisition was unknown for three. Malignancy diagnoses among those born abroad were more likely to be before or soon after entering HIV care, compared with those born within the country of cohort [12/31 (39%) vs. 17/106 (16%), $P=0.007$]. Five events (four non-Hodgkin lymphoma and one hepatocellular carcinoma) occurred after transfer to adult care, at a median (IQR) age of 24.4 (18.8–24.9) years.

A higher proportion of those with an ADM were diagnosed within 6 months of entry to HIV care [23% ($n=26$) vs. 7% ($n=2$), $P=0.106$] and a lower proportion

had previously been on ART [62% ($n=67$) vs. 96% ($n=26$), $P<0.001$], compared with those with an NADM. Age [9.9 (IQR 4.6–14.6) vs. 10.1 (8.0–16.4) years, $P=0.075$], $CD4^+$ [15 (6–23) vs. 19 (9–25), $P=0.273$] and the proportion with severe immunosuppression (58 vs. 36%, $P=0.133$) at malignancy diagnosis were similar between with ADM and NADM.

Of the 25 children diagnosed with Kaposi sarcoma, 44% were male, 40% were of black ethnicity, median age and $CD4^+$ at malignancy diagnosis were 10.4 (5.4–14.3) years and 6 (2–19) respectively. There were 11 events of Burkitts lymphoma, of which 73% were in males and 45% in those of white ethnicity, occurring at a median 14.3 (4.7–17.7) years of age and $CD4^+$ of 16 (10–30).

Overall, the rate of any malignancy was 1.18 [95% confidence interval (CI) 1.00–1.40] per 1000 person-years, and was 0.35 (0.18–0.68) among stable patients. Of ADM specifically, the rate was 0.94 (0.79–1.14) and of NADM was 0.23 (0.16–0.33). The rates of non-Hodgkin lymphoma, Kaposi sarcoma and Hodgkin lymphoma were 0.70 (0.57–0.87), 0.21 (0.14–0.31) and 0.13 (0.08–0.21), respectively. The rate of any malignancy, ADM and NADM by calendar time are shown in Fig. 1. The rate of ADM was high and steady over time between 1990/1991 and 1996/1997 ($P=0.210$; ranging between 2.37 and 3.02), after which it dropped dramatically, ranging between 0.31 and 1.09 between 1996/1997 and 2014/2015, with a slight continued decrease over time ($P=0.064$). The rate of NADM increased over time from 0.21 in 1994/1995 to 0.53 in 2014/2015 ($P=0.060$).

Risk factors for any malignancy are shown in Table 3. In multivariable analysis, female individuals [adjusted rate ratio (aRR) 0.63 (95% CI 0.45–0.89) vs. male individuals, $P=0.009$] and those from the Thai cohort [aRR 0.16 (0.04–0.69) vs. UK/Ireland, $P=0.017$] were at lower risk of a malignancy. Individuals who did not acquire HIV vertically were at lower risk than those who did so vertically [other modes of infection aRR 0.58 (0.31–0.98) vs. vertical, $P=0.049$]. Those with current severe immunosuppression [severe immunosuppression aRR 3.95 (2.48–6.29) vs. no severe immunosuppression, $P<0.001$] and viral load more than 400 copies/ml [aRR 1.80 (1.01–3.71) vs. ≤ 400 copies/ml, $P=0.027$] were more likely to have a malignancy. There was evidence of an interaction between current ART status and each of current calendar year and current age (Fig. 2). There was no change in risk over calendar time among those who were not on ART ($P=0.414$) or who had been on ART less than 6 months ($P=0.999$), but among those on ART for more than 6 months, the risk decreased over time [1996–2003 aRR 0.41 (0.23–0.74), 2004–2009 aRR 0.25 (0.12–0.51), ≥ 2010 aRR 0.22 (0.10–0.51) vs. <1996 , $P=0.001$]. There was a strong increase in risk with increasing current age among those not on




Fig. 1. Rates of malignancies over calendar time, overall and for AIDS-defining and non-AIDS-defining malignancies.

ART or on ART for less than 6 months, although less of an increase among those on ART at least 6 months. There was no evidence of an effect of current BMI-for-age z -score ($P=0.114$). In a supplementary analysis for individuals from the UK/Ireland and rest of Europe regions, there was no evidence of an effect of ethnicity or place of birth in the adjusted model ($P=0.588$, 0.955 , respectively, data not shown).

Risk factors for ADM and NADM are shown in Supplementary Tables 1 and 2, <http://links.lww.com/QAD/C163>. Risk factors for ADM were similar to those for any malignancy. For NADM, only immunosuppression and mode of infection were found to be associated; those with current severe immunosuppression were at higher risk [aRR 2.91 (1.06–7.97) vs. not severe, $P=0.038$] as were, in contrast to risk factors for any malignancy, those who did not acquire HIV vertically [aRR 3.13 (1.01–9.74) vs. vertical, $P=0.049$].

Overall, 58 (41%) patients with a malignancy event were reported to have died. Three deaths occurred in children with an unknown date of malignancy event, and the remaining 55 died at a median 2.4 (IQR 0.6–8.8) months (range 0.0–18.8 years) after malignancy diagnosis. Of 10 deaths among patients with an NADM, 8 (80%) were reported to be because of their NADM, and two (20%) because of an AIDS-defining event. Of 44 deaths among patients with an ADM, 27 (61%) were because of an AIDS-defining event, nine (20%) were reported as HIV-related, two (5%) because of an invasive bacterial infection, and the cause was unknown for six (14%).

The probability of death by 3 years after diagnosis was 41.1% (95% CI 33.1–50.3%). Those diagnosed with a malignancy before 1996 were more likely to die by 3 years [68.0% (51.9–83.1%)], with no change over time from

1996 onwards [30.6% (19.7–45.6%), 25.2% (12.1–47.8%), 33.7% (17.5–58.6%) for 1996–2003, 2004–2009 and after 2010, respectively] (Supplementary Figure 1, <http://links.lww.com/QAD/C163>). In multivariable analysis, only earlier calendar year of malignancy diagnosis and vertically acquired HIV were associated with higher risk of mortality (Supplementary Table 3, <http://links.lww.com/QAD/C163>).

Discussion

In this analysis of nearly 10 000 patients seen in paediatric HIV care across Eastern and Western Europe and Thailand, we assessed rates and risk factors for malignancies and malignancy outcomes among children and young people with HIV. The rate of ADM declined markedly following the introduction and widespread availability of combination ART for children in 1996, with some evidence of an increase in NADM, although rates were similar to those of any malignancy in the general population [1]. Increased risk of a malignancy was associated with older age among those not on treatment, male sex, being from a European cohort, vertical HIV acquisition, severe immunosuppression and viral load greater than 400 copies/ml. High mortality was observed following a malignancy diagnosis with no improvement in recent years.

Across the whole time period, the rate of any malignancy was 1.18 per 1000 person-years, approximately eight times higher than the estimated 0.14 per 1000 person-years for children and young people in the general population in Europe [1]. The rate of ADM decreased markedly in the late 1990s following the introduction and widespread availability of paediatric combination ART, in

Table 3. Risk factors for any malignancy.

	Univariable			Multivariable		
	Rate ratio	95% CI	P	Rate ratio	95% CI	P
Demographics, at entry to HIV care						
Sex						
Female	0.59	0.42–0.83	0.002	0.63	0.45–0.89	0.009
Male	1.00	–		1.00	–	
Region						
UK/Ireland	1.00	–	<0.001	1.00	–	0.017
Thailand	0.15	0.04–0.64		0.16	0.04–0.69	
Russia/Ukraine	0.39	0.18–0.86		0.66	0.29–1.54	
Rest of Europe	1.58	1.05–2.36		1.28	0.81–2.02	
Mode of HIV acquisition						
Vertical	1.00	–	0.055	1.00	–	0.049
Other	1.68	1.00–2.86		0.58	0.31–0.98	
Time-updated						
WHO immune stage						
Not severe	1.00	–	<0.001	1.00	–	<0.001
Severe	4.56	3.11–6.69		3.95	2.48–6.29	
Viral load						
Suppressed	1.00	–	0.003	1.00	–	0.027
Not suppressed	2.42	1.37–4.27		1.80	1.01–3.71	
BMI-for-age z-score						
>2	1.00	–	0.023	1.00	–	0.114
–2 to 2	2.29	0.38–13.75		2.01	0.32–12.53	
<–2	7.58	0.78–73.41		5.25	0.49–56.12	
ART status						
Not on ART	1.00	–	0.001			
<6 months on ART	3.59	1.78–7.23				
≥6 months on ART	1.68	1.10–2.57				
Calendar year						
<1996	1.00	–	<0.001	Test for interaction: $P = 0.144$		
1996–2003	0.60	0.39–0.94				
2004–2009	0.30	0.18–0.50				
≥2010	0.32	0.19–0.53				
Effect of time-updated calendar year among those not on ART						
			<1996	1.00	–	0.414
			1996–2003	1.41	0.52–3.84	
			2004–2009	0.63	0.17–2.42	
			≥2010	1.83	0.45–7.48	
Effect of time-updated calendar year among those on ART less than 6 months						
			<1996	1.00	–	0.999
			1996–2003	1.11	0.21–5.77	
			2004–2009	0.00	0.00 to >1000.00	
			≥2010	1.10	0.16–7.51	
Effect of time-updated calendar year among those on ART at least 6 months						
			<1996	1.00	–	0.001
			1996–2003	0.41	0.23, 0.74	
			2004–2009	0.25	0.12, 0.51	
			≥2010	0.22	0.10, 0.51	
Age (years)						
<5	1.00	–	0.033	Test for interaction: $P = 0.088$		
5 to <10	1.29	0.80–2.07				
10 to <15	1.46	0.90–2.38				
≥15	2.02	1.26–3.25				
Effect of time-updated age among those not on ART						
			<5	1.00	–	0.003
			5 to <10	2.52	0.98–6.46	
			10 to <15	7.17	2.55–20.14	
			≥15	5.43	0.63–47.07	
Effect of time-updated age among those on ART less than 6 months						
			<5	1.00	–	0.036
			5 to <10	3.63	0.60–21.80	
			10 to <15	7.95	1.38–45.88	
			≥15	17.65	2.261–137.72	
Effect of time-updated age among those on ART at least 6 months						
			<5	1.00	–	0.019
			5 to <10	0.95	0.52–1.70	
			10 to <15	1.09	0.56–2.12	
			≥15	1.18	1.18–4.83	

CI, confidence interval; ART, antiretroviral therapy.




Fig. 2. Rate ratios for association between any malignancy and current calendar year (a) and current age (b) by antiretroviral therapy status. Adjusted for sex, region, mode of HIV acquisition, WHO immune stage, viral suppression and BMI-for-age z-score.

line with findings from adult studies [4–6]. Conversely, the rate of NADM increased over time as reported elsewhere [15], which may be because of increased survival among those with HIV, although some NADM, including Hodgkin lymphoma, are known to be related to HIV [16]. Reassuringly, among virologically suppressed, nonimmunosuppressed patients on ART in our study, the rate of any malignancy was very low, 0.35 per 1000 person-years, although double the general population rate. There may be reasons for this higher rate beyond HIV status; key differences between our population and that in the general population study [1] are our inclusion of young people beyond 19 years of age and our wider calendar time period. Further, there is some evidence that the rate of malignancies is higher in children from black ethnic groups in the general population, and so higher than average rates may be expected in our study, regardless of HIV infection [17]. The potential carcinogenic effect of ART is unclear [18].

In adjusted analysis, children from the Thai cohort were at lower risk of a malignancy compared with those from Europe. One explanation for this may be differing exposures to oncogenic viruses. There is some evidence of a lower seroprevalence of HHV-8, which is associated

with Kaposi sarcoma, in Thailand compared with Europe and Africa [19,20]. A study estimating the risk of Kaposi sarcoma in children and adolescents reported no cases in Asia compared with rates between 11.4 and 85.8 per 100 000 person-years in sub-Saharan Africa and Europe [21]. The small number of Kaposi sarcoma events here precluded a similar analysis. Further, there is some evidence of lower rates of malignancies in the general paediatric population in Thailand compared with Europe (0.10 and 0.14 per 1000 person-years, respectively) [1,22].

The risk of a malignancy strongly increased with age among those not on or who had only recently started ART but there was less of an association, at least until adolescence, among those on treatment. Increasing risk among young people may be expected given the increased risk of malignancy in the general population in early adulthood [23]. Given that HIV infection is lifelong, the full elevated long-term risk may yet emerge, highlighting the importance of continued monitoring throughout adulthood of those with HIV acquired in childhood. Male individuals were at higher risk of ADM, which may be expected given an excess of non-Hodgkin lymphoma observed in the general male population [24].

Those with severe immune suppression and not virally suppressed were at higher risk, in line with other studies [9], reiterating the need for meeting UNAIDS targets for ART coverage, improving adherence, and reducing late diagnosis. Finally, those with vertically acquired HIV were shown to be at higher risk of ADM, yet lower risk of NADM, in adjusted analyses. Those with vertically acquired HIV are more likely to have been born abroad in our population, and so may have higher exposure to endemic oncogenic viruses, such as HHV-8 and EBV [25], although our analysis found no additional risk for those born abroad. Further, they had a longer duration of exposure to HIV. Reasons for the lower risk of NADM are unclear but may be related to their younger age, despite our adjustment for this.

Among those with a malignancy, all-cause mortality was high, with over 40% of children and young people reported to have died. In the general population, mortality by 3 years after a childhood cancer diagnosis in Europe has been estimated at 20% [26]. Although outcomes improved following the roll out of combination ART in 1996, worryingly, there was no improvement in mortality over time following this in our study, with approximately one-third of those diagnosed since 1996 dying within 3 years. This association was not explained by key characteristics, such as age at diagnosis, type of malignancy (ADM vs. NADM) or level of immunosuppression; vertically acquired HIV was the only other factor explored which was associated. In adults with HIV, malignancy outcomes have been shown to be broadly similar to the general population, although with poorer outcomes for some cancer types, including Hodgkin lymphoma [27,28]. Delayed malignancy diagnosis was believed to have contributed to poor outcomes in young adults diagnosed with a malignancy in the UK (some of whom are included here) [7]. Further, suboptimal treatment, perhaps because of concerns over the use of cancer treatment in children with immunosuppression [29], may have also contributed to the high mortality. No data on cancer treatment or stage at diagnosis, or any biological samples, were available to assess this here. Further, malignancy is associated with severe immunosuppression, which may have put patients at risk of dying from other causes. Among survivors, there may be an increased risk of secondary cancers, warranting regular screening [30].

Our study has several other limitations. Firstly, none of the cohorts were linked to cancer registries, so some cases may have been missed, with under-ascertainment and under diagnosis likely, especially in middle-income settings. However, estimated incidence rates were similar to other studies in children with HIV based on record linkage [8,31]. Secondly, limited data on co-infections with oncogenic viruses were available. Thirdly, despite the large size of our cohort, our ability to determine differences in risk factors for ADM and NADM was

limited by the relatively small number of NADM events, and low power resulted in wide confidence intervals when estimating rates over time. The major strength of this study is the long duration of follow-up and long calendar period covered as well as the inclusion of data following transfer to adult care.

In conclusion, in this analysis of nearly 10 000 children and young people with HIV across Europe and Thailand, we report a decline in the rates of ADM since the introduction of combination ART, with the rate of any malignancies among those on ART, virally suppressed and immunologically well similar to that in the general population. There was an increased risk with age, in particular, in those not currently on ART, which requires continued monitoring as young people with vertically acquired HIV progress into adulthood. Linkage of paediatric and adult registries, as well as HIV, malignancy and death registries, is crucial. Even in recent years, one-third of individuals with a malignancy died within 3 years of diagnosis, which is very concerning; possible causes should be explored and steps taken to address this.

Acknowledgements

We thank all the patients for their participation in these cohorts, and the staff members who cared for them.

Author contributors: Writing Group [consisting of Project Team first (ordered by contribution), followed by other Writing Group members (ordered alphabetically by country)].

Project Team: Elizabeth Chappell (EPPICC statistician), Anna Turkova (Irkutsk AIDS Centre, Russia), Tessa Goetghebuer (Hospital St Pierre Cohort, Brussels, Belgium), Charlotte Jackson (EPPICC epidemiologist), Elena Chiappini (Italian Register for HIV infection in children, Italy), Luisa Galli (Italian Register for HIV infection in children, Italy), Cosmina Gingaras ('Victor Babes' Hospital Cohort, Romania), Ali Judd (co-lead of EPPICC).

Other Writing Group members: Vana Spoulou (Greek cohort, Greece); Catuscia Lisi (Italian Register for HIV infection in children, Italy); Santa Anson (Latvian cohort, Latvia); Tom Wolfs (ATHENA paediatric cohort, Netherlands); Magdalena Marczyńska (Polish paediatric cohort, Poland); Luminita Ene ('Victor Babes' Hospital Cohort, Romania); Yulia Plotnikova (Irkutsk AIDS Centre, Russia); Evgeny Voronin (Republican Clinical Infectious Diseases Hospital' of the Ministry of Health of the Russian Federation, St Petersburg, Russia); Anna Samarina (The City HIV Centre, St Petersburg, Russia); Gonzague Jourdain, Nicole Ngo-Giang-Huong (Thailand Program for HIV Prevention and Treatment (PHPT) Study Group, Thailand); Claudia Fortuny

(CoRISPE-CAT, Catalonia cohort, Spain); Marisa Luisa Navarro, Jose Tomas Ramos (CoRISPE-S, rest of Spain cohort, Spain); Lars Navér (Karolinska University Hospital, Stockholm, Sweden); Pierre-Alex Crisinel (Swiss Mother and Child HIV Cohort Study, Switzerland); Heather Bailey, Ruslan Malyuta, Alla Volokha (Ukraine paediatric HIV cohort, Ukraine); Alasdair Bamford, Siobhan Crichton, Caroline Foster (Collaborative HIV Paediatric Study (CHIPS) and Integrated Screening Outcome Surveillance Service (ISOSS), UK and Ireland); Claire Thorne (European Collaborative Study); Intira Jeannie Collins, Carlo Giaquinto, Diana M. Gibb [EPPICC/Paediatric European Network for the Treatment of AIDS (PENTA)].

EPPICC/PENTA Co-ordinating Team: Elizabeth Chappell, Siobhan Crichton, Intira Jeannie Collins, Charlotte Duff, Carlo Giaquinto, Ruth Goodall, Daniel Gomez-pena, Charlotte Jackson, Ali Judd, Rebecca Lundin, Laura Mangiarini, Edith Milanzi, Alessandra Nardone, Claire Thorne.

Collaborating cohorts:

BELGIUM: Hospital St Pierre Cohort, Brussels: Tessa Goetghebuer, MD, PhD; Marc Hainaut, MD PhD; Evelyne Van der Kelen, Research nurse; Marc Delforge, data manager.

GREECE: Greek cohort: Vana Spoulou.

ITALY: Italian Register for HIV infection in Children. Coordinators: Maurizio de Martino, Luisa Galli (Florence), Pier Angelo Tovo, Clara Gabiano (Turin). Participants: Ines Carloni (Ancona), Domenico Larovere (Bari), Francesco Baldi, Angela Miniaci, Andrea Pession (Bologna) Raffaele Badolato (Brescia), Grazia Pantò (Catania), Elisa Anastasio (Catanzaro), Carlotta Montagnani, Elisabetta Venturini, Leila Bianchi (Florence), Alessandra Allodi, Antonio Di Biagio, Sara Grignolo (Genoa), Vania Giacommet, Paola Marchisio, Giuseppe Banderalli, Claudia Tagliabue (Milan), Monica Cellini (Modena), Eugenia Bruzzese, Pasquale Di Costanzo, Andrea Lo Vecchio (Naples), Carlo Giaquinto, Daniele Donà, Osvalda Rampon (Padua), Amelia Romano (Palermo), Icilio Dodi, Susanna Esposito (Parma), Valentina Zuccaro, Domenico Zanaboni (Pavia), Rita Consolini (Pisa), Stefania Bernardi, Orazio Genovese (Rome), Letizia Cristiano (Taranto), Antonio Mazza (Trento), Silvia Garazzino, Federica Mignone, Erika Silvestro (Turin), Vincenzo Portelli (Trapani).

LATVIA: Latvian Cohort (Santa Ansone).

NETHERLANDS: The ATHENA database is maintained by Stichting HIV Monitoring and supported by a grant from the Dutch Ministry of Health, Welfare and

Sport through the Centre for Infectious Disease Control of the National Institute for Public Health and the Environment.

Clinical centres (pediatric care)

Emma Kinderziekenhuis, Amsterdam, University Medical Centers: *HIV treating physicians:* D. Pajkrt, H.J. Scherpbier. *HIV nurse consultants:* A.M. Weijzenfeld, C.G. de Boer. *HIV clinical virologists/chemists:* S. Jurriaans, N.K.T. Back, H.L. Zaaijer, B. Berkhout, M.T.E. Cornelissen, C.J. Schinkel, K.C. Wolthers. Erasmus MC–Sophia, Rotterdam: *HIV treating physicians:* P.L.A. Fraaij, A.M.C. van Rossum, C.L. Vermont. *HIV nurse consultants:* L.C. van der Knaap, E.G. Visser. *HIV clinical virologists/chemists:* C.A.B. Boucher, M.P.G. Koopmans, J.J.A. van Kampen, S.D. Pas. Radboudumc, Nijmegen: *HIV treating physicians:* S.S.V. Henriët, M. van de Flier, K. van Aerde. *HIV nurse consultants:* R. Strik-Albers. *HIV clinical virologists/chemists:* J. Rahamat-Langendoen, F.F. Stelma. Universitair Medisch Centrum Groningen, Groningen: *HIV treating physicians:* E.H. Schölvinck. *HIV nurse consultants:* H. de Groot-de Jonge. *HIV clinical virologists/chemists:* H.G.M. Niesters, C.C. van Leer-Buter, M. Knoester. Wilhelmina Kinderziekenhuis, UMC Utrecht, Utrecht: *HIV treating physicians:* L.J. Bont, S.P.M. Geelen, T.F.W. Wolfs. *HIV nurse consultants:* N. Nauta. *HIV clinical virologists/chemists:* R. Schuurman, F. Verduyn-Lunel, A.M.J. Wensing.

Coordinating centre

Director: P. Reiss. *Deputy director:* S. Zaheri. *Data analysis:* D.O. Bezemer, A.I. van Sighem, C. Smit, F.W.M.N. Wit. *Data management and quality control:* M. Hillebrecht, A. de Jong, T. Woudstra. *Data monitoring:* D. Bergsma, S. Grivell, R. Meijering, M. Raethke, T. Rutkens. *Data collection:* L. de Groot, M. van den Akker, Y. Bakker, M. Bezemer, A. El Berkaoui, J. Geerlinks, J. Koops, E. Kruijne, C. Lodewijk, E. Lucas, R. van der Meer, L. Munjishvili, F. Paling, B. Peeck, C. Ree, R. Regtop, Y. Ruijs, L. van de Sande, M. Schoorl, P. Schnörr, E. Tuijn, L. Veenenberg, S. van der Vliet, A. Wisse, E.C. Witte. *Patient registration:* B. Tuk.

POLAND: Polish paediatric cohort: Head of the team: Professor Magdalena Marczyńska, MD, PhD. Members of the team: Jolanta Popielska, MD, PhD; Maria Pokorska-Śpiwak, MD, PhD; Agnieszka Ołdakowska, MD, PhD; Konrad Zawadka, MD, PhD; Urszula Coupland, MD, PhD; Administration assistant: Małgorzata Doroba. Affiliation: Medical University of Warsaw, Poland, Department of Children's Infectious Diseases; Hospital of Infectious Diseases in Warsaw, Poland.

ROMANIA: 'Victor Babes' Hospital Cohort, Bucharest: Dr Luminita Ene.

RUSSIA: Federal State-owned Institution 'Republican Clinical Infectious Diseases Hospital' of the Ministry of Health of the Russian Federation, St Petersburg: Liubov Okhonskaia, Evgeny Voronin, Milana Miloenko, Svetlana Labutina.

RUSSIA: The City HIV Centre, St Petersburg: Anna Samarina.

RUSSIA: Irkutsk AIDS Centre: Anna Turkova, Yulia Plotnikova.

SPAIN: CoRISPE-cat, Catalonia: financial support for CoRISPE-cat was provided by the Instituto de Salud Carlos III through the Red Temática de Investigación Cooperativa en Sida. Members: Hospital Universitari Vall d'Hebron, Barcelona [Pere Soler-Palacín, Maria Antoinette Frick and Santiago Pérez-Hoyos (statistician)], Hospital Universitari del Mar, Barcelona (Antonio Mur, Núria López), Hospital Universitari Germans Trias i Pujol, Badalona (María Méndez), Hospital Universitari Josep Trueta, Girona (Lluís Mayol), Hospital Universitari Arnau de Vilanova, Lleida (Teresa Vallmanya), Hospital Universitari Joan XXIII, Tarragona (Olga Calavia), Consorci Sanitari del Maresme, Mataró (Lourdes García), Hospital General de Granollers (Maite Coll), Corporació Sanitària Parc Taulí, Sabadell (Valentí Pineda), Hospital Universitari Sant Joan, Reus (Neus Rius), Fundació Althaia, Manresa (Núria Rovira), Hospital Son Espases, Mallorca (Joaquín Dueñas) and Hospital Sant Joan de Déu, Esplugues (Clàudia Fortuny, Antoni Noguera-Julian).

SPAIN: CoRISPE-S and Madrid cohort: María José Mellado, Luis Escosa, Milagros García Hortelano, Talía Sainz (Hospital La Paz); María Isabel González-Tomé, Pablo Rojo, Daniel Blázquez (Hospital Doce de Octubre, Madrid); José Tomás Ramos (Hospital Clínico San Carlos, Madrid); Luis Prieto, Sara Guillén (Hospital de Getafe); María Luisa Navarro, Jesús Saavedra, Mar Santos, María Angeles Muñoz, Beatriz Ruiz, Carolina Fernandez Mc Phee, Santiago Jimenez de Ory, Susana Alvarez (Hospital Gregorio Marañón); Miguel Ángel Roa (Hospital de Móstoles); José Beceiro (Hospital Príncipe de Asturias, Alcalá de Henares); Jorge Martínez (Hospital Niño Jesús, Madrid); Katie Badillo (Hospital de Torrejón); Miren Apilanez (Hospital de Donostia, San Sebastián); Itziar Pocheville (Hospital de Cruces, Bilbao); Elisa Garrote (Hospital de Basurto, Bilbao); Elena Colino (Hospital Insular Materno Infantil, Las Palmas de Gran Canaria); Jorge Gómez Sirvent (Hospital Virgen de la Candelaria, Santa Cruz de Tenerife); Mónica Garzón, Vicente Román (Hospital de Lanzarote); Abián Montesdeoca, Mercedes Mateo (Complejo Universitario de Canarias, La Laguna-Tenerife), María José Muñoz, Raquel Angulo (Hospital de Poniente, El Ejido); Olaf Neth, Lola Falcón (Hospital Virgen del Rocío, Sevilla); Pedro Terol

(Hospital Virgen de la Macarena, Sevilla); Juan Luis Santos (Hospital Virgen de las Nieves, Granada); David Moreno (Hospital Carlos Haya, Málaga); Francisco Lendínez (Hospital de Torrecárdenas, Almería); Ana Grande (Complejo Hospitalario Universitario Infanta Cristina, Badajoz); Francisco José Romero (Complejo Hospitalario de Cáceres); Carlos Pérez (Hospital de Cabueñes, Gijón); Miguel Lillo (Hospital de Albacete); Begoña Losada (Hospital Virgen de la Salud, Toledo); Mercedes Herranz (Hospital Virgen del Camino, Pamplona); Matilde Bustillo, Carmelo Guerrero (Hospital Miguel Servet, Zaragoza); Pilar Collado (Hospital Clínico Lozano Blesa, Zaragoza); José Antonio Couceiro (Complejo Hospitalario de Pontevedra); Amparo Pérez, Ana Isabel Piqueras, Rafael Bretón, Inmaculada Segarra (Hospital La Fe, Valencia); César Gavilán (Hospital San Juan de Alicante); Enrique Jareño (Hospital Clínico de Valencia); Elena Montesinos (Hospital General de Valencia); Marta Dapena (Hospital de Castellón); Cristina Álvarez (Hospital Marqués de Valdecilla, Santander); Ana Gloria Andrés (Hospital de León); Víctor Marugán, Carlos Ochoa (Hospital de Zamora); Santiago Alfayate, Ana Isabel Menasalvas (Hospital Virgen de la Arrixaca, Murcia); Elisa de Miguel (Complejo Hospitalario San Millán-San Pedro, Logroño) and Paediatric HIV-BioBank integrated in the Spanish AIDS Research Network and collaborating Centers.

Funding: This work has been partially funded by the Fundación para la Investigación y Prevención de SIDA en España (FIPSE) (FIPSE 3608229/09, FIPSE 240800/09, FIPSE 361910/10), Red Temática de Investigación en SIDA (RED RIS) supported by Instituto de Salud Carlos III (ISCIII) (RD12/0017/0035 and RD12/0017/0037), project as part of the Plan R+D+I and cofinanced by ISCIII-Subdirección General de Evaluación and Fondo Europeo de Desarrollo Regional (FEDER), Mutua Madrileña 2012/0077, Gilead Fellowship 2013/0071, FIS PI15/00694, CoRISpe (RED RIS RD06/0006/0035 y RD06/0006/0021).

SWEDEN: Karolinska University Hospital, Stockholm (Lars Navér, Sandra Soeria-Atmadja, Erik Belfrage, Vendela Hagås).

SWITZERLAND: *Members of the Swiss HIV Cohort Study (SHCS) and the Swiss Mother and Child HIV Cohort Study:* Aebi-Popp K, Anagnostopoulos A, Asner S, Battegay M, Baumann M, Bernasconi E, Böni J, Braun DL, Bucher HC, Calmy A, Cavassini M, Ciuffi A, Duppenhaler A, Dollenmaier G, Egger M, Elzi L, Fehr J, Fellay J, Francini K, Furrer H, Fux CA, Grawe C, Günthard HF (President of the SHCS), Haerry D (deputy of 'Positive Council'), Hasse B, Hirsch HH, Hoffmann M, Hösli I, Huber M, Kahlert CR (Chairman of the Mother & Child Substudy), Kaiser L, Keiser O, Klimkait T, Kottanattu L, Kouyos RD, Kovari H, Ledergerber B, Martinetti G, Martinez de Tejada B, Marzolini C, Metzner KJ, Müller

N, Nicca D, Paioni P, Pantaleo G, Perreau M, Polli Ch, Rauch A (Chairman of the Scientific Board), Rudin C, Scherrer AU (Head of Data Centre), Schmid P, Speck R, Stöckle M (Chairman of the Clinical and Laboratory Committee), Tarr P, Thanh Lecompte M, Trkola A, Vernazza P, Wagner N, Wandeler G, Weber R, Wyler CA, Yerly S.

Funding: This study has been financed within the framework of the Swiss HIV Cohort Study, supported by the Swiss National Science Foundation (grant #177499).

THAILAND: Program for HIV Prevention & Treatment (PHPT). Participating hospitals: Lamphun: Pornpun Wannarit; Phayao Provincial Hospital: Pornchai Techakunakorn; Chiangrai Prachanukroh: Rawiwan Hansudewechakul; Chiang Kham: Vanichaya Wanchaitanawong; Phan: Sookchai Theansavetrakul; Mae Sai: Sirisak Nanta; Prapokkiao: Chaiwat Ngampiyakul; Banglamung: Siriluk Phanomcheong; Chonburi: Suchat Hongsiwong; Rayong: Warit Karnchanamayul; Bhuddasothorn Chacheongsao: Ratchanee Kwanchaipanich; Nakornping: Suparat Kanjanavanit; Somdej Prapinkiao: Nareerat Kamonpakorn, Maneeratn Nantarukchaikul; Bhumibol Adulyadej: Prapaisri Layangool, Jutarat Mekmullica; Pranangkiao: Paiboon Lucksanapisitkul, Sudarat Watanayothin; Buddhachinaraj: Narong Lertpienthum; Hat Yai: Boonyarat Warachit; Regional Health Promotion Center 6, Khon Kaen: Sansanee Hanpinitak; Nong Khai: Sathit Potchalongsin; Samutsakhon: Pimpraphai Thanasiri, Sawitree Krikajornkitti; Phaholpolphayuhasena: Pornsawan Attaviniytrakarn; Kalasin: Sakulrat Sirojana; Nakhonpathom: Suthunya Bunjongpak; Samutprakarn: Achara Puangsombat; Maharakam: Sathaporn Na-Rajsima; Roi-et: Pornchai Ananpatharachai; Sanpatong: Noppadon Akarathum; Vachira Phuket: Weerasak Lawtongkum; Chiangdao: Prapawan Kheunjan, Thitiporn Suriyaboon, Airada Saipanya.

Data management team: Kanchana Than-in-at, Nirattiya Jaisieng, Rapeepan Suaysod, Sanuphong Chailoet, Naritsara Naratee, and Suttipong Kawilapat.

UKRAINE: Paediatric HIV Cohort: Dr T. Kaleeva, Dr Y. Baryshnikova (Odessa Regional Centre for HIV/AIDS); Dr S. Soloha (Donetsk Regional Centre for HIV/AIDS); Dr N. Bashkatova (Mariupol AIDS Center); Dr I. Raus (Kiev City Centre for HIV/AIDS); Dr O. Glutshenko, Dr Z. Ruban (Mykolaiv Regional Centre for HIV/AIDS); Dr N. Prymak (Kryvyi Rih); Dr G. Kiseleva (Simferopol); Dr Alla Volokha (Shupyk National Medical Academy of Postgraduate Education); Dr Ruslan Malyuta (Perinatal Prevention of AIDS Initiative, Odessa); Dr H. Bailey, Prof Claire Thorne (UCL, London, UK). Funding acknowledgement: PENTA Foundation.

UK AND IRELAND: Collaborative HIV Paediatric Study (CHIPS): CHIPS is funded by the NHS (London Specialised Commissioning Group) and has received additional support from Abbott, Boehringer Ingelheim, Bristol-Myers Squibb, GlaxoSmithKline, Gilead Sciences, Janssen and Roche. The MRC Clinical Trials Unit at UCL is supported by the Medical Research Council (<https://www.mrc.ac.uk>) programme number MC_UU_12023/26.

CHIPS Steering Committee: Hermione Lyall (chair), Alasdair Bamford, Karina Butler, Katja Doerholt, Conor Doherty, Caroline Foster, Ian Harrison, Julia Kenny, Nigel Klein, Gillian Letting, Paddy McMaster, Fungai Murau, Edith Nsangi, Katia Prime, Andrew Riordan, Fiona Shackley, Delane Shingadia, Sharon Storey, Gareth Tudor-Williams, Anna Turkova, Steve Welch. **MRC Clinical Trials Unit:** Intira Jeannie Collins, Claire Cook, Siobhan Crichton, Donna Dobson, Keith Fairbrother, Diana M. Gibb, Ali Judd, Marthe Le Prevost, Nadine Van Looy. **Integrated Screening Outcome Surveillance Service (ISOSS), UCL:** Helen Peters, Kate Francis, Claire Thorne.

Hospitals participating in CHIPS in 2019/20: University Hospitals Birmingham NHS Foundation Trust, Birmingham: L. Thrasyvoulou, S. Welch; Brighton and Sussex University Hospitals NHS Trust: K. Fidler; University Hospitals Bristol NHS Foundation Trust, Bristol: J. Bernatoniene, F. Manyika; Calderdale and Huddersfield NHS Foundation Trust, Halifax: G. Sharpe; Derby Teaching Hospitals NHS Foundation Trust: B. Subramaniam; Glasgow Royal Hospital for Children, Glasgow: R. Hague, V. Price; Great Ormond Street Hospital for Children NHS Foundation Trust, London: J. Flynn, A. Cardoso, M. Abou – Rayyah, N. Klein, A. Bamford, D. Shingadia; Oxford University Hospitals NHS Foundation Trust, Oxford: S. Yeadon, S. Segal; King's College Hospital NHS Foundation Trust, London: S. Hawkins; Leeds Teaching Hospitals NHS Trust, Leeds: M. Dowie; University Hospitals of Leicester NHS Trust, Leicester: S. Bandi, E. Percival; Luton and Dunstable Hospital NHS Foundation Trust, Luton: M. Eisenhut; K. Duncan; Milton Keynes General University Hospital NHS Foundation Trust, Milton Keynes: L. Anguava, L. Wren, Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle: T. Flood, A. Pickering; The Pennine Acute Hospitals NHS Trust, Manchester: P. McMaster, C. Murphy; North Middlesex University Hospital NHS Trust, London: J. Daniels, Y. Lees; Northampton General Hospital NHS Trust, Northampton: F. Thompson; London North West Healthcare NHS Trust, Middlesex: A. Williams, B. Williams, S. Pope; Barts Health NHS trust, London: Dr S. Libeschutz; Nottingham University Hospitals NHS Trust, Nottingham: L. Cliffe, S. Southall; Portsmouth Hospitals NHS Trust, Portsmouth: A. Freeman; Raigmore Hospital, Inverness: H. Freeman;

Royal Belfast Hospital for Sick Children, Belfast: S. Christie; Royal Berkshire NHS Foundation Trust, Reading: A. Gordon; Royal Children's Hospital, Aberdeen: D. Rosie Hague, L. Clarke; Royal Edinburgh Hospital for Sick Children, Edinburgh: L. Jones, L. Brown; Royal Free NHS Foundation Trust, London: M. Greenberg; Alder Hey Children's NHS Foundation Trust, Liverpool: C. Benson, A. Riordan; Sheffield Children's NHS Foundation Trust, Sheffield: L. Ibber-son, F. Shackley; University Hospital Southampton NHS Foundation Trust, Southampton: S. Patel, J. Hancock; St George's University Hospitals NHS Foundation Trust, London: K. Doerholt, K. Prime, M. Sharland, S. Storey; Imperial College Healthcare NHS Trust, London: E.G.H. Lyall, C. Foster, P. Seery, G. Tudor-Williams, N Kirkhope, S Raghunanan; Guy's and St Thomas' NHS Foundation Trust, London: Dr Julia Kenny, A. Callaghan; University Hospitals of North Midlands NHS Trust, Stoke On Trent: A Bridgwood, P. McMaster; University Hospital of Wales, Cardiff: J. Evans, E. Blake; NHS Frimley Health Foundation Trust, Slough: A. Yannoulis.

Funding statement: EPPICC receives funding from the PENTA Foundation (<http://penta-id.org>), and received support from the European Union's Horizon 2020 research and innovation programme under Grant Agreement No 825579 for the REACH study. The MRC Clinical Trials Unit at UCL is supported by the Medical Research Council (programme number MC_UU_12023/26).

Conflicts of interest

C.T. has received grant funding from ViiV Healthcare via the Penta Foundation (to UCL), and personally from ViiV for participation on an advisory board. C.F. has received funds for speaking at symposia organized on behalf of Gilead Sciences.

References

1. Steliarova-Foucher E, Fidler MM, Colombet M, Lacour B, Kaatsch P, Piñeros M, et al., ACCIS contributors. **Changing geographical patterns and trends in cancer incidence in children and adolescents in Europe, 1991–2010 (Automated Childhood Cancer Information System): a population-based study.** *Lancet Oncol* 2018; **19**:1159–1169.
2. Patel P, Hanson DL, Sullivan PS, Novak RM, Moorman AC, Tong TC, et al. **Incidence of types of cancer among HIV-infected persons compared with the general population in the United States, 1992–2003.** *Ann Intern Med* 2008; **148**:728–736.
3. Singh E, Naidu G, Davies M-A, Bohlius J. **HIV-associated malignancies in children.** *Curr Opin HIV AIDS* 2017; **12**:77–83.
4. Hessel NA, Whittemore H, Vittinghoff E, Hsu LC, Ma D, Scheer S, Schwarcz SK. **Incidence of first and second primary cancers diagnosed among people with HIV, 1985–2013: a population-based, registry linkage study.** *Lancet HIV* 2018; **5**:e647–e655.

5. Shiels MS, Pfeiffer RM, Gail MH, Hall HI, Li J, Chaturvedi AK, et al. **Cancer burden in the HIV-infected population in the United States.** *J Natl Cancer Inst* 2011; **103**:753–762.
6. Smith CJ, Ryom L, Weber R, Morlat P, Pradier C, Reiss P, et al., D:A:D Study Group. **Trends in underlying causes of death in people with HIV from 1999 to 2011 (D:A:D): a multicohort collaboration.** *Lancet* 2014; **384**:241–248.
7. Chhabra S, Fidler S, Ayers S, Bower M, Lyall H, Foster C. **Malignancy and all-cause mortality; incidence in adolescents and young adults living with perinatally acquired HIV.** *J Virus Erad* 2020; **6**:30–33.
8. Bohlius J, Maxwell N, Spoerri A, Wainwright R, Sawry S, Poole J, et al., for leDEA-Southern Africa. **Incidence of AIDS-defining and other cancers in HIV-positive children in South Africa: Record Linkage Study.** *Pediatr Infect Dis J* 2016; **35**:e164–e170.
9. Kest H, Brogly S, McSherry G, Dashefsky B, Oleske J, Seage GR 3rd. **Malignancy in perinatally human immunodeficiency virus-infected children in the United States.** *Pediatr Infect Dis J* 2005; **24**:237–242.
10. Irira M, Ngocho JS, Youze J, Shayo I, Komba V, Minja L, et al. **Prevalence and outcome of HIV-associated malignancies among HIV-infected children enrolled into care at Kilimanjaro Christian Medical Center 2006 to 2014: a hospital-based retrospective analytical study.** *J Pediatr Hematol Oncol* 2020; **42**:69–73.
11. Centers for Disease Control and Prevention. **Revised surveillance case definition for HIV infection—United States, 2014.** *MMWR Morbid Mortal Wly Rep* 2014; **63**:1–10.
12. World Health Organization. **WHO case definitions of HIV for surveillance and revised clinical staging and immunological classification of HIV-related disease in adults and children.** Geneva: World Health Organization; 2007.
13. World Health Organization. **WHO child growth standards: length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: methods and development.** Geneva: World Health Organization; 2006.
14. Rubin DB. **Multiple imputation for nonresponse in surveys.** Vol. 81. Hoboken, NJ: John Wiley & Sons; 2004.
15. Shiels MS, Engels EA. **Evolving epidemiology of HIV-associated malignancies.** *Current Opinion in HIV and AIDS* 2017; **12**:6–11.
16. Powles T, Robinson D, Stebbing J, Shamash J, Nelson M, Gazzard B, et al. **Highly active antiretroviral therapy and the incidence of non-AIDS-defining cancers in people with HIV infection.** *J Clin Oncol* 2009; **27**:884–890.
17. Sayeed S, Barnes I, Ali R. **Childhood cancer incidence by ethnic group in England, 2001–2007: a descriptive epidemiological study.** *BMC Cancer* 2017; **17**:570.
18. Borges AH. **Combination antiretroviral therapy and cancer risk.** *Curr Opin HIV AIDS* 2017; **12**:12–19.
19. Ayuthaya PIN, Katano H, Inagi R, Auwanit W, Sata T, Kurata T, et al. **The seroprevalence of human herpesvirus 8 infection in the Thai population.** *Southeast Asian J Trop Med Public Health* 2002; **33**:297–305.
20. Zhang T, Wang L. **Epidemiology of Kaposi's sarcoma-associated herpesvirus in Asia: challenges and opportunities.** *J Med Virol* 2017; **89**:563–570.
21. Pediatric AIDS-Defining Cancer Project Working Group for leDea Southern Africa TAPHOD Cohere in EuroCoord. Rohner E, Schmidlin K, Zwahlen M, Chakraborty R, Clifford G, et al. **Kaposi sarcoma risk in HIV-infected children and adolescents on combination antiretroviral therapy from sub-Saharan Africa, Europe, and Asia.** *Clin Infect Dis* 2016; **63**:1245–1253.
22. Bidwell SS, Peterson CC, Demanelis K, Zarins KR, Meza R, Sripilung H, et al. **Childhood cancer incidence and survival in Thailand: a comprehensive population-based registry analysis, 1990–2011.** *Pediatr Blood Cancer* 2019; **66**:e27428.
23. Cook MB, Dawsey SM, Freedman ND, Inskip PD, Wichner SM, Quraishi SM, et al. **Sex disparities in cancer incidence by period and age.** *Cancer epidemiology, biomarkers & prevention: a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* 2009; **18**:1174–1182.
24. Kaatsch P. **Epidemiology of childhood cancer.** *Cancer Treat Rev* 2010; **36**:277–285.
25. Rohner E, Wyss N, Trelle S, Mbulaiteye SM, Egger M, Novak U, et al. **HHV-8 seroprevalence: a global view.** *Syst Rev* 2014; **3**:11–111.

26. Gatta G, Botta L, Rossi S, Aareleid T, Bielska-Lasota M, Clavel J, *et al.* **Childhood cancer survival in Europe 1999–2007: results of EUROCare-5—a population-based study.** *Lancet Oncol* 2014; **15**:35–47.
27. Cingolani A, Cozzi Lepri A, Teofili L, Galli L, Mazzotta V, Baldin GM, *et al.*, ICONA Foundation Study group. **Survival and predictors of death in people with HIV-associated lymphoma compared to those with a diagnosis of lymphoma in general population.** *PloS One* 2017; **12**:e0186549–e1186549.
28. Trickey A, May MT, Gill MJ, Grabar S, Vehreschild J, Wit F, *et al.* **Cause-specific mortality after diagnosis of cancer among HIV-positive patients: a collaborative analysis of cohort studies.** *Int J Cancer* 2020; **146**:3134–3146.
29. Stefan DC. **Effect of HIV infection on the outcome of cancer therapy in children.** *Lancet Oncol* 2014; **15**:e562–e567.
30. Bohlius J, Foster C, Naidu G, Sengayi M, Turkova A. **Cancer in adolescents and young adults living with HIV.** *Curr Opin HIV AIDS* 2018; **13**:196–203.
31. Simard EP, Shiels MS, Bhatia K, Engels EA. **Long-term cancer risk among people diagnosed with AIDS during childhood.** *Cancer Epidemiol Biomarkers Prev* 2012; **21**: 148–154.