



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

The classical obstacle problem with Hölder continuous coefficients

Questa è la versione Preprint (Submitted version) della seguente pubblicazione:

Original Citation:

The classical obstacle problem with Hölder continuous coefficients / Andreucci G.; Focardi M.. - In: NONLINEAR ANALYSIS. - ISSN 0362-546X. - STAMPA. - 226:(2023), pp. 113139.1-113139.17. [10.1016/j.na.2022.113139]

Availability:

The webpage <https://hdl.handle.net/2158/1286023> of the repository was last updated on 2022-10-26T11:19:44Z

Published version:

DOI: 10.1016/j.na.2022.113139

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

Conformità alle politiche dell'editore / Compliance to publisher's policies

Questa versione della pubblicazione è conforme a quanto richiesto dalle politiche dell'editore in materia di copyright.

This version of the publication conforms to the publisher's copyright policies.

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

THE CLASSICAL OBSTACLE PROBLEM WITH HÖLDER CONTINUOUS COEFFICIENTS

GIOVANNA ANDREUCCI AND MATTEO FOCARDI

*This paper is dedicated to Prof. Emmanuele Di Benedetto
for having taught us, with his body of work, the taste for hard analysis problems.*

ABSTRACT. Weiss' and Monneau's type quasi-monotonicity formulas are established for quadratic energies having matrix of coefficients which are Dini, double-Dini continuous, respectively. Free boundary regularity for the corresponding classical obstacle problems under Hölder continuity assumptions is then deduced.

1. INTRODUCTION

In the last years several contributions have been devoted to the extension of the regularity theory for obstacle type problems to the case in which the involved linear elliptic operator in divergence form has coefficients with low regularity [1, 17, 2, 3, 12, 22, 23, 30, 18, 14, 20]. The aim of this note is to make another step in that direction for the classical obstacle problem by establishing Weiss' and Monneau's quasi-monotonicity formulas for quadratic forms having matrix of coefficients which are Dini, double-Dini continuous, respectively (for the sake of simplicity the lower order terms have the same regularity). Such results are instrumental to pursue the variational approach to establish the smoothness of the corresponding free boundaries. To keep the presentation as simple as possible the related free boundary analysis is performed only in the Hölder continuous case.

More precisely, in what follows we consider the functional $\mathcal{E} : W^{1,2}(\Omega) \rightarrow \mathbb{R}$ given by

$$\mathcal{E}(v) := \int_{\Omega} (\langle \mathbb{A}(x) \nabla v(x), \nabla v(x) \rangle + 2h(x)v(x)) \, dx, \quad (1.1)$$

and study regularity issues related to its unique minimizer w on the set

$$\mathcal{K}_{\psi,g} := \{v \in W^{1,2}(\Omega) : v \geq \psi \text{ } \mathcal{L}^n\text{-a.e. on } \Omega, \text{Tr}(v) = g \text{ on } \partial\Omega\}.$$

Existence and uniqueness of the minimizer is a standard issue for the problem under the ensuing hypotheses. For a bounded open set $\Omega \subset \mathbb{R}^n$, $n \geq 2$, we consider in what follows functions $\psi \in C_{\text{loc}}^{1,1}(\Omega)$ and $g \in H^{1/2}(\partial\Omega)$, such that $\psi \leq g$ \mathcal{H}^{n-1} -a.e on $\partial\Omega$, a matrix-valued field $\mathbb{A} : \Omega \rightarrow \mathbb{R}^{n \times n}$ and a function $f : \Omega \rightarrow \mathbb{R}$ satisfying:

(H1) $\mathbb{A}(x) = (a_{ij}(x))_{i,j=1,\dots,n} \in L^{\infty}(\Omega, \mathbb{R}^{n \times n})$ is symmetric and coercive, that is $a_{ij} = a_{ji}$ \mathcal{L}^n -a.e. in Ω for all $i, j \in \{1, \dots, n\}$, and for some $\Lambda \geq 1$

$$\Lambda^{-1}|\xi|^2 \leq \langle \mathbb{A}(x)\xi, \xi \rangle \leq \Lambda|\xi|^2 \quad (1.2)$$

2010 *Mathematics Subject Classification.* Primary: 35R35, 49N60.

Key words and phrases. Classical obstacle problem, free boundary, monotonicity formulas.

M. F. is a member of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

for \mathcal{L}^n -a.e. $x \in \Omega$, and for all $\xi \in \mathbb{R}^n$;

(H2) $f \in L^\infty(\Omega)$, $f := h - \operatorname{div}(\mathbb{A} \nabla \psi) > c_0$ \mathcal{L}^n -a.e. on Ω , for some $c_0 > 0$;

(H3) \mathbb{A} and f are α -Hölder continuous, for some $\alpha \in (0, 1]$,

(cf. Section 2 for the notation). Under assumptions (H1)-(H3) we establish a stratification result for the free boundary of solutions (see also Remark 4.1 for further possible extensions).

Theorem 1.1. *Assume (H1)-(H3), and let w be the (unique) minimizer of \mathcal{E} in (1.1) on $\mathcal{K}_{\psi,g}$.*

Then, w is $C_{\text{loc}}^{1,\alpha}(\Omega)$, and the free boundary decomposes as $\partial\{w = \psi\} \cap \Omega = \operatorname{Reg}(w) \cup \operatorname{Sing}(w)$, where $\operatorname{Reg}(w)$ and $\operatorname{Sing}(w)$ are called its regular and singular part, respectively. Moreover, $\operatorname{Reg}(w) \cap \operatorname{Sing}(w) = \emptyset$ and

- (i) *$\operatorname{Reg}(w)$ is relatively open in $\partial\{w = \psi\}$ and, for every point $x_0 \in \operatorname{Reg}(w)$, there exist $r = r(x_0) > 0$ such that $\partial\{w = \psi\} \cap B_r(x)$ is a $C^{1,\beta}$ $(n-1)$ -dimensional manifold, for some exponent $\beta \in (0, 1)$.*
- (ii) *$\operatorname{Sing}(w) = \cup_{k=0}^{n-1} S_k$, with S_k contained in the union of at most countably many sub-manifolds of dimension k and class C^1 .*

In the case of smooth matrix fields, Theorem 1.1 collects the fundamental contributions of Caffarelli to the classical obstacle problem (cf. for instance [4, 5, 7, 6] and the books [21, 8, 29] for more details and references also on related issues).

In the last years Theorem 1.1 has been extended to the case in which \mathbb{A} either is Lipschitz continuous in [12] or belongs to a fractional Sobolev space $W^{1+s,p}$ in [18], with $sp > 1$ and $p \geq n \wedge \frac{n^2}{n(1+s)-1}$, or belongs to the Sobolev space $W^{1,p}$ with $p > n$ in [14]. We point out that in all those cases the involved matrix fields \mathbb{A} turns out to be Hölder continuous in view of Sobolev type embeddings. Applications of the results in [12, 14] to the study of the obstacle problem for a large class of nonlinear energies were then given in [13]. In addition, counterexamples to smoothness of the free boundary are shown in case of the Laplacian if f is not Dini continuous in [1], and in case f is constant and \mathbb{A} is VMO in [3, Remark 4.2]. In this respect, if both the coefficients \mathbb{A} and f are assumed to be in VMO, Reifenberg vanishing flatness of $\operatorname{Reg}(u)$ has been established in [3].

The papers [12, 18, 14] follow the variational approach to free boundary analysis remarkably developed by Weiss [31] and by Monneau [27]. Central to this method are quasi-monotonicity formulas for suitable quantities, named in literature as Weiss' and Monneau's quasi-monotonicity formulas. The original papers by Weiss and Monneau are based on some key integral identities that hold true in the case of the Laplacian. The search for generalizations need several new insights and technical tools. Let us briefly recall the key ideas introduced in [12, 18, 14]. An extension of the Rellich and Nečas' inequality due to Payne and Weinberger (cf. [24]) is employed in the papers [12, 18]. On a technical side, the matrix field \mathbb{A} is differentiated, therefore enough smoothness of \mathbb{A} is needed. Instead, a different approach, leading to a weakening of the regularity assumptions on \mathbb{A} , has been pursued in [14] following an observation in [12, Remark 4.9]. The main difference in [14] with respect to [12, 18] concerns the monotone quantity itself: rather than considering the natural quadratic energy associated to the obstacle problem under study, a related constant coefficient quadratic form has been used. Higher order regularity of solutions, namely $W_{\text{loc}}^{2,p}$, in combination with their well-known quadratic growth from free boundary points then allow to conclude. In this note we push forward the ideas in [14] and extend the conclusions there to Hölder continuous matrix fields \mathbb{A} for which the above mentioned $W_{\text{loc}}^{2,p}$ regularity of solutions is not guaranteed.

More precisely, we prove Weiss' formula for Dini continuous matrix fields \mathbb{A} and Monneau's formula for double-Dini continuous ones. For simplicity we do not distinguish the degrees of regularity of \mathbb{A} and that of f . Let us stress that in the case of the Laplacian, low regularity assumptions on f have already been considered: Dini continuity has been deeply investigated in [1], and Dini continuity in L^p sense, p suitable, in [28] (cf. also [3] for both \mathbb{A} and f in VMO; [14] for Theorem 1.1 if \mathbb{A} is $W^{1,p}$, $p > n$, and f double-Dini continuous). A potential extension of the related free boundary analysis contained in Theorem 1.1 is discussed in Remark 4.1 (cf. also Remark 3.5). Note that if $\alpha = 1$ we recover the results in [12] with, partially, a different proof.

We close this introduction briefly resuming the structure of the paper: standard preliminaries for the classical obstacle problem are collected in Section 2. The mentioned generalizations of Weiss' and Monneau's quasi-monotonicity formulas are dealt with in Section 3, finally Section 4 contains the applications to the free boundary stratification problem.

2. PRELIMINARIES

2.1. Notation. Throughout the whole paper, the inner product in \mathbb{R}^n is denoted by $\langle \cdot, \cdot \rangle$, and the induced norm by $|\cdot|$. To distinguish it from the norm in $\mathbb{R}^{n \times n}$ we use the symbol $\|\cdot\|$ for the latter. We use standard notations for Lebesgue, Sobolev and Hölder spaces, quoting the necessary results when needed.

2.2. Reduction to the zero obstacle case. We first reduce ourselves to the zero obstacle problem. Let w be the unique minimizer of \mathcal{E} over $\mathcal{K}_{\psi,g}$, and define $u := w - \psi$. Then, u is the unique minimizer of

$$\mathcal{E}(v) := \int_{\Omega} (\langle \mathbb{A}(x) \nabla v(x), \nabla v(x) \rangle + 2f(x)v(x)) \, dx, \quad (2.1)$$

over $\mathbb{K}_{\psi,g} := \mathcal{K}_{0,\psi-g}$, where $f = h - \operatorname{div}(\mathbb{A} \nabla \psi)$. Clearly, $\partial\{w = \psi\} \cap \Omega = \partial\{u = 0\} \cap \Omega$, therefore we shall establish all the results in Theorem 1.1 for u . Notice that assumption (H2) is formulated exactly in terms of f , and that, for the zero obstacle problem, i.e. $\psi = 0$ the positivity assumption on f in (H2) involves only the lower order term h in the integrand and not the matrix field \mathbb{A} . It is also clear that given (H3), the L^∞ assumptions in (H1) and (H2) are redundant and the inequalities there hold in a pointwise sense. Despite this, we state (H1) and (H2) as in the Introduction because for some of the ensuing results we shall not need (H3) and we will sometimes substitute it with a weaker version (cf. (H4) and (H5) in what follows).

The first step is to prove that u satisfies a PDE both in the distributional sense and \mathcal{L}^n -a.e. on Ω . The next result has been established in [13, Proposition 3.2] more generally for variational inequalities inspired by some arguments in [31] (see also [15, 16]). The leading vector field in [13, Proposition 3.2] is assumed to be $C_{\text{loc}}^{1,1}$ in the full set of variables for the sake of existence of solutions in that general setting. On the contrary, such a regularity in the space variable is never used to deduce the conclusion of [13, Proposition 3.2]. We provide below the proof for the readers' convenience.

Proposition 2.1. *Assume (H1) and (H2). Let u be the minimizer of \mathcal{E} on $\mathbb{K}_{\psi,g}$. Then, there exists a function $\zeta \in L^\infty(\Omega)$ such that*

$$\operatorname{div}(\mathbb{A} \nabla u) = f - \zeta \quad (2.2)$$

\mathcal{L}^n -a.e. on Ω and in $\mathcal{D}'(\Omega)$, with

$$0 \leq \zeta \leq f\chi_{\{u=0\}}. \quad (2.3)$$

In particular, $u \in C_{\text{loc}}^{0,\beta}(\Omega)$ for some $\beta \in (0, 1)$.

Proof. Let $\varphi \in C_c^\infty(\Omega)$, and for all $\varepsilon > 0$ define $v_\varepsilon := (u + \varepsilon\varphi) \vee 0$. Note that v_ε belongs to the set $\mathbb{K}_{\psi,g}$. If $\varphi \geq 0$ then $v_\varepsilon = u + \varepsilon\varphi$ and, since u is the minimizer we have

$$-\frac{\varepsilon}{2} \int_{\Omega} \langle \mathbb{A}(x) \nabla \varphi, \nabla \varphi \rangle dx \leq \int_{\Omega} \langle \mathbb{A}(x) \nabla u, \nabla \varphi \rangle dx + \int_{\Omega} f(x) \varphi dx.$$

In turn, for $\varepsilon \rightarrow 0$ this implies that

$$0 \leq \int_{\Omega} \langle \mathbb{A}(x) \nabla u, \nabla \varphi \rangle dx + \int_{\Omega} f(x) \varphi dx \quad \forall \varphi \in C_c^\infty(\Omega), \varphi \geq 0.$$

The last inequality yields that the distribution $\mu := -\operatorname{div}(\mathbb{A}(\cdot) \nabla u) + f(\cdot) \mathcal{L}^n \llcorner \Omega$ is a non-negative Radon measure on Ω .

Let now φ and v_ε be as above without any assumption on the sign of φ , and set

$$\Omega_\varepsilon := \{x \in \Omega : u + \varepsilon\varphi < 0\}.$$

It is clear that $v_\varepsilon = 0$ and $\varphi < 0$ on Ω_ε , and that $v_\varepsilon = u + \varepsilon\varphi$ on $\Omega \setminus \Omega_\varepsilon$. Using again that u is minimizing, that $f \geq c_0 > 0$ on Ω by (H2), and that $v_\varepsilon \in \mathbb{K}_{\psi,g}$ we obtain

$$\begin{aligned} 0 \leq \mathcal{E}(v_\varepsilon) - \mathcal{E}(u) &= - \int_{\Omega_\varepsilon} \langle \mathbb{A} \nabla u, \nabla u \rangle dx + 2\varepsilon \int_{\Omega \setminus \Omega_\varepsilon} \langle \mathbb{A} \nabla u, \nabla \varphi \rangle dx \\ &\quad + \varepsilon^2 \int_{\Omega \setminus \Omega_\varepsilon} \langle \mathbb{A} \nabla \varphi, \nabla \varphi \rangle dx + 2 \int_{\Omega \setminus \Omega_\varepsilon} f(u + \varepsilon\varphi) dx - 2 \int_{\Omega} f u dx \\ &\leq 2\varepsilon \int_{\Omega} \langle \mathbb{A} \nabla u, \nabla \varphi \rangle dx + 2\varepsilon \int_{\Omega} f \varphi dx - 2\varepsilon \int_{\Omega_\varepsilon} \langle \mathbb{A} \nabla u, \nabla \varphi \rangle dx \\ &\quad + \varepsilon^2 \int_{\Omega \setminus \Omega_\varepsilon} \langle \mathbb{A} \nabla \varphi, \nabla \varphi \rangle dx - \int_{\Omega_\varepsilon} 2f(u + \varepsilon\varphi) dx. \end{aligned}$$

In the last inequality we have dropped the first term on the first line as it is negative (cf. (H1)). Thus, on account of the definition of μ , the last formula rewrites as

$$\int_{\Omega} \varphi d\mu \geq I_1(\varepsilon) + I_2(\varepsilon) + I_3(\varepsilon) \quad (2.4)$$

where

$$I_1(\varepsilon) = \int_{\Omega_\varepsilon} \langle \mathbb{A}(x) \nabla u, \nabla \varphi \rangle dx, \quad I_2(\varepsilon) = -\frac{\varepsilon}{2} \int_{\Omega \setminus \Omega_\varepsilon} \langle \mathbb{A} \nabla \varphi, \nabla \varphi \rangle dx, \quad I_3(\varepsilon) = \frac{1}{\varepsilon} \int_{\Omega_\varepsilon} f(u + \varepsilon\varphi) dx.$$

First, we note that the very definition of Ω_ε yields that

$$\mathcal{L}^n((\{u = 0\} \cap \{\varphi < 0\}) \setminus \Omega_\varepsilon) = \mathcal{L}^n(\Omega_\varepsilon \setminus (\{0 \leq u \leq \varepsilon \|\varphi\|_{L^\infty(\Omega)}\} \cap \{\varphi < 0\})) = 0,$$

and thus we may conclude that $\chi_{\Omega_\varepsilon} \rightarrow \chi_{\{u=0\} \cap \{\varphi < 0\}}$ in $L^1(\Omega)$. Hence, the dominated convergence theorem yields

$$\lim_{\varepsilon \rightarrow 0^+} I_1(\varepsilon) = \int_{\{u=0\} \cap \{\varphi < 0\}} \langle \mathbb{A} \nabla u, \nabla \varphi \rangle dx = 0, \quad (2.5)$$

by locality of the weak gradient.

Instead, to estimate $I_2(\varepsilon)$, we use (H1) as follows

$$|I_2(\varepsilon)| = \frac{\varepsilon}{2} \int_{\Omega \setminus \Omega_\varepsilon} \langle \mathbb{A} \nabla \varphi, \nabla \varphi \rangle dx \leq \frac{\varepsilon}{2} \Lambda \int_{\Omega} |\nabla \varphi|^2 dx,$$

so that

$$\lim_{\varepsilon \rightarrow 0^+} I_2(\varepsilon) = 0. \quad (2.6)$$

Lastly, we deal with $I_3(\varepsilon)$: note that $u \geq 0$ and $f \geq c_0 > 0$ on Ω by (H2), thus we have

$$\int_{\Omega_\varepsilon} f(u + \varepsilon \varphi) dx \geq \varepsilon \int_{\Omega_\varepsilon} f \varphi dx.$$

Hence, it is true that

$$\liminf_{\varepsilon \rightarrow 0^+} I_3(\varepsilon) \geq \int_{\{u=0\} \cap \{\varphi < 0\}} f \varphi dx. \quad (2.7)$$

Therefore, (2.4)-(2.7) yield that

$$\int_{\Omega} \varphi d\mu \geq \int_{\{u=0\} \cap \{\varphi < 0\}} f \varphi dx.$$

By repeating the same argument with $-\varphi$ we obtain the inequality

$$\int_{\Omega} \varphi d\mu \leq \int_{\{u=0\} \cap \{\varphi > 0\}} f \varphi dx.$$

Hence, by approximation for every $\varphi \in C_0^0(\Omega)$ it holds

$$\int_{\{u=0\} \cap \{\varphi < 0\}} f \varphi dx \leq \int_{\Omega} \varphi d\mu \leq \int_{\{u=0\} \cap \{\varphi > 0\}} f \varphi dx. \quad (2.8)$$

From this we infer that $\mu \ll \mathcal{L}^n \llcorner \Omega$, and thus $\mu = \zeta \mathcal{L}^n \llcorner \Omega$ with $\zeta \in L^1(\Omega)$. In conclusion, plugging this piece of information in (2.8) we conclude that $0 \leq \zeta \leq f \chi_{\{u=0\}}$ \mathcal{L}^n -a.e. on Ω .

The Hölder continuity of u follows from nowadays standard elliptic regularity [19, Theorem 8.22]. \square

Remark 2.2. In case \mathbb{A} is more regular, namely $W^{1,p}(\Omega; \mathbb{R}^{n \times n})$ for $p > n$, the minimizer u turns out to be $W_{\text{loc}}^{2,p}(\Omega)$ (cf. [14, Proposition 3.1]). In turn, from this one can easily prove that (2.2) rewrites as

$$\operatorname{div}(\mathbb{A} \nabla u) = f \chi_{\{u > 0\}} \quad (2.9)$$

\mathcal{L}^n -a.e. on Ω and in $\mathcal{D}'(\Omega)$ (cf. [13, Corollary 3.5], [14, Proposition 3.1]).

Here we limit ourselves to notice that, standing assumptions (H1) and (H2), by uniqueness of u and (2.2), ζ coincides with f on the interior $\{u = 0\}^\circ$ of $\{u = 0\}$, so that we may conclude the refined inequalities

$$f \chi_{\{u=0\}^\circ} \leq \zeta \leq f \chi_{\{u=0\}}.$$

This remark will be crucial in what follows in order to apply [2, Theorems 3.9]. In particular, we infer from this the quadratic non-degeneracy of the solution from free boundary points. Instead, a parabolic bound from above from free boundary points follows directly from (2.2) thanks to [2, Theorems 3.1]. We apply these properties to the blow up analysis of Section 3 (cf. Proposition 2.5).

Furthermore, observe that $\mathcal{L}^n(\Gamma_u) = 0$ in view of [2, Corollary 3.10], so that equation (2.9) is actually true \mathcal{L}^n -a.e. on Ω and in $\mathcal{D}'(\Omega)$. Despite this, we stress that we do not need such

a piece of information to establish the quasi-monotonicity formulas in Section 3: equation (2.2), but not (2.9), will be used in the proofs of Theorems 3.7 and 3.8.

We establish next two useful corollaries of Proposition 2.1. The first is simply a suitable version of Caccioppoli's inequality.

Corollary 2.3. *Assume (H1) and (H2). Let u be the minimizer of \mathcal{E} on $\mathbb{K}_{\psi,g}$. Then, there exists a constant $C = C(n, \Lambda) > 0$ such that for every $x_0 \in \Omega$ and for every $r \in (0, \frac{1}{4}\text{dist}(x_0, \partial\Omega))$*

$$\int_{B_r(x_0)} |\nabla u|^2 dx \leq \frac{C}{r^2} \int_{B_{2r}(x_0)} u^2 dx + C \|f\|_{L^\infty(B_{2r}(x_0))}^2 r^{n+2}. \quad (2.10)$$

Instead, the second corollary is an integration by parts formula which will be employed in the proof of the Monneau's quasi-monotonicity formula.

Corollary 2.4. *Assume (H1) and (H2). Let u be the minimizer of \mathcal{E} on $\mathbb{K}_{\psi,g}$. Then, for every $x_0 \in \Omega$ and for \mathcal{L}^1 -a.e. $r \in (0, \text{dist}(x_0, \partial\Omega))$*

$$\int_{B_r(x_0)} \langle \mathbb{A} \nabla u, \nabla \varphi \rangle dx + \int_{B_r(x_0)} (f - \zeta) \varphi dx = \int_{\partial B_r(x_0)} \langle \mathbb{A} \nabla u, \nu \rangle \varphi d\mathcal{H}^{n-1} \quad (2.11)$$

for every $\varphi \in W^{1,2}(\Omega)$, where $\nu(x) := \frac{x-x_0}{|x-x_0|}$.

Proof. For $x_0 \in \Omega$, $\varphi \in W^{1,2}(\Omega)$, $r \in (0, \text{dist}(x_0, \partial\Omega))$, and $s < r$ let

$$\psi_s(x) := 1 \wedge \frac{1}{r-s} \text{dist}(x, \partial B_r(x_0))$$

if $x \in B_r(x_0)$ and $\psi_s(x) := 0$ otherwise in Ω . Then $\psi_s \varphi \in W_0^{1,2}(\Omega)$ so that using it as test function in (2.2) we get

$$\int_{B_r(x_0)} \langle \mathbb{A} \nabla u, \nabla \varphi \rangle \psi_s dx + \int_{B_r(x_0)} (f - \zeta) \varphi \psi_s dx = - \int_{B_r(x_0)} \langle \mathbb{A} \nabla u, \nabla \psi_s \rangle \varphi dx := I_s.$$

It is clear that $\psi_s \rightarrow \chi_{B_r(x_0)}$ in $L^p(\Omega)$ for every $p \in [1, \infty)$ as $s \uparrow r$. Moreover, being $0 \leq \psi_s \leq 1$, the Lebesgue dominated convergence theorem implies that the left hand side converges to the left hand side in (2.11) as $s \uparrow r$. Computing explicitly the gradient of ψ_s , we evaluate I_s as follows:

$$I_s = \frac{1}{r-s} \int_{B_r(x_0) \setminus B_s(x_0)} \langle \mathbb{A} \nabla u, \frac{x-x_0}{|x-x_0|} \rangle \varphi dx = \frac{1}{r-s} \int_s^r dt \int_{\partial B_t(x_0)} \langle \mathbb{A} \nabla u, \nu \rangle \varphi d\mathcal{H}^{n-1},$$

where in the second equality we have used the coarea formula. The conclusion then follows at once (cf. for instance [11, Sections 3.4.3 and 3.4.4]). \square

We recall next the standard notations for the coincidence set and for the free boundary

$$\Lambda_u := \{x \in \Omega : u(x) = 0\}, \quad \Gamma_u := \partial \Lambda_u \cap \Omega. \quad (2.12)$$

For any point $x_0 \in \Gamma_u$, we introduce the family of rescaled functions

$$u_{x_0,r}(x) := \frac{u(x_0 + rx)}{r^2} \quad (2.13)$$

for $x \in \frac{1}{r}(\Omega - x_0)$. It is clear that changing variables in (2.1) implies that $u_{x_0,r}$ minimizes

$$\int_{\frac{1}{r}(\Omega - x_0)} (\langle \mathbb{A}(x_0 + rx) \nabla v(x), \nabla v(x) \rangle + 2f(x_0 + rx)v(x)) dx, \quad (2.14)$$

among all functions $v \geq 0$ on $\frac{1}{r}(\Omega - x_0)$, and with $v - u_{x_0,r} \in W_0^{1,2}(\frac{1}{r}(\Omega - x_0))$. Note that $\mathbb{A}(x_0 + r\cdot)$ and $f(x_0 + r\cdot)$ satisfy (H1)-(H2) uniformly in x_0 and r , i.e. the ellipticity constants of $\mathbb{A}(x_0 + r\cdot)$ are the same of those of \mathbb{A} , the L^∞ bound and the lower bound for $f(x_0 + r\cdot)$ are the same of those of f . Therefore, similarly to (2.2), from (2.14) we infer that

$$\operatorname{div}(\mathbb{A}(x_0 + rx)\nabla u_{x_0,r}) = f(x_0 + rx) - \zeta(x_0 + rx) \quad (2.15)$$

\mathcal{L}^n -a.e. on $\frac{1}{r}(\Omega - x_0)$ and in $\mathcal{D}'(\frac{1}{r}(\Omega - x_0))$, where ζ is the function in Proposition 2.1 (cf [19, Section 8]).

The first properties we recall on the family $(u_{x_0,r})_r$ follows from the fundamental quadratic growth and quadratic detachment of the solution from free boundary points established in [2, Theorems 3.1, 3.9] (cf. the discussion in Remark 2.2, see also [3, Theorems 2.3, 2.4]). We summarize the needed properties in the ensuing statement which is key for our approach. We remark that the estimates are uniform in x_0 and r .

Proposition 2.5. *Assume (H1) and (H2). Let u be the minimizer of \mathcal{E} over $\mathbb{K}_{\psi,g}$. There exists a constant $\vartheta = \vartheta(n, \Lambda, c_0, \|f\|_{L^\infty}) > 0$, such that for every $x_0 \in \Gamma_u$ and for every $r \in (0, \frac{1}{2}\operatorname{dist}(x_0, \partial\Omega))$, it holds*

$$\sup_{\partial B_1} u_{x_0,r} \geq \vartheta. \quad (2.16)$$

Moreover, for every $R > 0$ and for all compact sets $K \subset \Omega$ there exists a constant $C = C(n, \Lambda, c_0, \|f\|_{L^\infty}, R, K) > 0$ such that

$$\|u_{x_0,r}\|_{L^\infty(B_R)} \leq C, \quad (2.17)$$

for all $x_0 \in \Gamma_u \cap K$, and for all $r \in (0, \frac{1}{4R}\operatorname{dist}(K, \partial\Omega))$.

Remark 2.6. Notice that in view of the estimate in (2.10) in Corollary 2.3 and the estimate in (2.17), we may infer that for all $x_0 \in \Gamma_u \cap K$, and for all $r \in (0, \frac{1}{4R}\operatorname{dist}(K, \partial\Omega))$

$$\|u_{x_0,r}\|_{L^\infty(B_R)} + \|\nabla u_{x_0,r}\|_{L^2(B_R, \mathbb{R}^n)} \leq C. \quad (2.18)$$

In addition, if assumption (H3) holds, then the Hölder seminorms of $\mathbb{A}(x_0 + r\cdot)$ have uniform bounds with respect to x_0 and r . Thus, uniform Schauder estimates are deduced thanks to [19, Corollary 8.36]. Then the existence up to subsequences of $C^{1,\beta}$ -limits, $\beta < \alpha$, as $r \downarrow 0$ of the family $(u_{x_0,r})_r$ is standard.

Proposition 2.7. *Assume (H1)-(H3). Let u be the minimizer of \mathcal{E} over $\mathbb{K}_{\psi,g}$, and $K \subset \Omega$ a compact set. Then for every $x_0 \in K \cap \Gamma_u$ and for every $R > 0$ there exists a constant $C = C(n, \Lambda, \|f\|_{L^\infty}, \|\mathbb{A}\|_{C^{0,\alpha}}, R, K) > 0$ such that, for every $r \in (0, \frac{1}{4R}\operatorname{dist}(K, \partial\Omega))$*

$$\|u_{x_0,r}\|_{C^{1,\alpha}(B_R)} \leq C. \quad (2.19)$$

In particular, $(u_{x_0,r})_r$ is relatively compact in $C_{\operatorname{loc}}^{1,\beta}(A)$, for all $\beta \in (0, \alpha)$, and for every open set $A \subset \mathbb{R}^n$.

The functions arising in this limit process are called blow up limits. In particular, the blow up limits are non-trivial, i.e. not identically zero, in view of Proposition 2.5.

Corollary 2.8. *Assume (H1)-(H3). Let u be the minimizer of \mathcal{E} over $\mathbb{K}_{\psi,g}$, and let $x_0 \in \Gamma_u$. Then, for every sequence $r_k \downarrow 0$ there exists a subsequence $(r_{k_j})_j \subset (r_k)_k$ such that the rescaled functions $(u_{x_0,r_{k_j}})_j$ converge to a non trivial $C^{1,\alpha}(A)$ limit in $C_{\operatorname{loc}}^{1,\beta}(A)$, for all $\beta \in (0, \alpha)$, and for every open set $A \subset \mathbb{R}^n$.*

3. QUASI-MONOTONICITY FORMULAS

In this section we establish Weiss' and Monneau's type quasi-monotonicity formulas. The monotone quantities we consider in Section 3.1 are modeled upon the classical Dirichlet energy as in [14] under a pointwise normalization condition on the coefficients. We shall show in the subsequent Section 3.2 how to reduce to that formulation in a pointwise way via a suitable change of variables following [12]. The advantage of this approach is that the matrix field \mathbb{A} is not differentiated in deriving the quasi-monotonicity formulas contrary to [12, 18]. In those papers, instead, the natural quadratic energy \mathcal{E} associated to the obstacle problem under study had been considered. The $W_{\text{loc}}^{2,p}$ regularity of solutions and the quadratic growth from the free boundary were key properties to establish the quasi-monotonicity formulas in [14] for the Dirichlet based quantities. The new contribution of the current paper is to avoid the use of the former piece of information, which is not guaranteed in our setting, thanks to an elementary energy comparison argument to prove Weiss' formula (cf. Proposition 3.3), and in turn thanks to the latter and to Proposition 2.1 to prove Monneau's formula (cf. Proposition 3.6).

3.1. Weiss' and Monneau's quasi-monotonicity formulas under a normalization condition. We establish Weiss' and Monneau's quasi-monotonicity formulas for the minimizer u of \mathcal{E} on $\mathbb{K}_{\psi,g}$ under the normalization condition

$$x_0 = \underline{0} \in \Gamma_u, \quad \mathbb{A}(x_0) = \text{Id}, \quad f(x_0) = 1. \quad (3.1)$$

We will show in the next Section 3.2 how to reduce to (3.1) in each free boundary point thanks to a change of variables.

Moreover, for the sake of possible future generalizations of Theorem 1.1, we establish Weiss' and Monneau's formulas under Dini, double-Dini continuity assumptions, respectively (cf. Remark 3.5 for generalizations of Weiss' formula). Thus, we introduce some terminology. Given a uniformly continuous function $\zeta : \Omega \rightarrow \mathbb{R}^m$, $m \geq 1$, we consider a modulus of continuity of ζ namely an increasing function $\omega_\zeta : (0, \infty) \rightarrow (0, \infty)$ with $\lim_{t \rightarrow 0} \omega_\zeta(t) = 0$, and such that for all $t \in (0, \text{diam}\Omega)$

$$\sup_{x,y \in \Omega: |x-y| \leq t} \|\zeta(x) - \zeta(y)\| \leq \omega_\zeta(t).$$

With a slight abuse of notation with respect to our conventions we have denoted by $\|\cdot\|$ the norm in \mathbb{R}^m even in case $m = n$. In particular, ζ is said to be Dini continuous if for some ω_ζ as above

$$\int_0^{\text{diam}\Omega} \frac{\omega_\zeta(t)}{t} dt < \infty, \quad (3.2)$$

and double-Dini continuous if for some $a \geq 1$

$$\int_0^{\text{diam}\Omega} \frac{\omega_\zeta(t)}{t} |\log t|^a dt < \infty. \quad (3.3)$$

Therefore, we introduce the following weaker assumptions, each substituting (H3) in some of the results contained in this section:

- (H4) \mathbb{A} and f are Dini continuous;
- (H5) \mathbb{A} and f are double-Dini continuous.

We point out that under condition (H4), [25, Theorem 2.1 and Remark 2.2] imply that $u \in C^1(\Omega)$ (cf. also [26, 9, 10] for more on Schauder estimates for linear elliptic PDEs with Dini type continuity conditions on the matrix field).

Given $v \in W^{1,2}(\Omega)$ we consider the Weiss' energy

$$\Phi_v(r) := \frac{1}{r^{n+2}} \int_{B_r} (|\nabla v|^2 + 2v) \, dx - \frac{2}{r^{n+3}} \int_{\partial B_r} v^2 \, d\mathcal{H}^{n-1}, \quad (3.4)$$

and prove its quasi-monotonicity for $v = u$ in case (3.1) is satisfied. Let us also introduce the bulk energy

$$\mathcal{E}_v(r) := \int_{B_r} (|\nabla v|^2 + 2v) \, dx \quad (3.5)$$

and the boundary energy

$$\mathcal{H}_v(r) := \int_{\partial B_r} v^2 \, d\mathcal{H}^{n-1}, \quad (3.6)$$

so that

$$\Phi_v(r) = \frac{1}{r^{n+2}} \mathcal{E}_v(r) - \frac{2}{r^{n+3}} \mathcal{H}_v(r).$$

In the rest of the section to ease the notation we write u_r in place of $u_{0,r}$. Note then that

$$\mathcal{E}_{u_r}(1) = \frac{\mathcal{E}_u(r)}{r^{n+2}} \leq C, \quad \mathcal{H}_{u_r}(1) = \frac{\mathcal{H}_u(r)}{r^{n+3}} \leq C, \quad (3.7)$$

thanks to the bound (2.17) in Remark 2.6 (we stress that no continuity assumption on \mathbb{A} or f is needed).

We establish next two auxiliary results. The first is well-known, we prove it only for the readers' convenience being the fundamental identity from which Weiss' quasi-monotonicity is inferred.

Lemma 3.1. *Assume (H1), (H2) and (H4). Let u be the minimizer of \mathcal{E} over $\mathbb{K}_{\psi,g}$. If $\underline{0} \in \Omega$, for every $r \in (0, \text{dist}(\underline{0}, \partial\Omega))$ consider*

$$w_r(x) := |x|^2 u\left(r \frac{x}{|x|}\right). \quad (3.8)$$

Then, it is true that

$$\Phi'_u(r) = \frac{n+2}{r} (\Phi_{w_r}(1) - \Phi_{u_r}(1)) + \frac{1}{r} \int_{\partial B_1} (\langle \nabla u_r, \nu \rangle - 2u_r)^2 \, d\mathcal{H}^{n-1}. \quad (3.9)$$

Proof. By definition for every $r \in (0, \text{dist}(\underline{0}, \partial\Omega))$ we have

$$\Phi'_u(r) = \frac{\mathcal{E}'_u(r)}{r^{n+2}} - (n+2) \frac{\mathcal{E}_u(r)}{r^{n+3}} - 2 \frac{\mathcal{H}'_u(r)}{r^{n+3}} + 2(n+3) \frac{\mathcal{H}_u(r)}{r^{n+4}}. \quad (3.10)$$

A direct computation then gives

$$\mathcal{E}'_u(r) = \int_{\partial B_r} (|\nabla u|^2 + 2u) \, d\mathcal{H}^{n-1} = r^{n+1} \int_{\partial B_1} (|\nabla u_r|^2 + 2u_r) \, d\mathcal{H}^{n-1}. \quad (3.11)$$

By scaling it is easy to check that

$$\begin{aligned} \mathcal{H}'_u(r) &= \frac{d}{dr} \left(\int_{\partial B_1} u^2(r y) r^{n-1} \, d\mathcal{H}^{n-1} \right) \\ &= \frac{n-1}{r} \mathcal{H}_u(r) + 2r^{n+2} \int_{\partial B_1} u_r \langle \nabla u_r, y \rangle \, d\mathcal{H}^{n-1}, \end{aligned} \quad (3.12)$$

therefore

$$-\frac{2}{r^{n+3}}\mathcal{H}'_u(r) + \frac{2(n+3)}{r^{n+4}}\mathcal{H}_u(r) = \frac{8}{r^{n+4}}\mathcal{H}_u(r) - \frac{4}{r} \int_{\partial B_1} u_r \langle \nabla u_r, y \rangle d\mathcal{H}^{n-1}. \quad (3.13)$$

Plugging (3.13) and (3.11) in (3.9) we get

$$\begin{aligned} \Phi'_u(r) &= \frac{1}{r} \int_{\partial B_1} (|\nabla u_r|^2 + 2u_r) d\mathcal{H}^{n-1} - (n+2) \frac{\mathcal{E}_u(r)}{r^{n+3}} + \frac{8}{r^{n+4}}\mathcal{H}_u(r) - \frac{4}{r} \int_{\partial B_1} u_r \langle \nabla u_r, y \rangle d\mathcal{H}^{n-1} \\ &= -\frac{n+2}{r} \Phi_u(r) - \frac{2(n-2)}{r^{n+4}}\mathcal{H}_u(r) + \frac{1}{r} \int_{\partial B_1} (|\nabla u_r|^2 + 2u_r - 4u_r \langle \nabla u_r, y \rangle) d\mathcal{H}^{n-1} \\ &= -\frac{n+2}{r} \Phi_u(r) + \frac{1}{r} \int_{\partial B_1} (|\nabla u_r|^2 + 2u_r - 2(n-2)u_r^2 - 4u_r \langle \nabla u_r, y \rangle) d\mathcal{H}^{n-1} \\ &= -\frac{n+2}{r} \Phi_u(r) + \frac{1}{r} \int_{\partial B_1} ((\langle \nabla u_r, y \rangle - 2u_r)^2 + |\nabla_\tau u_r|^2 - 2nu_r^2 + 2u_r) d\mathcal{H}^{n-1} \\ &= \frac{n+2}{r} (\Phi_{w_r}(1) - \Phi_{u_r}(1)) + \frac{1}{r} \int_{\partial B_1} (\langle \nabla u_r, \nu \rangle - 2u_r)^2 d\mathcal{H}^{n-1}, \end{aligned}$$

where in the last equality we have used that

$$\Phi_{w_r}(1) = \frac{1}{n+2} \int_{\partial B_1} (|\nabla_\tau u_r|^2 - 2nu_r^2 + 2u_r) d\mathcal{H}^{n-1}.$$

which follows from a direct computation of the energy of the 2-homogeneous extension w_r . \square

The second result is a simple consequence of the continuity assumptions on the coefficients.

Lemma 3.2. *Assume (H4) and (3.1). For every $v \in W^{1,2}(B_r)$ with $v \geq 0$ \mathcal{L}^n -a.e. on B_r then*

$$\left| \int_{B_1} |\nabla v_r|^2 dx - \int_{B_1} \langle \mathbb{A}(rx) \nabla v_r, \nabla v_r \rangle dx \right| \leq \omega_{\mathbb{A}}(r) \int_{B_1} |\nabla v_r|^2 dx \quad (3.14)$$

$$\left| \int_{B_1} (1 - f(rx)) v_r dx \right| \leq \omega_f(r) \int_{B_1} v_r dx, \quad (3.15)$$

Proof. It suffices to take in to account the normalization assumption (3.1) and the definition of modulus of continuity. \square

Let us first establish Weiss' quasi-monotonicity under the normalization condition.

Proposition 3.3. *Assume (H1), (H2), (H4) and (3.1). Let u be the minimizer of \mathcal{E} over $\mathbb{K}_{\psi,g}$. There is a dimensional constant $C = C(n) > 0$ such that if for some $\gamma \geq 1$*

$$\|u_r\|_{L^\infty(B_2)} + \|\nabla u_r\|_{L^2(B_2; \mathbb{R}^n)} \leq \gamma \quad (3.16)$$

for every $r \in (0, \frac{1}{2} \text{dist}(\underline{0}, \partial\Omega))$, then

$$\frac{d}{dr} \left(\Phi_u(r) + C\gamma^2 \int_0^r \frac{\omega(t)}{t} dt \right) \geq \frac{1}{r} \int_{\partial B_1} (\langle \nabla u_r, x \rangle - 2u_r)^2 d\mathcal{H}^{n-1}, \quad (3.17)$$

for every $r \in (0, \frac{1}{2} \text{dist}(\underline{0}, \partial\Omega))$, where $\omega(r) := \omega_{\mathbb{A}}(r) + \omega_f(r)$.

In particular, Φ_u has a finite right limit in 0 denoted by $\Phi_u(0^+)$.

Proof. For $r \in (0, \text{dist}(\underline{0}, \partial\Omega))$, we use formula (3.9) for Φ'_u in combination with one of the following alternatives

- (a) $\Phi_{w_r}(1) \geq \Phi_{u_r}(1)$,
- (b) $\Phi_{w_r}(1) < \Phi_{u_r}(1)$.

In case (a) we conclude that

$$\Phi'_u(r) \geq \frac{1}{r} \int_{\partial B_1} (\langle \nabla u_r, \nu \rangle - 2u_r)^2 \, d\mathcal{H}^{n-1}. \quad (3.18)$$

Otherwise, being $u_r|_{\partial B_1} = w_r|_{\partial B_1}$, the inequality defining case (b) rewrites as

$$\mathcal{E}_{w_r}(1) < \mathcal{E}_{u_r}(1). \quad (3.19)$$

Thus we may estimate $\Phi_{w_r}(1) - \Phi_{u_r}(1)$ from below by taking into account Lemma 3.2 and that u_r minimizes the functional in (2.14) with respect to its boundary values, to conclude that

$$\begin{aligned} \Phi_{w_r}(1) - \Phi_{u_r}(1) &= \mathcal{E}_{w_r}(1) - \mathcal{E}_{u_r}(1) \\ &\geq \int_{B_1} (\langle \mathbb{A}(rx) \nabla w_r, \nabla w_r \rangle + 2f(rx)w_r) \, dx - \int_{B_1} (\langle \mathbb{A}(rx) \nabla u_r, \nabla u_r \rangle + 2f(rx)u_r) \, dx \\ &\quad - \omega(r)(\mathcal{E}_{w_r}(1) + \mathcal{E}_{u_r}(1)) \stackrel{(3.19)}{\geq} -2\omega(r)\mathcal{E}_{u_r}(1) \stackrel{(3.16)}{\geq} -C\gamma^2\omega(r), \end{aligned}$$

where we have set $\omega(r) = \omega_{\mathbb{A}}(r) + \omega_f(r)$ and $C = C(n) > 0$. Hence, in case (b) we infer that

$$\Phi'_u(r) \geq -C\gamma^2 \frac{\omega(r)}{r} + \frac{1}{r} \int_{\partial B_1} (\langle \nabla u_r, \nu \rangle - 2u_r)^2 \, d\mathcal{H}^{n-1}. \quad (3.20)$$

Inequalities (3.18) and (3.20) provide (3.17) for every $r \in (0, \frac{1}{2}\text{dist}(\mathbb{Q}, \partial\Omega))$. \square

Remark 3.4. Recalling that f and \mathbb{A} are Dini continuous by (H4), the modulus of continuity ω provided by Proposition 3.3 is in turn Dini continuous.

Remark 3.5. An inspection of the proof above shows that Weiss' formula can be deduced even for a weaker notion of Dini continuity, that is actually the one used in [25, Theorem 2.1 and Remark 2.2] to infer the mentioned C^1 regularity of solutions. In this respect, we need a different version of Lemma 3.2. To this aim, thanks to the mentioned Schauder estimates, in place of (3.14) and (3.15) we may consider for r sufficiently small the inequalities

$$\left| \int_{B_1} |\nabla u_r|^2 \, dx - \int_{B_1} \langle \mathbb{A}(rx) \nabla u_r, \nabla u_r \rangle \, dx \right| \leq \tilde{\omega}(r) \|\nabla u_r\|_{L^\infty(B_1; \mathbb{R}^n)}^2, \quad (3.21)$$

$$\left| \int_{B_1} (1 - f(rx))u_r \, dx \right| \leq \tilde{\omega}(r) \|u_r\|_{L^\infty(B_1)}, \quad (3.22)$$

where

$$\tilde{\omega}(r) := \mathcal{L}^n(B_1) \left(\sup_{y \in B_2} \int_{B_r(y)} \left(\|\mathbb{A}(x) - \mathbb{A}(y)\|^2 + |f(x) - f(y)|^2 \right) \, dx \right)^{1/2}.$$

Note that $\tilde{\omega}$ is not a modulus of continuity according to the definition given above as it is not increasing. Despite this, assume that it satisfies (3.2). Then, on one hand [25, Theorem 2.1 and Remark 2.2] provide C^1 regularity with a uniform modulus of continuity for the gradient of the solution; on the other hand (3.21) and (3.22) together with (3.27) below (rather than (3.16)) yield Weiss' quasi-monotonicity formula. As it will be discussed in Section 3.2 below, it is not restrictive to assume (3.27) thanks exactly to [25, Theorem 2.1 and Remark 2.2].

Finally, we note that even weaker notions of continuity are allowed to get C^1 regularity of solutions to divergence form elliptic equations (cf. [9, 10]). In this respect, Reifenberg vanishing flatness of $\text{Reg}(u)$ had already been proved in case of VMO coefficients in [3] by means of PDEs arguments rather than using the variational approach (cf. the introduction for the definition of $\text{Reg}(u)$).

For what Monneau's formula is concerned, let v be any positive 2-homogeneous polynomial solution of

$$\Delta v = 1 \quad \text{on } \mathbb{R}^n. \quad (3.23)$$

Then by 2-homogeneity, elementary calculations lead to

$$\Phi_v(r) = \Phi_v(1) = \int_{B_1} v \, dy, \quad (3.24)$$

for all $r > 0$. It is easy to prove that the value above is a dimensional constant independent of v , which we denote by θ . Then, being the space of polynomials of degree 2 finite dimensional, and being v 2-homogeneous we infer that

$$\|\nabla v\|_{L^2(B_1)} + \|v\|_{L^2(\partial B_1)} \leq C(n). \quad (3.25)$$

We prove next a quasi-monotonicity formula for solutions of the obstacle problem in case $x_0 \in \Gamma_u$ is a singular point of the free boundary, namely it is such that

$$\Phi_u(0^+) = \theta. \quad (3.26)$$

To prove Monneau's formula we need to strengthen condition (3.16) (cf. (3.27) below).

Proposition 3.6. *Assume (H1), (H2), (H5) with $a = 1$, and (3.1). Let u be the minimizer of \mathcal{E} over $\mathbb{K}_{\psi,g}$. There exists a dimensional constant $C = C(n) > 0$ such that if for some $\gamma \geq 1$*

$$\|u_r\|_{L^\infty(B_2)} + \|\nabla u_r\|_{L^\infty(B_2; \mathbb{R}^n)} \leq \gamma \quad (3.27)$$

for every $r \in (0, \frac{1}{2} \text{dist}(\underline{0}, \partial\Omega))$, then the function

$$(0, \frac{1}{2} \text{dist}(\underline{0}, \partial\Omega)) \ni r \longmapsto \int_{\partial B_1} (u_r - v_r)^2 \, dx + C\gamma^2 \int_0^r \frac{dt}{t} \int_0^t \frac{\omega(s)}{s} \, ds \quad (3.28)$$

is nondecreasing, where v is any positive 2-homogeneous polynomial solution of (3.23), and ω is the modulus of continuity provided by Proposition 3.3.

Proof. Let $w_r := u_r - v$, then by scaling and by taking into account the 2-homogeneity of v we get

$$\begin{aligned} \frac{d}{dr} \left(\frac{1}{r^{n+3}} \int_{\partial B_1} (u - v)^2 \, d\mathcal{H}^{n-1} \right) &= \frac{d}{dr} \left(\int_{\partial B_1} w_r^2 \, d\mathcal{H}^{n-1} \right) = \frac{2}{r} \int_{\partial B_1} w_r (\langle \nabla u_r, x \rangle - 2u_r) \, d\mathcal{H}^{n-1} \\ &\geq \frac{2}{r} \int_{\partial B_1} w_r (\langle \mathbb{A}(rx) \nabla u_r, x \rangle - 2u_r) \, d\mathcal{H}^{n-1} - \frac{\omega_{\mathbb{A}}(r)}{r} \|w_r\|_{L^2(\partial B_1)} \|\nabla u_r\|_{L^2(\partial B_1)} \\ &\geq \frac{2}{r} \int_{\partial B_1} w_r (\langle \mathbb{A}(rx) \nabla u_r, x \rangle - 2u_r) \, d\mathcal{H}^{n-1} - C\gamma^2 \frac{\omega_{\mathbb{A}}(r)}{r}, \end{aligned} \quad (3.29)$$

for some $C = C(n) > 0$, where we have used (3.25) and (3.27) in the last inequality.

We use next the integration by parts formula (2.11) in Corollary 2.4 to get

$$\int_{\partial B_1} w_r \langle \mathbb{A}(rx) \nabla u_r, x \rangle \, d\mathcal{H}^{n-1} = \int_{B_1} \langle \mathbb{A}(rx) \nabla u_r, \nabla w_r \rangle \, dx + \int_{B_1} (f(rx) - \zeta(rx)) w_r \, dx := I_1 + I_2.$$

We estimate the two addends above separately. We start off with I_1 :

$$\begin{aligned} I_1 &= \int_{B_1} \langle \mathbb{A}(rx) \nabla u_r, \nabla u_r \rangle dx - \int_{B_1} \langle \mathbb{A}(rx) \nabla u_r, \nabla v \rangle dx \\ &\geq \Phi_{u_r}(1) - \omega_{\mathbb{A}}(r) \int_{B_1} |\nabla u_r|^2 dx - 2 \int_{B_1} u_r dx + 2\mathcal{H}_{u_r}(1) - \int_{B_1} \langle \mathbb{A}(rx) \nabla u_r, \nabla v \rangle dx. \end{aligned} \quad (3.30)$$

By taking advantage of the 2-homogeneity of v and that $\Delta v = 1$ (cf. (3.23)) we get that

$$\begin{aligned} - \int_{B_1} \langle \mathbb{A}(rx) \nabla u_r, \nabla v \rangle dx &\geq - \int_{B_1} \langle \nabla u_r, \nabla v \rangle dx - \omega_{\mathbb{A}}(r) \|\nabla u_r\|_{L^2(B_1)} \|\nabla v\|_{L^2(B_1)} \\ &= \int_{B_1} u_r dx - 2 \int_{\partial B_1} u_r v d\mathcal{H}^{n-1} - \omega_{\mathbb{A}}(r) \|\nabla u_r\|_{L^2(B_1)} \|\nabla v\|_{L^2(B_1)} \\ &\geq \int_{B_1} u_r dx - 2 \int_{\partial B_1} u_r v d\mathcal{H}^{n-1} - C\gamma\omega_{\mathbb{A}}(r), \end{aligned}$$

for some $C = C(n) > 0$, where we have used (3.25) and (3.27) in the last inequality. Plugging the latter estimate in (3.30) we infer that

$$I_1 \geq \Phi_{u_r}(1) - \int_{B_1} u_r dx + 2 \int_{\partial B_1} u_r v d\mathcal{H}^{n-1} - C\gamma^2\omega_{\mathbb{A}}(r). \quad (3.31)$$

Next note that by (2.3) in Proposition 2.1

$$\begin{aligned} I_2 &= \int_{B_1} (f(rx) - \zeta(rx))u_r dx - \int_{B_1} (f(rx) - \zeta(rx))v dx \\ &\geq \int_{B_1} (f(rx) - f(rx)\chi_{\{u_r=0\}})u_r dx - \int_{B_1} f(rx)v dx \\ &= \int_{B_1} f(rx)\chi_{\{u_r>0\}}u_r - \int_{B_1} f(rx)v dx = \int_{B_1} f(rx)(u_r - v) dx. \end{aligned} \quad (3.32)$$

We now use (3.24), (3.31) and (3.32), to estimate (3.29)

$$\begin{aligned} \frac{d}{dr} \left(\frac{1}{r^{n+3}} \int_{\partial B_1} (u - v)^2 d\mathcal{H}^{n-1} \right) &\geq \frac{1}{r} (\Phi_{u_r}(1) - \Phi_v(1)) \\ &\quad + \frac{1}{r} \int_{B_1} (1 - f(rx))(v - u_r) dx - C\gamma^2 \frac{\omega_{\mathbb{A}}(r)}{r} \\ &\stackrel{(3.25)}{\geq} \frac{1}{r} (\Phi_{u_r}(1) - \Phi_u(0^+)) - C\gamma^2 \frac{\omega(r)}{r}, \end{aligned} \quad (3.33)$$

for some $C = C(n) > 0$. Therefore, we may finally use Proposition 3.3 to conclude that

$$\begin{aligned} \frac{d}{dr} \left(\frac{1}{r^{n+3}} \int_{\partial B_1} (u - v)^2 d\mathcal{H}^{n-1} \right) &\geq \frac{1}{r} \int_0^r \frac{1}{t} \int_{\partial B_1} (\langle \nabla u_t, x \rangle - 2u_t)^2 d\mathcal{H}^{n-1} - \frac{C}{r} \gamma^2 \int_0^r \frac{\omega(t)}{t} dt - C\gamma^2 \frac{\omega(r)}{r}. \end{aligned}$$

The conclusion then follows at once. \square

3.2. Weiss' and Monneau's quasi-monotonicity: general case. To establish Weiss' and Monneau's monotonicity in general, we follow [12] and show that by means of a change of variables one can always reduce to the normalized setting in (3.1) for every free boundary point $x_0 \in \Gamma_u$. Moreover, the new quantities appearing in the corresponding obstacle problems under such a transformation satisfy assumptions (H1), (H2) and either (H3) or (H4) or (H5), with uniform constants with respect to x_0 , according to the assumption imposed on \mathbb{A} and f .

Indeed, let $x_0 \in \Gamma_u$ be any point of the free boundary, consider the affine change of variables

$$x \mapsto x_0 + f^{-1/2}(x_0) \mathbb{A}^{1/2}(x_0) x =: x_0 + \mathbb{L}(x_0) x$$

Changing variables leads to

$$\mathcal{E}(u) = f^{1-\frac{n}{2}}(x_0) \det(\mathbb{A}^{1/2}(x_0)) \mathcal{E}_{\mathbb{L}(x_0)}(u_{\mathbb{L}(x_0)}), \quad (3.34)$$

where we have set $\Omega_{\mathbb{L}(x_0)} := \mathbb{L}^{-1}(x_0) (\Omega - x_0)$, and

$$\mathcal{E}_{\mathbb{L}(x_0)}(v) := \int_{\Omega_{\mathbb{L}(x_0)}} \left(\langle \mathbb{C}_{\mathbb{L}(x_0)}(x) \nabla v, \nabla v \rangle + 2 \frac{f_{\mathbb{L}(x_0)}}{f(x_0)} v \right) dx, \quad (3.35)$$

with

$$\begin{aligned} u_{\mathbb{L}(x_0)}(x) &:= u(x_0 + \mathbb{L}(x_0)x), \\ f_{\mathbb{L}(x_0)}(x) &:= f(x_0 + \mathbb{L}(x_0)x), \\ \mathbb{C}_{\mathbb{L}(x_0)}(x) &:= \mathbb{A}^{-1/2}(x_0) \mathbb{A}(x_0 + \mathbb{L}(x_0)x) \mathbb{A}^{-1/2}(x_0). \end{aligned} \quad (3.36)$$

Note that $f_{\mathbb{L}(x_0)}(\underline{0}) = f(x_0)$ and $\mathbb{C}_{\mathbb{L}(x_0)}(\underline{0}) = \text{Id}$. Moreover, the free boundary is transformed under this map into

$$\Gamma_{u_{\mathbb{L}(x_0)}} = \mathbb{L}^{-1}(x_0)(\Gamma_u - x_0),$$

and the energy \mathcal{E} in (1.1) is minimized by u on $\mathbb{K}_{\psi,g}$ if and only if $\mathcal{E}_{\mathbb{L}(x_0)}$ in (3.35) is minimized by $u_{\mathbb{L}(x_0)}$ in (3.36) on $\mathbb{K}_{\psi(\mathbb{L}^{-1}(x_0)(\cdot - x_0), g(\mathbb{L}^{-1}(x_0)(\cdot - x_0)))}$. In particular, the normalization assumption (3.1) is satisfied.

Moreover, $\mathbb{C}_{\mathbb{L}(x_0)}$ and $\frac{f_{\mathbb{L}(x_0)}}{f(x_0)}$ satisfy (H1)-(H5) with uniform constants. Indeed, for what (H1) is concerned, it is clear that $\mathbb{C}_{\mathbb{L}(x_0)}(\cdot)$ is symmetric, bounded and coercive, with

$$\Lambda^{-2} |\xi|^2 \leq \langle \mathbb{C}_{\mathbb{L}(x_0)}(x) \xi, \xi \rangle \leq \Lambda^2 |\xi|^2 \quad (3.37)$$

for \mathcal{L}^n -a.e. $x \in \Omega_{\mathbb{L}(x_0)}$, and for every $\xi \in \mathbb{R}^n$. Note that

$$\frac{c_0}{\|f\|_{L^\infty(\Omega)}} < \frac{f_{\mathbb{L}(x_0)}}{f(x_0)} \leq \frac{\|f\|_{L^\infty(\Omega)}}{c_0} \quad (3.38)$$

for \mathcal{L}^n -a.e. $x \in \Omega_{\mathbb{L}(x_0)}$, so that (H2) holds. Moreover, on setting $\bar{\omega}_{\mathbb{A}}(t) := (n\Lambda)^2 \omega_{\mathbb{A}} \left(\sqrt{\frac{n\Lambda}{c_0}} t \right)$, it is clear that for every $x, y \in \Omega_{\mathbb{L}(x_0)}$

$$\|\mathbb{C}_{\mathbb{L}(x_0)}(x) - \mathbb{C}_{\mathbb{L}(x_0)}(y)\| \leq \bar{\omega}_{\mathbb{A}}(|x - y|). \quad (3.39)$$

Analogously, $\frac{f_{\mathbb{L}(x_0)}}{f(x_0)}$ has modulus of continuity $\bar{\omega}_f(t) := c_0^{-1} \omega_f \left(\sqrt{\frac{n\Lambda}{c_0}} t \right)$. Therefore, either (H3) or (H4) or (H5) holds, according to the corresponding assumption on \mathbb{A} and f .

Furthermore, we note that in view of (3.37) and (3.38) formula (3.16) is satisfied uniformly in x_0 and r . More precisely, Remark 2.6 yields that if $R > 0$, $K \subset \Omega$ is compact and $x_0 \in K$,

then for every $r \in (0, \frac{1}{4R} \text{dist}(K, \partial\Omega))$, and for some constant $C = C(n, c_0, \Lambda, \|f\|_{L^\infty}, R, K) > 0$ it is true that

$$\|u_{\mathbb{L}(x_0),r}\|_{L^\infty(B_R)} + \|\nabla u_{\mathbb{L}(x_0),r}\|_{L^2(B_R;\mathbb{R}^n)} \leq C, \quad (3.40)$$

where we have set $u_{\mathbb{L}(x_0),r} := (u_{\mathbb{L}(x_0)})_{0,r}$ (notice that $\nabla u_{\mathbb{L}(x_0),r}(\cdot) = \mathbb{L}^t(x_0) \nabla u(x_0 + \mathbb{L}(x_0) \cdot)$).

We are now ready to establish Weiss' quasi-monotonicity formula by applying Proposition 3.3 to $u_{\mathbb{L}(x_0)}$ thanks to the discussion above.

Theorem 3.7 (Weiss' quasi-monotonicity formula). *Assume (H1), (H2) and (H4). Let u be the minimizer of \mathcal{E} over $\mathbb{K}_{\psi,g}$. If $K \subset \Omega$ is a compact set, there is a constant $C = C(n, c_0, \Lambda, \|f\|_{L^\infty}, K) > 0$ such that for all $x_0 \in K \cap \Gamma_u$*

$$\frac{d}{dr} \left(\Phi_{u_{\mathbb{L}(x_0)}}(r) + C \int_0^r \frac{\bar{\omega}(t)}{t} dt \right) \geq \frac{1}{r} \int_{\partial B_1} (\langle \nabla u_{\mathbb{L}(x_0),r}, x \rangle - 2u_{\mathbb{L}(x_0),r}^2) d\mathcal{H}^{n-1}, \quad (3.41)$$

for every $r \in (0, \frac{1}{4} \text{dist}(K, \partial\Omega))$, where $\bar{\omega}(r) := \bar{\omega}_{\mathbb{A}}(r) + \bar{\omega}_f(r)$.

In particular, $\Phi_{u_{\mathbb{L}(x_0)}}$ has finite right limit $\Phi_{u_{\mathbb{L}(x_0)}}(0^+)$ in zero, and for all $r \in (0, \frac{1}{4} \text{dist}(K, \partial\Omega))$,

$$\Phi_{u_{\mathbb{L}(x_0)}}(r) - \Phi_{u_{\mathbb{L}(x_0)}}(0^+) \geq -C \int_0^r \frac{\bar{\omega}(t)}{t} dt. \quad (3.42)$$

For what concerns Monneau's quasi-monotonicity formula we may apply Proposition 3.6 to $u_{\mathbb{L}(x_0)}$ on condition that (3.27) is satisfied. This follows from the discussion above, and thanks to [25, Theorem 2.1 and Remark 2.2] which provide a modulus of continuity for $\nabla u_{\mathbb{L}(x_0)}$ depending only on $n, c_0, \Lambda, \bar{\omega}_{\mathbb{A}}$, and $\|f\|_{L^\infty(\Omega)}$ (cf. (3.37)-(3.39)).

Theorem 3.8 (Monneau's quasi-monotonicity formula). *Assume (H1), (H2) and (H5) with $a = 1$. Let u be the minimizer of \mathcal{E} over $\mathbb{K}_{\psi,g}$. If $K \subset \Omega$ is a compact set and (3.24) holds for $x_0 \in K \cap \Gamma_u$, then there exists a constant $C = C(n, c_0, \Lambda, \|f\|_{L^\infty}, K) > 0$ such that the function*

$$(0, \frac{1}{4} \text{dist}(K, \partial\Omega)) \ni r \longmapsto \int_{\partial B_1} (u_{\mathbb{L}(x_0),r} - v)^2 dx + C\gamma^2 \int_0^r \frac{dt}{t} \int_0^t \frac{\bar{\omega}(s)}{s} ds. \quad (3.43)$$

is nondecreasing, where v is any 2-homogeneous polynomial solution of (3.23), and $\bar{\omega}$ is the modulus of continuity provided by Theorem 3.7.

4. FREE BOUNDARY ANALYSIS

The regularity of the free boundary of the minimizer u of \mathcal{E} on $\mathbb{K}_{\psi,g}$ can be established thanks to the Weiss' and Monneau's quasi-monotonicity formulas proved in Section 3 at least if assumption (H3) is satisfied. In doing this we follow the approach introduced in [31, 27] for the classical obstacle problem related to the Dirichlet energy, and developed in [12, 18, 13, 14] both for linear elliptic operators in divergence form and in the nonlinear setting with suitable smoothness assumptions (see also [29] for a systematic presentation).

In particular, in this section we improve upon [12, Theorems 4.12 and 4.14], [18, Theorem 1.3] and [14, Theorem 2.1], since in all those cases the matrix field \mathbb{A} turns out to be in particular Hölder continuous due to Sobolev type embeddings.

In the ensuing proof we will highlight only the substantial changes since the arguments are essentially those given in [12].

Proof of Theorem 1.1. First recall that we may establish the conclusions for the function $u = w - \psi$ introduced in Section 2. Given this, the only minor change to be done to the arguments in [12, Section 4] is related to the freezing of the energy where the regularity of the coefficients plays a substantial role. More precisely, under assumption (3.1) we have in view of Lemma 3.2

$$\left| \int_{B_1} (\mathbb{A}(rx) \nabla v, \nabla v) + 2f(rx)v \, dx - \mathcal{E}_v(1) \right| \leq \omega(r) \mathcal{E}_v(1)$$

for all $v \in W^{1,2}(B_1)$.

We then describe shortly how to infer all the conclusions. We start off recalling that the quasi-monotonicity formulas established in [12, Section 3] are to be substituted by those in Section 3. Then the 2-homogeneity of blow up limits in [12, Proposition 4.2] now follows from Theorem 3.7. Nondegeneracy of blow up limits is contained in Proposition 2.5 (see [12, Lemma 4.3]). The classification of blow up limits is obtained exactly as in [12, Proposition 4.5]. Uniqueness of blow up limits at regular points, that follows from [12, Lemma 4.8], can be obtained with essentially no difference. The proofs of [12, Propositions 4.10, 4.11, Theorems 4.12, 4.14] remain unchanged. \square

Remark 4.1. Thanks to the quasi-monotonicity formulas in Section 3 we expect to be possible to deduce results analogous to Theorem 1.1 under assumption (H5) with $a > 2$ (cf. [14, Theorem 2.1] for such a statement if $\mathbb{A} \in W^{1,p}(\Omega; \mathbb{R}^{n \times n})$, $p > n$, and f satisfies (H5) with $a > 2$). Moreover, in view of Remark 3.5 it is likely that analogous results hold even in case $\tilde{\omega}$ there satisfies a double-Dini continuity condition. We do not insist on this issue here since several arguments should be carefully checked along the proofs of [12].

REFERENCES

- [1] I. BLANK. Sharp results for the regularity and stability of the free boundary in the obstacle problem. *Indiana Univ. Math. J.* 50 (2001), no. 3, 1077–1112.
- [2] I. BLANK, Z. HAO. The mean value theorem and basic properties of the obstacle problem for divergence form elliptic operators. *Communications in Analysis and Geometry* 23 (2015), no. 1, 129–158.
- [3] I. BLANK, Z. HAO. Reifenberg flatness of free boundaries in obstacle problems with VMO ingredients. *Calc. Var. Partial Differential Equations* 53 (2015), no. 3-4, 943–959.
- [4] L. A. CAFFARELLI. The regularity of free boundaries in higher dimensions. *Acta Math.* 139 (1977), no. 3-4, 155–184.
- [5] L. A. CAFFARELLI. Compactness methods in free boundary problems. *Comm. Partial Differential Equations* 5 (1980), no. 4, 427–448.
- [6] L. A. CAFFARELLI. The obstacle problem revisited. *J. Fourier Anal. Appl.* 4 (1998), no. 4-5, 383–402.
- [7] L. A. CAFFARELLI. The obstacle problem revisited. *Lezioni Fermiane. [Fermi Lectures]* Accademia Nazionale dei Lincei, Rome; Scuola Normale Superiore, Pisa, 1998. ii+54 pp.
- [8] L. A. CAFFARELLI, S. Salsa. A geometric approach to free boundary problems. Graduate Studies in Mathematics, 68. American Mathematical Society, Providence, RI, 2005. x+270 pp.
- [9] K. DONG, S. KIM. On C^1 , C^2 , and weak type-(1, 1) estimates for linear elliptic operators. *Comm. Partial Differential Equations* 42 (2017), no. 3, 417–435.
- [10] K. DONG, L. ESCAURIAZA, S. KIM. On C^1 , C^2 , and weak type-(1, 1) estimates for linear elliptic operators: part II. *Math. Ann.* 370 (2018), no. 1-2, 447–489.
- [11] L.C. EVANS, R.F. GARIEPY. Measure theory and fine properties of functions. Revised edition. Textbooks in Mathematics. CRC Press, Boca Raton, FL, 2015. xiv+299 pp.

- [12] M. FOCARDI, M. S. GELLI, E. SPADARO. Monotonicity formulas for obstacle problems with Lipschitz coefficients. *Calc. Var. Partial Differential Equations* 54 (2015), no. 2, 1547–1573.
- [13] M. FOCARDI, F. GERACI, E. SPADARO. The classical obstacle problem for nonlinear variational energies. *Nonlinear Anal.* 154 (2017), 71–87.
- [14] M. FOCARDI, F. GERACI, E. SPADARO. Quasi-monotonicity formulas for classical obstacle problems with Sobolev coefficients and applications. *J. Optim. Theory Appl.* 184 (2020), no. 1, 125–138.
- [15] M. FUCHS. Hölder continuity of the gradient for degenerate variational inequalities. *Nonlinear Anal. TMA* 15 (1) (1990), 85–100.
- [16] M. FUCHS, G. MINGIONE. Full $C^{1,\alpha}$ -regularity for free and constrained local minimizers of elliptic variational integrals with nearly linear growth. *Manuscripta Math.* 102 (2000), 227–250.
- [17] N. GAROFALO, M. SMIT VEGA GARCIA. New monotonicity formulas and the optimal regularity in the Signorini problem with variable coefficients. *Adv. Math.* 262 (2014), 682–750.
- [18] F. GERACI. The classical obstacle problem with coefficients in fractional Sobolev spaces. *Ann. Mat. Pura Appl.* (4) 197 (2018), no. 2, 549–581.
- [19] D. GILBARG, N.S. TRUDINGER. Elliptic partial differential equations of second order. Reprint of the 1998 edition. *Classics in Mathematics*. Springer-Verlag, Berlin, 2001. xiv+517 pp.
- [20] S. JEON, A. PETROSYAN, M. SMIT VEGA GARCIA. Almost minimizers for the thin obstacle problem with variable coefficients. arXiv:2007.07349
- [21] D. KINDERLEHRER, G. STAMPACCHIA. An introduction to variational inequalities and their applications. *Pure and Applied Mathematics*, 88. Academic Press, Inc., New York-London, 1980. xiv+313 pp.
- [22] H. KOCH, A. RÜLAND, W. SHI. The variable coefficient thin obstacle problem: Carleman inequalities. *Adv. Math.* 301 (2016), 820–866.
- [23] H. KOCH, A. RÜLAND, W. SHI. The variable coefficient thin obstacle problem: optimal regularity and regularity of the regular free boundary. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 34 (2017), no. 4, 845–897.
- [24] I. KUKAVICA. Quantitative uniqueness for second-order elliptic operators. *Duke Math. J.* 91 (1998), no. 2, 225–240.
- [25] Y.Y. LI. On the C^1 regularity of solutions to divergence form elliptic systems with Dini-continuous coefficients. *Chin. Ann. Math. Ser. B* 38 (2017), no. 2, 489–496.
- [26] G. LIEBERMAN. Hölder continuity of the gradient of solutions of uniformly parabolic equations with conormal boundary conditions. *Ann. Mat. Pura Appl.* 148 (1987), 77–99.
- [27] R. MONNEAU. On the number of singularities for the obstacle problem in two dimensions. *J. Geom. Anal.* 13 (2003), no. 2, 359–389.
- [28] R. MONNEAU. Pointwise estimates for Laplace equation. Applications to the free boundary of the obstacle problem with Dini coefficients. *J. Fourier Anal. Appl.* 15 (2009), no. 3, 279–335.
- [29] A. PETROSYAN, H. SHAHGHOLIAN, N. URALTSEVA. Regularity of free boundaries in obstacle-type problems. *Graduate Studies in Mathematics*, 136. American Mathematical Society, Providence, RI, 2012. x+221 pp.
- [30] A. RÜLAND, W. SHI. Optimal regularity for the thin obstacle problem with $C^{0,\alpha}$ coefficients *Calc. Var. Partial Differential Equations* 56 (2017), no. 5, Art. 129, 41 pp.
- [31] G. S. WEISS. A homogeneity improvement approach to the obstacle problem. *Invent. Math.* 138 (1999), no. 1, 23–50.

UNIVERSITÀ DI ROMA “LA SAPIENZA”

Current address: P.le Aldo Moro 5, Roma (Italy)

Email address: giovanna.andreucci@uniroma1.it

DiMAI, UNIVERSITÀ DEGLI STUDI DI FIRENZE

Current address: Viale Morgagni 67/A, 50134 Firenze (Italy)

Email address: matteo.focardi@unifi.it