




ORIGINAL ARTICLE

Frailty, psychological well-being, and social isolation in older adults with cognitive impairment during the SARS-CoV-2 pandemic: data from the GeroCovid initiative

Camilla TERZIOTTI,¹ Chiara CEOLIN ,¹ Maria DEVITA ,^{1,2} Cecilia RAFFAELLI,¹ Sara ANTENUCCI,³ Salvatore BAZZANO,⁴ Andrea CAPASSO,⁵ Manuela CASTELLINO,⁶ Stefania Del SIGNORE,⁷ Francesca LUBIAN,⁸ Mariangela MAIOTTI,⁹ Fiammetta MONACELLI,¹⁰ Maria Teresa MORMILE,¹¹ Claudia SGARITO,¹² Filomena VELLA,¹³ Giuseppe SERGI,¹ Pietro GARERI,¹⁴ Caterina TREVISAN,^{1,15} Andrea BELLIO,¹ Filippo FINI,¹ Alba MALARA,¹⁶ Enrico MOSSELLO,¹⁷ Stefano FUMAGALLI,¹⁷ Stefano VOLPATO,¹⁵ Fabio MONZANI,¹⁸ Giuseppe BELLELLI ,¹⁹ Gianluca ZIA,⁷ Raffaele Antonelli INCALZI²⁰ and Alessandra COIN¹

Department of ¹Medicine (DIMED), Geriatrics Division and ²General Psychology, University of Padova and ⁴Geriatrics Unit, Azienda ULSS 16, Padova, ³Psychogeriatric Outpatient Clinic, Ortona, ⁵Territorial Care Department, ASL NA2 Nord and ¹¹CDCD DS 46 and 47, ASL Napoli 2 Nord, Naples, ⁶“B.V. Consolata” Rehabilitation Hospital-Fatebenefratelli, San Maurizio Canavese, ⁸Geriatrics Unit, Memory Clinic, Bozen Hospital, Bozen, ⁹Clinical Neuropsychology, S.Giovanni Battista Hospital, Foligno, ¹⁰Section of Geriatrics, Department of Internal Medicine and Medical Specialties (DIMI), University of Genoa, Genoa, ¹²UOC Involutive Degenerative Diseases, Territorial Psychogeriatrics, ASP of Agrigento, Agrigento, ¹³CDCD, District 2, Trieste, ¹⁴Center for Cognitive Disorders and Dementia-Catanzaro Lido ASP, Catanzaro, ¹⁵Department of Medical Sciences, University of Ferrara, Ferrara, ¹⁶ANASTE Humanitas Foundation and ²⁰Unit of Geriatrics, Department of Medicine, Campus Bio-Medico University and Teaching Hospital, Rome, ¹⁷Geriatric Intensive Care Unit, Department of Experimental and Clinical Medicine, University of Firenze, Firenze, ¹⁸Geriatrics Unit, Department of Clinical and Experimental Medicine, University of Pisa, Pisa and ¹⁹School of Medicine and Surgery, Acute Geriatric Unit, University of Milano-Bicocca, San Gerardo Hospital, Monza, Italy and ⁷Bluecompanion, Ltd, London, UK

Correspondence: Chiara Ceolin, Department of Medicine (DIMED) Geriatrics Division, University of Padua, Via Giustiniani 2, 35128 Padua, Italy. Email: chiara.ceolin.1@gmail.com

Camilla Terziotti and Chiara Ceolin are equally contributed to this work.

Received 9 February 2023; revision received 26 June 2023; accepted 16 August 2023.

Key words: frailty, NPI, older adults, SARS-CoV-2, social isolation, UCLA.

Abstract

Background: The containment measures linked to the COVID-19 pandemic negatively affected the phyco-physical well-being of the population, especially older adults with neurocognitive disorders (NCDs). This study aims to evaluate whether the frailty of NCD patients was associated with different changes in multiple health domains, in particular in relation to loneliness and social isolation, pre- and post-lockdown.

Materials and Methods: Patients were recruited from 10 Italian Centers for Cognitive Disorders and Dementia. Data were collected in the pre-pandemic period (T0), during the pandemic lockdown (T1), and 6–9 months post-lockdown (T2). The UCLA Loneliness Scale-3, Activities of Daily Living (ADL), Instrumental ADL (IADL), Mini-Mental State Examination, and Neuropsychiatric Inventory (NPI) were administered. Caregivers' burden was also tested. Patients were categorized as non-frail, pre-frail, and frail according to the Fatigue, Resistance, Ambulation, Illness, and Loss of Weight scale.

Results: The sample included 165 subjects (61.9% women, mean age 79.5 ± 4.9 years). In the whole sample, the ADL, IADL, and NPI scores significantly declined between T0 and T2. There were no significative variations in functional and cognitive domains between the frail groups. During lockdown we recorded higher Depression Anxiety Stress Scales and Perceived Stress Scale scores in frail people. In multivariable logistic regression, frailty was associated with an increase in social isolation, and a loss of IADL.

Conclusions: We observed a global deterioration in functional and neuropsychiatric domains irrespective of the degree of frailty. Frailty was associated with the worsening of social isolation during lockdown. Frail patients and their caregivers seemed to experience more anxiety and stress disorders during SARS-CoV-2 pandemic.

INTRODUCTION

The COVID-19 pandemic has required important clinical, social and economic interventions in order to limit escalation of the disease and safeguard individuals' health. Considering the well-known impact of lockdown and forced isolation on physical and psychological well-being, particularly in older adults,^{1–4} many studies conducted during the COVID-19 pandemic focused on describing the prevalence during this period of frailty, underlying increased vulnerability and compromised ability to cope with everyday or acute stressors.^{5–8} Pandemic and containment measures exacerbated self-reported loneliness in a large part of the adult population,⁹ increasing the risk of 'social frailty', known as the loss of social resources, social activities, or abilities and general resources that are important for fulfilling basic social needs during one's lifetime.¹⁰ The pandemic's effects on psychological well-being were more intensive in patients with severe cognitive impairment.¹¹ Although the increased prevalence of loneliness and social isolation in older cognitively compromised adults has already been reported before and after the COVID-19 pandemic,^{12,13} studies that considered the trend during the pandemic phases, with particular attention to neurocognitive disorders, are still lacking. In fact, previous findings suggest that the association of frailty and poor global cognition describes the 'frailest among the frail' individuals, who experienced different psychological and/or behavioural effects of COVID-19 and, for these reasons, deserve particular attention.^{2,14–17} Given these premises, we could hypothesize an increase of psycho-cognitive and behavioural disorders, with an increasing gradient according to the degree of fragility of the patient. Despite this, to our knowledge no studies have focused on trends of neuropsychiatric disorders according to the degree of frailty during the SARS-CoV-2 pandemic.

The aims of this study were (i) to evaluate the association between frailty and different health domains, particularly in relation to loneliness and social isolation, in a sample of older adults affected by neurocognitive disorders (NCDs) before and after the COVID-19 first wave lockdown; and (ii) to estimate whether the lockdown impacted the psychological and affective well-being of older adults with NCDs and their caregivers' burden differently according to the presence of frailty.

MATERIALS AND METHODS

Study design

The GeroCovid CDCD study ('GeroCovid dementia—psychological health cohort') was part of a multi-purpose, multicentre, and multi-setting initiative¹¹ called the GeroCovid Observational project. The retrospective GeroCovid CDCD study involves 10 Italian Centers for Cognitive Decline and Dementia (CDCDs) and considers three phases during 2020: from January to February (pre-pandemic, T0), from March to May (lockdown, T1), and from July to December (post-lockdown, T2). In Italy lockdown started from March 2020 with the closure of schools and all types of commercial activities, with the exception of basic necessities, and with strict restrictions on moving.

All participants (or their caregivers) gave informed consent to their involvement in the study. The study was approved by the BIO-CAMPUS Ethics Committee, University of Rome—Prot. Number: 22.5 (20).20 OSS ComEtUCBM. Each centre received approval from the local Ethics Committee to perform the study.

Participants

Participants were recruited from Italian CDCDs, using the following inclusion criteria: (i) a last routine cognitive evaluation between January and March 2020, and a next follow-up ambulatory visit expected between September and December 2020; (ii) a diagnosis of Mild Cognitive Impairment (MCI), or of mild-to-moderate dementia (through a complete evaluation that considered DSM-5 criteria, Mini-Mental State Examination [MMSE] score and neuroimaging); and (iii) an MMSE score of >16. Exclusion criteria were the following: (i) inability to undergo psychometrics tests for any reason; and (ii) history of psychiatric illness, according to clinical anamnesis.

Of the 496 individuals who met the inclusion criteria and were originally contacted, 275 agreed to participate in the study and were assessed at T0 and T1. At T2, the comprehensive geriatric assessment (CGA) was administered to 214 individuals (35 refused, nine patients were no longer at home with the same caregiver, six were institutionalized, and 11 died). Of the 214 patients who performed both T0 and T2 CGA evaluations, we selected only those evaluated with the Fatigue, Resistance, Ambulation, Illness, and Loss of Weight (FRAIL) scale,

resulting in 165 participants with MMSE score recorded from 18 (the lowest MMSE score recorded) to 30 and aged 70–91 years (Fig. 1).

Procedure

Trained physicians gathered the following information on the patient characteristics for each participant from medical records.

We considered participants' sociodemographic data (age, sex, cohabiting status, years of education), information on risk behaviours (smoking and alcohol consumption), medical history (including diagnosis of

cognitive impairment, depressive mood, and other coexisting chronic diseases and comorbidities), and drug treatments.

At T0 and T2, the CGA¹⁸ was performed during ambulatory evaluations and included the following scales: MMSE,¹⁹ Neuropsychiatric Inventory (NPI),²⁰ Activities of Daily Living (ADL),²¹ and Instrumental Activities of Daily Living (IADL).²² The NPI is used to assess the presence and severity of specific behavioural and psychological symptoms of dementia (BPSD) experienced by each participant, as rated by a reliable study partner. BPSD symptoms assessed




Figure 1 STROBE flowchart of patient selection

in the NPI questionnaire include delusions, hallucinations, agitation/aggression, depression/dysphoria, anxiety, euphoria, apathy, disinhibition, irritability/lability, aberrant motor activity, night-time behaviour, and appetite disturbance. For each symptom endorsed, the caregiver rated both the severity (1 to 3, for mild, moderate, and marked) and frequency (1 = occasionally, less than once per week; 2 = often, about once per week; 3 = frequently, several times per week but less than every day; 4 = very frequently, once or more per day or continuously) of each symptom. Frequency and severity scores were multiplied to yield an overall severity score (maximum of 12) for each domain, which was summed across all domains to yield a total NPI score.

To assess social isolation and loneliness, moreover, the UCLA Loneliness Scale-3²³ was administered. The three items of the UCLA scale were (i) lack of companionship, (ii) feeling left out, and (iii) feeling isolated, in the past 2 weeks. For each question, a 4-point Likert scale (1 = never, 2 = rarely, 3 = sometimes, and 4 = always) was used; scores ranged from 3 to 12, with higher scores indicating higher levels of loneliness. Concerning participants' social environment, we considered the presence of a formal or informal caregiver, and the number of informal visits received on average by each participant before the pandemic period.

T1 evaluations were carried-out by telephonic interviews to the patients, and included the Depression Anxiety Stress Scales-21 (DASS-21),²⁴ Perceived Stress Scale (PSS),²⁵ and Coping Orientation to Problems Experienced (COPE),²⁶ described below.

The DASS-21²⁴ is composed of a set of three self-report scales designed to measure the emotional states of depression, anxiety, and stress. Each of the three DASS-21 scales contains seven items, divided into subscales with similar content. Cut-off scores for depression, anxiety, and stress were 10, 8, and 15, respectively. The depression scale (including items 3, 5, 10, 13, 16, 17, and 21) assesses dysphoria, hopelessness, devaluation of life, self-deprecation, lack of interest/involvement, anhedonia, and inertia. The anxiety scale (including the items 2, 4, 7, 9, 15, 19, and 20) assesses autonomic arousal, skeletal muscle effects, situational anxiety, and subjective experience of anxious affect. The stress scale is sensitive to levels of chronic non-specific arousal and included the items 1, 6, 8, 11, 12, 14, and 18). It assesses difficulty

relaxing, nervous arousal, and being easily upset/agitated, irritable/over-reactive, and impatient. Scores for depression, anxiety, and stress are calculated by summing the scores for the relevant items, then multiplied by two. The cut-offs were used to detect the presence of symptoms of depression.

The PSS²⁵ is the most frequently used psychological measure to assess perceptions of stress. The degree to which the situations in a person's life are rated as stressful are evaluated by 10 items constructed to capture the level at which respondents perceive their lives as unpredictable, uncontrollable, or overloaded. The scale also contains a series of direct questions about current levels of perceived stress. The PSS was designed to be used in samples drawn from the general population with an educational level at least equal to lower middle school. The items and the response alternatives are easy to understand: for each item, respondents are asked to indicate how often they felt a certain way in the last month (0 = Never, to 4 = Very often). The PSS scores are obtained by reverse-scoring the responses to the four positively formulated items (items 4, 5, 7, and 8), then adding together the scores for all items. A short 4-item scale can be obtained using questions 2, 4, 5, and 10 of the 10 items in the PSS scale.

The COPE²⁶ is a multidimensional coping inventory to assess the different ways in which people respond to stress. Five scales (of four items each) measure conceptually distinct aspects of problem-focused coping (active coping, planning, suppression of competing activities, restraint coping, seeking of instrumental social support); five scales measure aspects of what might be viewed as emotion-focused coping (seeking of emotional social support, positive reinterpretation, acceptance, denial, turning to religion); and three scales measuring coping responses that arguably are less useful (focus on and venting of emotions, behavioural disengagement, mental disengagement).

The subscales drawn from the COPE for use in the current study are calculated as follows: the subscale 'social support' is indicated by the sum of items 1, 10, 15, 18, and 25; 'positive attitude' is indicated by the sum of items 2, 6, 12, 16, 23, and 24; 'orientation to problem' is indicated by the sum of items 3, 5, 9, 13, 20), and 'transcendent orientation' (indicated by the sum of items 8, 11, 14, 19). The sum of the remaining scales refers to "avoidance strategies".

The presence of a caregiver was always required during telephone interviews in order to limit potential biases due to patients' cognitive impairment and their ability to answer questions.^{11,27} Finally, the caregivers were evaluated through the Caregiver Burden Inventory (CBI)²⁸ during the same telephone call. The CBI is a 24-item self-report questionnaire for assessing the burden of caregivers caring for people with chronic disease. The items are rated on a 5-point Likert scale from 0 (Never) to 4 (Nearly always). The questions cover five dimensions of caregiver burden: objective burden; time-dependence, referring to time demands for assistance; psychological burden, understood as the caregiver's feelings of exclusion from expectations and opportunities; physical burden, which describes the caregiver's feelings of fatigue and health problems; social burden, which describes the caregiver's feelings of role conflict; and emotional burden, which describes the caregiver's feelings of shame or embarrassment caused by the patient. Time spent for assistance, social involvement, physical involvement, and relational involvement are represented, respectively by the sum of the items from 1 to 5, from 6 to 10, 11 to 14, and 15 to 19.

We categorized participants according to the FRAIL scale,²⁹ which consists in a brief assessment of five items: fatigue, resistance, ambulation, illnesses, and loss of weight. For this study, 'fatigue' was measured by asking respondents how much time during the past 4 weeks they felt tired, 'resistance' was assessed by asking participants if they had any difficulty walking up 10 steps alone without resting and without aids, and 'ambulation' by asking if they had any difficulty walking several hundred yards alone and without aids; yes responses were each scored as 1 point. 'Illness' was scored 1 for respondents who reported five or more illnesses in their pathological anamnesis. Loss of weight was scored 1 for respondents with a weight decline of 5% or greater within the past 12 months based on self-report. We performed frailty evaluations with patients and their caregivers, so that the caregiver could help to reconstruct the response when the patient was unable to provide an answer or provided it uncertainly. Each FRAIL scale characteristic was scored 0–1, and scores ranged from 0 (best) to 5 (worst). According to the FRAIL scale, participants were categorized as frail (score ≥ 3), pre-frail

(score = 1–2) and non-frail (score = 0). The FRAIL scale was administrated at both T0 and T2.

Analyses

Descriptive characteristics of the sample are expressed as means \pm standard deviations (SDs) or as count (%), as appropriate.

The comparison of the sociodemographic and health-related characteristics of non-frail, pre-frail, and frail participants was performed by using chi-squared or Fisher exact tests for categorical variables and generalized linear model testing for homoscedasticity through the Levene's test for quantitative ones. Post hoc analyses with Bonferroni adjustment for multiple comparisons were applied. The comparison of quantitative variables between T0 and T2 was performed through paired-samples *t*-tests. Delta (Δ) values for functional and psycho-cognitive variables were obtained from the difference of T2 and T0 values of the same items. In particular, Δ values for ADL and IADL tests were considered as 'loss of functional autonomy', 'loss of ADL', or 'loss of IADL' when there was a loss of at least 1 point in the final score compared to the previous evaluation.

A multivariable logistic regression was performed to test the association between frailty and changes in social isolation feeling, functional and cognitive status, and depressive or behavioural symptoms after T2. The model was corrected for age, sex, and education status. Analyses were performed in IBM SPSS version 28.0.1.0 (IBM Corp., Armonk, NY).

RESULTS

We evaluated 165 individuals (61.9% women), including 47 individuals who were not frail, 92 who were pre-frail, and 26 who were frail. The most common cognitive disorders observed were Alzheimer's disease (28.7%), vascular dementia (24.7%), MCI (24.3%), and mixed dementia (14.9%).

Table 1 and Fig. 2 show the participants' functional, cognitive, and neuropsychiatric characteristics, expressed as variations between T0 and T2, and as single scores at T0 and T2. The sample's mean age was 79.49 ± 4.89 years and the frail patients were the oldest group (82.08 ± 4.17 vs 78.21 ± 4.65 for no-frail and 79.96 ± 4.87 for pre-frail, $P = 0.004$). Subjects were stable in functional and psycho-cognitive domains during the observational period

between frailty groups (Table 1). In pre-frail patients we observed significant reductions of functional autonomy parameters during the 6 months of follow-

up (Fig. 2, $P < 0.01$). Differences in the MMSE and NPI total scores between the frailty groups before and after the lockdown were not observed, while

Table 1 Descriptive functional and psycho-cognitive characteristic as variations between T0 and T2 in the total sample and sorted by frailty levels (FRAIL scale)

| N item (18) | All patients (n = 165) | Not frail (FRAIL = 0) (n = 47) | Pre-frail (FRAIL = 1–2) (n = 92) | Frail (FRAIL >2) (n = 26) | P-value |
|----------------------------|------------------------|--------------------------------|----------------------------------|---------------------------|------------------|
| Age (years) | 79.59 ± 4.89 | 78.21 ± 4.65 | 79.96 ± 4.87 | 82.08 ± 4.17 | 0.01 |
| Sex (F) | 61.9% | 55.3% | 65.2% | 65.4% | 0.49 |
| Education (years) | 7.36 ± 3.74 | 7.23 ± 3.98 | 7.04 ± 3.380 | 7.00 ± 3.55 | 0.94 |
| Lived alone: yes | 22.4% | 27.7% | 17.4% | 30.8% | 0.16 |
| Caregiver: yes | 78.5% | 80.9% | 76.1% | 92.3% | 0.18 |
| Smoking habit | | | | | 0.69 |
| Never | 76.4% | 76.0% | 74.2% | 84.6% | |
| Former | 17.6% | 18.0% | 20.2% | 7.7% | |
| Current | 6.0% | 6.0% | 5.6% | 7.7% | |
| Use of antipsychotics | 15.7% | 14.9% | 14.1% | 23.8% | 0.54 |
| Depressive mood | 19.4% | 20.0% | 14.3% | 34.6% | 0.72 |
| Hearing deficits | 12.5% | 8.0% | 15.5% | 19.2% | 0.36 |
| Vision deficits | 11.3% | 4.0% | 13.1% | 11.5% | 0.12 |
| Diabetes mellitus | 17.6% | 12.2% | 17.9% | 26.9% | 0.28 |
| Hypertension | 68.9% | 46.0% | 81.2% | 73.1% | <0.001 |
| Atrial fibrillation | 13.1% | 8.0% | 13.1% | 23.1% | 0.18 |
| COPD | 4.4% | 0 | 6.0% | 7.7% | 0.17 |
| Number of chronic diseases | 3.28 ± 1.72 | 2.35 ± 1.02 | 3.49 ± 1.73 | 4.30 ± 1.95 | <0.001 |
| Functional domains | | | | | |
| ΔADL | −0.23 ± 1.01 | −0.04 ± 0.59 | −0.27 ± 0.98 | −0.36 ± 1.75 | 0.35 |
| ΔIADL | −0.49 ± 1.37 | −0.40 ± 1.09 | −0.71 ± 1.48 | −0.31 ± 1.49 | 0.27 |
| Psycho-cognitive domains | | | | | |
| ΔMMSE | −0.17 ± 2.35 | −0.32 ± 1.91 | −0.07 ± 2.46 | −0.92 ± 2.79 | 0.27 |
| ΔUCLA | 0.49 ± 0.21 | 0.61 ± 1.50 | 0.45 ± 1.18 | 0.70 ± 1.14 | 0.65 |
| ΔNPI | 2.35 ± 11.28 | 1.42 ± 8.04 | 2.86 ± 12.02 | 1.92 ± 15.12 | 0.87 |

Note: Numbers are mean ± standard deviation, or count (%), as appropriate. P-values refer to the comparisons between FRAIL groups. Post hoc analyses with Bonferroni adjustment for multiple comparisons were applied. Abbreviations: FRAIL, Fatigue, Resistance, Ambulation, Illness, and Loss of Weight scale; COPD, chronic obstructive pulmonary disease; ADL, Activities of Daily Living; IADL, Instrumental ADL; MMSE, Mini-Mental State Examination; UCLA, UCLA Loneliness Scale-3; NPI, Neuropsychiatric Inventory. The values of $p < 0.05$ are in bold.




Figure 2 T0 and T2 descriptive cognitive, functional, neuropsychiatric, and social characteristics sorted by frailty levels (FRAIL scale). ADL, Activities of Daily Living; IADL, Instrumental ADL; MMSE, Mini-Mental State Examination; NPI, Neuropsychiatric Inventory; NPI, Neuropsychiatric Inventory; UCLA, UCLA Loneliness Scale-3. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

Table 2 T0 and T2 descriptive neuropsychiatric characteristics in the total sample and sorted by frailty levels

| N item (26) | All patients evaluated with FRAIL scale (<i>n</i> = 165) | Non-frail (FRAIL = 0) (<i>n</i> = 47) | Pre-frail (FRAIL = 1–2) (<i>n</i> = 92) | Frail (FRAIL >2) (<i>n</i> = 26) | <i>P</i> -value |
|---------------------------------------|---|--|--|-----------------------------------|------------------|
| NPI T0 | 9.59 ± 10.62 | 8.36 ± 9.95 | 8.29 ± 9.75 | 18.36 ± 12.38 | 0.01 |
| NPI T2 | 12.52 ± 11.65* | 10.46 ± 11.49 | 11.22 ± 10.42* | 21.71 ± 13.23 | 0.01 |
| Delusions | | | | | |
| NPI T0 Delusions F × S | 0.38 ± 1.78 | 0.47 ± 1.52 | 0.21 ± 1.49 | 0.93 ± 3.19 | 0.37 |
| NPI T2 Delusions F × S | 0.51 ± 1.98* | 0.43 ± 1.73 | 0.53 ± 2.02 | 0.60 ± 2.32 | 0.95 |
| Hallucination | | | | | |
| NPI T0 Hallucination F × S | 0.09 ± 0.46 | 0.19 ± 0.74 | 0.05 ± 0.27 | 0.00 ± 0.00 | 0.23 |
| NPI T2 Hallucination F × S | 0.10 ± 0.57 | 0.14 ± 0.75 | 0.10 ± 0.53 | 0.07 ± 0.25 | 0.90 |
| Agitation/aggression | | | | | |
| NPI T0 Agitation/aggression F × S | 0.50 ± 1.28 | 0.61 ± 1.29 | 0.48 ± 1.37 | 0.36 ± 0.74 | 0.79 |
| NPI T2 Agitation/aggression F × S | 1.27 ± 2.35** | 1.29 ± 2.29* | 1.44 ± 2.59* | 0.53 ± 1.12 | 0.41 |
| Depression/dysphoria | | | | | |
| NPI T0 Depression/dysphoria F × S | 2.06 ± 3.32 | 1.97 ± 3.59 | 1.75 ± 2.70 | 3.79 ± 4.75 | 0.11 |
| NPI T2 Depression/dysphoria F × S | 3.01 ± 3.78 | 2.79 ± 3.95 | 2.49 ± 3.20* | 5.56 ± 4.84 | 0.01 |
| Anxiety | | | | | |
| NPI T0 Anxiety F × S | 1.60 ± 2.74 | 1.26 ± 1.97 | 1.24 ± 2.45 | 4.14 ± 4.24 | <0.001 |
| NPI T2 Anxiety F × S | 1.86 ± 2.75 | 1.45 ± 2.37 | 1.67 ± 2.16 | 3.50 ± 4.73 | 0.03 |
| Euphoria/elation | | | | | |
| NPI T0 Euphoria/elation F × S | 0.16 ± 0.84 | 0.31 ± 1.34 | 0.11 ± 0.50 | 0.00 ± 0.00 | 0.39 |
| NPI T2 Euphoria/elation F × S | 0.10 ± 0.57* | 0.18 ± 0.77 | 0.10 ± 0.53* | 0.00 ± 0.00 | 0.61 |
| Apathy/indifference | | | | | |
| NPI T0 Apathy/indifference F × S | 1.58 ± 2.83 | 1.03 ± 2.23 | 1.40 ± 2.43 | 3.86 ± 4.65 | 0.01 |
| NPI T2 Apathy/indifference F × S | 1.62 ± 2.76 | 1.23 ± 2.17 | 1.44 ± 2.56 | 3.06 ± 4.02 | 0.07 |
| Disinhibition | | | | | |
| NPI T0 Disinhibition F × S | 0.23 ± 1.29 | 0.14 ± 0.68 | 0.33 ± 1.63 | 0.00 ± 0.00 | 0.60 |
| NPI T2 Disinhibition F × S | 0.31 ± 1.21 | 0.43 ± 1.37 | 0.27 ± 1.23 | 0.27 ± 0.79 | 0.84 |
| Irritability/lability | | | | | |
| NPI T0 Irritability/lability F × S | 1.03 ± 2.38 | 0.61 ± 1.55 | 1.03 ± 2.35 | 2.14 ± 3.75 | 0.12 |
| NPI T2 Irritability/lability F × S | 1.22 ± 1.93 | 1.04 ± 2.13 | 1.36 ± 1.91 | 0.93 ± 1.62 | 0.62 |
| Aberrant motor behaviour | | | | | |
| NPI T0 Aberrant motor behaviour F × S | 0.22 ± 1.03 | 0.22 ± 0.76 | 0.27 ± 1.25 | 0.00 ± 0.00 | 0.68 |
| NPI T2 Aberrant motor behaviour F × S | 0.20 ± 0.93 | 0.36 ± 1.33 | 0.16 ± 0.81 | 0.07 ± 0.25 | 0.55 |
| Nighttime behaviours | | | | | |
| NPI T0 Nighttime behaviours F × S | 1.44 ± 2.53 | 0.97 ± 2.45 | 1.55 ± 2.28 | 2.14 ± 3.61 | 0.30 |
| NPI T2 Nighttime behaviours F × S | 2.41 ± 3.23** | 1.72 ± 3.25 | 2.33 ± 2.87* | 4.13 ± 4.29* | 0.05 |
| Appetite/eating | | | | | |
| NPI T0 Appetite/eating F × S | 0.41 ± 1.46 | 0.37 ± 2.03 | 0.30 ± 0.75 | 1.00 ± 2.18 | 0.26 |
| NPI T2 Appetite/eating F × S | 0.61 ± 1.66 | 0.32 ± 0.81 | 0.33 ± 0.86 | 2.33 ± 3.57 | <0.001 |

Note: Numbers are mean ± standard deviation, or count (%), as appropriate. *P*-values refer to the comparisons between FRAIL groups. Frequencies do not sum to 100% due to missing values. Post-hoc analyses with Bonferroni adjustment for multiple comparisons were applied. Abbreviations: FRAIL, Fatigue, Resistance, Ambulation, Illness, and Loss of Weight scale; NPI, Neuropsychiatric Inventory; F, frequency; S, severity. The values of *p* < 0.05 are in bold. * < 0.05; ** < 0.01.

UCLA scores increased from T0 to T2 in all groups ($P < 0.01$).

The analysis of the descriptive items of the NPI scale sorted by frailty levels at T0 and T2 is reported in Table 2. Frail people had higher depression (5.56 ± 4.84 vs 2.79 ± 3.95 in not frail and 2.49 ± 3.20 in pre-frail, $P = 0.01$), anxiety (3.50 ± 4.73 vs 1.45 ± 2.37 in not frail and 1.67 ± 2.16 in pre-frail, $P = 0.03$), and eating disorders (2.33 ± 3.57 vs 0.32 ± 0.81 in not frail and 0.33 ± 0.86 in pre-frail, $P < 0.001$) scores at T2. Between T0 and T2, there was an increase in scores for agitation (0.48 ± 1.37 vs 1.44 ± 2.59 , $P < 0.05$), depression (1.75 ± 2.70 vs 2.49 ± 3.20 , $P < 0.05$), and sleep disorders (1.55 ± 2.28 vs 2.33 ± 2.87 , $P < 0.05$) in the pre-frail group.

The evaluation during the lockdown period (T1) showed that frail individuals had higher total DASS (19.76 ± 13.99 vs 11.53 ± 11.57 in not frail and 14.99 ± 10.90 in pre-frail, $P = 0.02$) and PSS (18.88 ± 7.72 vs 14.07 ± 7.41 in not frail and 15.93 ± 6.74 in pre-frail, $P = 0.02$) scores (Table 3), and worse 'positive attitude' coping strategies compared to the other groups (13.77 ± 3.07 vs 16.98 ± 4.12 in not frail and 15.87 ± 4.09 in pre-frail).

Higher caregiving burden, in particular for social and relational involvement (items 15–19 and items 6–10, respectively) were observed in caregivers of frail subjects.

In the multivariable logistic regression, after adjusting for age, sex, and schooling, we found that frailty at T2 was associated with the worsening of social isolation during lockdown (odds ratio (OR): 1.24, 95% confidence interval (CI) 1.060–1.950,

Table 4 Logistic multivariable regression: association between frailty at T2 and covariates

| Variable | OR | 95% CI | | P-value |
|---------------|------|--------|-------|-------------|
| | | Lower | Upper | |
| Age | 0.84 | 0.55 | 1.29 | 0.43 |
| Sex M | 0.19 | 0.01 | 6.09 | 0.36 |
| Schooling | 1.45 | 0.88 | 2.37 | 0.14 |
| Δ IADL | 5.72 | 1.03 | 1.89 | 0.04 |
| Δ MMSE | 0.48 | 0.20 | 1.15 | 0.10 |
| Δ NPI | 0.87 | 0.65 | 1.15 | 0.31 |
| Δ UCLA | 1.24 | 1.06 | 1.95 | 0.04 |

Abbreviations: OR, odds ratio; CI, confidence interval; IADL, Instrumental Activities of Daily Living; MMSE, Mini-Mental State Examination; NPI, Neuropsychiatric Inventory; UCLA, UCLA Loneliness Scale-3. The values of $p < 0.05$ are in bold.

Table 3 Results of the evaluation through DASS, COPE and CBI scales at T1

| N Item (15) | All patients ($n = 165$) | Non-frail (FRAIL = 0) ($n = 47$) | Pre-frail (FRAIL = 1–2) ($n = 92$) | Frail (FRAIL >2) ($n = 26$) | P-value |
|-------------------------------|----------------------------|------------------------------------|--------------------------------------|-------------------------------|-------------|
| DASS total (sum of all items) | 14.73 \pm 11.83 | 11.53 \pm 11.57 | 14.99 \pm 10.90 | 19.76 \pm 13.99 | 0.02 |
| DASS stress | 11.41 \pm 9.33 | 8.77 \pm 8.58 | 12.11 \pm 9.26 | 13.69 \pm 10.19 | 0.05 |
| DASS anxiety | 6.04 \pm 7.21 | 4.60 \pm 6.83 | 5.63 \pm 6.58 | 10.08 \pm 8.73 | 0.01 |
| DASS depression | 12.11 \pm 9.77 | 9.70 \pm 9.24 | 12.24 \pm 9.29 | 16.16 \pm 11.35 | 0.01 |
| PSS total | 15.89 \pm 7.21 | 14.07 \pm 7.41 | 15.93 \pm 6.74 | 18.88 \pm 7.72 | 0.02 |
| COPE | | | | | |
| COPE social support | 11.68 \pm 3.94 | 12.16 \pm 3.69 | 11.60 \pm 3.95 | 11.15 \pm 4.36 | 0.56 |
| COPE avoidance strategies | 8.35 \pm 2.66 | 7.96 \pm 2.83 | 8.45 \pm 2.47 | 8.69 \pm 3.01 | 0.46 |
| COPE positive attitude | 15.85 \pm 4.07 | 16.98 \pm 4.12 | 15.87 \pm 4.09 | 13.77 \pm 3.07 | 0.01 |
| COPE orientations to problem | 12.55 \pm 3.47 | 13.20 \pm 3.33 | 12.30 \pm 3.54 | 12.23 \pm 3.45 | 0.32 |
| COPE transcendent orientation | 10.52 \pm 4.26 | 10.44 \pm 4.48 | 10.44 \pm 4.20 | 10.92 \pm 4.22 | 0.87 |
| CBI | | | | | |
| CBI time-dependence | 7.09 \pm 5.29 | 5.97 \pm 5.62 | 6.88 \pm 4.62 | 9.33 \pm 6.12 | 0.04 |
| CBI psychological burden | 4.83 \pm 4.91 | 4.71 \pm 5.20 | 3.96 \pm 3.82 | 7.48 \pm 6.35 | 0.01 |
| CBI physical burden | 3.57 \pm 3.70 | 3.26 \pm 3.39 | 3.18 \pm 3.37 | 5.08 \pm 4.68 | 0.07 |
| CBI social burden | 2.40 \pm 3.28 | 2.88 \pm 3.43 | 1.54 \pm 2.39 | 4.13 \pm 4.40 | 0.01 |
| CBI emotional burden | 1.45 \pm 2.22 | 1.51 \pm 2.10 | 1.21 \pm 2.07 | 2.00 \pm 2.72 | 0.31 |

Note: Numbers are mean \pm standard deviation, or count (%), as appropriate. P-values refer to the comparisons between FRAIL groups. Frequencies do not sum to 100% due to missing values. Post hoc analyses with Bonferroni adjustment for multiple comparisons were applied. Abbreviations: FRAIL, Fatigue, Resistance, Ambulation, Illness, and Loss of Weight scale; DASS, Depression Anxiety Stress Scales; PSS, Perceived Stress Scale; COPE, Coping Orientation to Problems Experienced; CBI, Caregiver Burden Inventory. The values of $p < 0.05$ are in bold.

$P = 0.042$), and loss of IADL (OR: 5.72, 95% CI 1.025–1.889, $P = 0.047$) (Table 4).

DISCUSSION

In our study, we observed more anxiety and stress symptoms in frail patients and their caregivers during lockdown; frailty was associated with increased feelings of social isolation and loneliness after the first wave of the COVID-19 pandemic. Nevertheless, the worsening in functional and psycho-cognitive domains was not exacerbated in frail individuals, but occurred similarly in all involved individuals. Pre-frail patients seemed to experience more neuropsychiatric disorders (especially agitation, aggression, and sleep disorder domains) during lockdown.

During containment measures, we observed a slight increase in anxiety, depression, and stress symptoms in frail patients. Looking at caregiver burden, in the context of the COVID-19 pandemic we confirmed increases in especially social and relational involvement, and less but still significant in time spent for assistance and physical involvement, for caregivers of frail individuals, in line with previous studies.^{30–32} Moreover, our data confirmed the association between social isolation and frailty in patients affected by cognitive impairment in the post-pandemic phase.^{17,33} High prevalence of loneliness and social isolation in older adults have already been reported before the COVID-19 pandemic, with loss of social partners,³⁴ reduction of social contacts,³⁵ and physical limitations as some of the mechanisms involved.¹² The mobility limitations induced by the lockdown could both limit communication with family and friends, and could erase the routine, confusing the circadian rhythm and flattening the days of older people, resulting in reduced cognitive stimulation. All of these may impact negatively on cognitive and affective well-being.¹³ We hope that in the future greater attention will be given to early detection of social isolation as a modifiable risk factor for frailty, especially considering patients affected by NCDs; part of primary care may involve taking charge of the psychological well-being of the patient and his or her caregiver, an aspect of attention that today is unfortunately reserved only for cases where required.

Regarding the health domains analysed, surprisingly persons affected by NCDs remained broadly stable, save for some expected minimal within-group

variation in functional and neuro-psychiatric areas. The UCLA scores were higher after lockdown in all subjects, regardless of frailty, while there was a deterioration especially in the not frail and pre-frail patients in terms of the IADL scores. A possible explanation could be that, while frail patients were more impaired even before the start of the pandemic, the reduction in mobility due to the lockdown could have significantly compromised the residual autonomy of the remaining two groups of subjects; physical limitations could be responsible for an exacerbation of joint pain, which translates into greater stiffening and increasing difficulty in performing even the simplest movements.³⁶ In our sample we did not observe significant changes in the MMSE scores, in contrast with previous studies^{37–39}; however, the methodological strategies were different, i.e. the follow-up periods and the way in which the MMSE scale was administered. Our evaluations at T0 and T2 were both performed during ambulatory visits, in contrast with other studies, in which the first clinic appointment was followed by a telephone interview. Moreover, our last evaluation was at 69 months, suggesting that minimal cognitive changes could not be highlighted with the MMSE. This is supported also in the study of Sardella *et al.*, in which no MMSE change between baseline and the first follow-up was reported.

An interesting result was the NPI trend, in which scores increased in all groups from T0 to T2, but no significant variations between the three type of patients considered were reported. There was a further worsening of agitation/aggression in the pre-frail and not frail groups, and of night-time behaviour in frail and pre-frail individuals, supporting the hypothesis that restrictive measures adopted during the COVID-19 lockdown could increase BPSD even in older adults with cognitive impairment despite of their frailty status.⁴⁰ Previous studies conducted both in the periods before and during COVID-19 showed a substantial stability of the NPI, showing an increase only in specific sub-categories^{41–43}; however, they were conducted with different scales and covering shorter periods than our study.^{2,44} Although not perfectly comparable to other papers actually in the literature, due to different methods or timing of evaluation, we support the hypothesis that the restrictive measures adopted during the COVID-19 waves and the pandemic itself could increase

especially the agitation, anxiety, and depression of patients due to social isolation and concern for their own health status and that of their families, perhaps explaining our results. The unique value of our results lies in the longer observation time of the patients, which was more able to highlight the pandemic lockdown's effect in subjects with NCDs. To our knowledge, this is the first study that considered variations of functional, psycho-cognitive, and behavioural disorders based on the degree of frailty. Overall, we can say that pre-frail patients were apparently the most vulnerable to minimal changes during the pandemic, as if the lockdown had somehow moved the threshold for defining a frail subject. In line with this, the impairment of coping ability inherent in the concept of fragility^{7,45} fails, as no significant differences were detected in the COPE scale during the lockdown.

Some limitations should be acknowledged. First, the scales used for the psycho-affective evaluation at T1 have not been validated for remote administration. However, these tools have been previously administered remotely,⁴⁶ although originally not designed in this format. Second, we do not know the psycho-affective profile of the patients with cognitive impairment before the pandemic; we only have reference population data. Another limit is that our study did not consider a control group of individuals without cognitive impairment. Also, due to the strict criteria of our study, the sample size was small, limiting the statistical power of analyses, especially considering the inter-group and intra-group comparisons that were conducted.

Our study also has some strengths. It was in fact a multicentre study that investigated the topic of cognitive-behavioural changes during the COVID-19 pandemic on a specific population of elderly people affected by cognitive impairment reevaluated at different time points. Finally, we evaluated multiple health domains, considering different aspects of psychological well-being, including the caregiver's point of view.

CONCLUSIONS

This study found that during the Covid-19 pandemic, people affected by NCDs with a frail status seemed to experience more anxiety and stress disorders than pre-frail and not frail individuals, although a worsening of BPSDs was observed in all patients regardless

of their degree of frailty. Also, frail patients' caregivers seem to have suffered the effects of the pandemic the most. Our study points out the role of social isolation in worsening the feeling of loneliness of frail patients with cognitive impairment.

FUNDING INFORMATION

This research did not receive any specific grants from funding agencies in the public, commercial, or not-for-profit sectors.

DISCLOSURE

There are no known conflicts of interest associated with this publication, and there has been no significant financial support for this work that could have influenced its outcome. The manuscript has been read and approved by all named authors, and there are no other persons who satisfy the criteria for authorship that are not listed. The order of authors listed in the manuscript has been approved by all of the authors. The Corresponding Author is the sole contact for enquiries, and we confirm that we have provided a current, correct email address which is accessible by the Corresponding Author.

STUDY APPROVAL STATEMENT

The study protocol was conducted according to good clinical practice guidelines and the ethical standards of the 1964 Declaration of Helsinki as revised in 2000 and was reviewed and approved by the BIO-CAMPUS Ethics Committee, University of Rome—Prot. Number: 22.5 (20).20 OSS ComEtUCBM.

CONSENT TO PARTICIPATE STATEMENT

The subjects participating in this study received a thorough explanation of the risks and benefits of inclusion and gave their oral and written informed consent to publish the data.

DATA AVAILABILITY STATEMENT

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

REFERENCES

- Pancani L, Marinucci M, Riva AN. Forced social isolation and mental health: a study on 1006 Italians under COVID-19 lockdown. *Front Psychol* 2021; **12**: 663799. <https://doi.org/10.3389/fpsyg.2021.663799>.
- Cagnin A, Di LR, Marra C *et al*. Behavioral and psychological effects of coronavirus Disease-19 quarantine in patients with dementia. *Art Ther* 2020; **11**: 1. <https://doi.org/10.3389/fpsyg.2020.578015>.
- Mendoza-Ruvalcaba NM, Gutiérrez-Herrera R, López C *et al*. Impact of quarantine due to COVID-19 pandemic on health and lifestyle conditions in older adults from Centro American countries. *PloS One* 2022; **17**: e0264087. <https://doi.org/10.1371/JOURNAL.PONE.0264087>.
- Orrù G, Ciacchin R, Gemignani A, Conversano C. Psychological intervention measures during the COVID-19 pandemic. *Clin Neuropsychiatry* 2020; **17**: 76–79. Accessed November 2, 2022. <https://arpi.unipi.it/retrieve/handle/11568/1039146/592302/PsychologicalinterventionmeasuresduringtheCOVID-19pandemic.pdf>.
- Müller I, Mancinetti M, Renner A *et al*. Frailty assessment for COVID-19 follow-up: a prospective cohort study. *BMJ open Respir Res* 2022; **9**(1): e001227. <https://doi.org/10.1136/BMJRESP-2022-001227>.
- Fried LP, Tangen CM, Walston J *et al*. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001; **56**: M146–M157. <https://doi.org/10.1093/GERONA/56.3.M146>.
- Fried LP, Hadley EC, Walston JD *et al*. From bedside to bench: research agenda for frailty. *Sci Aging Knowledge Environ* 2005; **2005**(31): pe24. <https://doi.org/10.1126/SAGEKE.2005.31.PE24>.
- Leng S, Chen X, Mao G. Frailty syndrome: an overview. *Clin Interv Aging* 2014; **9**: 433–441. <https://doi.org/10.2147/CIA.S45300>.
- Killgore WDS, Cloonan SA, Taylor EC, Dailey NS. Loneliness: a signature mental health concern in the era of COVID-19. *Psychiatry Res* 2020; **290**: 113117. <https://doi.org/10.1016/J.PSYCHRES.2020.113117>.
- Bunt S, Steverink N, Olthof J, van der Schans CP, Hobbelen JSM. Social frailty in older adults: a scoping review. *Eur J Ageing* 2017; **14**: 323–334. <https://doi.org/10.1007/S10433-017-0414-7>.
- Coin A, Devita M, Trevisan C *et al*. Psychological well-being of older adults with cognitive deterioration during quarantine: preliminary results from the GeroCovid initiative. *Front Med* 2021; **8**: 8. <https://doi.org/10.3389/FMED.2021.715294>.
- Pinquart M, Sörensen S. Influences on loneliness in older adults: a meta-analysis. *Basic Appl Soc Psych* 2001; **23**: 245–266. https://doi.org/10.1207/S15324834BASP2304_2.
- Smith L, Boska J, Jacob L *et al*. Is loneliness associated with mild cognitive impairment in low- and middle-income countries? *Int J Geriatr Psychiatry* 2021; **36**: 1345–1353. <https://doi.org/10.1002/GPS.5524>.
- Vernuccio L, Sarà D, Inzerillo F *et al*. Effect of COVID-19 quarantine on cognitive, functional and neuropsychiatric symptoms in patients with mild cognitive impairment and dementia. *Aging Clin Exp Res* 2022; **34**: 1187–1194. <https://doi.org/10.1007/s40520-022-02113-z>.
- Simonetti A, Pais C, Jones M *et al*. Neuropsychiatric symptoms in elderly with dementia during COVID-19 pandemic: definition, treatment, and future directions. *Front Psychiatry* 2020; **11**: 579842. <https://doi.org/10.3389/FPSYT.2020.579842/FULL>.
- Devita M, Bordignon A, Sergi G, Coin A. The psychological and cognitive impact of Covid-19 on individuals with neurocognitive impairments: research topics and remote intervention proposals. *Aging Clin Exp Res* 2021; **33**: 733–736. <https://doi.org/10.1007/S40520-020-01637-6>.
- Kotwal AA, Holt-Lunstad J, Newmark RL *et al*. Social isolation and loneliness among San Francisco Bay Area older adults during the COVID-19 shelter-in-place orders. *J Am Geriatr Soc* 2021; **69**: 20–29. <https://doi.org/10.1111/JGS.16865>.
- Schippinger W. Comprehensive geriatric assessment. *Wien Med Wochenschr* 2022; **172**: 122–125. <https://doi.org/10.1007/S10354-021-00905-Y>.
- Magni E, Binetti G, Bianchetti A, Rozzini R, Trabucchi M. Mini-mental state examination: a normative study in Italian elderly population. *Eur J Neurol* 1996; **3**: 198–202. <https://doi.org/10.1111/J.1468-1331.1996.TB00423.X>.
- Cummings J. The neuropsychiatric inventory: development and applications. *J Geriatr Psychiatry Neurol* 2020; **33**: 73–84. <https://doi.org/10.1177/0891988719882102>.
- Katz S, Downs TD, Cash HR, Grotz RC. Progress in development of the index of ADL. *Gerontologist* 1970; **10**: 20–30. https://doi.org/10.1093/geront/10.1_Part_1.20.
- Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist* 1969; **9**: 179–186. https://doi.org/10.1093/geront/9.3_Part_1.179.
- Russell DW. UCLA loneliness scale (version 3): reliability, validity, and factor structure. *J Pers Assess* 1996; **66**: 20–40. https://doi.org/10.1207/S15327752JPA6601_2.
- Bottesi G, Ghisi M, Altoè G, Conforti E, Melli G, Sica C. The Italian version of the depression anxiety stress Scales-21: factor structure and psychometric properties on community and clinical samples. *Compr Psychiatry* 2015; **60**: 170–181. <https://doi.org/10.1016/J.COMPSPSYCH.2015.04.005>.
- The Social Psychology of Health. Open Library, Accessed November 6, 2022. https://openlibrary.org/books/OL2404670M/The_social_psychology_of_health.
- Carver CS. You want to measure coping but your protocol's too long: consider the brief COPE. *Int J Behav Med* 1997; **4**: 92–100. https://doi.org/10.1207/S15327558IJBMO401_6.
- Savla J, Roberto KA, Blieszner R, McCann BR, Hoyt E, Knight AL. Dementia caregiving during the “stay-at-home” phase of COVID-19 pandemic. *J Gerontol B Psychol Sci Soc Sci* 2021; **76**: E241–E245. <https://doi.org/10.1093/GERONB/GBAA129>.
- Novak M, Guest C. Application of a multidimensional caregiver burden inventory. *Gerontologist* 1989; **29**: 798–803. <https://doi.org/10.1093/GERONT/29.6.798>.
- Morley JE, Malmstrom TK, Miller DK. A simple frailty questionnaire (FRAIL) predicts outcomes in middle aged African Americans. *J Nutr Health Aging* 2012; **16**: 601–608. <https://doi.org/10.1007/S12603-012-0084-2>.
- Cohen SA, Kunicki ZJ, Drohan MM, Greaney ML. Exploring changes in caregiver burden and caregiving intensity due to COVID-19. *Gerontol Geriatr Med* 2021; **7**: 7. <https://doi.org/10.1177/2333721421999279>.
- Canevelli M, Valletta M, Toccaceli Blasi M *et al*. Facing dementia during the COVID-19 outbreak. *J Am Geriatr Soc* 2020; **68**: 1673–1676. <https://doi.org/10.1111/JGS.16644>.
- Park SS. Caregivers' mental health and somatic symptoms during COVID-19. *J Gerontol B Psychol Sci Soc Sci* 2021; **76**: E235–E240. <https://doi.org/10.1093/GERONB/GBAA121>.
- Koyama Y, Nawa N, Yamaoka Y *et al*. Interplay between social isolation and loneliness and chronic systemic inflammation

- during the COVID-19 pandemic in Japan: results from U-CORONA study. *Brain Behav Immun* 2021; **94**: 51–59. <https://doi.org/10.1016/J.BBI.2021.03.007>.
- 34 Lang PJ, Bradley MM, Cuthbert BN. Emotion, motivation, and anxiety: brain mechanisms and psychophysiology. *Biol Psychiatry* 1998; **44**: 1248–1263.
 - 35 Okun MA, Keith VM. Effects of positive and negative social exchanges with various sources on depressive symptoms in younger and older adults. *Journals Gerontol-Ser B Psychol Sci Soc Sci* 1998; **53**: P4–P20. <https://doi.org/10.1093/GERONB/53B.1.P4>.
 - 36 Smith TO, Belderson P, Dainty JR et al. Impact of COVID-19 pandemic social restriction measures on people with rheumatic and musculoskeletal diseases in the UK: a mixed-methods study. *BMJ Open* 2021; **11**: 1–12. <https://doi.org/10.1136/bmjopen-2021-048772>.
 - 37 Sardella A, Chiara E, Alibrandi A et al. Changes in cognitive and functional status and in quality of life of older outpatients during the COVID-19 pandemic. *Gerontology* 2022; **68**: 1285–1290. <https://doi.org/10.1159/000525041>.
 - 38 Ismail IL, Kamel WA, Al-Hashel JY. Association of COVID-19 pandemic and rate of cognitive decline in patients with dementia and mild cognitive impairment: a cross-sectional study. *Gerontol Geriatr Med* 2021; **7**: 23337214211005224. <https://doi.org/10.1177/23337214211005223>.
 - 39 Chen Z-C, Liu S, Gan J et al. The impact of the COVID-19 pandemic and lockdown on mild cognitive impairment, Alzheimer's disease and dementia with Lewy bodies in China: a 1-year follow-up study. *Front Psych* 2021; **12**: 711658. <https://doi.org/10.3389/fpsy.2021.711658>.
 - 40 Sugimoto T, Ono R, Kimura A et al. Physical frailty correlates with behavioral and psychological symptoms of dementia and caregiver burden in Alzheimer's disease. *J Clin Psychiatry* 2018; **79**: 8633. <https://doi.org/10.4088/JCP.17M11991>.
 - 41 Bergh S, Engedal K, Røen I, Selbæk G. The course of neuropsychiatric symptoms in patients with dementia in Norwegian nursing homes. *Int Psychogeriatr* 2011; **23**: 1231–1239. <https://doi.org/10.1017/S1041610211001177>.
 - 42 Eikelboom WS, van den Berg E, Singleton EH et al. Neuropsychiatric and cognitive symptoms across the Alzheimer disease clinical spectrum: cross-sectional and longitudinal associations. *Neurology* 2021; **97**: e1276–e1287. <https://doi.org/10.1212/WNL.00000000000012598>.
 - 43 Vik-Mo AO, Gjøll LM, Borda MG, Ballard C, Aarsland D. The individual course of neuropsychiatric symptoms in people with Alzheimer's and Lewy body dementia: 12-year longitudinal cohort study. *Br J Psychiatry* 2020; **216**: 43–48. <https://doi.org/10.1192/bjp.2019.195>.
 - 44 Manini A, Brambilla M, Maggiore L, Pomati S, Pantoni L. The impact of lockdown during SARS-CoV-2 outbreak on behavioral and psychological symptoms of dementia. *Neurol Sci* 2021; **42**: 825–833. <https://doi.org/10.1007/S10072-020-05035-8>.
 - 45 Fried LP, Tangen CM, Walston J et al. Frailty in older adults: evidence for a phenotype. *J Gerontol Med Sci Copyr* 2001; **56**: 146–156. Accessed November 2, 2022. <https://academic.oup.com/biomedgerontology/article/56/3/M146/545770>.
 - 46 Scriven H, Doherty DP, Ward EC. Evaluation of a multisite telehealth group model for persistent pain management for rural/remote participants. *Rural Remote Health* 2019; **19**(1): 4710. <https://doi.org/10.22605/RRH4710>.