



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Synthesis of nitroarenes and azoxyarenes through the selenium-mediated on water oxidation of aryl amines

Questa è la versione Preprint (Submitted version) della seguente pubblicazione:

Original Citation:

Synthesis of nitroarenes and azoxyarenes through the selenium-mediated on water oxidation of aryl amines / Capperucci, A., Dalia, C., Cenni, A., Tanini, D.. - In: PHOSPHORUS SULFUR AND SILICON AND THE RELATED ELEMENTS. - ISSN 1042-6507. - ELETTRONICO. - 198:(2023), pp. 526-531. [10.1080/10426507.2023.2166044]

Availability:

The webpage <https://hdl.handle.net/2158/1353875> of the repository was last updated on 2024-03-25T17:32:26Z

Published version:

DOI: 10.1080/10426507.2023.2166044

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

Conformità alle politiche dell'editore / Compliance to publisher's policies

Questa versione della pubblicazione è conforme a quanto richiesto dalle politiche dell'editore in materia di copyright.

This version of the publication conforms to the publisher's copyright policies.

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

Synthesis of nitroarenes and azoxyarenes through the selenium-mediated *on water* oxidation of aryl amines


Antonella Capperucci, Camilla Dalia, Alessio Cenni, Damiano Tanini*

Department of Chemistry "Ugo Schiff", University of Florence, Via Della Lastruccia 3-13, Sesto Fiorentino, Firenze, Italy. E-mail: damiano.tanini@unifi.it

Abstract

Selenium-promoted oxidations enable a broad range of valuable oxidative functional group interconversions. Anilines are oxidised with hydrogen peroxide in water in the presence of selenium catalysts. The selenium catalyst determines the nature of the oxidation product; whereas nitroarenes are obtained employing diphenyl diselenide or benzeneseleninic acid, azoxyarenes are formed using selenium dioxide or sodium seleninate. The reaction mechanism relies on the reactivity of Se(IV) active oxidants, while Se(VI) species have been demonstrated not to be involved in.

Graphical Abstract



Selenium-promoted oxidative functional group interconversions occupy a privileged position in organic chemistry.¹⁻⁴ Both organic and inorganic selenium species are employed as catalysts for these reactions. While diselenides, seleninic acids, anhydrides, as well as other derivatives are used as organoselenium catalysts,⁵ selenium(IV) oxide is arguably the most widely employed inorganic selenium species.⁶ Hydrogen peroxide and tertbutyl hydroperoxide (TBHP) are the oxidant of choice in selenium-mediated oxidations.

A wide variety of organoselenium-catalysed methodologies for the oxidation of a number of organic compounds, including alkenes,^{5,7,8} alkynes,⁵ alcohols,^{5,7-9} aldehydes, and ketones,⁵ have been developed. Several selenium-catalysed versions of the Baeyer-Villiger oxidation,¹⁰ as well as selenium-mediated α -oxyfunctionalization of alkenes¹¹ and ring contraction of cycloalkanones,^{12,13} have also been reported. Additionally, a range of valuable selenium(IV) oxide-mediated protocols for the allylic oxidation of alkenes, cycloalkenes and alkynes, as well as for the trans-dihydroxylation of olefins have emerged as powerful transformations.⁶

The synthetic potential of such selenium-mediated routes is showcased by their use in the total synthesis of a number of natural products. For example, the SeO_2 -mediated oxidation of the α -methylene group of a cyclohexanone derivative to the corresponding α -keto group is one of the steps involved in the total synthesis of 6-*epi*-(-)-hamigeran B.¹⁴ The organoselenium-catalysed oxidation of indolines to indoles has been applied to the final step of the total synthesis of ergot alkaloids, such as ($\pm$)-lysergol.¹⁵ The well-

known SeO_2 /TBHP-promoted allylic oxidation was successfully employed for the preparation of a key intermediate in the total synthesis of (–)-okilactomycin (Figure 1).¹⁶




Figure 1. Structures of selected natural products whose total synthesis involves selenium-mediated oxidation steps.

Besides the above-described transformations, the selenium-mediated thiol-disulfide oxidative interconversion has also received considerable attention because of its importance in biology. Indeed, the oxidation of glutathione (GSH) to the corresponding disulfide (GSSG) using peroxides (*i.e.*, H_2O_2), which in turn is reduced to safe species (*i.e.*, H_2O), plays a key role in the detoxification of cells from reactive oxygen species (ROS).¹⁷⁻²⁰ This simple but essential reaction is catalysed by glutathione peroxidases (GPxs), which are selenoenzymes and feature a selenocysteine residue in their catalytic pocket. Considering the potential importance of this transformation, a steadily growing number of selenium-containing small molecules capable to catalyse the oxidation of thiols to disulfides have been reported over the past decades.²¹⁻³³


Oxidative transformations of alkenes and amines are broadly employed in the synthesis of biologically important small molecules. However, while the selenium-mediated epoxidation and dihydroxylation of alkenes have been widely explored, the oxidation of amines has received far less attention. Indeed, while a very few methodologies for the conversion of anilines to nitrosoarenes³⁴ or azoxyarenes³⁵ have been described, a selenium-based route for the oxidation of anilines to nitroarenes has been only recently developed by our group.³⁶ Nitro derivatives occupy an important position in organic synthesis³⁷ and are employed as versatile building blocks for the preparation of pharmaceutical,^{38,39} dye,^{40,41} materials,⁴² and perfumes.⁴³ In some cases, the synthesis of nitroarenes *via* oxidation of anilines represents an attractive alternative to the variety of available nitration methodologies.⁴⁴⁻⁴⁷

During our studies in the chemistry of chalcogens, we became interested in the development of selenium-catalysed methodologies for the oxidation of sulfur- and nitrogen-containing functional groups.^{26,28,29,31,36,48}

Very recently, we focused our attention on anilines and envisaged the possibility to develop a sustainable route for their conversion to nitroarenes by exploiting the redox chemistry of organoselenium compounds.³⁶

During our initial optimisation studies, a series of selenium-containing compounds at different oxidation state were evaluated as catalysts in the oxidation of aniline with hydrogen peroxide. Dialkyl- and diaryl-selenides and diselenides, as well as benzeneseleninic acid were employed. The study was also enlarged to include inorganic selenium species, such as selenium dioxide (SeO_2), sodium seleninate (Na_2SeO_3), and potassium selenonate (K_2SeO_4). This screening led us to find that both diphenyl diselenide (PhSeSePh) and benzeneseleninic acid (PhSeOOH) were effective in promoting the oxidation of aniline to nitrobenzene **1a** with high selectivity (Scheme 1, *reaction a*). Notably, the selectivity of the reaction could be controlled by simply switching the nature of the catalyst; indeed, azoxyarenes **2a** were formed upon using selenium(IV) oxide or sodium seleninate (Scheme 1, *reaction b*). Besides selenium-based catalysts, other chalcogen-


containing compounds – including disulfides, tellurides and ditellurides – were evaluated and found to be ineffective in promoting the oxidation herein described.



Scheme 1. selenium-promoted synthesis of nitroarenes and azoxyarenes.

As reported in the Scheme 1, the use of H_2O as the solvent and the possibility to efficiently conduct the reaction at ambient temperature, represent additional attractive features of these selenium-promoted oxidation routes.


The substrate scope of both these processes was wide and a range of nitroarenes **1** and azoxyarenes **2** were easily prepared in high yields under the optimised reaction conditions. Anilines bearing both electron-donating and electron-withdrawing substituents were amenable to these Se-mediated methodologies and were efficiently converted into the corresponding substituted nitro- and azoxy- derivatives **1** and **2**. (Schemes 2 and 3). Further functionalisable halo-substituted derivatives were also achieved.




Scheme 2. Se-mediated oxidation of anilines to the corresponding nitroarenes (selected examples). Isolated yields are given. ^aObtained from 4-nitroaniline. ^bThe reaction mixture was warmed at 60 °C.

The reaction mechanism of selenium-mediated oxidation reactions generally proceeds through Se(IV) species (*i.e.*, benzeneseleninic acid and peroxybenzeneseleninic acid). However, the involvement of Se(VI) species (*i.e.*, benzeneselenonic acid and peroxybenzeneselenonic acid) in the epoxidation of alkenes has

been recently demonstrated.⁴⁹ Some Se(IV) and Se(VI) species involved in selenium-promoted oxidation reactions, along with their interconversion equilibria, are reported in the Scheme 4.



Scheme 3. Se-mediated oxidation of anilines to the corresponding azoxyarenes (selected examples). Isolated yields are given.




Scheme 4. Some Se(IV) and Se(VI) species involved in selenium-promoted oxidation processes.


Our studies suggested that while Se(IV) species play a key role in the mechanism of the *on water* selenium-mediated oxidation of anilines to nitroarenes, Se(VI) species do not behave as effective oxidants in this transformation. Indeed, benzeneselenonic acid **6** – easily prepared upon treatment of benzeneselenenyl chloride with hydrogen peroxide⁵⁰ (Scheme 5, *reaction a*) – does not promote the oxidation of aniline to nitrobenzene using hydrogen peroxide (Scheme 5, *reaction c*). On the other hand, peroxybenzeneseleninic acid **5** – obtained according to the literature⁵¹ *via* oxidation of benzeneselenenic acid with H_2O_2 (Scheme 5, *reaction b*) – quickly oxidises aniline to provide nitrobenzene (Scheme 5, *reaction d*), thus demonstrating the key role of Se(IV) oxidants in this reaction mechanism.

Spurred by these observations, we proposed the mechanism reported in the Scheme 6. Oxidation of diphenyl diselenide with hydrogen peroxide provides benzeneselenenic acid **9**, which is further oxidised to give benzeneseleninic acid **3**.⁵² Oxidation of **3** with hydrogen peroxide affords the peroxyselenurane **4**,

which can undergo dehydration to afford peroxybenzeneseleninic acid **5** or benzeneselenonic acid **6**. The latter has been demonstrated ineffective in promoting the conversion of aniline to nitrobenzene using hydrogen peroxide as the oxidant and water as the solvent (Scheme 6, *part a*). Conversely, **5** proved to effectively oxidise aniline to nitrobenzene; therefore, the proposed mechanism relies on the reactivity of peroxybenzeneseleninic acid **5** as the oxidant (Scheme 6, *part b*). Reaction of **5** with anilines reasonably provides the hydroxylamine **10**, which is further oxidised by **5** to afford the dihydroxylamine derivative **11**. In both these reactions, benzeneseleninic acid **3** is also formed. The active oxidant **5** is regenerated upon oxidation of **3** with hydrogen peroxide. The dihydroxylamine **11** probably undergoes an acid-promoted water elimination to provide the corresponding nitroso derivative **12**. Both benzeneseleninic acid and benzeneselenonic acid can promote the conversion of **11** to **12**. A final oxidation step is required to oxidise the nitrosoarene **12** to the nitroarene **1**.



Scheme 5. Synthesis of benzeneselenonic acid **6** and peroxybenzeneseleninic acid **5** and their reactivity in the oxidation of aniline.



Scheme 6. Selenium species formed upon oxidation of diphenyl diselenide (*Part a*). Proposed mechanism for the oxidation of anilines to nitroarenes and selenium species involved (*Part b*).

Notably, these results highlight that the mechanism operating for the selenium-mediated oxidation of anilines to nitroarenes proceeds through different species with respect to that involved in the mechanism of the epoxidation of alkenes. Particularly, while Se(VI) species (*i.e.*, selenonic acid, selenonium benzeneselenonate) have been showed to behave as catalysts in the epoxidation of olefins,⁴⁹ we recently demonstrated that such compounds do not play a catalytic role in the *on water* oxidation of anilines to nitroarenes with H₂O₂,³⁶ which plausibly relies on Se(IV) species (*i.e.*, benzeneseleninic acid). The active oxidant involved in the oxidation of anilines is reasonably peroxybenzeneseleninic acid **5** (or the peroxyselecurane derivative **4**). The above-described findings suggest that the active oxidants and the catalytic pathway of selenium-promoted oxidative interconversions might depend on the nature of the substrate and on the reaction conditions, thus opening new avenues for the development of selective oxidation protocols.

References.

1. Thurow, S.; Abenante, L.; Anghinoni, J. M.; Lenardão, E. J. Selenium as a Versatile Reagent in Organic Synthesis: More than Allylic Oxidation. *Curr. Org. Synth.* **2021**, *19*, 331-365.
2. Lenardão, E.J.; Santi, C.; Sancineto, L. *New Frontiers in Organoselenium Compounds*; Springer: Gewerbestrasse, Switzerland, 2018.
3. Singh, F.V.; Wirth, T. *Organoselenium Chemistry. Synthesis and Reactions*; Wirth, T. Ed.; Wiley-VCH: Weinheim, Germany, 2012; Chapter 8.
4. Rathore, V.; Jose, C.; Kumar, S. Organoselenium small molecules as catalysts for the oxidative functionalization of organic molecules. *New J. Chem.* **2019**, *43*, 8852–8864.
5. Młochowski, J.; Brząszcz, M.; Giurg, M.; Palus, J.; Wójtowicz, H. Selenium-promoted oxidation of organic compounds: Reactions and mechanisms. *Eur. J. Org. Chem.* **2003**, 4329–4339.
6. Młochowski, J.; Wójtowicz-Młochowska, H. Developments in synthetic application of selenium(IV) oxide and organoselenium compounds as oxygen donors and oxygen-transfer agents. *Molecules* **2015**, *20*, 10205–10243.
7. Yu, L.; Wang, J.; Chen, T.; Wang, Y.; Xu, Q. Recyclable 1,2-bis[3,5-bis(trifluoromethyl)phenyl]diselane-catalyzed oxidation of cyclohexene with H₂O₂: A practical access to *trans*-1,2-cyclohexanediol. *Appl. Organomet. Chem.* **2014**, *28*, 652–656.
8. Jensen, H.P.; Sharpless, K.B. Selenium dioxide oxidation of d-limonene. Reinvestigation. *J. Org. Chem.* **1975**, *40*, 264–265.
9. Back, T.G. Oxidation of azasteroid lactams and alcohols with benzeneseleninic anhydride. *J. Org. Chem.* **1981**, *46*, 1442–1446.
10. ten Brink, G.-J.; Martijn, J.-M.; Vis, J.-H.; Arends, I. W. C.; Sheldon, R. A. Selenium-Catalyzed Oxidations with Aqueous Hydrogen Peroxide. 2. Baeyer–Villiger Reactions in Homogeneous Solution. *J. Org. Chem.*, **2001**, *66*, 2429-2433.

11. Barton, D. H.; Wang, T.-L. Novel allylic oxidation reagents. *Tetrahedron Lett.*, **1994**, *35*, 5149-5152.
12. Giurg, M.; Młochowski, J. Oxidative Ring Contraction of Cycloalkanones: A Facile Method for Synthesis of Medium Ring Cycloalkanecarboxylic Acids. *Synth. Commun.*, **1999**, *29*, 2281-2291.
13. Karimi, S.; Ma, S.; Ramig, K.; Greer, E. M.; Szalda, D. J.; Subramaniam, G. Oxidative ring-contraction of 3*H*-1-benzazepines to quinoline derivatives. *Tetrahedron Lett.*, **2015**, *56*, 6886-6889.
14. Mehta, G.; Shinde, H.M. Enantiospecific total synthesis of 6-epi(-)-hamigeran B. Intramolecular Heck reaction in a sterically constrained environment. *Tetrahedron Lett.* **2003**, *44*, 7049–7053
15. Ninomiya, I.; Hashimoto, C.; Kiguchi, T.; Naito, T.; Barton, D.H.R.; Luchinchi, X.; Milliet, P. Dehydrogenation with Benzeneseleninic Anhydride in Total Synthesis of Ergot Alkaloids. *J. Chem. Soc. Perkin. Trans. 1*, **1990**, 707–713.
16. Smith III, A. B.; Bosanac, T.; Basu, K. Evolution of the Total Synthesis of (-)-Okilactomycin Exploiting a Tandem Oxy-Cope Rearrangement/Oxidation, a Petasis-Ferrier Union/Rearrangement, and Ring-Closing Metathesis. *J. Am. Chem. Soc.* **2009**, *131*, 2348–2358.
17. Flohé, L.; Brigelius-Flohé, R. Selenoproteins of the Glutathione Peroxidase Family. In *Selenium. Its Molecular Biology and Role in Human Health*, 3rd ed.; Hatfield, D.L., Berry, M.J., Gladyshev, V.N., Eds.; Springer: London, UK, 2012; pp. 167–180.
18. Brigelius-Flohé, R.; Flohé, L. Regulatory Phenomena in the Glutathione Peroxidase Superfamily. *Antioxid. Redox Signal.* **2020**, *33*, 498–516.
19. Flohé, L. Looking Back at the Early Stages of Redox Biology. *Antioxidants* **2020**, *9*, 1254.
20. Barchielli, G.; Capperucci, A.; Tanini, D. The Role of Selenium in Pathologies: An Updated Review. *Antioxidants* **2022**, *11*, 251.
21. Bhabak, K. P.; Mughesh, G. Functional Mimics of Glutathione Peroxidase: Bioinspired Synthetic Antioxidants. *Acc. Chem. Res.*, **2010**, *43*, 1408-1419.
22. Pacula, A. J.; Mangiavacchi, F.; Sancineto, L.; Lenardao, E. J.; Scianowski, J.; Santi, C. An Update on “Selenium Containing Compounds from Poison to Drug Candidates: A Review on the GPx-like Activity”. *Curr. Chem. Biol.*, **2015**, *9*, 97-112.
23. Singh, F. V.; Wirth, T. in *Organoselenium Compounds in Biology and Medicine: Synthesis, Biological and Therapeutic Treatments*, ed. V. K. Jain and K. I. Priyadarsini, Royal Society of Chemistry, London, 2018, Ch. 3.
24. Capperucci, A.; Coronello, M.; Salvini, F.; Tanini, D.; Dei, S.; Teodori, E.; Giovannelli, L. Synthesis of functionalised organochalcogenides and in vitro evaluation of their antioxidant activity. *Bioorg. Chem.* **2021**, *110*, 104812.
25. Yoshida, S.; Kumakura, F.; Komatsu, I.; Arai, K.; Onuma, Y.; Hojo, H.; Singh, B. G.; Priyadarsini, K. I.; Iwaoka, M. Antioxidative Glutathione Peroxidase Activity of Selenogluthathione. *Angew. Chem., Int. Ed.*, **2011**, *50*, 2125-2128.

26. Capperucci, A.; Petrucci, A.; Faggi, C.; Tanini, D. Click reaction of selenols with isocyanates: Rapid access to selenocarbamates as peroxide-switchable reservoir of thiol-peroxidase-like catalysts. *Adv. Synth. Catal.* **2021**, *363*, 4256–4263.
27. Arai, K.; Sato, Y.; Nakajima, I.; Saito, M.; Sasaki, M.; Kanamori, A.; Iwaoka, M. Glutathione peroxidase-like functions of 1, 2-diselenane-4, 5-diol and its amphiphilic derivatives: Switchable catalytic cycles depending on peroxide substrates. *Bioorg. Med. Chem.* **2021**, *29*, 115866.
28. Tanini, D.; Capperucci, A. Synthesis and applications of organic selenols. *Adv. Synth. Catal.* **2021**, *363*, 5360–5385.
29. Tanini, D.; Lupori, B.; Malevolti, G.; Ambrosi, M.; Lo Nostro, P.; Capperucci, A. Direct biocatalysed synthesis of first sulfur-, selenium- and tellurium- containing L-ascorbyl hybrid derivatives with radical trapping and GPx-like properties. *Chem. Commun.* **2019**, *55*, 5705–5708.
30. Sands, K.N.; Tuck, T.A.; Back, T.G. Cyclic Seleninate esters, spirodioxyselenuranes and related compounds: New classes of biological antioxidants that emulate glutathione peroxidase. *Chem. Eur. J.* **2018**, *24*, 9714–9728.
31. Tanini, D.; Grechi, A.; Ricci, L.; Dei, S.; Teodori, E.; Capperucci, A. Novel functionalized organotellurides with enhanced thiol peroxidase catalytic activity. *New. J. Chem.* **2018**, *42*, 6077–6083.
32. Kumakura, F.; Mishra, B.; Priyadarsini, K.I.; Iwaoka, M. A watersoluble cyclic selenide with enhanced glutathione peroxidase-like catalytic activities. *Eur. J. Org. Chem.* **2010**, *3*, 440–445.
33. Mugesh, G.; Singh, H.B. Synthetic organoselenium compounds as antioxidants: Glutathione peroxidase activity. *Chem. Soc. Rev.* **2000**, *29*, 347–357.
34. Gebhardt, C.; Priewisch, G. B.; Irran, E.; Rück-Braun, K. Oxidation of Anilines with Hydrogen Peroxide and Selenium Dioxide as Catalyst. *Synthesis*, **2008**, 1889–1894.
35. Zhao, D.; Johansson, M.; Bäckvall, J.-E. In Situ Generation of Nitroso Compounds from Catalytic Hydrogen Peroxide Oxidation of Primary Aromatic Amines and Their One-Pot Use in Hetero-Diels–Alder Reactions. *Eur. J. Org. Chem.*, **2007**, 4431–4436.
36. Tanini, D.; Dalia, C.; Capperucci, A. The polyhedral nature of selenium-catalysed reactions: Se(IV) species instead of Se(VI) species make the difference in the on water selenium-mediated oxidation of arylamines. *Green Chem.*, **2021**, *23*, 5680–5686.
37. Ono, N. *The Nitro Group in Organic Synthesis*; Wiley-VCH: New York, NY, USA, 2001.
38. Muller, W.E. *The Benzodiazepine Receptor*; Cambridge University Press: New York, NY, USA, 1988.
39. Belciug, M.; Ananthanarayanan, V.S. Interaction of calcium channel antagonists with calcium: Structural studies on nifedipine and its Ca²⁺ complex. *J. Med. Chem.* **1994**, *37*, 4392–4399.
40. Zollinger, H. *Color Chemistry*; Wiley-VCH: New York, NY, USA, 1987.
41. Koh, J.; Kim, S.; Kim, J.P. Synthesis and spectral properties of azohydroxypyridone disperse dyes containing a fluorosulphonyl group. *Color. Technol.* **2004**, *120*, 241–246.

42. Fan, F.-R.F.; Yao, Y.; Cai, L.; Cheng, L.; Tour, J.M.; Bard, A.J. Structure-dependent charge transport and storage in self-assembled monolayers of compounds of interest in molecular electronics: Effects of tip material, headgroup, and surface concentration. *J. Am. Chem. Soc.* **2004**, *126*, 4035–4042.
43. Luckenbach, T.; Epel, D. Nitromusk and polycyclic musk compounds as long-term inhibitors of cellular xenobiotic defense systems mediated by multidrug transporters. *Environ. Health Perspect.* **2005**, *113*, 17–24.
44. Yan, G.; Yang, M. Recent advances in the synthesis of aromatic nitro compounds. *Org. Biomol. Chem.* **2013**, *11*, 2554–2566.
45. Weissmehl, K.; Arpe, H.-J. *Industrial Organic Chemistry*; Wiley-VCH: Weinheim, Germany, 2008.
46. Prakash, G.K.S.; Mathew, T. *ipso*-Nitration of arenes. *Angew. Chem. Int. Ed.* **2010**, *49*, 1726–1728.
47. Capperucci, A.; Tanini, D. Synthesis of Nitroarenes by Oxidation of Aryl Amines. *Chemistry*, **2022**, *4*, 77–97.
48. Młostoń, G.; Capperucci, A.; Tanini, D.; Hamera-Fałdyga, R.; Heimgartner, H. Dialkyl Dicyanofumarates as Oxidizing Reagents for the Conversion of Thiols into Disulfides and Selenols into Diselenides. *Eur. J. Org. Chem.*, 2017, 6831–6839.
49. Sands, K. N.; Rengifo, E. M.; George, G. N.; Pickering, I. J.; Gelfand, B. S.; Back, T. G. The Unexpected Role of Se^{VI} Species in Epoxidations with Benzeneseleninic Acid and Hydrogen Peroxide. *Angew. Chem. Int. Ed.*, **2020**, *59*, 4283–4287.
50. Wei, D.; Han, M.; Yu, L. Solid-State ⁷⁷Se NMR of Organoselenium Compounds through Cross Polarization Magic Angle Spinning (CPMAS) Method. *Sci. Rep.* **2017**, *7*, 6376.
51. Syper, L.; Młochowski, J. Benzeneperoxseleninic acids-synthesis and properties. *Tetrahedron*, **1987**, *43*, 207–213.
52. Ribaud, G.; Bellanda, M.; Menegazzo, I.; Wolters, L. P.; Bortoli, M.; Ferrer-Sueta, G.; Zagotto, G.; Orian, L. Mechanistic Insight into the Oxidation of Organic Phenylselenides by H₂O₂. *Chem. Eur. J.*, **2017**, *23*, 2405–2422.