



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

On the strong maximum principle for fully nonlinear parabolic equations of second order

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

On the strong maximum principle for fully nonlinear parabolic equations of second order / Goffi, A.. - In: PROCEEDINGS OF THE AMERICAN MATHEMATICAL SOCIETY. - ISSN 1088-6826. - ELETTRONICO. - 153:(2025), pp. 1575-1583. [10.1090/proc/17050]

Availability:

The webpage <https://hdl.handle.net/2158/1401918> of the repository was last updated on 2025-02-21T13:10:40Z

Published version:

DOI: 10.1090/proc/17050

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

ON THE STRONG MAXIMUM PRINCIPLE FOR FULLY NONLINEAR PARABOLIC EQUATIONS OF SECOND ORDER

ABSTRACT. We provide a proof of strong maximum and minimum principles for fully nonlinear uniformly parabolic equations of second order. The approach is of parabolic nature and does not exploit the parabolic Harnack inequality.

1. INTRODUCTION

In this note we prove a strong maximum principle for fully nonlinear uniformly parabolic equations of the form

$$(1) \quad F(x, t, u, Du, D^2u) - \partial_t u = 0 \text{ in } D \subset \mathbb{R}^{n+1},$$

where $F : D \times \mathbb{R} \times \mathbb{R}^n \times \mathcal{S}_n \rightarrow \mathbb{R}$, $\mathcal{S}_n$ being the space of $n \times n$ symmetric matrices, is assumed to satisfy the following structure condition: there exist $b \geq 0$, $c \leq 0$ continuous such that for $M, N \in \mathcal{S}_n$, $N \geq 0$, and every $p, q \in \mathbb{R}^n$, $(x, t) \in D$ we have

$$(2) \quad \lambda \|N\| - b|p - q| + c(r - s) \leq F(x, t, r, p, M + N) - F(x, t, s, q, N) \leq \Lambda \|N\| + b|p - q| + c(r - s)$$

and $F(x, t, 0, 0, 0) = 0$ on D , $\|N\| = \text{Tr}(N)$.

By strong maximum (resp. minimum) principle for an evolution operator $\mathcal{P}u = 0$ in $\Omega_T := \Omega \times (0, T)$ we mean the following property:

Any continuous viscosity subsolution (resp. supersolution) to $\mathcal{P}u = 0$ in Ω_T that attains a nonnegative maximum (minimum) at $(x_0, t_0) \in \Omega_T$ is constant in $\overline{\Omega} \times (0, t_0)$.

L. Nirenberg [17], see also [9], proved the strong maximum principle for classical solutions of linear parabolic equations, extending the proof due to E. Hopf [18] for linear elliptic equations. This property was later investigated in the framework of fully nonlinear parabolic equations in [5, 8]. The known proofs require to show a vertical propagation of maximum points, which in turn use the elliptic propagation on suitable small rectangles, see e.g. [9]. As for the strong maximum principle for viscosity solutions of linear and nonlinear, even degenerate, elliptic equations we refer to [2, 3, 6, 10, 11, 18] and the references therein. We mention that F. Da

Date: August 18, 2024.

2020 Mathematics Subject Classification. 35B50, 35D40, 35K10.

Key words and phrases. Strong maximum principle, Strong comparison principle, Fully nonlinear equation, Bellman parabolic equation, Isaacs parabolic equation, Uniqueness of solutions.

The author is member of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). He was partially supported by the INdAM-GNAMPA Projects 2023 and 2024, by the King Abdullah University of Science and Technology (KAUST) project CRG2021-4674 “Mean-Field Games: models, theory and computational aspects” and by the project funded by the European Union – NextGenerationEU under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.1 - Call PRIN 2022 No. 104 of February 2, 2022 of Italian Ministry of University and Research; Project 2022W58BJ5 (subject area: PE - Physical Sciences and Engineering) “PDEs and optimal control methods in mean field games, population dynamics and multi-agent models”.

Lio [8] proved the result for rather more general equations than (1), even degenerate, following the route of [9, 17] in the context of viscosity solutions. The paper [5] used the argument in [17], proving the result when $F \in C^1$ in all arguments in the form of a strong comparison principle: in this case F was not assumed uniformly parabolic and the derivatives $|F_p|, |F_r|$ are not necessarily uniformly bounded.

Our proof follows the approach of [12, 15] for classical solutions of linear parabolic, nondivergence structure, equations and slightly differs from that of L. Nirenberg [17], L. Caffarelli-Y. Li-L. Nirenberg [5] and F. Da Lio [8]. The main difference with respect to the aforementioned works relies on the choice of a space-time barrier-like function, which results in a vertical propagation of maximum points on inclined cylinders whose axes are broken lines starting from the maximum points. We recall that other different proofs of the strong maximum principle for linear and nonlinear parabolic equations involve the weak Harnack inequality [13, 16], see also Proposition 4.9 in [4] for the elliptic case.

We emphasize that this strategy, at least in a nonlinear setting, allows to avoid different sets of assumptions on the nonlinearity to prove the horizontal and vertical propagation of maximum points. Moreover, since it does not make use of the parabolic Harnack inequality as in [4, 13], it could be particularly useful in those settings where the latter tool is not available in full generality. This is the case of fully nonlinear subelliptic equations [3] and those posed on Riemannian manifolds [10].

We conclude the paper with some remarks about the applications of the strong maximum principle to various qualitative properties of nonlinear evolution equations, such as the strong comparison principle for fully nonlinear parabolic equations, a Tychonoff uniqueness principle for Hessian evolution equations and a positivity result. Notably, the result also gives a proof of the strong maximum principle for fully nonlinear elliptic equations [2, 3, 6, 10] by means of a different method. Some new examples to which the results apply will be presented in the last section of the manuscript.

2. NOTATIONS AND PRELIMINARIES

Let $D \subset \mathbb{R}^{n+1}$ be a bounded domain and b, c, f be continuous and bounded functions in $D \subset \mathbb{R}^{n+1}$, $b \geq 0$, $c \leq 0$ and $\lambda \leq \Lambda$ be two positive constants (the ellipticity constants). We denote for $M \in \mathcal{S}_n$ by

$$\begin{aligned}\mathcal{M}_{\lambda, \Lambda}^+(M) &= \sup_{\lambda I_n \leq A \leq \Lambda I_n} \text{Tr}(AM) = \Lambda \sum_{e_i > 0} e_i(M) + \lambda \sum_{e_i < 0} e_i(M), \\ \mathcal{M}_{\lambda, \Lambda}^-(M) &= \inf_{\lambda I_n \leq A \leq \Lambda I_n} \text{Tr}(AM) = \lambda \sum_{e_i > 0} e_i(M) + \Lambda \sum_{e_i < 0} e_i(M),\end{aligned}$$

e_i being the eigenvalues of M , the Pucci's extremal operators. We denote by $\underline{S}(\lambda, \Lambda, b, c, f)$ the space of continuous functions $u \in C(D)$ such that

$$\mathcal{M}_{\lambda, \Lambda}^+(D^2u) + b(x, t)|Du| + c(x, t)u - \partial_t u \geq f(x, t)$$

in the viscosity sense in D . Similarly, we denote by $\overline{S}(\lambda, \Lambda, b, c, f)$ the space of continuous functions $u \in C(D)$ such that

$$\mathcal{M}_{\lambda, \Lambda}^-(D^2u) - b(x, t)|Du| + c(x, t)u - \partial_t u \leq f(x, t).$$

Note that if F in (1) satisfies (2) and $F(x, t, 0, 0, 0) = 0$, then it satisfies in the viscosity sense (3) $\mathcal{M}_{\lambda, \Lambda}^+(D^2u) + b(x, t)|Du| + c(x, t)u - \partial_t u \geq 0 \geq \mathcal{M}_{\lambda, \Lambda}^-(D^2u) - b(x, t)|Du| + c(x, t)u - \partial_t u$.

For more details on these classes and the proof of the results we refer to [4].

We follow the notation in [15]. We denote by $\mathbb{Q}_{x_0, R}^{t_1, t_2} := \{(x, t) \in \mathbb{R}^{n+1} : |x - x_0| < R, t_1 < t < t_2\}$ a cylinder in $\mathbb{R}^{n+1}$. Moreover, $S_{x_0, R}^{t_1, t_2}$ is the lateral (closed) surface, $Q_{x_0, R}^{t_1}$ is the lower (open) base (i.e. the n -dimensional open ball of radius R centered at (x_0, t_1) on the hyperplane $t = t_1$) and $Q_{x_0, R}^{t_2}$ is the upper (open) base of the cylinder $\mathbb{Q}_{x_0, R}^{t_1, t_2}$. Here, the t -axis is directed upward.

By the upper base of the domain D , denoted with $\gamma(D)$, we mean the set of those points $(x, t) \in \partial D$ satisfying the following property: for each point of $\gamma(D)$ there exists $h > 0$ such that the cylinder $\mathbb{Q}_{x, h}^{t-h, t}$ belongs to D , while the cylinder $\mathbb{Q}_{x, h}^{t, t+h}$ lies outside D . Let $(x_0, t_0) \in \overline{D}$. A set D' is called subdomain of D subordinate to (x_0, t_0) if (x_0, t_0) can be joined with every point of D' by a broken line that lies in D (except at most (x_0, t_0)), is uniquely projected to the t -axis (i.e. two points of the broken line are projected to different points of the t -axis), and has (x_0, t_0) as upper end-point.

3. MAIN RESULT

We consider the following equation

$$(4) \quad F(x, t, u, Du, D^2u) - \partial_t u = 0 \text{ in } D.$$

The next result is a fully nonlinear version of Theorem 2.3 in Chapter 3 of [15], cf. also Theorem 6 in [12], and applies to viscosity solutions. Note also that the next property holds for a general parabolic domain in $\mathbb{R}^{n+1}$ rather than cylinders of the form $\Omega_T := \Omega \times (0, T)$.

Theorem 3.1 (Strong maximum/minimum principles). *Let D be a domain in $\mathbb{R}^{n+1}$ with upper base $\gamma(D)$. Let u be a continuous viscosity subsolution (resp. supersolution) in $D \cup \gamma(D)$ to (4), with F satisfying (2), such that $c(x, t) \leq 0$, $b \geq 0$ and b, c are bounded, attaining a nonnegative maximum (nonpositive minimum) at some point $(x_0, t_0) \in D \cup \gamma(D)$. Then, u is constant in a subdomain D' of D subordinate to the point (x_0, t_0) .*

To prove the strong maximum principle we need some preliminary lemmas. The next is the crucial step towards the propagation of maximum/minimum points vertically in the parabolic cylinder.

Lemma 3.2. *Under the same assumptions of Theorem 3.1, let u be a continuous viscosity subsolution (resp. supersolution) of (4) in the cylinder $\mathbb{Q}_{x_0, R}^{t_1, t_2}$ and on the upper base $Q_{x_0, R}^{t_2}$. If u attains the nonnegative maximum (nonpositive minimum) M at (x_0, t_2) , then the maximum (minimum) propagates, namely $u = M$, on the axis of the cylinder.*

Proof. We prove the statement for viscosity subsolutions, the other being similar using that $-u \in \underline{\mathcal{S}}$. First, since F satisfies (3), it is enough to prove the claim for viscosity solutions belonging to $\underline{\mathcal{S}}$, i.e. to the viscosity inequality

$$\mathcal{M}_{\lambda, \Lambda}^+(D^2u(x, t)) + b(x, t)|Du| + c(x, t)u - \partial_t u \geq 0.$$

Suppose by contradiction that there exists a point (x_0, t') on the axis of the cylinder at which $u < M - \alpha$ for some $\alpha > 0$. By continuity, we can find r_0 small enough such that

$r_0 < \min\{R, 1\}$ and $u(x, t') < M - \alpha$ for $|x - x_0| < r_0$. In the inner closed cylinder $\mathbb{Q}_{x_0, r_0}^{t', t_2}$ we consider the function

$$v(x, t) = M - \alpha e^{-\beta(t-t')} [r_0^2 - |x - x_0|^2]^2.$$

We assume for the moment that there exists $\beta > 0$ such that v is a classical (strict) supersolution to

$$\mathcal{M}_{\lambda, \Lambda}^+(D^2 u(x, t)) + b(x, t)|Du| + c(x, t)u - \partial_t u = 0.$$

We first show how to obtain a contradiction and conclude the proof. We first observe that $u - v$ is nonpositive on the lower base and the lateral surface of the cylinder, since $v > M - \alpha$ on the lower base and equals M on the lateral surface $S_{x_0, R}^{t', t_2}$. By the comparison principle, see e.g. Theorem 8.2 in [7], $u \leq v$ in the cylinder, and we have that

$$M = u(x_0, t_2) \leq v(x_0, t_2) = M - \alpha r_0^2 e^{-\beta(t_2-t')} < M,$$

which gives the desired contradiction.

We are now left to prove the existence of β . We define $\varphi(x) = [r_0^2 - |x - x_0|^2]^2$. Then

$$\partial_{x_i} \varphi(x) = -4[r_0^2 - |x - x_0|^2](x_i - (x_0)_i),$$

$$D^2 \varphi(x) = 8|x - x_0|^2 \frac{x - x_0}{|x - x_0|} \otimes \frac{x - x_0}{|x - x_0|} - 4[r_0^2 - |x - x_0|^2]I_n,$$

where $(x \otimes x)_{ij} = x_i x_j$. We recall that the eigenvalues of a matrix M of the form

$$M = \nu I_n + \xi v \otimes v, \quad v \in \mathbb{R}^n, \quad |v| = 1, \quad \nu, \xi \in \mathbb{R},$$

are ν with multiplicity $n - 1$ and $\nu + \xi$. Therefore, the eigenvalues of $D^2 \varphi$ are $e_1(D^2 \varphi) = -4[r_0^2 - |x - x_0|^2]$ with multiplicity $n - 1$ and $e_2(D^2 \varphi) = 8|x - x_0|^2 - 4[r_0^2 - |x - x_0|^2] = 12|x - x_0|^2 - 4r_0^2$, which is simple. In particular, $e_1 \leq 0$. As for e_2 , note that when $|x - x_0|^2 \leq \frac{r_0^2}{3}$ we have $e_2 \leq 0$, hence

$$\begin{aligned} \mathcal{M}_{\lambda, \Lambda}^+(D^2 \varphi) &= \lambda\{-4(n-1)[r_0^2 - |x - x_0|^2] + 8|x - x_0|^2 - 4[r_0^2 - |x - x_0|^2]\} \\ &= \lambda\{-4n[r_0^2 - |x - x_0|^2] + 8|x - x_0|^2\}. \end{aligned}$$

Thus, by the properties of Pucci's extremal operators we get

$$\mathcal{M}_{\lambda, \Lambda}^+(D^2 v) = -\alpha e^{-\beta(t-t')} \{8\lambda|x - x_0|^2 - 4n\lambda[r_0^2 - |x - x_0|^2]\}.$$

If instead $|x - x_0|^2 \geq \frac{r_0^2}{3}$ (and $|x - x_0| < r_0$) we have $e_2 \geq 0$, and we conclude

$$\begin{aligned} \mathcal{M}_{\lambda, \Lambda}^+(D^2 \varphi) &= \lambda\{-4(n-1)[r_0^2 - |x - x_0|^2]\} + \Lambda\{8|x - x_0|^2 - 4[r_0^2 - |x - x_0|^2]\} \\ &= 8\Lambda|x - x_0|^2 - [4\lambda(n-1) + 4\Lambda]\{r_0^2 - |x - x_0|^2\}. \end{aligned}$$

Then

$$\mathcal{M}_{\lambda, \Lambda}^+(D^2 v) = -\alpha e^{-\beta(t-t')} \{8\Lambda|x - x_0|^2 - [4\lambda(n-1) + 4\Lambda](r_0^2 - |x - x_0|^2)\}.$$

In both cases we end up with

$$\mathcal{M}_{\lambda, \Lambda}^+(D^2 v) \leq -\alpha e^{-\beta(t-t')} \{8\lambda|x - x_0|^2 - c_{n, \lambda, \Lambda}(r_0^2 - |x - x_0|^2)\}$$

for $c_{n, \lambda, \Lambda} := 4\lambda(n-1) + 4\Lambda$. Since

$$\partial_t v = \beta \alpha e^{-\beta(t-t')} [r_0^2 - |x - x_0|^2]^2$$

we get, using $c \leq 0$,

$$\begin{aligned} & \mathcal{M}_{\lambda,\Lambda}^+(D^2v) + b(x,t)|Dv(x,t)| + c(x,t)v(x,t) - \partial_t v \\ & \leq Mc - \alpha e^{-\beta(t-t')} \{8\lambda|x-x_0|^2 - r(x,t)[r_0^2 - |x-x_0|^2] + \beta[r_0^2 - |x-x_0|^2]^2\} \\ & \leq -\alpha e^{-\beta(t-t')} \{8\lambda|x-x_0|^2 - K[r_0^2 - |x-x_0|^2] + \beta[r_0^2 - |x-x_0|^2]^2\} =: -\alpha e^{-\beta(t-t')} \Psi(|x-x_0|), \end{aligned}$$

where $r(x,t)$ is a bounded function, i.e. $|r(x,t)| \leq K$, for some $K > 0$ depending on n, λ, Λ and on the coefficients b, c , due to their boundedness assumptions. Setting $s := r_0^2 - |x-x_0|^2$ we have

$$\Psi(|x-x_0|) = \beta s^2 - (K + 8\lambda)s + 8\lambda r_0^2,$$

which is quadratic in s . Therefore, $\Psi > 0$ if $\beta > 0$ is chosen large enough, since $(K + 8\lambda)^2 - 32\beta\lambda r_0^2 < 0$. $\square$

We now prove that maximum/minimum points propagate along the axes of inclined cylinders.

Lemma 3.3. *Consider the inclined cylinder $D^I := \{(x,t) : |x - [x_0 + \eta(t - t_1)]| < R, t_1 < t \leq t_2\}$, $\eta \in \mathbb{R}^n$. Let u be a continuous viscosity subsolution (resp. supersolution) of (4) up to the upper base of D^I . If u attains the nonnegative maximum (nonpositive minimum) M at $(x_0 + \eta(t_2 - t_1), t_2)$, then $u = M$ at the points $x = x_0 + \eta(t - t_1)$, $t \in (t_1, t_2)$.*

Proof. It is enough to make the change of variables

$$\tilde{t} = t, \tilde{x} = x - \eta(t - t_1),$$

which leads to the viscosity inequality

$$\mathcal{M}_{\lambda,\Lambda}^+(\tilde{D}^2u) + b|\tilde{D}u| - \eta \cdot \tilde{D}u + cu - \partial_{\tilde{t}}u \geq 0,$$

where $\tilde{D}, \tilde{D}^2$ stands for the gradient and Hessian operators with respect to the variable $\tilde{x}$, which has the same form of the one treated in the previous Lemma 3.2 with the addition of a term involving a first-order power of the quantity $r_0^2 - |x-x_0|^2$. The presence of this new term does not affect the proof of Lemma 3.2. Then, the inclined cylinder becomes $t_1 < t < t_2$, $|\tilde{x} - x_0| < R$. We are thus in position to apply Lemma 3.2 from which $u \equiv M$ on $t_1 < t < t_2$ and $x = x_0 + \eta(t - t_1)$. $\square$

Proof of Theorem 3.1. It is only geometrical and the same of the linear case. If u attains its maximum at $(x_0, t_0) \in D \cup \gamma(D)$, assume that the point (x', t') can be joined to (x_0, t_0) by a broken line in $D \cup \gamma(D)$ having (x_0, t_0) as upper end-point. For every segment of the broken line one can construct a cylinder, which is in general inclined as in Lemma 3.3, such that its lower base is orthogonal to the t -axis, the broken line is the axis of the cylinder and it belongs to $D \cup \gamma(D)$. We then apply repeatedly Lemma 3.3 to these cylinders, starting from the upper one, to find $u \equiv M$ at every point of the broken line, and hence at (x', t') . $\square$

In the case of an equation without lower-order terms of the form

$$(5) \quad F(x, t, D^2u) - \partial_t u = 0 \text{ in } D \subset \mathbb{R}^{n+1},$$

in the statement of Theorem 3.1 we can drop the sign property of the maximum/minimum point by simply saying that it attains a maximum/minimum, as the operator does not change by the addition of a constant. The result then reads as follows:

Theorem 3.4. *Let D be a domain in $\mathbb{R}^{n+1}$ with upper base $\gamma(D)$. Let u be a continuous viscosity subsolution (resp. supersolution) in $D \cup \gamma(D)$ to (5), with F satisfying (2) and attaining a maximum (minimum) at some point $(x_0, t_0) \in D \cup \gamma(D)$. Then, u is constant in a subdomain D' of D subordinate to the point (x_0, t_0) .*

Some remarks are now in order:

Remark 3.5. The strong maximum and minimum principles involve only subdomains subordinate to the maximum/minimum point (x_0, t_0) and fail for larger subdomains. Moreover, the strong maximum/minimum principles hold only for boundary points belonging to the upper base, while for an arbitrary point it may fail. In this direction, we refer to [15] for some counterexamples in the linear case.

Remark 3.6. Theorem 3.1 implies the validity of strong maximum and minimum principles for the elliptic problem

$$F(x, u, Du, D^2u) = 0 \text{ in } \Omega,$$

where Ω is an open connected set in $\mathbb{R}^n$, namely any viscosity subsolution (resp. supersolution) attaining a nonnegative maximum (nonpositive minimum) in Ω is constant. It is sufficient to notice that any solution $u(x)$ of the elliptic equation solves the parabolic problem

$$F(x, u, Du, D^2u) - \partial_t u = 0 \text{ in } \Omega \times (0, 1).$$

Remark 3.7. Theorem 3.4 implies a strong comparison principle (also known as tangency theorem, cf. Chapter 2 of [18]) for the uniformly parabolic equation $F(D^2u) - \partial_t u = 0$. This property means that if $u - v \leq 0$ in Ω_T and there exists a point $z = (\tilde{x}, \tilde{t}) \in \Omega_T$ such that $(u - v)(z) = 0$, then $u - v = 0$ in $\Omega \times (0, \tilde{t}]$. An example for strictly parabolic equations can be found in Theorem 4.1 of [5]. This follows from the fact that $u - v$ solves a parabolic Pucci's extremal equation by means of Theorem 5.3 of [4].

Remark 3.8. The strong maximum principle also implies a Tychonoff uniqueness principle for exponentially growing solutions to the uniformly parabolic equation

$$F(x, t, D^2u) - \partial_t u = 0$$

by means of the three-curve property in [14]. Indeed, it is sufficient to note that if u, v are two continuous solutions of the above equation in the strip $\mathbb{R}^n \times (0, T)$ with $u(x, 0) = v(x, 0) = g(x)$ and satisfying the growth condition

$$|u|, |v| \leq c_1 e^{c_2 |x|^2}, \quad c_1, c_2 > 0$$

uniformly in $t \in (0, T)$, then $w = u - v$ solves

$$\mathcal{M}_{\lambda, \Lambda}^+(D^2w) - \partial_t w \geq 0,$$

by the results in [4]. Therefore, we can apply Theorem 1.1 in [14] and the argument of Theorem 1.3 therein along with the strong maximum principle to conclude $u \equiv v$ in $\mathbb{R}^n \times (0, T)$.

Remark 3.9. If the strong minimum principle holds for (4) in $\overline{D} = \overline{\Omega}_T$, then we have the following conservation of positivity result: if $u \in C(\overline{\Omega}_T)$ with $u(x, 0) = u_0(x) \geq 0$ and $u_0 \not\equiv 0$, then it is strictly positive in the whole cylinder.

4. EXAMPLES

Examples of operators to which the strong maximum and/or minimum principles proved in the previous section apply are those of the form

$$E(x, t, u, \partial_t u, Du, D^2 u) = a(x, t)G(Du, D^2 u) + b(x, t) \cdot Du + c(x, t)u - \partial_t u$$

when b, c are bounded, $c \leq 0$, $a > 0$ continuous and bounded. G can be of the following form:

- A Pucci's extremal operator or, more generally, a Bellman or an Isaacs operator defined respectively as

$$G(x, t, D^2 u) = \sup_{\alpha} \text{Tr}(A_{\alpha}(x, t)D^2 u), \quad G(D^2 u) = \inf_{\alpha} \text{Tr}(A_{\alpha}(x, t)D^2 u),$$

$$G(x, t, D^2 u) = \sup_{\beta} \inf_{\alpha} \text{Tr}(A_{\alpha, \beta}(x, t)D^2 u),$$

where the linear operators are uniformly parabolic. These evolution operators satisfy both the strong maximum and minimum principles, as already discussed in [8] using a different proof.

- The time-dependent normalized (1-homogeneous) p -Laplacian, $p > 1$, so that E corresponds to the evolution

$$|Du|^{2-p} \text{div}(|Du|^{p-2} Du) - \partial_t u = 0.$$

It is known that such operator belongs to the classes $\mathcal{S}$ defined above for $\lambda = \min\{1, p-1\}$, $\Lambda = \max\{1, p-1\}$, $b = c = 0$. We recall that the strong maximum and minimum principle fails for the standard parabolic p -Laplacian, cf. Example 2.6 in [8].

- The truncated (degenerate) Pucci's operator defined for $k < n$ by

$$\mathcal{M}_{\lambda, \Lambda; k}^{-}(D^2 u) = \lambda \sum_{\substack{i=0 \\ e_i > 0}}^k e_i + \Lambda \sum_{\substack{i=0 \\ e_i < 0}}^k e_i$$

satisfies the strong maximum principle by comparison with the Pucci's extremal operator $\mathcal{M}_{\lambda, \Lambda}^{-}$. Indeed, a subsolution to $\mathcal{M}_{\lambda, \Lambda; k}^{-}(D^2 u) - \partial_t u = 0$ is also a subsolution to $\mathcal{M}_{\lambda', \Lambda'}^{-}(D^2 u) - \partial_t u = 0$, where λ', Λ' are possibly different ellipticity constants, via the inequality

$$\mathcal{M}_{\frac{n}{k}\lambda, \frac{n}{k}\Lambda; k}^{-}(X) \leq \mathcal{M}_{\lambda, \Lambda; n}^{-}(X) = \mathcal{M}_{\lambda, \Lambda}^{-}(X), \quad X \in \mathcal{S}_n.$$

It is known that the strong minimum principle for this operator fails even for the simpler case $\lambda = \Lambda = 1$, where $\mathcal{M}_{\lambda, \Lambda; k}^{-}(D^2 u) = \sum_{i=1}^k e_i(D^2 u)$. Symmetrically, by duality one observes that $\mathcal{M}_{\lambda, \Lambda; k}^{+}(D^2 u) - \partial_t u$ satisfies the strong minimum principle, but not the strong maximum principle (a counterexample is Example 4 of [1] for the case $\lambda = \Lambda = 1$, $k < n$).

- Consider now the operator $G(D^2 u) = \sum_{i=1}^n \arctan e_i(D^2 u)$ and the equation

$$\sum_{i=1}^n \arctan e_i(D^2 u) - \partial_t u = 0.$$

Such an equation is called potential equation for the Lagrangian Mean Curvature Flow. It is known that when

$$\Theta > (n-2)\frac{\pi}{2} \text{ and } \Theta = \sum_{i=1}^n \arctan e_i(D^2u)$$

the operator is quasiconcave. Solutions of such equations with bounded Hessian are uniformly parabolic, and hence any subsolution (resp. supersolution) belongs to the class $\underline{S}$ (resp. $\overline{S}$). Therefore, the operator satisfies both the strong maximum and minimum principles. The same continues to hold for its elliptic counterpart via Remark 3.6.

REFERENCES

- [1] M. Alfaro and I. Birindelli. Evolution equations involving nonlinear truncated Laplacian operators. *Discrete Contin. Dyn. Syst.*, 40(6):3057–3073, 2020.
- [2] M. Bardi and F. Da Lio. On the strong maximum principle for fully nonlinear degenerate elliptic equations. *Arch. Math. (Basel)*, 73(4):276–285, 1999.
- [3] M. Bardi and A. Goffi. New strong maximum and comparison principles for fully nonlinear degenerate elliptic PDEs. *Calc. Var. Partial Differential Equations*, 58(6):Paper No. 184, 20, 2019.
- [4] L. A. Caffarelli and X. Cabré. *Fully nonlinear elliptic equations*, volume 43 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 1995.
- [5] L. A. Caffarelli, Y. Li, and L. Nirenberg. Some remarks on singular solutions of nonlinear elliptic equations III: viscosity solutions including parabolic operators. *Comm. Pure Appl. Math.*, 66(1):109–143, 2013.
- [6] E. Calabi. An extension of E. Hopf’s maximum principle with an application to Riemannian geometry. *Duke Math. J.*, 25:45–56, 1958.
- [7] M. G. Crandall, H. Ishii, and P.-L. Lions. User’s guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc. (N.S.)*, 27(1):1–67, 1992.
- [8] F. Da Lio. Remarks on the strong maximum principle for viscosity solutions to fully nonlinear parabolic equations. *Commun. Pure Appl. Anal.*, 3(3):395–415, 2004.
- [9] A. Friedman. *Partial differential equations of parabolic type*. Prentice-Hall, Inc., Englewood Cliffs, N.J., 1964.
- [10] A. Goffi and F. Pediconi. A note on the strong maximum principle for fully nonlinear equations on Riemannian manifolds. *J. Geom. Anal.*, 31(8):8641–8665, 2021.
- [11] F. R. Harvey and H. B. j. Lawson. Characterizing the strong maximum principle for constant coefficient subequations. *Rend. Mat. Appl., VII. Ser.*, 37(1-2):63–104, 2016.
- [12] A. M. Il’in, A. S. Kalashnikov, and O. A. Oleĭnik. Second-order linear equations of parabolic type. *J. Math. Sci. (New York)*, (108):435–542, 4, 2002.
- [13] C. Imbert and L. Silvestre. An introduction to fully nonlinear parabolic equations. In *An introduction to the Kähler-Ricci flow*, volume 2086 of *Lecture Notes in Math.*, pages 7–88. Springer, Cham, 2013.
- [14] J. Kovats. A three-curves theorem for viscosity subsolutions of parabolic equations. *Proc. Amer. Math. Soc.*, 131(5):1509–1514, 2003.
- [15] E. M. Landis. *Second order equations of elliptic and parabolic type*, volume 171 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1998.
- [16] G. M. Lieberman. *Second order parabolic differential equations*. World Scientific Publishing Co., Inc., River Edge, NJ, 1996.
- [17] L. Nirenberg. A strong maximum principle for parabolic equations. *Comm. Pure Appl. Math.*, 6:167–177, 1953.
- [18] P. Pucci and J. Serrin. *The maximum principle*, volume 73 of *Progress in Nonlinear Differential Equations and their Applications*. Birkhäuser Verlag, Basel, 2007.