



UNIVERSITÀ  
DEGLI STUDI  
FIRENZE

## FLORE

# Repository istituzionale dell'Università degli Studi di Firenze

### **Carotid intimal-medial thickness and stiffness are not affected by hypercholesterolemia in uncomplicated essential hypertension**

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

*Original Citation:*

Carotid intimal-medial thickness and stiffness are not affected by hypercholesterolemia in uncomplicated essential hypertension / P.S. SABA; M.J. ROMAN; C. LONGHINI; D. SCORZONI; R. PINI; R.B. DEVEREUX; A. GANAU. - In: ARTERIOSCLEROSIS, THROMBOSIS, AND VASCULAR BIOLOGY. - ISSN 1079-5642. - STAMPA. - 19:(1999), pp. 2788-2794.

*Availability:*

The webpage <https://hdl.handle.net/2158/219234> of the repository was last updated on

*Terms of use:*

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

*Publisher copyright claim:*

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

# Arteriosclerosis, Thrombosis, and Vascular Biology



JOURNAL OF THE AMERICAN HEART ASSOCIATION

## **Carotid Intimal-Medial Thickness and Stiffness Are Not Affected by Hypercholesterolemia in Uncomplicated Essential Hypertension**

Pier Sergio Saba, Mary J. Roman, Carlo Longhini, Daniela Scorzoni, Riccardo Pini, Richard B. Devereux and Antonello Ganau

*Arterioscler Thromb Vasc Biol.* 1999;19:2788-2794

doi: 10.1161/01.ATV.19.11.2788

*Arteriosclerosis, Thrombosis, and Vascular Biology* is published by the American Heart Association, 7272  
Greenville Avenue, Dallas, TX 75231

Copyright © 1999 American Heart Association, Inc. All rights reserved.

Print ISSN: 1079-5642. Online ISSN: 1524-4636

The online version of this article, along with updated information and services, is located on the  
World Wide Web at:

<http://atvb.ahajournals.org/content/19/11/2788>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Arteriosclerosis, Thrombosis, and Vascular Biology* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:

<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Arteriosclerosis, Thrombosis, and Vascular Biology* is online at:

<http://atvb.ahajournals.org/subscriptions/>

# Carotid Intimal-Medial Thickness and Stiffness Are Not Affected by Hypercholesterolemia in Uncomplicated Essential Hypertension

Pier Sergio Saba, Mary J. Roman, Carlo Longhini, Daniela Scorzoni, Riccardo Pini, Richard B. Devereux, Antonello Ganau

**Abstract**—The combined effects of hypertension and hypercholesterolemia on carotid anatomy and stiffness were studied in 62 normotensives, 141 uncomplicated essential hypertensives with a total cholesterol level <240 mg/dL, and 60 essential hypertensives with a total cholesterol level  $\geq$ 240 mg/dL. Carotid ultrasonography was performed to evaluate intimal-medial thickness (IMT), relative wall thickness, and the presence of plaque. Carotid pressure waveforms were recorded by applanation tonometry to measure carotid stiffness ( $\beta$ ) and pressure wave reflection (ie, augmentation index). After adjusting for age, body mass index, and smoking habit by analysis of covariance, no significant differences were found between normocholesterolemic hypertensives and hypercholesterolemic hypertensives in terms of IMT ( $0.79 \pm 0.19$  versus  $0.81 \pm 0.19$  mm), relative wall thickness ( $0.27 \pm 0.07$  versus  $0.28 \pm 0.07$ ), carotid stiffness ( $6.1 \pm 3.2$  versus  $5.6 \pm 2.7$ ), augmentation index ( $18.7 \pm 12.9\%$  versus  $17.3 \pm 12.8\%$ ), and prevalence of plaque (30.8% versus 30.7%). In the whole population, carotid IMT was significantly related to age ( $r=0.43$ ), systolic ( $r=0.35$ ) and diastolic ( $r=0.35$ ) blood pressures, body surface area ( $r=0.22$ ), and cholesterol levels ( $r=0.22$ ) (all  $P<0.05$ ). Carotid stiffness was significantly related to age, blood pressure, body mass index, and body surface area but not to cholesterol levels. In multivariate analyses, age, body surface area, and systolic blood pressure, but not cholesterol, smoking habit, or sex, were independent correlates of IMT (multiple  $R=0.54$ ,  $P<0.0001$ ), whereas carotid stiffness was independently associated with age, body surface area, and sex ( $R=0.38$ ,  $P<0.0001$ ). In conclusion, hypertension is a potent stimulus of vascular hypertrophy. The superimposition of hypercholesterolemia does not substantially augment these changes or further increase arterial stiffness in uncomplicated hypertensive subjects. (*Arterioscler Thromb Vasc Biol.* 1999;19:2788-2794.)

**Key Words:** hypertension ■ hypercholesterolemia ■ carotid arteries ■ atherosclerosis ■ arterial tonometry

Hypertension and hypercholesterolemia are major risk factors for cardiovascular morbidity and mortality in Western countries.<sup>1,2</sup> Both conditions are common in the general population and are frequently associated, thereby steeply increasing the risk of overall and coronary heart disease mortality.<sup>1</sup> Identification of patients at high risk by detection of preclinical disease, when target-organ damage may be in an early and potentially reversible stage, has been emphasized in recent years as a key step in the prevention of major clinical events.<sup>3</sup> Previous reports have shown a close relationship between carotid atherosclerosis and both cerebrovascular and coronary artery disease,<sup>4,5</sup> suggesting that carotid artery evaluation may help to identify high-risk subjects. However, carotid plaque may represent a late stage in the evolution of atherosclerotic disease in several vascular beds.<sup>6-8</sup>

The recent availability of high-resolution ultrasound techniques allows the detection of early arterial modifications,

such as intimal-medial thickening, which may precede the development of atherosclerotic plaques. Previous reports have indicated that carotid intimal-medial thickening is associated with coronary artery disease<sup>9,10</sup> and cardiovascular morbidity and mortality.<sup>11-15</sup> Moreover, carotid intimal-medial thickening has been described in association with hypertension<sup>16,17</sup> and hypercholesterolemia.<sup>18,19</sup> Studies of hypercholesterolemia have, however, included subjects with mild hypertension and/or symptomatic cardiovascular disease, and thus, they could not assess the independent contributions of uncomplicated hypertension and hypercholesterolemia to intimal-medial thickening.

Although arterial stiffening has been described in animals with diet-induced hypercholesterolemia in association with structural arterial modifications,<sup>20,21</sup> few data are presently available on the effect of hypercholesterolemia on arterial viscoelastic properties in humans. Because an increased pulse

Received December 15, 1998; revision accepted April 19, 1999.

From the Department of Medicine (M.J.R., R.B.D.), New York Hospital-Cornell Medical Center, New York, NY; the Institute of Clinical Medicine (P.S.S., A.G.), University of Sassari, Sassari, Italy; the Institute of Internal Medicine (C.L., D.S.), University of Ferrara, Ferrara, Italy; and the Division of Gerontology and Geriatrics (R.P.), University of Firenze, Firenze, Italy.

Correspondence to Mary J. Roman, MD, Division of Cardiology, New York Hospital-Cornell Medical Center, 525 E 68th St, New York, NY 10021. E-mail mroman@mail.med.cornell.edu

© 1999 American Heart Association, Inc.

*Arterioscler Thromb Vasc Biol.* is available at <http://www.atvbaha.org>

wave velocity has been described in hypercholesterolemic monkeys,<sup>21</sup> an earlier return of reflected pressure waves to the central aorta in hypercholesterolemia may augment central systolic blood pressure<sup>22</sup> and thereby ventricular afterload. This study evaluated a population of uncomplicated hypertensive patients with normal and high cholesterol levels to assess the relative contribution of hypercholesterolemia to carotid structural and functional modifications associated with hypertension.

## Methods

### Study Population

The study is a retrospective evaluation of a population of 395 normotensive and uncomplicated hypertensive subjects enrolled in a cooperative study of preclinical cardiovascular disease in hypertension conducted at Cornell University Medical Center, New York, NY; the Institute of Clinical Medicine, University of Sassari, Sassari, Italy; and the Institute of Internal Medicine, University of Ferrara, Ferrara, Italy. Exclusion criteria were a history or clinical signs of coronary artery disease, heart failure, or valvular heart disease; secondary hypertension; diabetes mellitus; and renal insufficiency. Hypertension was defined as a cuff blood pressure  $\geq 140$  mm Hg systolic and/or  $\geq 90$  mm Hg diastolic on repeated measurements taken by mercury sphygmomanometry. Sixty-two normotensive, normocholesterolemic subjects and 201 uncomplicated essential hypertensives were selected from this population, based on the availability of cholesterol data. All antihypertensive drug therapy was stopped at least 3 weeks before the study in the 116 (58%) patients who had been on antihypertensive treatment. None of the patients were on lipid-lowering therapy.

The effect of hypercholesterolemia in association with hypertension on carotid structure and stiffness was evaluated by dividing the study population into 3 groups: normal-cholesterol normotensives, normal-cholesterol hypertensives, and high-cholesterol hypertensives, on the basis of a normal to borderline-high ( $< 6.22$  mmol/L, or 240 mg/dL) or a high ( $\geq 6.22$  mmol/L) fasting total serum cholesterol level obtained within 1 month of the vascular examination.<sup>23</sup> Body habitus was evaluated by body mass index (BMI, weight in kilograms per square of height in meters squared) and body surface area (BSA, obtained from height and weight measurements by standard nomograms). The study protocol was approved by the Committee on Human Rights in Research of Cornell University Medical College in 1989 and at intervals thereafter and by the Ethics Committees of the Universities of Sassari and Ferrara.

### Carotid Ultrasonography

Carotid ultrasonography was performed in subjects recruited in New York (n=199) by using a Biosound Genesis II system (OTE Biomedica) equipped with a 7.5-MHz probe; subjects recruited in Sassari (n=43) and Ferrara (n=21) were examined with a Toshiba 270 HG system equipped with a 7.5-MHz probe. With the subject in the supine position and the neck in slight hyperextension, the common carotid artery, carotid bulb, and the extracranial portions of internal and external carotid arteries were identified. Two-dimensionally guided M-mode tracings of the distal common carotid artery ( $\approx 1$  cm proximal to the carotid bulb) were obtained with simultaneous ECG and contralateral carotid pressure waveform tracings (see below) and recorded on 1/2-in. Super VHS videotape. After the videotape was reviewed, suitable frames for measurement of M-mode images were obtained by using a frame grabber (Imaging Technology Inc) interfaced with a high-resolution (640 $\times$ 480 pixels) video monitor and stored on diskette.

Carotid measurements were performed on the stored images by using mouse-driven software written by 1 of the investigators (R.P.), after calibration for depth and time. The simultaneous carotid pressure tracing (see below) was used to time the carotid artery measurement at end diastole (minimum arterial pressure) and at the time of peak systolic pressure. Measurements included the combined intimal-medial thickness (IMT) of the far wall at end diastole, as have been validated in animal and human studies, and end dia-

stolic and peak systolic internal dimensions obtained by continuous tracing of the intima-lumen interface of the near and far walls. All measurements were performed on several cycles and averaged.

Relative wall thickness (RWT) of the artery was calculated according to the formula  $2 \times \text{far-wall diastolic thickness} / \text{end-diastolic diameter}$ . Because increased distending pressure, as occurs in hypertensive patients, could decrease the IMT due to vessel wall stretching, carotid intimal-medial cross-sectional area was also calculated as  $\{[(\text{diastolic diameter}/2) + \text{far-wall diastolic thickness}]^2 \times \pi\} - [(\text{diastolic diameter}/2)^2 \times \pi]$  as an estimate of circumferential intimal-medial tissue mass.<sup>17,25</sup> Both carotid arteries were scanned to identify the presence and size of atherosclerotic plaques, defined as focal increases in IMT  $> 50\%$  of the surrounding wall; standard wall thickness measurements were never obtained at the level of a discrete plaque.

To minimize operator variability, ultrasonography and tonometry were always performed in each center under the guidance of the same operator (M.J.R., P.S.S., D.S.), and measurements of carotid anatomy were carried out by 2 investigators (M.J.R. for the New York data and P.S.S. for the Sassari and Ferrara data). As previously reported, intraobserver ( $r=0.98$ ,  $SEE=0.04$  mm for both) and interobserver ( $r=0.97$ ,  $SEE=0.05$  mm) reproducibility of blinded wall thickness measurements between the 2 readers was high.<sup>16</sup>

### Carotid Stiffness

In association with carotid ultrasonography, carotid pressure waveforms were obtained as described elsewhere<sup>26-28</sup> by using a high-fidelity external pressure transducer (Millar Instruments, Inc) applied to the skin overlying the pulse of the contralateral common carotid artery.

The transducer is internally calibrated (0.2 V/100 mm Hg) and registers absolute changes in applied pressure over a range of 300 mm Hg. To obtain actual carotid blood pressure values, the waveforms require additional external calibration. Based on the observation that mean arterial pressure is nearly identical in all capacitance vessels,<sup>29-31</sup> brachial artery pressure was measured by the cuff and mercury sphygmomanometer method with the patient in the supine position, and mean pressure was calculated as diastolic blood pressure +  $(1/3 \times \text{pulse pressure})$ . The resultant value was assigned to the planimetrically computer-derived mean blood pressure of the carotid waveforms with software written by 1 of the investigators (R.P.). Carotid artery stiffness was estimated by Young's modulus and the stiffness index. Young's modulus (E; wall tension per centimeter of thickness for a 100% diameter increase) was calculated according to the formula  $E = [(P_s - P_d) / (D_s - D_d)] \times (D/h)$ , where  $P_s$  and  $P_d$  are systolic and diastolic pressures, respectively;  $D_s$  and  $D_d$  are systolic and diastolic carotid dimensions, respectively; and  $D$  and  $h$  are carotid mean diameter and wall thickness, respectively.<sup>32</sup> Carotid stiffness was also calculated by the pressure-independent stiffness index  $\beta$  according to the formula<sup>33,34</sup>  $\beta = [\ln(P_s/P_d)] / [(D_s - D_d)/D_d]$ . This index takes into account the logarithmic relation between arterial pressure and diameter.

### Pressure Wave Reflection

The systolic portion of the carotid waveform was analyzed to calculate the augmentation index,<sup>35,36</sup> a means of quantifying the contribution of reflected pressure waves to the central pulse pressure. After identification of the early and late systolic peaks and the inflection that separates them, pressures were determined at peak systolic pressure ( $P_{pk}$ ) and at the inflection point ( $P_i$ ) by using the previously described calibrated computer system. The augmentation index was calculated as  $(P_{pk} - P_i) / \text{pulse pressure}$  (when  $P_{pk}$  occurred in late systole) or as  $(P_i - P_{pk}) / \text{pulse pressure}$  (when  $P_{pk}$  occurred in early systole).<sup>36</sup> Pulse wave analysis was performed by a single investigator (P.S.S.); measurements were repeated on several cycles and averaged.

### Plasma Lipid Determinations

Total cholesterol and triglycerides were determined by standard enzymatic methods; the HDL cholesterol fraction was determined after precipitation of apolipoprotein B-containing lipoproteins, and LDL cholesterol was calculated as described by Friedewald et al.<sup>37</sup>

**TABLE 1. Characteristics of Normotensive Subjects and of Normal- and High-Cholesterol Hypertensive Patients**

	Normotensive Subjects (Chol<6.22 mmol/L)	Hypertensive Patients	
		Chol<6.22 mmol/L	Chol≥6.22 mmol/L
n	62	141	60
Age, y	49±19	53±11	58±12*†
Sex, % male	61.3	61.7	55.0
Body weight, kg	69.2±17.9	77.1±16.3*	75.0±17.1‡
Body height, m	1.69±0.10	1.70±0.10	1.67±0.10
BMI, kg/m <sup>2</sup>	24.0±4.4	26.4±4.0*	26.5±3.7*
BSA, m <sup>2</sup>	1.79±0.25	1.88±0.23‡	1.84±0.25
Systolic BP, mm Hg	126±16	156±19*	160±20*
Diastolic BP, mm Hg	76±9	95±12*	92±12*
Total chol, mmol/L	5.10±0.75	5.15±0.67	7.12±0.80*†
HDL chol, mmol/L	1.42±0.41	1.35±0.41	1.45±0.41
LDL chol, mmol/L	3.13±0.83	3.21±0.67	4.79±0.73*†
Total HDL ratio	4.2±1.1	4.2±1.4	5.3±1.5*†
Former or current smokers, %	29.5	40.0	56.7*†

Chol indicates cholesterol; BP, blood pressure.

\* $P<0.01$  vs normotensive subjects.

† $P<0.01$  vs hypertensive subjects with cholesterol<6.22 mmol/L.

‡ $P<0.05$  vs normotensive subjects.

Hypercholesterolemia was defined as total cholesterol  $\geq 6.22$  mmol/L ( $\geq 240$  mg/dL).<sup>23</sup>

### Statistical Analysis

Data were stored and analyzed by using the Crunch<sup>4</sup> Statistical Package (Crunch Software Corp). Relationships between continuous variables were evaluated by linear regression analyses. Comparisons between normotensive and hypertensive subjects were performed by Student's *t* test. Carotid anatomy and stiffness were compared among normal-cholesterol normotensives and normal- and high-cholesterol hypertensives by ANOVA. ANCOVA was also performed to compare groups after controlling for differences in age, BSA, and smoking habit. Both analyses were followed by the Ryan, Einot, Gabriel, and Welsch *F* test multiple-comparison procedure. Power analysis was performed to assess the minimal difference in IMT detectable in our study. Based on the size of our groups and within-group SDs, differences as small as 0.07 mm for the IMT were detectable between normal and high-cholesterol hypertensives with an  $\alpha$  error of 0.05 and a statistical power of 0.90; this detectable difference is lower than the absolute between-group difference in IMT associated with a significantly higher risk of cardiovascular and/or cerebrovascular events.<sup>38,39</sup> Multiple regression analyses were performed to assess independent determinants of carotid intimal-medial cross sectional area and the stiffness index in normotensive and hypertensive subjects. Data are expressed as mean±SD. A 2-tailed *P* value <0.05 was considered significant.

## Results

### Characteristics of Normotensive and Hypertensive Subjects

Among the 201 hypertensive patients, 141 had total cholesterol levels <6.22 mmol/L (240 mg/dL), while 60 had a total cholesterol level  $\geq 6.22$  mmol/L (Table 1). Hypertensive patients with higher cholesterol levels were significantly older than normocholesterolemic hypertensive and normotensive subjects. Both hypertensive groups had significantly greater body weights and BMIs than did normotensives, while the distribution of sex and height was similar among the groups. BSA was significantly greater in normocholesterolemic hypertensives than in normo-

tensive subjects. Both hypertensive groups had similar blood pressure levels, and the normocholesterolemic groups did not differ in total and LDL cholesterol. The total-HDL ratio was significantly increased in hypercholesterolemic hypertensives. Hypercholesterolemic hypertensives were more likely to be current or former smokers.

### Carotid Anatomy and Stiffness in Normal- and High-Cholesterol Hypertensive Patients

Carotid wall thickness, RWT, cross-sectional area, and prevalence of plaque were significantly greater in both hypertensive groups than in the normotensive subjects, while normal- and high-cholesterol hypertensives did not differ in these parameters (Table 2). Neither the stiffness index nor Young's elastic modulus differed significantly among the 3 groups, whereas the augmentation index was similarly higher in both hypertensive groups than in normotensive subjects. ANCOVA, after controlling for differences in age, BSA, and smoking habit, yielded similar results (Table 2, values in parentheses). To avoid a potentially inadequate adjustment for age in ANCOVA, each hypertensive group was also stratified by age into 2 subsets (<50 and  $\geq 50$  years); carotid parameters were then analyzed in relation to the cholesterol levels. There were no significant differences between the 2 subgroups comparable by age but different in cholesterol level. For example, mean carotid IMT was  $0.75\pm 0.17$  and  $0.71\pm 0.16$  mm in hypercholesterolemic and normocholesterolemic patients <50 years old and  $0.88\pm 0.19$  and  $0.85\pm 0.19$  mm in older hypercholesterolemic and normocholesterolemic patients ( $P=0.38$  and  $0.40$ , respectively).

### Univariate Correlates of Carotid Anatomy and Stiffness Parameters in Normotensive and Hypertensive Subjects

In the entire population (upper portion of Table 3), carotid IMT, RWT, and cross-sectional area were directly and

**TABLE 2. Carotid Anatomy and Stiffness in Normal- and High-Cholesterol Hypertensive Subjects**

	Normotensive Subjects	Hypertensive Patients	
		Chol<6.22 mmol/L	Chol≥6.22 mmol/L
IMT, mm	0.66±0.17 (0.70)	0.80±0.19* (0.79)*	0.84±0.19* (0.81)*
Internal diameter, mm	5.7±0.7 (5.8)	5.9±0.8 (5.9)	5.9±0.7 (5.8)
RWT	0.24±0.06 (0.25)	0.27±0.07* (0.27)*	0.29±0.07* (0.28)*
Cross-sectional area, mm <sup>2</sup>	13.3±4.4 (14.5)	16.9±5.6* (17.0)*	17.5±4.8* (16.6)*
Cross-sectional area/BSA, mm <sup>2</sup> /m <sup>2</sup>	7.5±2.3 (7.8)	9.0±2.8* (9.1)*	9.7±2.9* (9.3)*
Plaques, %	16.1 (24.3)	27.7* (30.8)*	28.3* (30.7)*
Stiffness index, β	6.2±2.3 (6.5)	6.1±3.2 (6.1)	6.0±2.7 (5.6)
Young's elastic modulus, dynes/cm <sup>2</sup> ×mm×10 <sup>-6</sup>	0.94±0.46 (0.96)	0.96±0.45 (0.97)	0.93±0.48 (0.89)
Augmentation index, %	8.8±15.1 (10.5)	18.2±12.9* (18.7)*	19.7±12.8* (17.3)*

Chol indicates cholesterol. Values in parentheses are means corrected for age, BMI, and smoking history by ANCOVA. \*P<0.01 vs normotensive subjects.

significantly related to total and LDL cholesterol, age, BMI, BSA, and systolic and diastolic blood pressures. Similarly, RWT was directly related to cholesterol levels, age, BMI, and systolic pressure but not to BSA or diastolic blood pressure (Table 3). Carotid lumen diameter was most closely related to older age and higher blood pressure. Young's modulus was significantly related to age and systolic blood pressures and negatively to BMI and BSA. The stiffness index was directly related to age and negatively to BSA; no stiffness index showed a significant relationship with cholesterol levels. Similar results were obtained after repeating the analysis in the hypertensive population only (lower portion of Table 3).

**Independent Predictors of Carotid IMT and Stiffness Index**

Multiple regression analyses were performed to assess independent predictors of carotid IMT and stiffness (Table 4). Age, BSA, and systolic blood pressure were independent predictors of carotid IMT (multiple R=0.54, P<0.0001), while total chole-

sterol, smoking history, and sex did not enter the model. Age, BSA, and male sex, but not cholesterol level, smoking history, and systolic blood pressure, were independent predictors of arterial stiffness (multiple R=0.38, P<0.0001).

In the subset of hypertensive patients, age and BSA were independent predictors of carotid IMT (multiple R=0.42, P<0.01), and no residual correlation was found with cholesterol levels (Figure 1). Similarly, Young's elastic modulus of the carotid was independently predicted by systolic blood pressure and BSA, while the stiffness index was independently positively correlated with age and smoking history and negatively with cholesterol levels (Figure 2).

**Discussion**

The present study indicates that mild to moderate hypercholesterolemia does not play a primary role in increasing common carotid artery IMT in otherwise-healthy hypertensive patients. In fact, although significant, positive, multivariate relationships were found between carotid structure and

**TABLE 3. Univariate Correlates of Carotid Anatomy and Stiffness Measures With Cholesterol Levels and Demographic Characteristics**

	Total Chol	LDL Chol	Age	BMI	BSA	Systolic BP	Diastolic BP
Total population							
IMT	0.22*	0.21*	0.43*	0.25*	0.22†	0.35*	0.18*
Internal diameter	0.01	0.05	0.24*	0.14†	0.23*	0.29*	0.24*
Cross-sectional area	0.16*	0.16†	0.45*	0.26*	0.26*	0.40*	0.22*
RWT	0.18*	0.15†	0.33*	0.20*	0.11	0.20*	0.04
Young's modulus	-0.03	0.02	0.19*	-0.13†	-0.21*	0.21*	0.19*
Stiffness index β	0.01	0.03	0.33*	-0.07	-0.17*	0.12	0.03
Hypertensive patients							
IMT	0.15†	0.14	0.36*	0.13	0.15†	0.16†	-0.08
Internal diameter	-0.04	-0.03	0.24*	0.07	0.15†	0.29*	0.20*
Cross-sectional area	0.09	0.07	0.40*	0.14	0.19*	0.26*	-0.01
RWT	0.12	0.12	0.23*	0.10	0.06	-0.01	-0.22*
Young's modulus	-0.06	-0.03	0.19*	-0.15†	-0.22*	0.28*	0.22*
Stiffness index β	-0.04	-0.03	0.31*	-0.09	-0.18†	0.17†	0.02

Chol indicates cholesterol; BP, blood pressure.

\*P<0.01.

**TABLE 4. Independent Predictors of Carotid IMT and Stiffness Index  $\beta$**

	Multiple		Partial Correlation	F	P
	R	R <sup>2</sup>	Coefficient*		
Dependent variable: IMT					
Age	0.44	0.19	0.39	69.2	0.0001
BSA	0.52	0.27	0.27	27.5	0.0001
Systolic BP	0.54	0.29	0.19	10.6	0.0013
Total cholesterol	Not entered		...	1.3	0.24
Smoking history†	Not entered		...	0.2	0.69
Sex‡	Not entered		...	0.03	0.86
Dependent variable: stiffness index $\beta$					
Age	0.33	0.11	0.32	30.3	0.0001
BSA	0.36	0.13	-0.25	5.5	0.02
Sex‡	0.38	0.15	-0.18	5.4	0.02
Total cholesterol	Not entered		...	2.4	0.12
Smoking history†	Not entered		...	2.1	0.14
Systolic BP	Not entered		...	0.05	0.83

BP indicates blood pressure.

\*After the last step.

†Smoking history: never=1; former or current=2.


‡Sex: male=1, female=2.

cholesterol levels after controlling for age and other potentially confounding parameters, hypertensive patients with normal and high cholesterol levels did not differ in carotid wall thickness, RWT, and cross-sectional area. In contrast, the association of hypertension with increased IMT was evident in both groups. Accordingly, independent predictors of carotid cross-sectional area included older age and higher BSA and systolic blood pressure but not total serum cholesterol (Table 4).


These data are partially in contrast with previous reports that indicated a role of hypercholesterolemia in inducing carotid intimal-medial thickening.<sup>12,19,40-44</sup> However, 2 of these studies<sup>19,42</sup> included hypercholesterolemic patients with mild or moderate hypertension; moreover, diabetic patients and/or subjects with signs or symptoms of cardiovascular disease were not excluded. Nevertheless, although Poli et al<sup>42</sup> detected a significant increase in IMT in their hypercholesterolemic group, there was no significant correlation between carotid IMT and plasma cholesterol levels. In contrast, in the population described by Wendelhag et al,<sup>19</sup> the distribution of

IMT in hypercholesterolemic patients appeared to be similar to that in normocholesterolemic subjects except for a few subjects in whom IMT values suggested the inclusion of plaques in the calculation. In addition, studies conducted in normotensive children with familial hypercholesterolemia and very high cholesterol levels showed only modest associations between carotid IMT and cholesterol levels.<sup>43,44</sup>

Although it is unclear whether it is more clinically and prognostically meaningful to exclude from the calculation of IMT all discrete areas of wall thickening, measurements performed in arterial segments including discrete plaques could overemphasize the association between risk factors for atherosclerotic disease and the IMT of representative segments of the arterial tree. Data from our study suggest that when IMT is measured in areas free of discrete atherosclerotic lesions, common carotid wall thickness in hypertensive patients is not strongly affected by mild to moderate hypercholesterolemia. Moreover, our results are in accordance with experimental studies<sup>45</sup> that suggest that intimal-medial thick-



**Figure 1.** Residual correlation between carotid IMT and cholesterol levels in hypertensive patients after controlling for age and BSA in a multiple regression analysis (multiple  $R=0.42$ ). Variables introduced in the model were age, sex, systolic blood pressure, and smoking history.



**Figure 2.** An inverse, significant, residual correlation was present between carotid stiffness index and cholesterol levels in hypertensive patients after controlling for age and smoking history in a multiple regression analysis (multiple  $R=0.34$ ). Variables introduced in the model were age, sex, systolic blood pressure, and BSA.

ening is mainly related to increased deformational arterial stresses due to blood pressure elevation, whereas hypercholesterolemia may be more influential in the genesis of discrete atherosclerotic plaques by focal deposition of lipids in the subintimal layers of the arterial wall. This concept is in agreement with observations that normotensive hypercholesterolemic children had mean wall thicknesses similar to those of normocholesterolemic controls, whereas the maximum IMT was significantly higher in the hypercholesterolemic group.<sup>43</sup>

A potential limitation of ultrasonographic evaluation of the arterial intimal-medial complex is that it does not allow separation of increased medial mass due to hypertrophy from increased intimal mass due to atherosclerosis and arteriosclerosis. Therefore, early atherosclerotic lesions involving only the intimal layer may cause only small increases in the composite wall thickness measured by vascular ultrasonography that might be confused with medial hypertrophy in the presence of hypertension. Although this could be a major limitation of ultrasonographic studies of preclinical atherosclerosis, the consistent ability of increased IMT to predict cardiovascular events in multiple populations<sup>11–15</sup> suggests that the different processes that increase overall carotid wall thickness are all prognostically adverse. Furthermore, since the atherosclerotic process typically involves the arterial wall in a focal fashion, whereas hypertension induces diffuse thickening of the intimal-medial complex,<sup>46</sup> ultrasonographic identification of focal intimal-medial thickening can be considered a reliable indicator of atherosclerosis per se.

Surprisingly, carotid stiffness, which is commonly considered an early indicator of vascular damage, was not increased by hypercholesterolemia. In fact, the stiffness index, which takes into account the nonlinearity of the pressure-diameter relation, and Young's elastic modulus, which considers adaptive wall thickening, were not elevated in either hypertensive group, and in particular, were not increased in the hypercholesterolemic patients. Multiple regression analysis conducted in the whole population indicated that age, BSA, and sex, but not cholesterol levels, were independent predictors of the stiffness index. However, in a subanalysis of the hypertensive population, an unexpected modest but significant independent inverse relationship was found between stiffness index and cholesterol level. This surprising finding, which needs verification by other studies, confirms the lack of positive correlations between carotid stiffness and cholesterol levels after controlling for age and smoking history. Moreover, the augmentation index, which is related to aortic pulse wave velocity, was equally elevated in normocholesterolemic and hypercholesterolemic hypertensives, suggesting that neither carotid nor aortic compliance was substantially modified by mild-to-moderate hypercholesterolemia. The evidence for altered arterial stiffness in hypertensive patients when the pressure-sensitive augmentation index was used but not with the pressure-insensitive stiffness index is in accordance with previous reports that did not detect an impact of hypertension on arterial stiffness evaluated at a standardized pressure level.<sup>47,48</sup> Although studies in hypercholesterolemic animals indicated increased arterial stiffness,<sup>20,21</sup> results may have been affected by use of the pressure-dependent pulse wave velocity as the index of vascular stiffness. Other human observations did not detect increased arterial compliance in

nonfamilial hypercholesterolemia<sup>49</sup> or an effect of high cholesterol levels on age-related increases in arterial stiffness.<sup>50,51</sup> Moreover, a recent study reported a lack of association between aortic stiffening and extracoronary atherosclerotic disease, after controlling for the affect of age.<sup>52</sup>

Although ultrasonographic evaluation of uncomplicated hypertensive subjects allows the assessment of vascular damage in a preclinical phase of atherosclerotic disease, the lack of cholesterol-induced arterial modifications in our study may be due to selection of hypertensive subjects at relatively low risk. In fact, the prevalence of carotid plaques, although significantly higher than in normocholesterolemic subjects, was relatively low in the high-cholesterol hypertensive group (31%).

In conclusion, this study indicates that in patients with relatively mild, uncomplicated essential hypertension, sporadic hypercholesterolemia does not substantially affect carotid wall thickness or stiffness. However, although the relatively small sample size of this study had adequate power to detect a moderate intimal-medial thickening induced by high cholesterol levels, lesser changes might have gone undetected in our moderately sized population. Additionally, in the present study, the impact of hypercholesterolemia may have been overwhelmed by hypertension-induced arterial modifications. Further larger studies are needed to assess the independent effect of high cholesterol levels on arterial anatomy and stiffness in normotensive subjects.

### Acknowledgments

This study was supported in part by grants HL-18323 and HL-47540 from the National Heart, Lung, and Blood Institute, Bethesda, Md; the Michael Wolk Heart Foundation; and grant 97.04129.CT04 from Consiglio Nazionale delle Ricerche and grant ex-40% 1997 from MURST, Italy (to A.G.). P.S.S. has been the recipient of a research grant from Regione Autonoma della Sardegna, Italy.

### References

1. Kannel WB, Neaton JD, Wentworth D, Thomas HE, Stamler J, Hulley SB, Kjelsberg MO. Overall and coronary heart disease mortality rates in relation to major risk factors in 325,348 men screened for the MRFIT Study. *Am Heart J*. 1986;112:825–836.
2. Kannel WB, Castelli WP, Gordon T, McNamara PM. Serum cholesterol, lipoproteins and the risk of coronary heart disease: the Framingham Study. *Ann Intern Med*. 1971;74:1–12.
3. Devereux RB, Alderman MH. Role of preclinical cardiovascular disease in the evolution from risk factor exposure to development of morbid events. *Circulation*. 1993;88:1444–1455.
4. Hennerici M, Hulsbomer HM, Hefter H, Lammerts D, Rautemberg W. Natural history of asymptomatic extracranial arterial disease. *Brain*. 1987;110:777–791.
5. Craven TE, Ryu JE, Espeland MA, Kahl FR, McKinney WM, Toole JF, McMahan MR, Thompson CJ, Heiss G, Crouse JR III. Evaluation of the associations between carotid artery atherosclerosis and coronary artery stenosis: a case-control study. *Circulation*. 1990;82:1230–1242.
6. Giral P, Pithois-Merli I, Filitti V, Levenson J, Plainfosse MC, Mainardi C, Simon AC. Risk factors and early extracoronary atherosclerotic plaques detected by three-site ultrasound imaging in hypercholesterolemic men. *Arch Intern Med*. 1991;151:950–956.
7. Suurkula M, Agewall S, Fagenberg B, Wendelhag I, Widgren B, Wikstrand J. Ultrasound evaluation of atherosclerotic manifestations in the carotid artery in high-risk hypertensive patients. *Arterioscler Thromb*. 1994;14:1297–1304.
8. Young W, Gofman JW, Tandy R, Malamud N, Waters ESG. The quantification of atherosclerosis, III: the extent of correlation of degrees of atherosclerosis within and between the coronary and cerebral vascular beds. *Am J Cardiol*. 1960;6:300–308.
9. Mitchell JRA, Schwartz CJ. Relationship between arterial disease in different sites: a study of the aorta and coronary, carotid and iliac arteries. *BMJ*. 1963;2:123–126.

10. Geroulakos G, O'Gorman DJ, Kalodiki E, Sheridan DJ, Nicolaides AN. The carotid intima-media thickness as a marker of severe symptomatic coronary artery disease. *Eur Heart J.* 1994;15:781-785.
11. Burke GL, Evans GW, Riley WA, Sharrett AR, Howard G, Barnes RW, Rosamond W, Crow RS, Rautaharju PM, Heiss G. Arterial wall thickness is associated with prevalent cardiovascular disease in middle-aged adults: the Atherosclerosis Risk in Communities (ARIC) Study. *Stroke.* 1995;26:386-391.
12. Salonen JT, Salonen R. Arterial wall thickness, carotid atherosclerosis and the risk of myocardial infarction and cerebrovascular stroke. In: Touboul PJ, Crouse JR, eds. *Intima-Media Thickness and Atherosclerosis.* New York, NY: Parthenon Publishing Group; 1997.
13. Kuller LH, Shemanski L, Psaty BM, Borhani NO, Gardin J, Haan MN, O'Leary DH, Savage PJ, Tell GS, Tracy R. Subclinical disease as an independent risk factor for cardiovascular disease. *Circulation.* 1995;92:720-726.
14. O'Leary DH, Polak JF, Kronmal RA, Kittner SJ, Bond MG, Wolfson SK Jr, Bommer W, Price TR, Gardin JM, Savage PJ. Distribution and correlates of sonographically detected carotid artery disease in the Cardiovascular Health Study. *Stroke.* 1992;23:1752-1760.
15. Belcaro G, Nicolaides AN, Laurora G, Cesarone MR, De Sanctis M, Incandela L, Barsotti A. Ultrasound morphology classification of the arterial wall and cardiovascular events in a 6-year follow-up study. *Arterioscler Thromb Vasc Biol.* 1996;16:851-856.
16. Roman MJ, Saba PS, Pini R, Spitzer M, Pickering TG, Rosen S, Alderman MH, Devereux RB. Parallel cardiac and vascular adaptation in hypertension. *Circulation.* 1992;86:1909-1918.
17. Lemne C, Jøgestrand T, deFaire U. Carotid intima-media thickness and plaque in borderline hypertension. *Stroke.* 1995;26:34-39.
18. Poli A, Tremoli E, Colombo A, Sirtori M, Pignoli P, Paoletti R. Ultrasonographic measurement of the common carotid artery wall thickness in hypercholesterolemic patients: a new model for the quantitation and follow-up of preclinical atherosclerosis in living human subjects. *Atherosclerosis.* 1988;70:253-261.
19. Wendelhag I, Wiklund O, Wikstrand J. Arterial wall thickness in familial hypercholesterolemia: ultrasound measurement of intima-media thickness in the common carotid artery. *Arterioscler Thromb.* 1992;12:70-77.
20. Burattini R, Montanari L, Mulligan LJ, Cannon MS, Gross DR. Evaluation of hypercholesterol diet-induced changes in viscoelastic properties of carotid circulation in pigs. *Am J Physiol.* 1992;263:H1919-H1926.
21. Farrar DJ, Bond MG, Riley WA, Sawyer JK. Anatomic correlates of aortic pulse wave velocity and carotid artery elasticity during atherosclerosis progression and regression in monkeys. *Circulation.* 1991;83:1754-1763.
22. Yaginuma T, Cohn JN. Introduction to wave reflection, wave travel and changes with modification of geometric and physical parameters. *J Hypertens.* 1992;10(suppl):S71-S72.
23. Working Group on Management of Patients with Hypertension and High Blood Cholesterol. National Education Programs Working Group report on the management of patients with hypertension and high blood cholesterol. *Ann Intern Med.* 1991;114:224-237.
24. Pignoli P, Tremoli E, Poli A, Oreste P, Paoletti R. Intimal plus medial thickness of the arterial wall: a direct measurement with ultrasound imaging. *Circulation.* 1986;74:1399-1406.
25. Roman MJ, Pickering TG, Pini R, Schwartz JE, Devereux RB. Prevalence and determinants of cardiac and vascular hypertrophy in hypertension. *Hypertension.* 1995;26:369-373.
26. Roman MJ, Pini R, Pickering TG, Devereux RB. Noninvasive measurement of arterial compliance in hypertensive compared with normotensive adults. *J Hypertens.* 1992;10(suppl 6):S115-S118.
27. Saba PS, Roman MJ, Pini R, Spitzer M, Ganau A, Devereux RB. Relation of pressure waveform to left ventricular and carotid anatomy in normotensive subjects. *J Am Coll Cardiol.* 1993;22:1873-1880.
28. Saba PS, Scorzoni D, Cavallini C, Longhini C, Pini R, Ganau A. Arterial tonometry: principles and clinical applications in hypertension. *High Blood Press.* 1996;5:241-250.
29. Hamilton WF, Dow P. An experimental study of the standing waves in the pulse propagated through the aorta. *Am J Physiol.* 1939;125:48-59.
30. Schnabel TG Jr, Fitzpatrick HF, Peterson LH, Rashkind WJ, Talley D, Raphael RL. A technic of vascular catheterization with small plastic catheters: its utilization to measure the arterial pulse wave velocity in man. *Circulation.* 1952;5:257-262.
31. Kroeker EJ, Wood EH. Comparison of simultaneously recorded central and peripheral arterial pressure pulses during rest, exercise and tilted position in man. *Circ Res.* 1955;3:623-632.
32. O'Rourke MF. Arterial stiffness, systolic blood pressure and logical treatment of arterial hypertension. *Hypertension.* 1990;15:339-347.
33. Kawasaki T, Sasayama S, Yagi SI, Asakawa T, Hirai T. Non-invasive assessment of the age related changes in stiffness of major branches of the human arteries. *Cardiovasc Res.* 1987;21:678-687.
34. Hirai T, Sasayama S, Kawasaki T, Yagi SI. Stiffness of systemic arteries in patients with myocardial infarction: a noninvasive method to predict severity of coronary atherosclerosis. *Circulation.* 1989;80:78-86.
35. Kelly R, Daley J, Avolio A, O'Rourke M. Arterial dilation and reduced wave reflection: benefit of diltiazem in hypertension. *Hypertension.* 1989;14:14-21.
36. Murgu JP, Westerhof N, Giolma JP, Altabelli SA. Aortic input impedance in normal man: relationship to pressure wave forms. *Circulation.* 1980;62:105-116.
37. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem.* 1972;18:499-502.
38. O'Leary DH, Polak JF, Kronmal RA, Manolio TA, Burke GL, Wolfson SK Jr. Carotid-artery intima and media thickness as a risk factor for myocardial infarction and stroke in older adults: Cardiovascular Health Study Collaborative Research Group. *N Engl J Med.* 1999;340:14-22.
39. Hodis HN, Mack WJ, LaBree L, Selzer RH, Liu C-r, Liu C-h, Azen SP. The role of carotid arterial intima-media thickness in predicting clinical coronary events. *Ann Intern Med.* 1998;128:262-269.
40. Crouse JR, Toole JF, McKinney WM, Dignan MB, Howard G, Kahl FR, McMahan MR, Harpold GH. Risk factors for extracranial carotid artery atherosclerosis. *Stroke.* 1987;18:990-996.
41. Handa N, Matsumoto M, Maeda H, Hougaku H, Ogawa S, Fukunaga R, Yoneda S, Kimura K, Kamada T. Ultrasonic evaluation of early carotid atherosclerosis. *Stroke.* 1990;21:1567-1572.
42. Poli A, Tremoli E, Colombo A, Sirtori M, Pignoli P, Paoletti R. Ultrasonographic measurement of the common carotid artery wall thickness in hypercholesterolemic patients: a new model for the quantitation and follow-up of preclinical atherosclerosis in living human subjects. *Atherosclerosis.* 1988;70:253-261.
43. Paucullio P, Iannuzzi A, Sartorio R, Irace C, Covetti G, Di Costanzo A, Rubba P. Increased intima-media thickness of the common carotid artery in hypercholesterolemic children. *Arterioscler Thromb.* 1994;14:1075-1079.
44. Tonstad S, Joakimsen O, Stensland-Bugge E, Leren TP, Ose L, Russel D, Bønaa KH. Risk factors related to carotid intima-media thickness and plaque in children with familial hypercholesterolemia and control subjects. *Arterioscler Thromb Vasc Biol.* 1996;16:984-991.
45. Dobrin PB. Mechanical factors associated with the development of intimal and medial thickening in vein grafts subjected to arterial pressure: a model of arteries exposed to hypertension. *Hypertension.* 1995;26:38-43.
46. Stary HC, Chandler AB, Glagov S, Guyton JR, Insull W, Rosenfeld ME, Schaffer SA, Schwartz CJ, Wagner WD, Wissler RW. A definition of initial, fatty streak, and intermediate lesions of atherosclerosis: a report from the Committee on Vascular Lesions of the Council on Arteriosclerosis, American Heart Association. *Circulation.* 1994;89:2462-2478.
47. Devereux RB, Waeber B, Roman MJ. Conclusions on the measurement of arterial wall thickness: anatomic, physiologic and methodologic considerations. *J Hypertens.* 1992;10(suppl 6):S119-S121.
48. Laurent S, Caviezel B, Beck L, Girerd X, Billaud E, Boutouyrie P, Hoeks A, Safar M. Carotid artery distensibility and distending pressure in hypertensive humans. *Hypertension.* 1994;23:878-883.
49. Giannattasio C, Mangoni AA, Carugo S, Bombelli M, Stefanoni P, Failla M, Stella ML, Sega R, Grassi G, Vergani C, Mancia G. Arterial compliance in familial hypercholesterolemia: a preliminary report. *J Hypertens.* 1993;11(suppl 5):S82-S83.
50. Dart AM, Lacombe F, Yeoh JK, Cameron JD, Jennings GL, Laufer E, Esmore DS. Aortic distensibility in patients with isolated hypercholesterolemia, coronary artery disease, or cardiac transplant. *Lancet.* 1991;338:270-273.
51. O'Rourke MF, Avolio AP, Clyde K, Simons L, Ho K, Bains D. High serum cholesterol and atherosclerosis do not contribute to increased arterial stiffening with age. *J Am Coll Cardiol.* 1986;7(suppl A):247A. Abstract.
52. Megnen JL, Simon A, Denarie N, Del Pino M, Garipey J, Segond P, Levenson J. Aortic stiffening does not predict coronary and extracoronary atherosclerosis in asymptomatic men at risk from cardiovascular disease. *Am J Hypertens.* 1998;11:293-301.