UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Dendritic cells in T- and B-cell proliferation in the skin

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Dendritic cells in T- and B-cell proliferation in the skin / PIMPINELLI N.; M. SANTUCCI; ROMAGNOLI P.; GIANNOTTI B.. - In: DERMATOLOGIC CLINICS. - ISSN 0733-8635. - STAMPA. - 12:(1994), pp. 255-270.

Availability:

This version is available at: 2158/223194 since:

Publisher:

W B Saunders Company:Fulfillment Department, The Curtis Center, Independence Square West:

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

DENDRITIC CELLS IN T- AND B-CELL PROLIFERATION IN THE SKIN

Nicola Pimpinelli, MD, PhD, Marco Santucci, MD, Paolo Romagnoli, MD, and Benvenuto Giannotti, MD

ANATOMY, PHYSIOLOGY, AND DISTRIBUTION OF DENDRITIC CELLS IN NORMAL SKIN

Dendritic cells (DC), otherwise known as *dendritic leukocytes* or *dendrocytes*, represent a relatively homogeneous population of non-lymphocytic cells within lymphoid organs, epithelia, connective tissue, and lymph.⁸⁵ The main, common features of DC are their typical dendritic shape (from which their name, although devoid of functional implication, is derived); bone-marrow origin; a constitutive expression of major histocompatibility complex (MHC) class II antigens; usually low phagocytic activity; and high efficiency in stimulating (hence the definition *antigen-presenting cells*, APC) and regulating cell-mediated immune responses.¹¹⁸ Ultrastructurally, DC share some basic features: a roundish to oval body and variably long, thin cytoplasmic branches (dendrites); an indented nucleus with pale chromatin (excluding a thin peripheral rim); and a cytoplasm containing cytoskeletal filaments (but no fibrils), mitochondria, few stacks of rough endoplasmic

reticulum, many smooth vesicles and tubules, a large Golgi apparatus, and few dense bodies (primary lysosomes).⁸⁵ This group comprises interdigitating reticulum cells (T-cell areas) and dendritic reticulum cells ([DRC] B-cell areas) of secondary lymphoid organs, so-called *veiled cells* in the afferent lymph, Langerhans cells (LC) and other dendritic leukocytes of squamous and nonsquamous stratified epithelia, and interstitial dendritic leukocytes.

Cutaneous LC⁶⁹ are the most well-known and extensively studied DC. Their histogenetic, morphologic, antigenic, and functional features, as well as their alteration by aging and physical-chemical stimuli and their involvement in pathology, have been extensively reviewed elsewhere^{19, 24, 95, 100, 105} and will not be the subject of this article. In extreme synthesis, the cardinal features of LC are the presence of typical cytoplasmic inclusions (Birbeck granules, shown by electron microscopy)¹⁷ and the expression of CD1a antigen on their plasma membranes. According to a strict definition, *only cells containing Birbeck granules should be considered LC*. It has been widely demonstrated that LC acquire their complete morphologic features, as judged from the presence and number of Birbeck granules, only following their establishment in the epidermis^{10, 23, 81, 91}; for this reason, epidermal and dermal DC not containing Birbeck

The original research of the authors in this field was supported by grants from the Italian Ministry of University and Scientific and Technologic Research (University funds, 40% and 60%), and Italian National Research Council (grant no. 92.01087.04).

From the Dermatology Clinic II (NP, BG), Institute of Anatomic Pathology (MS), and Departments of Histology, Cytology, and Embryology (PR), University of Florence School of Medicine, Florence, Italy

DERMATOLOGIC CLINICS

granules have been interpreted as putative precursors of LC and termed *indeterminate cells*, for almost 15 years.^{23, 51, 101} Indeed, the apparent lack of Birbeck granules may be related either to a loss of these typical structures following the migration of LC from the epidermis into the dermis or to an absent formation due to the lack of—or minimal—homing in the epidermal microenvironment.^{10, 23, 51, 101} The findings of recent reports,¹² however, suggest that the identification of Birbeck granules may also depend on the careful evaluation of many serial ultrathin sections. Consequently, the term *indeterminate cells* seems today a little bit confusing, and many researchers prefer to avoid its use.

In addition to their identificative antigenic marker (i.e., CD1a), LC constitutively express many other membrane molecules that may be related to their functional role in physiology and pathology (e.g., antigen presentation and T-cell activation, cytokine production, adhesion to the extracellular matrix, and migration).^{19, 74, 95, 100, 105} The expression of these functionally relevant molecules by LC can be either up regulated or down regulated in specific conditions, with possible clinical consequences. These features support the interpretation of these cells as the "sentinels" of the skin-associated lymphoid tissue (SALT)¹¹⁹ or skin immune system (SIS).¹⁸

A novel type of DC has been described in the skin and increasingly investigated in the last years, that is, dermal dendritic cells. Some common, basic features of these dermal DC, variably termed as *dermal dendrocytes* ([DD] the most popular definition),^{3, 30, 40, 58, 59} *perivascular veil cells*,²² or *perivascular dendritic macrophages*,¹³ are the expression of class II MHC antigens, a mainly perivascular location (predominantly around the superficial vascular plexus), the lack of expression of CD1a antigen and vascular markers (factor VIII-RA, Ulex Europaeus lectin), and the ultrastructural features of macrophages. The antigenic phenotype of DD, however, is actually a matter of controversy; in fact, despite the substantial agreement concerning their interpretation as a subset of CD68+ dermal macrophages, they have been variably defined as factor XIII+^{3, 30, 40, 58, 59} CD11b- / D11c+ / D36-¹¹³ CD11b+ / CD11c+ / CD14+ / KiM8+ / D36-¹²⁹ or CD11b- / CD11c+ / CD14+ / D36+^{30, 58, 59, 125} Thy-1+^{43, 44} (Thy-1 is a human antigen, possibly related to the immunoglobulin supergene family as well as its murine counterpart⁷⁷; however, human Thy-1

is expressed by dermal perivascular DC and not by epidermal DC in both normal and pathologic skin.^{43, 44}) The role of these resident dermal DC in murine and human SIS is also controversial: some experimental evidences suggest that they are a further cell type, in addition to LC, capable of initiating delayed-type hypersensitivity responses.^{34, 120, 125} Other studies, on the contrary, support the hypothesis that DD may be responsible for the activation of suppressor pathways,^{43, 72} similar to a CD36+ subset of blood monocytes.¹⁰⁹ At the present state of our knowledge, it is not possible to hypothesize whether the CD36+ DC identified in normal dermis correspond to those demonstrated in UV-irradiated skin,^{4, 5, 32, 33} which are capable of activating suppressor cell responses.

A specific subset of DC, with features closely resembling those of cutaneous DD, has been described in the lamina propria of the oral mucosa.⁷⁴ These cells show a typical perivascular location and a CD36+ / CD1a- / CD11b- / CD11c- antigenic phenotype; their ultrastructural profile is characterized by the presence of many lysosomes (both primary and secondary, often containing melanin), well-developed rough endoplasmic reticulum, and focal adhesion sites to the extracellular matrix.

CD36+ / CD1a- DC have been found in normal epidermis,^{114, 114} although not regularly.⁷² It remains to be established whether and how these CD36+ epidermal DC, when present, are related to those found in the dermis. The identification of all CD36 expressing cells as monocyte-derived APC seems at present to be unwise. In fact, CD36+ cells—variably dendritic in shape but not expressing class II MHC antigens or CD45 antigen (the latter indicative of bone marrow origin)—have been found by some authors at the dermoepidermal junction.¹²⁹

THE IMPORTANCE OF STUDYING DENDRITIC CELLS IN LYMPHOPROLIFERATIVE DISORDERS OF THE SKIN

In lymphoproliferative disorders of the skin, lymphocytes are the proliferating cells and DC may be considered their "living room," that is, the framework in which lymphocytes live and proliferate. Therefore, why study DC? This is an obvious, reasonable

question. First of all, DC have a key role in T- and B-cell homing.^{85, 118} Second, DC can be the specific target of living and physical and chemical agents, and their alterations may have a pathogenic role in certain cutaneous lymphoproliferative diseases.^{50, 74, 75, 110} Third, the morphologic, antigenic, and topographic features of DC may have diagnostic and histogenetic relevance in specific conditions⁹⁵ as discussed later on. Finally, DC are crucial for both effector and regulatory immune responses,¹¹⁸ and might play a role in antitumoral lymphocyte reactions.⁵⁷ In addition, investigating the morphologic, antigenic, and functional features and the distribution of DC in different cutaneous pathologic skin models characterized by aberrant lymphoid cell migration and proliferation may greatly contribute to a better understanding of the differentiation and function of SIS and the pathogenesis of SIS-related disorders.⁹⁵

Concerning cutaneous lymphomas, that is, non-Hodgkin's lymphomas primarily presenting in the skin, the significance and possible role of DC in the natural history of the disease have been subjects of investigation for almost 20 years, with no definitive conclusions reached at present.

DENDRITIC CELLS IN T-CELL PROLIFERATIVE DISORDERS

Since 1976,^{20, 55, 56} the presence of DC in mycosis fungoides (MF), a classic type of cutaneous T-cell lymphoma (CTCL), has been described by means of electron microscopy and considered the specific expression of lymphocyte homing. This finding has been widely confirmed by numerous immunohistochemical and ultrastructural studies* in MF, Sézary syndrome (SS, which is interpreted as the leukemic variant of MF), and localized pagetoid reticulosis.

In fact, different hypotheses have been proposed concerning the significance and possible role of DC in the pathogenesis of MF. The presence of viruslike particles in the cytoplasm of LC induced some authors to hypothesize a viral infection of these APC, with the consequence of their persistent stimulatory action on CD4+ T cells, eventually leading to the malignant transformation of CD4+ T cells.^{50, 74, 75, 110, 127} The report that only intraepi-

dermal T cells are cycling in MF⁸³ supports the hypothesis of a possible significant role of epidermal LC in the pathophysiology of MF. Other studies, however, have stressed that close contacts between LC and T cells are occasional, even in initial patches, notwithstanding the high numbers of epidermal CD1a+ LC⁷⁻¹¹; these findings raise doubts on this hypothesis of a persistent stimulation of T cells by LC as a crucial event in the pathogenesis of MF. Conversely, a clear-cut correlation between the number, immunomorphology, and ultrastructure of dermal CD1a+ DC (indeterminate cells) and the clinical stage-related histopathologic modifications has been documented.^{9, 10} In early patches of MF (flat lesions, with a scarce, mostly perivascular lymphoid infiltrate), indeterminate cells are relatively few, intermingled with high numbers of CD11c+ /CD1a- monocytoïd cells (putative precursors?),¹⁰ and form a rudimentary cellular network, establishing close contacts between each other and with cerebriform lymphocytes: according to our initial paradox, lymphocytes "reclaim their living room." In the plaque stage, indeterminate cells are more numerous, form an extended cellular network, show morphologic signs of enhanced functional activity (abundant cytoplasm with large amounts of mitochondria, multiple Golgi stacks, and numerous cisternae of rough and smooth endoplasmic reticulum), and establish close contacts with CD4+ T cells, often showing blastic morphology and an activated-proliferating phenotype (HLA-DR+ /CD25+ /CD71+ /Ki-67+)⁹ (Figs. 1 to 4). In the tumor stage, blastic T cells markedly increase in number and lose their mature CD4+ phenotype, whereas indeterminate cells are sparse and less rich in organelles and show only occasional contacts with T cells.⁹ On the basis of these findings, it has been suggested that dermal CD1a+ DC are possibly involved in the homing, proliferation, and neoplastic progression of T cells.⁹ On the other hand, CD1a+ DC may differentiate and persist only under the influence of a CD4+ T-cell microenvironment.^{9, 10}

If the paucity of close contacts between lymphocytes and epidermal LC seems to speak against a crucial role of epidermal LC in the pathogenesis of MF, how can the constant finding of large numbers of these cells in the epidermis of MF patches and plaques be interpreted? It could be considered an epiphenomenon related to the migration of CD11c+ and CD1a- monocytoïd precursors

*References 9-11, 21, 26, 31, 36, 41, 50, 61, 63, 65, 67, 68, 74-76, 99, 110, 112, 123

Figure 1. MF (plaque stage). Epidermotropic, bandlike infiltration of CD4⁺ T cells. The phenotype of these cells is HLA-DR⁺, CD25⁺, CD71⁺, Ki-67⁺. Immunohistochemical technique used (APAAP) on frozen section (original magnification $\times 160$).

to DC from the dermis into the epidermis,^{8,10} possibly due to the increased release of cytokines¹²¹ (the role of cytokines is extensively dealt with in the article by Dummer and Schwarz). In the epidermis, these precursor cells eventually acquire the typical antigenic (CD1a expression), morphologic (Birbeck granules), and, probably, functional features of fully mature LC.^{8,10} According to the aforementioned data, the recent report of a positive correlation between higher numbers of epidermal CD1a⁺ LC and a better prognosis in patients with MF⁷⁹ might simply depend on a correlation between the numbers of epidermal LC and the disease stage. The presence of skin tumors, in fact, is accompanied by a paucity of epidermal CD1a⁺ DC and is related to a poor prognosis. The hypothesis that epidermal LC alterations are a consequence, rather than a cause, of MF lesions is supported by the finding that there are no significant differences in the number and distribu-

tion of these cells between the clinically normal skin of MF patients and that of healthy control subjects.

Similar to S100⁺/CD1a⁺ DC, factor XIIIa⁺ DD coexpressing Thy-1 antigen are constantly found in the papillary dermis of MF lesions and progressively increase in numbers from early patches to plaques.^{43,44} The increase of factor XIIIa⁺ and Thy-1⁺ DC, possibly neodifferentiated from a pool of CD34⁺ progenitors that have migrated into the papillary dermis from the reticular dermis—where they are usually found in perivascular location⁴⁴—and recruited from the circulation via adhesion to ELAM-1⁺/ICAM-1⁺/VCAM-1⁺ endothelium,⁴⁴ seems to counterbalance the increase of CD1a⁺ DC in MF. This same situation has been observed in some widely studied inflammatory dermatoses, such as lichen planus (LP) or atopic dermatitis.³⁰ Factor XIIIa⁺, CD36⁺ DD could be crucial in the selective activation of sup-

Figure 2. MF (plaque stage). Extended network of CD1a+ dendritic cells. These cells are very numerous in both the epidermis and the dermis. Immunohistochemical technique used (APAAP) on frozen section (original magnification $\times 160$).

pressor-inducer pathways that, if overstimulated, may down regulate host immune responses leading to suppression of antitumor immunity.^{43, 72} Factor XIIIa+ and CD1a+ DC³⁹ and CD36+ /CD1a- DC^{57, 71} have been found in MF lesional epidermis; no definite hypothesis can be proposed yet on a role of these cells in the disease. The possible significance of DC in the pathogenesis of CTCL is discussed in detail in another article of this issue.

The presence and possible significance of DC in non-MF/S CTCL have not been extensively studied. CD1a+ DC have been observed in low numbers and scattered distribution in both CD30(Ki-1)+^{6, 13} and CD30- large T-cell lymphomas^{13, 47} primarily presenting in the skin, but no conclusion can be drawn concerning the possible relationship between the characteristics of DC and the clinicobiologic features of these distinctive

subtypes of CTCL (e.g., the DC pattern of regressing skin lesions in CD30+ CTCL is presently not known).

Some recent reports specifically address clinical, histologic, and immunohistologic features possibly useful for the differential diagnosis between CTCL and pseudo-CTCL.⁹⁸ Specifically concerning dendritic cells, the dermal infiltrate of MF-like pseudo-CTCL (characterized by a bandlike, epidermotropic infiltration of the skin by cerebriform T cells) seems to contain a percentage of CD1a+ DC lower than that commonly found in plaque-stage MF.⁹⁸ It should be interesting to compare MF and drug-induced pseudo-MF on electron microscopy to verify whether some differences do exist comparable to those found between MF plaques and well-developed lesions of LP, a classic model of immune cell-mediated, benign dermatosis.⁸⁹ In both cases, DC form an extended cellular network

Figure 3. MF (plaque stage). An indeterminate cell (ic, dendritic cell of Langerhans lineage, but without identifiable Birbeck granules) is in close contact with a lymphoid cell undergoing clear-cut blastic transformation (bc). Electron micrograph (original magnification $\times 11,000$).

in the dermis and establish close appositions with activated, cerebriform T cells. In LP, however, neither DC showing cytomorphologic signs of activation nor lymphocytes with blastic phenotype are found, thus suggesting that a "physiologic" mechanism of lymphocyte activation is operating in LP. A crucial difference between pseudo-CTCL, histologically characterized by a nodular pattern mimicking that found in pleomorphic CTCL, and true pleomorphic CTCL is the abundance of histiocytic cells in the former;⁸⁸ no data on the immunophenotype of these cells are currently available.

DENDRITIC CELLS IN B-CELL PROLIFERATIVE DISORDERS

Dendritic Cells in Normal and Neoplastic Lymphoid Tissue

DRC, whose myelomonocytic origin has been only recently demonstrated unequivocally,⁴⁵

are DC restricted to the B-cell areas of secondary lymphoid tissue (e.g., lymph nodes, spleen, tonsils, Peyer's patches), namely, primary and secondary follicles.¹²⁷ For this reason, they are also commonly known as *follicular dendritic cells*.^{88, 116, 127} They are not identifiable in extrafollicular areas of the lymphoid tissue or in normal extranodal tissues (thymus, liver, skin, or kidney).¹³⁸ The detection of these cells by conventional light microscopy is extremely difficult and is made possible by enzyme cytochemistry (5-nucleotidase activity⁸⁰), immunohistochemistry⁸⁸ (see following), and electron microscopy.^{46, 87} The immunophenotype of DRC may be difficult to assess in situ, owing to the close association between them and surrounding B cells.¹⁰¹ Therefore, immunophenotyping of DRC has been performed in single cell suspensions, giving controversial results that may be related to the heterogeneity of these cells,⁸⁸ as discussed further on. In this article, the anatomy and physiology of B-cell areas of

Figure 4. MF (plaque stage). Dermal indeterminate cell showing clear-cut morphologic signs of enhanced functional activity: abundant cytoplasm with large amounts of mitochondria (arrow), multiple Golgi stacks (g), and very high density of tubules and vesicles (arrowhead). Electron micrograph (original magnification $\times 16,000$).

lymphoid tissue cannot be analyzed extensively, and more details can be found in recently published reviews⁸⁸; nevertheless, a simplified description seems necessary to clarify as much as possible the discussion concerning the significance and possible role of DRC in B-cell lymphoproliferative disorders.

In normal lymphoid tissue, DRC form a dendritic network in both primary follicles (in which no antigen-driven processes are taking place) and secondary follicles ("sanctuary" of B-cell differentiation, activation, and proliferation).⁷³ Concerning their functional significance, a specific role of DRC in B-cell homing is suggested by the expression of VCAM-1 antigen by these cells and of VLA-4 antigen (ligand to VCAM-1) by B lymphocytes.^{48, 49} In addition, recent studies⁸⁹ suggest that DRC have a significant role in the "trapping" and presentation of antigens (in the form of membrane-bound immune complexes) to antigen-specific B cells, with consequent differentiation, activation, and proliferation.

In primary follicles, DRC form a rudimentary cell network.^{73, 88, 127} In secondary follicles, DRC have different features according to their localization. In the germinal center, DRC form a dense, sharply demarcated dendritic web, closely associated with CD10+ B cells (follicle-

ular center cells, [FCC]).^{73, 88, 116, 127} On electron microscopy, in addition to the aforementioned common features of DC, DRC show characteristic narrow plasma membrane infoldings coated with electron dense material, presumably representing fixed immunocomplexes^{46, 87} (see following). The immunophenotype of DRC in the germinal center is fairly constant⁸⁸: DRC-1 (R4/23) +, CD9+, CD11b (C3bi receptor) +, CD14+, CD21 (C3d receptor) +, CD23 (low-affinity receptor for IgE) +, CD35 (C3b receptor) +, CD54 (ICAM-1) +, VCAM-1+, HLA-DR+, acid cysteine-proteinase inhibitor (ACPI) +,² nerve growth factor receptor (NGFr) +,¹²¹ and IgG, M, and A+. In non-Hodgkin's lymphomas of FCC origin (known as FCC lymphoma or follicular lymphoma^{70, 88, 116}), showing bcl-2 gene rearrangement^{1, 126} related to a chromosomal translocation (t(14;18)),¹³² DRC are a constant finding. They usually retain most of the morphologic and antigenic features found in the normal germinal center, but specifically lack some of them: the electron dense coating of membrane infoldings,⁸⁷ membrane-bound immunoglobulins,^{88, 116} and ACPI- and NGFr expression.¹²¹ Notwithstanding these morphologic and antigenic aberrations, the capacity to favor B-cell homing and to create an ade-

Figure 5. Dermal infiltrate of non-Hodgkin's B-cell lymphoma. Two dendritic reticulum cells (stars), one of which presumably binucleated (black star), establish close contact with blast cells showing a centrocytelike (cc) or centroblastlike (cb) nuclear morphology. Note the absence of electron dense material between dendritic cells and neoplastic lymphoid cells. Electron micrograph (original magnification $\times 8,500$).

quate microenvironment for the proliferation of B cells seems to be retained in neoplastic conditions, confirming that DRC are a necessary component for B-cell proliferation and suggesting that they can provide a costimulatory signal for neoplastic lymphocytes.⁵⁸

In the follicle mantle (the external portion of secondary follicles), both normal and neoplastic DRC are more loosely arranged^{58, 116} and show some significant immunophenotypic differences as compared with those found in the germinal center. In fact, they are CD9-, CD11b+/-, CD14-, CD23-, and IgD+.^{28, 29, 54} No lack of specific antigenic markers has been reported so far in lymphomas of follicle mantle origin.^{28, 29, 54, 88, 116, 130, 131} In this external portion of secondary follicles, DRC are associated with two main B-cell populations: CD5+ / CD10- B cells, and CD5- / CD10- B cells. The former—mantle zone lymphocytes,^{130, 131} or mantle cells^{70, 58}—may represent the step of B-cell differentiation next to FCC.⁷³ The CD5- and CD10- cells are particularly well represented in the spleen, where they (*marginal cells*⁷³) form a distinct area of the follicle mantle (*marginal zone*), external to that populated by mantle cells (mantle zone). In the lymph node, the CD5- /

CD10- B-cell population (parafollicular cells^{35, 37, 82}) grow and become clearly evident in some reactive (immature sinus histiocytosis^{36, 86, 117}) and neoplastic conditions (known as *parafollicular lymphoma*^{35, 37} or *monocytoid B-cell lymphoma*^{82, 96, 107, 108}). This parafollicular cell might be interpreted as a B cell that has undergone T-cell-dependent antigen priming and is putatively capable of differentiating into various morphologic subtypes (FCC, mantle cells, plasmacytic cells). This hypothesis is supported by some characteristic features of parafollicular-monocytoid lymphomas. In this peculiar type of lymphoma, typical parafollicular-monocytoid cells are very frequently admixed with other, often monoclonal, cell types: plasmacytic cells (sometimes with transitional forms to typical parafollicular-monocytoid cells) and/or FCC cells and/or mantle cells (often in small clusters), indeed a spectrum of distinct cytologic types originating from one cell, the parafollicular-monocytoid cell.^{35, 37, 82, 96, 107, 108} These characteristic features, along with the typical finding of "lymphoepithelial lesions" (clusters of neoplastic B cells infiltrating glandular ducts), have been widely described also in primary extranodal B-cell lymphomas,

Figure 6. Primary CBCL. Folliclelike nodules of CD22+ B cells in the superficial dermis of recently developed lesions. These cells constantly show a CD5⁺, CD10⁻, slg⁺ (monoclonal) phenotype. Immunohistochemical technique used (APAAP) on frozen section (original magnification $\times 250$).

namely, the so-called MALT lymphoma (lymphoma of the mucosa-associated lymphoid tissue).^{66, 84}

Dendritic Cells in Cutaneous B-cell Proliferative Disorders

Since 1981, numerous studies have demonstrated that cutaneous B-cell proliferative disorders are often characterized by the presence of a specialized microenvironment of DRC. The presence of DRC in the skin has been considered specifically related to the FCC homing and, consequently, to the development of "pseudolymphomatous" FCC reactions* or to the localization (primary† or secondary^{25, 26, 51, 63, 90, 112, 133}) of neoplastic FCC (FCC lymphoma, follicular lymphoma), independent of the histologic evidence of follicles. According to their ultrastructure, DRC found in the skin infiltrate of FCC lymphomas show

a limited degree of morphologic differentiation⁹⁰ (Fig. 5), similar to that described in nodal lymphomas.⁸⁷

In a recently published study,¹⁰² we have analyzed—retrospectively and prospectively—a large group of patients with unequivocal diagnosis of primary cutaneous B-cell lymphoma (CBCL). The diagnosis was based on negative staging, positive staining for B-cell restricted antigens CD19/CD20/CD22, and the surface Ig light chain monoclonality of neoplastic B cells, this latter feature being essential for the differentiation from pseudolymphoma.⁹⁷ According to our data, CBCL show a constellation of features strikingly similar to those described in parafollicular-monocytoid lymphoma^{35, 37, 82, 96, 107, 108} and MALT lymphoma^{66, 84}: (1) nonaggressive clinical behavior (preferentially locoregional extension, good response to local treatments, low tendency to spread, and excellent prognosis despite frequent relapses); (2) uniform immunophenotypic (CD5⁻, CD10⁻)^{14-16, 53, 93, 102, 104} and genotypic features of neoplastic B cells (lack of bcl-2 gene rearrangement),^{14, 15, 139} despite the wide variability of cytomorpho-

*References 21, 25, 26, 41, 52, 64, 97, 112, 136, 137, 139

†References 21, 25, 26, 41, 51, 63, 92, 97, 112, 133-135, 139

Figure 7. CBCL. DRC-1+ dendritic reticulum cells are clustered with neoplastic B cells (see Figure 6, serial section) in the superficial dermis of recently developed lesions. These cells show an aberrant "centrifugal" pattern and characteristically have a CD14- phenotype. Immunohistochemical technique used (APAAP) on frozen section (original magnification $\times 250$).

logic appearances (the whole spectrum of perifollicular cells, admixed with small lymphocytes, plasma and lymphoplasmacytoid cells); and (3) lymphoepithelial lesions. These features suggest the interpretation of CBCL as the cutaneous counterpart of MALT lymphoma, that is, *skin-associated lymphoid tissue-related B-cell lymphoma*,^{53, 93, 102, 104} and lead to hypothesizing their perifollicular origin.¹⁰⁴

The systematic analysis of biopsies sequentially taken from these patients allows delineation of the histologic and immunohistologic spectrum of CBCL, whose different aspects are largely related to the age and growth rates of skin lesions,¹⁰² without any correlation with clinical course and prognosis (which are good overall). This spectrum ranges from the sometimes inconspicuous "top heavy" infiltrate of intermediate-sized neoplastic B cells (resembling small perifollicular cells^{35, 82}) and many reactive T cells observed in early lesions ("low-grade" malignancy picture, classically considered typical of cutaneous pseudolymphomas^{21, 25-27, 42, 63, 64, 136, 137}) to the uniformly heavy dermal infiltrate of large neoplastic B

cells ("high-grade" malignancy picture) characteristic of late, rapidly growing lesions.

In CBCL, DRC are present in three main fashions^{92, 93, 103-104}: (1) associated with CD5- and CD10-, monoclonal, proliferating (Ki-67+ and CD71+) B cells to form folliclelike clusters. These B-cell DRC clusters are mainly found in the superficial dermal infiltrate of recently developed lesions and are irregularly surrounded by reactive T cells and associated CD1a+ DC, thus delineating a quite distinct compartmentalization of the infiltrate not identifiable on histologic grounds alone (Figs. 6 to 8). In this instance, DRC have a DRC-1+ /NGFr+ /CD14- /CD21(+)- /CD35+ phenotype¹⁰²⁻¹⁰⁴ and often show an aberrant "centrifugal" pattern (loosely arranged in ill-defined meshworks with blurred and radiating contours, and skipping the nodule centers)¹⁰² (Fig. 7); (2) scattered among neoplastic B cells, which are intermingled with high numbers of reactive T cells without a clear-cut compartmentalization in the upper and mid-dermis. In this instance, typical of slowly grown, long-standing lesions, DRC

Figure 8. CBCL. CD3+ T cells are abundant and irregularly surround the DRC-B-cell clusters (Figures 6, 7, serial section) in the superficial dermis of recently developed lesions. Immunohistochemical technique used (APAAP) on frozen section (original magnification $\times 250$).

have the same DRC-1+ /NGFr+ /CD14- /CD21(+)- /CD35+ phenotype; (3) sited in reactive, polyclonal lymphoid follicles, mainly located in the mid-lower reticular dermis of slowly grown, long-standing lesions. These follicles more frequently show a morphologically and immunologically typical organization, that is, a lightly staining germinal center (composed of CD10+ B cells and CD14+ DRC) and a darkly staining mantle (composed of CD5+ /CD10- and CD5- /CD10- B cells and CD14- DRC)¹⁰²⁻¹⁰⁴ (Fig. 9) and are surrounded by broad strands of neoplastic B cells in a fashion reminiscent of a parafollicular and interfollicular pattern typically observed in parafollicular-monocytoid lymphoma.¹⁰² Parafollicular cells are frequently associated with variable numbers of plasma cells and lymphoplasmacytoid cells, often monoclonal, and a morphologic continuum between parafollicular cells and plasmacytic cells is sometimes evident.^{93,102,104} Features resembling lymphoepithelial lesions observed in MALT B-cell lymphoma⁶⁶ and other extranodal parafollicular-monocytoid lymphoma,^{82, 84} namely, neoplastic cells infiltrat-

ing the ductal portion of eccrine sweat glands and the pilosebaceous units, are typically, yet rarely, found.^{93, 102} Summarizing, the presence of reactive, polyclonal, typically organized secondary follicles is not necessarily indicative of a reactive, pseudolymphomatous process, and particular attention has to be paid to the features of surrounding B cells: if the latter are virtually all polyclonal, the finding is indicative of pseudolymphoma; on the contrary, if B cells outside the polyclonal follicles are monoclonal (associated or not with sparse polyclonal cells), the finding is definitely suggestive of CBCL.

From a speculative point of view, the evolutionary modifications of skin lesions support the hypothesis that—at the very beginning—DRC favor the homing and proliferation of neoplastic B cells. The neoplastic growth of these B cells is long contrabanced by a heavy T-cell reaction (also supported by a specific, CD1a+ DC microenvironment) and only late does this control becomes ineffective. Interestingly, however, this neoplastic progression is not accompanied by a parallel worsening of the clinical behavior, which is

Figure 9. CBCL. CD35+ DRC in the germinal center of a reactive, polyclonal follicle in the mid-lower reticular dermis of a slowly grown, long-standing lesion. These follicles more frequently show a morphologically and immunologically typical organization: CD10+ B cells and CD14- DRC in the germinal center, and CD5+ /CD10- (mantle cells) or CD5- /CD10- B cells (marginal cells) in the mantle. Immunohistochemical technique used (APAAP) on fixed-embedded material (original magnification $\times 250$).

and remains overall very good. The possible explanation for this discrepancy is given by the putative parafollicular origin of CBCL; in fact, parafollicular-monocytoid nodal lymphoma, MALT lymphoma, and CBCL share nonaggressive clinical behavior and uniform immunophenotype and genotype of neoplastic B cells (CD5- /CD10-; lack of bcl-2 rearrangement).

SUMMARY

In lymphoproliferative diseases of the skin, DC have a key role in T- and B-cell homing. Furthermore, DC alterations may have a pathogenic role in the natural history of specific disorders, either in the neoplastic lymphoid cell progression or in antitumoral lymphocyte reaction. Finally, the morphoantigenic and topographic features of DC may have diagnostic and histogenetic relevance in specific conditions.

In CTCL, dermal CD1a+ DC ("indeterminate cells") seem to play a significant role in the neoplastic progression of MF, whereas the possible pathogenetic role of specific alterations of epidermal LC is yet to be proven. Recently, a possible implication of DD (resident, perivascular factor XIIIa+ /CD1a- DC)

in the pathogenesis of MF has been also suggested. The presence and possible significance of DC in CTCL non-MF are presently poorly studied. At present, DC number, distribution, and phenotype seem possibly useful in the differential diagnosis between CTCL and pseudo-CTCL, but this hypothesis has to be adequately confirmed.

CBCL has been recently proposed as a unique type of clinically low-grade lymphoma, namely, skin-associated lymphoid tissue (SALT)-related B-cell lymphoma. Both SALT- and mucosa-associated lymphoid tissue (MALT)-related B-cell lymphoma share with a peculiar nodal lymphoma of follicle mantle origin (parafollicular-monocytoid lymphoma) the nonaggressive clinical behavior and the uniform phenotype (CD5-, CD10-) and genotype (lack of bcl-2 gene rearrangement) of neoplastic B cells, despite the wide variability of cytomorphologic appearances. The putative origin of CBCL is further supported by the typical CD14-, nerve growth factor receptor (NGFr) immunophenotype of DRC. Moreover, the immunophenotype and architectural fashion of DRC are interesting clues to the differentiation between neoplastic and true reactive folliclelike nodules and may be of help in the differential diagnosis between CBCL and B-cell pseudo-lymphoma as well as in the correct interpre-

tation of lesions showing monoclonal proliferations of B cells accompanied by polyclonal follicular reactions.

References

1. Aisenberg AC, Wilkes BM, Jacobson L: The bcl-2 gene is rearranged in many diffuse B-cell lymphomas. *Blood* 71:969, 1988
2. Alavajkko M, Ajne R, Rinne A, et al: Behaviour of dendritic reticulum cells possessing immunoreactive acid cysteine-proteinase inhibitor in human lymphoid secondary follicles and in follicular centre-cell lymphomas. *Int J Cancer* 35:319, 1985
3. Arrese Estrada J, Piérard GE: Factor-XIIIa-positive dendrocytes and the dermal microvascular unit. *Dermatologica* 180:51, 1990
4. Baadsgaard O, Fox DA, Cooper KD: Human epidermal cells from ultraviolet light-exposed skin preferentially activate autoreactive CD4+2H4+ suppressor-inducer lymphocytes and CD8+ suppressor/cytotoxic lymphocytes. *J Immunol* 140:1738, 1988
5. Baadsgaard O, Salvo B, Mannie A, et al: In vivo ultraviolet-exposed human epidermal cells activate T suppressor cell pathways that involve CD4+CD45RA+ suppressor-inducer T cells. *J Immunol* 145:2854, 1990
6. Banerjee SS, Heald J, Harris M: Twelve cases of Ki-1 positive anaplastic large cell lymphoma of skin. *J Clin Pathol* 4:117, 1991
7. Bani D, Moretti S: Are Langerhans cells usual components of the dermal infiltrate of mycosis fungoides? *Arch Dermatol Res* 279:561, 1987
8. Bani D, Moretti S, Pimpinelli N, et al: Differentiation of monocytes into Langerhans cells in human epidermis. An ultrastructural study. *In* Thivolet J, Schmitt D (eds): *The Langerhans Cell*. London, Montrouge, John Libbey Eurotext, Coll. INSERM vol. 172, 1988, p 75
9. Bani D, Moretti S, Pimpinelli N, et al: Interdigitating reticulum cells in the dermal infiltrate of mycosis fungoides. An ultrastructural and immunohistochemical study. *Virchows Arch A (Pathol Anat Histopathol)* 412:451, 1988
10. Bani D, Giannotti B: Differentiation of interdigitating reticulum cells and Langerhans cells in the human skin with T-lymphoid infiltrate. An immunocytochemical and ultrastructural study. *Arch Histol Cytol* 52:361, 1989
11. Bani D, Pimpinelli N, Moretti S, et al: Langerhans cells and mycosis fungoides: A critical overview of their role in the disease. *Clin Exp Dermatol* 15:7, 1990
12. Bartosik J: The non-keratinocytes in normal epidermis. *Eur J Dermatol* 1:131, 1991
13. Beljaards RC, Meijer CJLM, Scheffer E, et al: Prognostic significance of CD30 (Ki-1/Ber-H2) expression in primary cutaneous large cell lymphomas of T-cell origin. A clinicopathologic and immunohistochemical study of 20 patients. *Am J Pathol* 135:1169, 1989
14. Berti E, Alessi E, Caputo R, et al: Reticulohistiocytoma of the dorsum. *J Am Acad Dermatol* 19:259, 1988
15. Berti E, Gianotti R, Alessi E, et al: Primary cutaneous follicular center cell lymphoma: Immunophenotypic and immunogenotypical aspects. *In* van Vloten WA, Willemze R, Lange Vojlsgaard G, et al (eds): *Cutaneous Lymphoma*. Basel, Karger, 1990, p 196
16. Berti E, Alessi E, Caputo R: Reticulohistiocytoma of the dorsum (Crosti's disease) and other B-cell lymphomas. *Semin Diagn Pathol* 8:82, 1991
17. Birbeck MSC, Breathnach AS, Everall JD: An electron microscopic study of basal melanocytes and high level clear cells (Langerhans cells) in vitiligo. *J Invest Dermatol* 37:51, 1961
18. Bos JD, Kapsenberg ML: The skin immune system (SIS): Its cellular constituents and their interactions. *Immunol Today* 7:235, 1986
19. Bos JD, Teunissen MBM, Kapsenberg ML: Dendritic cells of the skin immune system (SIS). *In* Thivolet J, Schmitt D (eds): *The Langerhans cell*. London, Montrouge, John Libbey Eurotext, Coll. INSERM, vol. 172, 1988, p 9
20. Braun-Falco O, Schmoeckel C, Wolff HH: The ultrastructure of mycosis fungoides, of Sézary syndrome and of Wöringer-Kolopp's disease (pagetoid reticulosis). *Bull Cancer* 64:191, 1977
21. Braun-Falco O, Burg G, Schmoeckel C: Recent advances in the understanding of cutaneous lymphoma. *Clin Exp Dermatol* 6:89, 1981
22. Braverman JM, Sibley J, Keh-Yen A: A study of the veil cells around normal, diabetic, and aged cutaneous microvessels. *J Invest Dermatol* 86:57, 1986
23. Breathnach AS: Variations in ultrastructural appearance of Langerhans cells of normal human epidermis. *Br J Dermatol* 97(suppl 15):14, 1977
24. Breathnach AS: The Langerhans cell. *Br J Dermatol* 119:463, 1988
25. Burg G, Braun-Falco O, Hoffmann-Fezer G, et al: Differentiation between pseudolymphomas and malignant B-cell lymphomas of the skin. *In* Goos M, Christophers E (eds): *Lymphoproliferative Disease of the Skin*. Berlin, Springer-Verlag, 1982, p 101
26. Burg G, Braun-Falco O: *Cutaneous Lymphomas, Pseudolymphomas, and Related Disorders*. Berlin, Springer-Verlag, 1983
27. Burke J: Malignant lymphomas of the skin: Their differentiation from lymphoid and non-lymphoid cutaneous infiltrates that simulate lymphoma. *Semin Diagn Pathol* 2:169, 1985
28. Carbone A, Manconi R, Poletti A, et al: Heterogeneous immunostaining patterns of follicular dendritic reticulum cells in human lymphoid tissues with selected antibodies reacting with different cell lineages. *Hum Pathol* 19:51, 1988
29. Carbone A, Poletti A, Manconi R, et al: In situ immunophenotyping of follicular dendritic reticulum cells in malignant lymphomas of B-cell origin. *Cancer* 60:2919, 1987
30. Cerio R, Griffiths CEM, Cooper KD, et al: Characterization of factor XIIIa and dermal dendritic cells in normal and inflamed skin. *Br J Dermatol* 121:421, 1989
31. Chu A, Berger LC, Kung P, et al: In situ identification of Langerhans cells in the dermal infiltrate of cutaneous T-cell lymphoma. *J Am Acad Dermatol* 6:350, 1982
32. Cooper KD, Fox P, Neises C, et al: Effects of ultraviolet radiation on human epidermal cell alloantigen presentation: Initial depression of Langerhans cell-dependent function is followed by appearance of T6-DR+ cells that enhance epidermal alloantigen presentation. *J Immunol* 134:129, 1985
33. Cooper KD, Neises C, Katz SI: Antigen-presenting OKM5+ melanophages appear in human epidermis

- after ultraviolet radiation. *J Invest Dermatol* 86:363, 1986
34. Cooper KD, Duraiswamy N, Kang S, et al: Murine dermal cells in suspension contain T cell-activating presenting cells. *J Invest Dermatol* 88:482, 1987
 35. Cousar JB, McGinn DL, Glick AD, et al: Report of an unusual lymphoma arising from perifollicular B-lymphocytes (PBLs) or so called monocytoid lymphocytes. *Am J Clin Pathol* 87:121, 1987
 36. Cox NH, Turbitt ML, Ashworth J, et al: Distribution of T-cell subsets and Langerhans cells in mycosis fungoides, and the effect of PUVA therapy. *Clin Exp Dermatol* 11:564, 1986
 37. Davis G, York J, Glick A, et al: Plasmacytic differentiation in perifollicular (monocytoid) B-cell lymphoma: A study of eleven cases. *Lab Invest* 64:70A, 1991
 38. DeAlmeida PC, Harris NL, Bhan AK: Characterization of immature sinus histiocytes (monocytoid cells) in reactive lymph nodes by use of monoclonal antibodies. *Hum Pathol* 15:330, 1984
 39. Dréno B, Fleischmann M, Célrier P, et al: Dermal F XIIIa, CD1 positive cells in mycosis fungoides. *J Invest Dermatol* 99:113s, 1992
 40. Drijkoningen M, De Wolf-Peeters C, Van der Steen K, et al: Epidermal Langerhans' cells and dermal dendritic cells in human fetal and neonatal skin: An immunohistochemical study. *Pediatr Dermatol* 4:11, 1987
 41. Drijkoningen M, De Wolf-Peeters C, Snauwaert J, et al: Immunohistochemical study of epidermal Langerhans cells and dermal dendritic cells in benign and malignant skin lesions characterized by a dermal lymphoid infiltrate consisting either of B-cells or T-cells. *Virchows Arch A (Pathol Anat Histopathol)* 411:337, 1987
 42. Evans HL, Winkelmann RK, Banks PM: Differential diagnosis of malignant and benign cutaneous lymphoid infiltrates. *Cancer* 44:699, 1979
 43. Fivenson DP, Rheins LA, Nordlund JJ, et al: Thy-1 and T-cell receptor antigen expression in mycosis fungoides and benign inflammatory dermatoses. *J Natl Cancer Inst* 83:1088, 1991
 44. Fivenson DP, Douglass MC, Nickoloff BJ: Cutaneous expression of Thy-1 in mycosis fungoides. *Am J Pathol* 141:1373, 1992
 45. Flidner A, Parwaresch MR, Feller AC: Induction of antigen expression of follicular dendritic cells in a monoblastic cell line. A contribution to its cellular origin. *J Pathol* 161:71, 1990
 46. Fossum S, Ford WL: The organization of cell populations within lymph nodes: Their origin, life history and functional relationships. *Histopathology* 9:469, 1985
 47. Franchi F, Seminara P, Calvieri S: Interdigitating reticulum cell and lymphoproliferative disease: Possible correlations between skin and thymus-dependent lymphocyte system revisited. *Acta Haematol* 70:154, 1983
 48. Freedman AS, Munro JM, Rice GE, et al: Adhesion of human B-cells to germinal centres in vitro involves VLA-4 and ICAM-110. *Science* 249:1030, 1990
 49. Freedman AS, Munro JM, Rhyndart KR, et al: Adhesion of follicular lymphoma cells to neoplastic and normal germinal centres in vitro involves VLA-4 and VCAM-1. *Blood* 76(suppl 1):349a, 1990
 50. Fullbrandt U, Meissner K, Loning Th, et al: A second look to intraepithelial Langerhans cells in mycosis fungoides and related disorders. Ultrastructural study with special reference to Langerhans granules and virus-like particles. *Virchows Arch A (Pathol Anat Histopathol)* 402:47, 1983
 51. Garcia CF, Weiss LM, Warnke RA, et al: Cutaneous follicular lymphoma. *Am J Surg Pathol* 10:454, 1986
 52. Geerts M, Kaiserling E: A morphologic study of lymphadenosis benigna cutis. *Dermatologica* 170:121, 1985
 53. Giannotti B, Santucci M: Skin-associated lymphoid tissue (SALT)-related B-cell lymphoma (primary cutaneous B-cell lymphoma): A concept and a clinicopathologic entity. *Arch Dermatol* 129:353, 1993
 54. Gloghini A, Carbone A: The nodal lymphoid microenvironment of reactive follicles and lymphomas of follicular origin as defined by immunohistology on paraffin-embedded tissues. *Hum Pathol* 24:67, 1993
 55. Goos M: T-cell specific microenvironment in mycosis fungoides. *Arch Dermatol Res* 255:215, 1976
 56. Goos M, Kaiserling E, Lennert K: Mycosis fungoides: A model for T lymphocyte homing to the skin? *Br J Dermatol* 94:221, 1976
 57. Hansen ER, Baadsgaard O, Lisby S, et al: Cutaneous T-cell lymphoma lesional epidermal cells activate autologous CD4+ T lymphocytes: Involvement of both CD1+OKM5+ and CD1+OKM5- antigen-presenting cells. *J Invest Dermatol* 94:485, 1990
 58. Headington JT: The dermal dendrocyte. In Callen JP, Dahl MV, Golitz ME, et al (eds): *Advances in Dermatology*, Chicago, Year Book Medical Publishers, 1986, p 159
 59. Headington JT, Cerio R: Dendritic cells and the dermis: 1990. *Am J Dermatopathol* 12:217, 1990
 60. Holbert JM Jr, Chesney T, McC: Malignant lymphoma of the skin: A review of recent advances in diagnosis and classification. *J Cutan Pathol* 9:133, 1982
 61. Holden CA, Morgan EW, MacDonald DM: The cell population in the cutaneous infiltrate of mycosis fungoides: In situ studies using monoclonal antisera. *Br J Dermatol* 106:385, 1982
 62. Kaudewitz P, Burg G, Majewski G, et al: Cell populations in pagetoid reticulosis: an immunological study using cell activation associated monoclonal antibodies. *Acta Derm Venereol (Stockh)* 67:24, 1987
 63. Kerl H, Rauch HJ, Hödl S: Cutaneous B-cell lymphomas. In Goos M, Christophers E (eds): *Lymphoproliferative Disease of the Skin*. Berlin, Springer-Verlag, 1982, p 179
 64. Kerl H, Smolle J: Classification of cutaneous pseudolymphomas. Cutaneous lymphoma. *Curr Probl Dermatol* 19:167, 1990
 65. Igisu K, Watanabe S, Shimozato Y, et al: Langerhans cells and their precursors with S-100 protein in mycosis fungoides. *Jpn J Clin Oncol* 13:693, 1983
 66. Isaacson PG, Spencer J: Malignant lymphoma of mucosa-associated lymphoid tissue. *Histopathology* 11:445, 1987
 67. Iwahara K, Hashimoto K: T-cell subsets and nuclear contour index of skin-infiltrating T-cells in cutaneous T-cell lymphoma. *Cancer* 54:440, 1984
 68. Jimbow K, Chiba M, Horikishi T: Electron microscope identification of Langerhans cells in the dermal infiltrates of mycosis fungoides. *J Invest Dermatol* 78:102, 1982
 69. Langerhans P: Über die Nerven der menschlichen Haut. *Virchows Arch* 44:325, 1868
 70. Lennert K, Feller AC: *Histopathology of Non-Hodgkin's Lymphomas (Based on the Updated Kiel Classification)*, ed 2. Berlin, Springer-Verlag, 1992
 71. Lisby S, Baadsgaard O, Cooper KD, et al: Phenotype, ultrastructure, and function of CD1+DR+ epider-

- mal cells that express CD36 (OKM5) in cutaneous T-cell lymphoma. *Scand J Immunol* 32:111, 1990
72. Lisby S, Ralfkiaer E, Hansen ER, et al: Keratinocyte and epidermal leukocyte expression of CD36 (OKM5) in benign and malignant skin diseases. *Acta Dermatovenereol* 70:18, 1990
 73. Liu YJ, Johnson GD, Gordon J, et al: Germinal centres in T-cell-dependent antibody responses. *Immunol Today* 13:17, 1992
 74. van der Loo EM, van Muijen GNP, van Vloten WA, et al: C-type virus-like particles specifically localized in Langerhans cells and related cells of skin and lymph nodes of patients with mycosis fungoides and Sézary syndrome. *Virchows Arch B (Cell Pathol)* 31:193, 1979
 75. MacKie R: Initial event in mycosis fungoides of the skin is viral infection of epidermal Langerhans cells. *Lancet* 2:283, 1981
 76. MacKie RM, Turbitt ML: The use of a double-label immunoperoxidase monoclonal antibody technique in the investigation of patients of mycosis fungoides. *Br J Dermatol* 106:379, 1982
 77. McKenzie JL, Fabre JW: Human Thy-1: Unusual localization and possible functional significance in lymphoid tissues. *J Immunol* 126:843, 1980
 78. McMillan EM, Beeman K, Wasik R, et al: Demonstration of OKT6-reactive cells in mycosis fungoides. *J Am Acad Dermatol* 6:880, 1982
 79. Meissner K, Michaelis K, Rehpenning W, et al: Epidermal Langerhans cell densities influence survival in mycosis fungoides and Sézary syndrome. *Cancer* 65:2069, 1990
 80. Muller-Hermelink HK: Characterization of the B cell and T cell regions of human lymphatic tissue through enzyme cytochemical demonstration of ATPase and 5'-nucleotidase activities. *Virchows Arch B (Cell Pathol)* 16:371, 1974
 81. Murphy GF, Bhan AK, Harriot JT, et al: In situ identification of T6-positive cells in normal human dermis by immunoelectron microscopy. *Br J Dermatol* 108:423, 1983
 82. Nathwani B, Morhmann RL, Brynes RK, et al: Monocytoid B-cell lymphomas: An assessment of diagnostic criteria and a perspective on histogenesis. *Hum Pathol* 23:1061, 1992
 83. Nickoloff BJ, Griffiths CEM: Intraepidermal but not dermal T lymphocytes are positive for a cell-cycle-associated antigen (Ki-67) in mycosis fungoides. *Am J Pathol* 136:261, 1990
 84. Nizze H, Cogliatti SB, Von Schilling C, et al: Monocytoid B-cell lymphoma: Morphologic variants and relationship to low-grade B-cell lymphoma of the mucosa-associated lymphoid tissue. *Histopathology* 18:403, 1991
 85. Nussenzweig MC, Witner MD, Nogueira N, et al: A comparison of dendritic cells and macrophages. In Forster O, Landy M (eds): *Heterogeneity of Mononuclear Phagocytes*. San Diego, Academic Press, 1981, p 189
 86. Oord van den JJ, De Wolf-Peters C, De Vos R, et al: Immature sinus histiocytosis: Light and electron microscopic features, immunologic phenotype, and relationship to marginal zone lymphocytes. *Am J Pathol* 118:266, 1985
 87. Peters JP, Rademakers LHPM, Roelofs JMM, et al: Distribution of dendritic reticulum cells in follicular lymphoma and reactive hyperplasia. *Virchows Arch B (Cell Pathol)* 46:215, 1984
 88. Petrasch S, Stein H, Kosco MH, et al: Follicular dendritic cells in non-Hodgkin lymphomas: Localization, characterisation and pathophysiological aspects. *Eur J Cancer* 27:1052, 1991
 89. Pimpinelli N, Bani D: Non-lymphoid accessory cells in lichen planus and mycosis fungoides. *Ital Gen Rev Dermatol* 25:82, 1988
 90. Pimpinelli N, Romagnoli P, Moretti S, et al: Non-lymphoid accessory cells in the cutaneous infiltrate of B cell lymphomas. An immunohistochemical and ultrastructural study. *Br J Dermatol* 118:353, 1988
 91. Pimpinelli N, Bani D: Langerhans cells and epidermal microenvironment. *Am J Dermatopathol* 11:188, 1989
 92. Pimpinelli N, Santucci M, Bosi A, et al: Primary cutaneous follicular center-cell lymphoma: A lymphoproliferative disease with favourable prognosis. *Clin Exp Dermatol* 14:12, 1989
 93. Pimpinelli N, Santucci M, Carli P, et al: Primary cutaneous follicular center cell lymphoma: Clinical and histological aspects. In van Vloten WA, Willemze R, Lange Veilgaard G, et al (eds): *Cutaneous Lymphoma*. Basel, Karger, 1990, p 203
 94. Pimpinelli N, Borgognoni L, Riccardi R, et al: CD36(OKM5) - dendritic cells in the oral mucosa of HIV - and HIV + subjects. *J Invest Dermatol* 97:537, 1991
 95. Pimpinelli N, Romagnoli P, Santucci M, et al: Morpho-antigenic features of dendritic cells as a clue to the understanding of skin immune system related disorders. In van Vloten WA, Lambert WC, Giannotti B (eds): *Basic mechanisms of physiologic and aberrant lymphoproliferation in the skin*. New York, Plenum Press, 1993 (in press)
 96. Piris MA, Rivas C, Morente M, et al: Monocytoid B-cell lymphoma: A tumor related to the marginal zone. *Histopathology* 12:383, 1987
 97. Rijlaarsdam JU, Meijer CJLM, Willemze R: Differentiation between lymphadenosis benigna cutis and primary cutaneous follicular center cell lymphomas. A comparative clinicopathologic study of 57 patients. *Cancer* 65:2301, 1990
 98. Rijlaarsdam JU, Scheffer E, Meijer CJLM, et al: Cutaneous pseudo-T-cell lymphomas: A clinicopathologic study of 20 patients. *Cancer* 69:717, 1992
 99. Romagnoli P, Moretti S, Fattorossi A, et al: Dendritic cells in the dermal infiltrate of Sézary syndrome. *Histopathology* 10:25, 1986
 100. Romagnoli P, Pimpinelli N, Lotti T: Dendritic cells in the skin and oral mucosa. *Ital Gen Rev Dermatol* 28:49, 1991
 101. Rowden G, Phillips TM, Lewis MC: Ia antigens on indeterminate cells in the epidermis. Immunoelectron microscopic studies on surface antigens. *Br J Dermatol* 108:531, 1979
 102. Santucci M, Pimpinelli N, Arganini L: Primary cutaneous B-cell lymphoma: A unique type of low-grade lymphoma. *Cancer* 67:2311, 1991
 103. Santucci M, Mori M, Pimpinelli N, et al: Nerve growth factor receptor expression on dendritic reticulum cells in cutaneous B-cell lymphoma. *Hum Pathol* 23:1088, 1992
 104. Santucci M, Pimpinelli N: Cutaneous B-cell lymphoma: A SAT-related tumor? In van Vloten WA, Lambert WC, Giannotti B (eds): *Basic Mechanisms of Physiologic and Aberrant Lymphoproliferation in the Skin*. New York, Plenum Press, 1993 (in press)
 105. Schuler G: *Epidermal Langerhans Cells*. Boca Raton, CRC Press, 1991
 106. Sheibani K, Fritz RM, Winberg CD, et al: Monocytoid cells in reactive follicular hyperplasia with and without multifocal histiocytic reactions: An immu-

- nohistochemical study of 21 cases including suspected cases of toxoplasmic lymphadenitis. *Am J Clin Pathol* 81:453, 1984
107. Sheibani K, Sohn CC, Burke JS, et al: Monocytoid B-cell lymphoma: A novel B-cell neoplasm. *Am J Pathol* 124:310, 1986
 108. Sheibani K, Burke JS, Swartz WG, et al: Monocytoid B-cell lymphoma: Clinicopathologic study of 21 cases of a unique type of low-grade lymphoma. *Cancer* 62:1531-1538, 1988
 109. Shen HH, Talle MA, Goldstein G, et al: Functional subsets of human monocytes defined by monoclonal antibodies: a distinct subset of monocytes contains the cells capable of inducing the autologous mixed lymphocyte culture. *J Immunol* 130:698, 1983
 110. Slater DN, Bleehen S, Rooney N, et al: Type C retrovirus-like particles in mycosis fungoides. *Br J Dermatol* 109:120, 1983
 111. Smolle J, Ehall R, Kerl H: Inflammatory cell types in normal human epidermis: An immunohistochemical and morphometric study. *Acta Dermatovenereol* 65:479, 1985
 112. Smolle J, Kaudewitz P, Burg G, et al: Significance of non-lymphoid (accessory) cells in malignant lymphomas and pseudolymphomas of the skin. *Br J Dermatol* 113:677, 1985
 113. Sontheimer RD: Perivascular dendritic macrophages as immunobiological constituents of the human dermal microvascular unit. *J Invest Dermatol* 93:s96, 1989
 114. Soyer HP, Smolle J, Kerl H: Distribution patterns of the OKM5 antigen in normal and diseased human epidermis. *J Cutan Pathol* 16:60, 1988
 115. Spencer J, Finn T, Pulford KAF, et al: The human gut contains a novel population of B lymphocytes which resemble marginal zone cells. *Clin Exp Immunol* 62:607, 1985
 116. Stein H, Gerdes J, Mason DY: The normal and malignant germinal centre. *Clin Haematol* 11:531, 1982
 117. Stein H, Lennert K, Mason DY, et al: Immature sinus histiocytes: Their identification as a novel B-cell population. *Am J Pathol* 117:44, 1984
 118. Steinman RM: The dendritic cell system and its role in immunogenicity. *Annu Rev Immunol* 9:271, 1991
 119. Streilein JW: Skin-associated lymphoid tissue (SALT): Origins and functions. *J Invest Dermatol* 80:12, 1983
 120. Streilein JW: Antigen-presenting cells in the induction of contact hypersensitivity in mice—evidence that Langerhans cells are sufficient but not required. *J Invest Dermatol* 93:443, 1989
 121. Strobach RS, Nakamine H, Masih AS, et al: Nerve growth factor receptor expression on dendritic reticulum cells in follicular lymphoid proliferations. *Hum Pathol* 22:481, 1991
 122. Tan RSH, Butterworth CM, McLaughlin H, et al: Mycosis fungoides, a disease of antigen persistence. *Br J Dermatol* 91:607, 1974
 123. Tan RSH, MacLeod HF, Dean SG: Pagetoid reticulis, epidermotropic mycosis fungoides and mycosis fungoides: a disease spectrum. *Br J Dermatol* 116:67, 1987
 124. Tron VA, Rosenthal D, Sauder DN: Epidermal interleukin-1 is increased in cutaneous T-cell lymphoma. *J Invest Dermatol* 90:378, 1988
 125. Tse Y, Cooper KD: Cutaneous dermal Ia+ cells are capable of initiating delayed type hypersensitivity responses. *J Invest Dermatol* 94:267, 1990
 126. Tsujimoto Y, Cossman J, Jaffe E, et al: Involvement of the bcl-2 gene in human follicular lymphoma. *Science* 228:1440, 1985
 127. van der Valk P, van der Loo EM, Jansen J, et al: Analysis of lymphoid and dendritic cells in human lymph node, tonsil and spleen. *Virchows Arch B (Cell Pathol)* 45:169, 1984
 128. van der Valk P, Meijer CJLM: The histology of reactive lymph nodes. *Am J Surg Pathol* 11:866, 1987
 129. Weber-Matthiesen K, Sterry W: Organization of the monocyte/macrophage system of normal human skin. *J Invest Dermatol* 95:83, 1990
 130. Weisenburger DD, Nathwani BN, Diamond LW, et al: Malignant lymphoma, intermediate lymphocytic type: A clinico-pathologic study of 42 cases. *Cancer* 48:1415, 1981
 131. Weisenburger DD, Kim H, Rappaport H: Mantle zone lymphoma: A follicular variant of intermediate lymphocytic lymphoma. *Cancer* 49:1429, 1982
 132. Weiss LM, Warnke RA, Sklar J, et al: Molecular analysis of the t(14;18) chromosomal translocation in malignant lymphomas. *N Engl J Med* 317:1185, 1987
 133. Willemze R, De-Graaff Reitsma GB, van Vloten WA, et al: The cell population of cutaneous B-cell lymphomas. *Br J Dermatol* 108:395, 1983
 134. Willemze R, Meijer CJLM, Sentis H, et al: Primary cutaneous large cell lymphomas of follicular center cell origin. *J Am Acad Dermatol* 16:518, 1987
 135. Willemze R, Meijer CJLM, Scheffer E, et al: Diffuse large cell lymphomas of follicular center cell origin. *Am J Pathol* 126:325, 1987
 136. Winkelmann RK, Dabski K: Large cell lymphocytoma: Follow-up, immunopathology studies, and comparison to cutaneous follicular and Crosti lymphoma. *Arch Dermatol Res* 279:s81, 1987
 137. Wirt DP, Grogan TM, Jolley CS, et al: The immunorearchitecture of cutaneous pseudolymphoma. *Hum Pathol* 16:492, 1985
 138. Wood GS, Turner RR, Shiurba RA, et al: Human dendritic cells and macrophages: In situ immunophenotypic definition of subsets that exhibit specific morphologic and microenvironmental characteristics. *Am J Pathol* 119:73, 1985
 139. Wood GS, Ngan B-Y, Tung R, et al: Clonal rearrangement of immunoglobulin genes and progression to B-cell lymphoma in cutaneous lymphoid hyperplasia. *Am J Pathol* 135:13, 1989

Address reprint requests to

Nicola Pimpinelli, MD, PhD
 Clinica Dermatologica II
 Università degli Studi di Firenze
 Via della Pergola, 58
 I-50121 Firenze
 Italy