UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

IFNgamma and TNFalpha account for a pro-clonogenic activity secreted by activated murine peritoneal macrophages.

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

IFNgamma and TNFalpha account for a pro-clonogenic activity secreted by activated murine peritoneal macrophages / L. Calorini; F. Bianchini; A. Mannini; G. Mugnai; M. Balzi; A. Becciolini; S. Ruggieri. - In: CLINICAL & EXPERIMENTAL METASTASIS. - ISSN 0262-0898. - STAMPA. - 19:(2002), pp. 259-264. [10.1023/A:1015583322354]

Availability:

This version is available at: 2158/2251 since: 2020-10-23T10:58:44Z

Published version:

DOI: 10.1023/A:1015583322354

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

Research paper

IFN γ and TNF α account for a pro-clonogenic activity secreted by activated murine peritoneal macrophages

Lido Calorini¹, Francesca Bianchini¹, Antonella Mannini¹, Gabriele Mugnai¹, Manuela Balzi², Aldo Becciolini² & Salvatore Ruggieri¹

¹Department of Experimental Pathology and Oncology, ²Department of Clinical Physiopathology, Radiation Biology Laboratory, University of Florence, Italy

Key words: F10-M3 cells, *Corynebacterium parvum*-, BCG- or *Listeria monocytogenes*-elicited macrophages, MHC class I antigens, lung colonization, macrophage pro-clonogenic activity, IFN γ , TNF α

Abstract

In the present study, we found that murine peritoneal macrophages elicited by BCG or *Listeria monocytogenes* release into the media an activity capable of stimulating the lung colonization as well as the expression of MHC class I antigens in B16 melanoma cells. A similar activity has previously been found in media conditioned by *Corynebacterium parvum*-elicited macrophages. Analysis by gel filtration chromatography of media conditioned by *Corynebacterium parvum*-, BCG- or *Listeria monocytogenes*-elicited macrophages revealed that the material responsible for the pro-clonogenic activity concentrated in chromatographic fractions corresponding to molecular weights (25 to 52 kDa) which are characteristic of certain cytokines. Thus, we challenged the various macrophage-conditioned media with polyclonal antibodies against IFN γ and TNF α , and found that the macrophage pro-clonogenic activity was completely abolished in the presence of anti-IFN γ antibodies, but only partially inhibited by anti-TNF α antibodies. This finding suggests a cooperative participation of the two cytokines to the pro-clonogenic activity of the media conditioned by *Corynebacterium parvum*-, BCG- or *Listeria monocytogenes*-elicited macrophages.

Introduction

In a previous study, we demonstrated that cultures of peritoneal macrophages elicited *in vivo* with *Corynebacterium parvum* contained a biological activity which stimulated the lung colonization in B16-F10 murine melanoma cells [1]. We also found that the pro-clonogenic activity stimulated the adhesiveness to endothelium, invasiveness through Matrigel and growth rate in melanoma cells. Moreover, melanoma cells exposed to the macrophage pro-clonogenic activity showed an increased expression of the major histocompatibility complex (MHC) class I antigens (K^b and D^b) [2].

In the present study, we tried to gain insight into the nature of the pro-clonogenic activity released by the elicited macrophages into their growth medium. Conditioned media of *C. parvum*- as well as of BCG- or *Listeria monocytogenes*-elicited macrophages were used as sources of a macrophage pro-clonogenic activity. These media were fractionated according to molecular weight by the use of gel filtration chromatography. The data obtained through this

procedure suggested that the macrophage pro-clonogenic activity might derive from certain cytokines, a suggestion that was confirmed by the use of antibodies against IFN γ and TNF α .

Materials and methods

Reagents

Corynebacterium parvum (Coparwax) was a gift from Wellcome Foundation (London, UK); BCG (OncoTICE) was supplied by Organon Teknika (Boxtel, The Netherlands), and *Listeria monocytogenes* was provided by Dr R. Dei (Department of Microbiology, University of Florence). AF6-88.5.3 mAb, specific for K^b determinant of MHC class I antigen (H-2K^b antigen) was a gift from Dr S. Gattoni-Celli (Medical University of South Carolina, Charleston, South Carolina), rabbit polyclonal antibodies anti-murine TNF α and polyclonal antibodies anti-murine IFN γ were purchased from PeproTech EC (London, UK). Gel filtration calibration kit for molecular weight determination was purchased from Pharmacia (Uppsala, Sweden).

Correspondence to: Dr Lido Calorini, Department of Experimental Pathology and Oncology, Florence University, Viale G.B. Morgagni 50, 50134 Florence, Italy. Tel: +39-055-4282322; Fax: +39-055-4282333; E-mail: lcalorini@unifi.it

Table 1. Change of lung-colonizing potential and expression of H-2K^b antigen in F10-M3 melanoma cells grown in media conditioned by *C. parvum*-, BCG-, or *L. monocytogenes*-elicited macrophages.¹

Growth conditions	No. of animals in the experiment	Lung colonies	H-2K ^b antigen expression
Standard medium	5	4 ± 1 ²	6–15 ³
Macrophage-conditioned medium:			
BCG	3	> 300 ⁴	72–96
<i>L. monocytogenes</i>	5	158 ± 35 ⁴	54–88
<i>C. parvum</i>	3	220 ± 15 ⁴	76–98

¹Data reported in the Table are derived from a typical experiment.

²Values represent the mean ± SEM of lung colonies found in the experimental animals.

³% of cells positive for H-2K^b antigen as assayed by flow cytometry using a specific mAb.

⁴Significantly different at *P* < 0.03 from cells grown in a standard medium.

Cell line and culture conditions

F10-M3 cells, a clone isolated from B16-F10 cell line [3], were kindly provided by Dr S. Gattoni-Celli (Medical University of South Carolina, Charleston, South Carolina). Cells were grown in Dulbecco's modified Eagle medium containing 4,500 mg/l glucose (DMEM 4500) (GIBCO, Life Technologies, Italy) supplemented with 10% fetal calf serum (FCS) (Boehringer Mannheim, Germany), at 37 °C in a 10% CO₂-humidified atmosphere. 5.0 × 10⁵ cells were seeded in 100 mm Falcon dishes and propagated every three days by incubation with a trypsin solution (GIBCO).

Cultures were periodically monitored for mycoplasma contamination using Chen's fluorochrome test [4].

Preparation of macrophage-conditioned media

As previously reported, monolayers were established from macrophages isolated from a lavage of the peritoneal cavities of female C57Bl/6 or CBA mice which had received an intraperitoneal injection of 1 ml of a suspension containing 0.7 mg/ml of killed *C. parvum*, or 0.4 ml of a suspension of BCG containing 5 × 10⁶ CFU/ml or 1 ml of a suspension of *L. monocytogenes* containing 15 × 10⁴ CFU/ml. Macrophage-conditioned media were prepared from monolayers of *C. parvum*-, BCG- or *L. monocytogenes*-elicited macrophages (3–5 × 10⁴ macrophages/cm²) incubated for 24 h in DMEM 4500 supplemented with 250 µg/ml of bovine serum albumin (BSA) [1, 2].

Media conditioned by BCG- or *L. monocytogenes*-elicited macrophages were found to contain amounts of NO and TNFα comparable to those found in media conditioned by *C. parvum*-elicited macrophages.

Determination of the pro-clonogenic activity in media conditioned by *C. parvum*-, BCG-, or *L. monocytogenes*-elicited macrophages

Melanoma cells were grown in a standard medium for 24 h and then for a further 24 h period in macrophage-conditioned media that were supplemented with FCS to a final concentration of 10%. Melanoma cells grown for 48 h in a standard medium were used as a control. Cultures were harvested by trypsinization, washed by centrifugation in PBS, and then resuspended at 5 × 10⁵ cells/ml in serum free DMEM 4500.

0.2 ml of this suspension were injected intravenously into female C57Bl/6 mice, which were sacrificed 21 days later. Metastatic nodules on the lung surfaces were counted using a dissecting microscope.

Expression of H-2K^b antigen in melanoma cells grown in a standard medium or in a medium conditioned by *C. parvum*-, BCG- or *L. monocytogenes*-elicited macrophages

The expression of H-2K^b antigen on melanoma cells grown in a standard medium or in a macrophage conditioned medium was measured by FACS analysis (FACScan, Becton Dickinson, Mountain View, California) using a AF6-88.5.3 mAb.

Analysis of macrophage-conditioned media by size-exclusion chromatography

The conditioned media were concentrated by lyophilization, dissolved in water and loaded on a Sephacryl S-300 column (Pharmacia) (1.5 × 40 cm) equilibrated with DMEM 4500. Chromatography was performed at RT with a flow-rate of 0.4 ml/min, and the eluted material was monitored at 280 nm. The chromatographic fractions were collected, supplemented with FCS to a final concentration of 10%, and then transferred into 24 multiwell dishes, where melanoma cells had been previously layered. After a 24-h period of incubation at 37 °C, melanoma cells were removed and tested for the expression of H-2K^b antigen.

The molecular weights of the bioactive material eluted from the Sephacryl column was determined by the use of a calibration kit containing ribonuclease A (13.7 kDa), chymotrypsinogen A (25 kDa), ovalbumin (43 kDa), and BSA (67 kDa).

Challenge of macrophage-conditioned media with anti-IFNγ or -TNFα antibodies

The macrophage-conditioned media, intact or diluted with fresh medium (DMEM 4500 plus BSA), were challenged with various concentrations of rabbit polyclonal antibodies against murine TNFα (1:64,000-fold dilution of a 1 mg/ml antibody reacts with 1 µg/ml TNFα in a ELISA test) or with polyclonal antibodies against murine IFNγ (ND₅₀ of

Figure 1. Lung-colonizing potential (A, C) and expression of H-2K^b antigen (B, D) in F10-M3 melanoma cells grown in media conditioned by BCG- (A, B) or *L. monocytogenes*- (C, D) elicited macrophages, and then subcultivated in a standard medium for two passages. Tumor cells grown in a standard medium for the entire experimental protocol (empty columns) served as a control. Values represent the mean $\pm$ SEM of lung colonies found in the experimental animals (groups of 3–4 mice). *Significantly different at $P < 0.02$ from cells grown in a standard medium.

the biological activity of 0.3 ng/ml of mIFN γ requires a concentration of 0.03–0.05 μ g/ml of antibody). The conditioned media-antibody mixtures were incubated at 37 °C, for 1.5 h.

Statistical analysis

The statistical significance of the differences between the lung colonization of melanoma cells grown in a standard medium and in a medium conditioned by *C. parvum*-, BCG-, or *L. monocytogenes*-elicited macrophages was determined by the use of the Mann–Whitney test.

Results

As shown in Table 1, growth in media conditioned by BCG- or *L. monocytogenes*-elicited macrophages enhanced the lung-colonizing potential and stimulated the expression of H-2K^b antigen in F10-M3 melanoma cells to levels comparable to that of melanoma cells grown in media conditioned

by *C. parvum*-elicited macrophages. As previously found with B16 melanoma cells grown in media conditioned by *C. parvum*-elicited macrophages [2], the lung-colonizing potential and expression of H-2K^b antigen in melanoma cells stimulated by the pro-clonogenic activity released by BCG- or *L. monocytogenes*-elicited macrophages declined to the same level as that found in unstimulated cells after two subcultivations in a standard medium (Figure 1).

Figure 2 gives the profiles of the pro-clonogenic activity detected in the various fractions obtained by gel filtration chromatographic analysis of media conditioned by *C. parvum*-, BCG- or *L. monocytogenes*-elicited macrophages. Regardless of the eliciting agent, the highest levels of the macrophage pro-clonogenic activity were found in chromatographic fractions corresponding to molecular weights ranging between 25 and 52 kDa (maximum peak of activity at 35–38 kDa.). These values are compatible with various cytokines, in particular IFN γ (34 kDa) [5] and TNF α (51 kDa) [6]. At this point, we investigated

Figure 2. Gel filtration chromatography of media conditioned by macrophages elicited with *C. parvum* (A), BCG (B), or *L. monocytogenes* (C). The histograms correspond to the % of F10-M3 melanoma cells which expressed the H-2K^b antigen after exposure to the various chromatographic fractions. The chromatographic elution positions of chymotrypsinogen A (25 kDa), ovalbumin (43 kDa) and BSA (67 kDa) are reported as size standards.

Figure 3. Change of the lung-colonizing potential (A, C) and expression of H-2K^b antigen (B, D) in F10-M3 cells grown in media conditioned by BCG- (A, B) or *L. monocytogenes*- (C, D) elicited macrophages (light grey columns), and in the same media challenged with anti-IFN γ (dark grey columns) or anti-TNF α antibodies (black columns). Tumor cells grown in a standard medium for the entire experimental protocol (empty columns) served as a control. Data reported in the Figure are derived from a typical experiment. Values represent the mean $\pm$ SEM of lung colonies found in the experimental animals (groups of 4–5 mice). Significantly different at * $P = 0.02$, ** $P = 0.008$, *** $P = 0.01$ from cells grown in a standard medium.

Figure 4. Change of the lung-colonizing potential in F10-M3 melanoma cells treated with exogenous IFN γ and/or TNF α . Melanoma cells were grown in media supplemented with IFN γ (25 U/ml) or TNF α (25 μ g/ml) following the indicated sequence. Tumor cells grown in a standard medium for the entire experimental protocol (empty columns) served as a control. Values represent the mean $\pm$ SEM of lung colonies found in the experimental animals (groups of 5–6 mice). *Significantly different at $P = 0.008$ from cells grown in a standard medium.

whether the treatment of macrophage-conditioned media with anti-IFN γ and anti-TNF α antibodies affected the pro-clonogenic activity present in these media. As shown in Figure 3, the capacity of the media conditioned by BCG- or *L. monocytogenes*-elicited macrophages of up-regulating H-2K^b antigen as well as enhancing the lung-colonizing potential in melanoma cells was lost after treatment with anti-IFN γ antibodies. Treatment of macrophage-conditioned media with anti-TNF α antibodies only partially affected their pro-clonogenic activity.

Discussion

The present study revealed that a pro-clonogenic activity, previously demonstrated in media conditioned by *C. parvum*-elicited macrophages [1, 2], is also present in media conditioned by BCG- or *L. monocytogenes*-elicited macrophages. Moreover, as in the case of *C. parvum*-elicited macrophages [2], the pro-clonogenic activity generated by BCG- or *L. monocytogenes*-elicited macrophages enhanced the expression of MHC class I antigens in F10-M3 melanoma cells.

The gel filtration chromatographic analysis of macrophage-conditioned media revealed that the molecular weight of the pro-clonogenic activity present in these media ranged between values compatible with certain cytokines, such as IFN γ and TNF α . Indeed, the challenge of the macrophage-conditioned media with anti-IFN γ antibodies abolished their capacity of increasing the lung-colonizing potential as well as up-regulating the expression of H-2K^b antigen

in melanoma cells. The recent finding that macrophages, besides lymphocytes, can secrete IFN γ [7–9] sustains the participation of IFN γ in the pro-clonogenic activity released by the elicited macrophages. Our observation that IFN γ is involved in the macrophage pro-clonogenic activity is analogous to the finding that treatment of tumor cells with IFN γ enhances their colonization in secondary organs [10–15]. The pro-clonogenic effect of IFN γ has been attributed to an inhibition of the anti-metastatic NK cell activity caused by the IFN γ -mediated enhancement of MHC class I antigens [16–18].

The limited inhibitory effect of anti-TNF α antibodies on the macrophage pro-clonogenic activity suggests a different contribution of TNF α as compared to IFN γ in promoting organ colonization of melanoma cells. It is possible that a prior stimulation by IFN γ is required for TNF α to produce the maximal pro-clonogenic activity. This hypothesis is sustained by our observation, reported in Figure 4, that the lung-colonizing potential is enhanced in melanoma cells treated with exogenous TNF α provided the cells were previously exposed to IFN γ , while the reversal of this sequence was not effective. The possibility that TNF α displays a pro-clonogenic activity implies the presence of specific receptors whose expression, as recently reported [19], is promoted by IFN γ .

Our observation that TNF α and IFN γ contribute to the pro-clonogenic activity generated by elicited macrophages is in contrast with the *in vivo* therapeutic use of IFN γ and TNF α as anti-metastatic agents [20–23]. This discrepancy might be explained on the basis of the differences between the *in vivo* and *in vitro* conditions. Indeed, *in vivo* treatment

with $\text{INF}\gamma$ and $\text{TNF}\alpha$ may evoke a complex array of effects in different homeostatic systems of the host, due to the pleiotropic properties of these cytokines [7–24]. Nevertheless, use of *in vitro* experimental protocols offers the advantage of exploring specific tumor cell/host cell interactions under well controlled conditions.

Acknowledgements

This study was funded by grants from MURST 40% – Cofin1999 and MURST ex 60%. The authors wish to thank Prof. Alberto Fonnesu for his interest in this work.

References

- Cecconi O, Calorini L, Mannini A et al. Enhancement of lung-colonizing potential of murine tumor cell lines co-cultivated with activated macrophages. *Clin Exp Metastasis* 1997; 15: 94–101.
- Calorini L, Mannini A, Bianchini F et al. Biological properties associated with the enhanced lung-colonizing potential in a B16 murine melanoma line grown in a medium conditioned by syngeneic *Corynebacterium parvum*-elicited macrophages. *Clin Exp Metastasis* 1999; 17: 889–95.
- Gattoni-Celli S, Calorini L, Simile MM et al. Modulation by MHC Class I antigens of the biology of melanoma cells. Non-immunological mechanisms. *Melanoma Res* 1993; 3: 285–9.
- Chen TR. *In situ* detection of *Mycoplasma* contamination in cell cultures by fluorescent Hoechst 33258 stain. *Exp Cell Res* 1977; 104: 255–62.
- Havell EA, Spitalny GL. Two molecular weight species of murine gamma interferon. *Virology* 1983; 129: 508–13.
- Wingfield P, Pain RH, Craig S. Tumor necrosis factor is a compact trimer. *FEBS Lett* 1987; 211: 179–84.
- Billiau A. Interferon- γ : Biology and role in pathogenesis. *Adv Immunol* 1996; 62: 61–130.
- Puddu P, Fantuzzi L, Borghi P et al. IL-12 induces IFN-gamma expression and secretion in mouse peritoneal macrophages. *J Immunol* 1997; 159: 3490–7.
- Guillemard E, Geniteau-Legendre M, Kergot R et al. Simultaneous production of IFN-gamma, IFN-alpha/beta and nitric oxide in peritoneal macrophages from TDM-treated mice. *J Biol Regul Homeost Agents* 1998; 12: 106–11.
- Taniguchi K, Petersson M, Hoglund P et al. Interferon γ induces lung colonization by intravenously inoculated B16 melanoma cells in parallel with enhanced expression of class I major histocompatibility complex antigens. *Proc Natl Acad Sci USA* 1987; 84: 3405–9.
- McMillan TJ, Rao J, Everett CA et al. Interferon-induced alterations in metastatic capacity, class-I antigen expression and natural killer cell sensitivity of melanoma cells. *Int J Cancer* 1987; 40: 659–63.
- Zoller M, Strubel A, Hammerling G et al. Interferon-gamma treatment of B16 melanoma cells: Opposing effects for non-adaptive and adaptive immune defense and its reflection by metastatic spread. *Int J Cancer* 1988; 41: 256–66.
- Ramani P, Balkwill FR. Enhanced metastases of a mouse carcinoma after *in vitro* treatment with murine interferon gamma. *Int J Cancer* 1987; 40: 830–4.
- Kelly SA, Gschmeissner S, East N et al. Enhancement of metastatic potential by γ -interferon. *Cancer Res* 1991; 51: 4020–7.
- Lollini PL, De Giovanni C, Nicoletti G et al. Enhancement of experimental metastatic ability by tumor necrosis factor-alpha alone or in combination with interferon-gamma. *Clin Exp Metastasis* 1990; 8: 215–24.
- Piontek G, Taniguchi K, Ljunggren H et al. YAC-1 MHC class I variants reveal association between decreased NK sensitivity and increased H-2 expression after interferon treatment or *in vivo* passage. *J Immunol* 1985; 135: 4281–8.
- Taniguchi K, Karre K, Klein G. Lung colonization and metastasis by disseminated B16 melanoma cells: H-2 associated control at the level of the host and the tumor cells. *Int J Cancer* 1985; 36: 503–10.
- Kawano Y-I, Taniguchi K, Toshitani A et al. Synergistic defense system by cooperative natural effectors against metastasis of B16 melanoma cells in H-2-associated control: Different behaviour of H-2⁺ and H-2⁻ cells in metastatic process. *J Immunol* 1986; 136: 4729–34.
- Carrel S, Hartmann F, Salvi S et al. Expression of type A and B tumor necrosis factor (TNF) receptors on melanoma cells can be regulated by dbc-AMP and IFN γ . *Int J Cancer* 1995; 62: 76–83.
- Saiki I, Maeda H, Murata J et al. Antimetastatic effect of endogenous tumor necrosis factor induced by the treatment of recombinant interferon gamma followed by an analogue (GLA-60) to synthetic lipid A subunit. *Cancer Immunol Immunother* 1989; 30: 151–157.
- Schultz RM, Altom MG. Protective activity of recombinant murine tumor necrosis factor-alpha and interferon-gamma against experimental murine lung carcinoma metastases. *J Interferon Res* 1990; 10: 229–36.
- Ruegg C, Yilmaz A, Bieler G et al. Evidence for the involvement of endothelial cell integrin alphaVbeta3 in the disruption of the tumor vasculature induced by TNF and IFN-gamma. *Nat Med* 1998; 4: 408–14.
- van Moorselaar RJA, Hendriks BT, van Stratum P et al. Synergistic antitumor effects of rat γ -interferon and human tumor necrosis factor α against androgen-dependent and -independent rat prostatic tumors. *Cancer Res* 1991; 51: 2329–34.
- Tracey KJ, Cerami A. Tumor necrosis factor: A pleiotropic cytokine and therapeutic target. *Annu Rev Med* 1994; 45: 491–503.