



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Androgen receptor and prostate cancer invasion.

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Androgen receptor and prostate cancer invasion / L. BONACCORSI ; M. MURATORI ; V. CARLONI ; S. ZECCHI-ORLANDINI ; L. FORMIGLI ; G. FORTI ; E. BALDI. - In: INTERNATIONAL JOURNAL OF ANDROLOGY. - ISSN 0105-6263. - STAMPA. - 26:(2003), pp. 21-25.

Availability:

The webpage <https://hdl.handle.net/2158/252370> of the repository was last updated on

Publisher:

Blackwell Science Limited:PO Box 88, Oxford OX2 0NE United Kingdom:011 44 1865 776868, 011 44 1865

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

REVIEW

Androgen receptor and prostate cancer invasion¹

LORELLA BONACCORSI,* MONICA MURATORI,* VINICIO CARLONI,† SANDRA ZECCHI,‡ LUCIA FORMIGLI,‡ GIANNI FORTI* and ELISABETTA BALDI*

*Dipartimento di Fisiopatologia Clinica, Unità di Andrologia, †Dipartimento di Medicina Interna e Immunologia, and ‡Dipartimento di Anatomia Umana, Università di Firenze, Firenze, Italy

Summary

Evidence indicates that androgen-sensitive prostate cancer cells have a lower malignant potential. We previously demonstrated that expression of androgen receptor (AR) by transfection of the androgen-independent prostate cancer cell line PC3 decreases invasion and adhesion of these cells through modulation of $\alpha 6 \beta 4$ expression. Treatment with the androgen further reduced adhesion and invasion of the cells without, however, modifying $\alpha 6 \beta 4$. Here we investigated whether the androgen has a direct effect on $\alpha 6 \beta 4$ -EGF receptor (EGFR) interaction and signalling leading to invasion of these cells. Immunoconfocal microscopy demonstrated that in control cells (PC3-Neo), $\alpha 6 \beta 4$ and EGFR colocalize and redistribute in response to epidermal growth factor (EGF). In PC3-AR cells colocalization and redistribution between the two molecules was reduced and abolished by pre-treatment with R1881. Co-immunoprecipitation studies demonstrated that tyrosine phosphorylation of $\beta 4$ in response to EGF was reduced in PC3-AR cells compared to PC3-Neo. Immunoconfocal and co-immunoprecipitation studies demonstrated colocalization at membrane level and co-immunoprecipitation of EGFR and AR, indicating an interaction between the two proteins. PI3K activity, a key signalling pathway for invasion of these cells, was decreased in PC3-AR cells in response to EGF and further reduced by treatment with R1881. EGFR internalization was strongly reduced in PC3-AR compared with PC3-Neo cells and was reduced by treatment with R1881. In conclusion, the expression of AR by transfection in PC3 cells confers a less malignant phenotype by interfering with EGFR- $\alpha 6 \beta 4$ interaction and signalling leading to invasion through a mechanism involving an interaction between the classic AR and EGFR.

Keywords: epidermal growth factor, integrin, prostate cancer

Introduction

Prostate cancer (PC) is one of the most common cancers and the second leading cause of death among

Correspondence: Dr. Lorella Bonaccorsi, Andrology Unit, Department of Clinical Physiopathology, University of Florence, Viale Pieraccini, 6, 50139 Florence, Italy. Fax: +39-055-4271371; e-mail: l.bonaccorsi@dfc.unifi.it

¹This is a review article for the Second European Congress of Andrology in Malmö, Sweden.

other populations than American men only (Amanatullah *et al.*, 2000). As PC cell growth is enhanced by androgens, in the advanced stages of the disease androgen ablation therapy represents a valuable tool for the treatment of these patients. However, the development in most patients after few years of treatment of androgen-independent clones characterized by higher invasiveness and metastatic properties, has focused attention on the molecular mechanisms that lead to loss of androgen dependence as well as

on the pathways that are regulated by androgens in these cells. At this stage no valuable tools for the treatment of the patients are available. Studies performed with patient specimens have shown that the androgen receptor (AR) is expressed in almost all cancers of the prostate after androgen ablation therapy indicating that loss of androgen-dependence is not related to the expression of AR (Van der Kwast *et al.*, 1993). Indeed, maintenance of androgen-sensitivity appears to have a more differentiated and less malignant phenotype of PC cells. The ability to produce tumours in nude mice, for instance, is higher in androgen-insensitive cell lines (such as PC3 and DU145) with respect to the androgen-sensitive (LNCaP) (Witkowski *et al.*, 1993). Moreover, invasion ability of androgen-independent prostate carcinoma cells in response to epidermal growth factor (EGF) is decreased by transfection with an AR expression vector (Bonaccorsi *et al.*, 2000; Cinar *et al.*, 2001). In this light, the role of androgens and AR in the regulation of the signalling pathways involved in invasion and metastasis represents a primary task in studies on PC biology. As a result, some androgen-regulated genes involved in signalling pathways leading to invasion has been recently identified and their role in lower invasion ability of androgen-sensitive PC cells indicated (Bonaccorsi *et al.*, 2000; Sumitomo *et al.*, 2000; Manos *et al.*, 2001; Trusolino *et al.*, 2001), suggesting that loss of androgen regulation leads to increased invasion through multiple pathways.

Multiple pathways resulting from extracellular and intracellular signals regulate invasion of carcinoma cells. Indeed, cell migration results from merging of signalling pathways, which employ growth factors and their receptors, adhesion receptors (integrins) and cytoskeletal elements.

Integrins are heterodimers composed of no covalently associated alpha and beta subunits (Meredith *et al.*, 1996), and have a key role in the regulation of proliferation and invasion of carcinomas (Juliano & Verner, 1993). In particular, $\alpha 6 \beta 4$, one of the main receptors of laminins, is a primary actor in the process of migration and invasion (Dans *et al.*, 2001). In normal epithelia $\alpha 6 \beta 4$ mediates the formation of stable adhesive structures termed hemidesmosomes, that link intermediate filaments to extracellular matrix, whereas in carcinoma cells, it promotes migration on laminin through association with actin cytoskeleton (Dans *et al.*, 2001). It has been demonstrated that $\alpha 6 \beta 4$ co-immunoprecipitates with ErbB-2 and EGF receptor (EGFR) to promote cell migration and invasion in response to EGF (Gambaletta *et al.*, 2000; Dans *et al.*, 2001). EGF stimulates phosphorylation and activation of $\beta 4$ cytoplasmic domain, which leads to phosphoinositide-3 kinase (PI3K) activation and migration (Gambaletta *et al.*, 2000; L. Bonaccorsi, V. Carloni, M. Muratori, L. Formigli, S. Zecchi, C. Forti & E. Baldi, unpublished data). In addition, it has been recently shown that $\alpha 6 \beta 4$ may also represent a functional amplifier of biochemical signals leading to invasion by

potentiating growth factor activation of Ras- and PI3K-dependent pathways (Trusolino *et al.*, 2001).

In a previous study we demonstrated that the expression of $\alpha 6 \beta 4$ is higher in androgen-independent prostate carcinoma cell lines (PC3, DU145) with respect to androgen-sensitive PC cells, LNCaP and PC3 transfected with a full length hAR expression vector (PC3-AR) (Bonaccorsi *et al.*, 2000). PC3-AR cells were characterized by lower binding to laminin-1, anchorage-independent growth and EGF-mediated invasion compared with androgen-independent cells (Bonaccorsi *et al.*, 2000). The treatment with the synthetic androgen R1881 determined a further decrease of the invasion ability of these cells, without however, modifying the surface expression of $\alpha 6 \beta 4$ (Bonaccorsi *et al.*, 2000) and prospecting an effect of the androgen on EGF-mediated signalling related to invasion. In agreement with our results, in a different cell model, Cinar *et al.* (2001) have recently demonstrated, that invasion of PC cells is decreased by transfection with AR and by treatment with androgens. In order to study if androgens and AR have a direct effect on EGFR- $\alpha 6 \beta 4$ interaction and signalling leading to invasion in response to EGF, we investigated the effect of androgens on EGFR- $\alpha 6 \beta 4$ interaction, association of AR with EGFR and PI3K activation leading to invasion in PC3-AR cells in response to EGF (L. Bonaccorsi, V. Carloni, M. Muratori, L. Formigli, S. Zecchi, C. Forti & E. Baldi, unpublished data).

Results

EGFR and $\alpha 6 \beta 4$ colocalization in response to EGF is disrupted in PC3-AR cells

As mentioned above the interaction between EGFR and $\alpha 6 \beta 4$ is essential for carcinoma cell invasion (Mainiero *et al.*, 1996; Gambaletta *et al.*, 2000). By laser confocal immunofluorescence studies, we demonstrated that $\alpha 6$ and $\beta 4$ colocalized with the EGFR in basal conditions in PC3-Neo cells (transfected only with the vector) on laminin-1. In response to EGF $\alpha 6$, $\beta 4$ and EGFR redistributed and colocalized in discrete areas at the level of membrane ruffles. This pattern of colocalization of $\alpha 6 \beta 4$ and EGFR in response to EGF was not substantially affected by pre-treatment with androgens. On the contrary, in PC3-AR cells colocalization between $\alpha 6$, $\beta 4$ and EGFR was reduced with respect to PC3-Neo both in basal conditions and in response to EGF. In addition, pre-treatment with androgens further reduced redistribution of $\alpha 6 \beta 4$ and EGFR suggesting an interference of AR and androgens with the mechanisms leading to interaction between the two proteins. Immunoprecipitation studies conducted on PC3-Neo and PC3-AR cells confirmed that this colocalization was caused by an interaction between $\alpha 6 \beta 4$ and EGFR. After immunoprecipitation of EGFR, $\beta 4$ was detected in both cell lines and after immunoprecipitation of $\beta 4$, EGFR was detected in both cell lines indicating that the two proteins are constantly associated in our cell models. By probing the blot with an

antibody against phosphotyrosine residues we detected a tyrosine phosphorylated band at 200 kDa exactly migrating as $\beta 4$ and a phosphorylated band at 180 kDa corresponding to EGFR. Tyrosine phosphorylation of both bands increased in response to EGF in both cell lines, although EGF-induced tyrosine phosphorylation of $\beta 4$ was much less evident in PC3-AR cells. Moreover, in PC3-AR cells autophosphorylation of EGFR in response to EGF was lower compared with PC3-Neo suggesting a reduced activation of EGFR and $\beta 4$ integrin subunit by EGF in the presence of the AR.

The androgen receptor colocalizes with EGFR at the plasma membrane

Recent data indicate that besides its classical localization at the nuclear level, the AR may be targeted at the membrane level where interactions with molecules involved in growth factor signalling, such as src kinase family members (Migliaccio *et al.*, 2000), caveolin-1 (Lu *et al.*, 2001) and PI3K (Simoncini *et al.*, 2000) have been demonstrated. We investigated the localization of AR and EGFR by laser confocal immunofluorescence experiments. The AR localizes both to the nucleus and the cytoplasm of PC3-AR cells, although after stimulation with androgens, increased location to the nuclei was evident. Interestingly, a marked colocalization of AR with EGFR at the plasma membrane level was present. In response to EGF, colocalization between AR and EGFR was also evident at the level of intracellular granules, reflecting internalization of the EGFR-AR complex. When PC3-AR cells were stimulated with EGF in the presence of androgens, the pattern of colocalization between the two proteins was different compared with EGF alone because no discrete granules were present. After immunoprecipitation with an antibody against EGFR, a band at 110 kDa corresponding to AR was detected in PC3-AR cells. Tyrosine phosphorylated bands corresponding to EGFR and $\beta 4$ were also detected. Similarly, following immunoprecipitation of AR in PC3-AR cells, a tyrosine phosphorylated band corresponding to EGFR was also detected.

EGFR and PI-3 kinase are essential for EGF-mediated invasion in PC3-AR

An association between the EGFR and $\alpha 6\beta 4$ promotes the activation of several signalling pathways including PI3K (Shaw, 2001), which finally leads to cell migration and invasion. To evaluate whether PI3K activation is involved in Matrigel invasion of PC3-Neo cells, invasion assays were performed in the absence or presence of the PI3K inhibitor LY294002. Matrigel invasion in response to EGF was inhibited by the treatment with LY294002. EGF-mediated Matrigel invasion was also suppressed by ZD1839 a tyrosine kinase inhibitor selective for EGFR (Moasser *et al.*, 2001). To investigate whether PI3K activity was altered in PC3-AR cells, a PI3K activity assay was performed in both cell lines. We observed that basal and EGF-stimulated PI3K activity was reduced in PC3-AR cells with respect to

PC3-Neo. Pre-treatment with androgens determined a further decrease of EGF-stimulated PI3K activity in PC3-AR cells, whereas it was ineffective in PC3-Neo cells.

EGFR internalization is reduced in PC3-AR with respect to PC3-Neo

Recent evidences indicate that a correct endocytotic pathway is important for EGFR signalling by controlling the specificity of the response to the agonist (Wiley & Burke, 2001). It has also been demonstrated that internalized EGFR is enzymatically active, is still phosphorylated and maintains association with many adaptor proteins (Burke *et al.*, 2001). To investigate whether EGFR internalization following addition of the growth factor is affected by androgens, the expression of EGFR on cell surface by flow cytometry was studied. After EGF treatment (30 min), a significant decrease in cell surface expression of the receptor was detected in PC3-Neo cells. In PC3-AR cells this decrease of EGFR expression was less marked indicating a decreased internalization. Treatment with androgens further reduced EGFR internalization in response to EGF in PC3-AR but did not affect EGFR internalization in PC3-Neo cells.

Discussion

Androgen-insensitive PC cells are characterized by increased growth, adhesion, invasion and migration (Witkowski *et al.*, 1993; Bonaccorsi *et al.*, 2000; Cinar *et al.*, 2001). We show here that an interaction between AR and EGFR occurs in androgen-sensitive PC cell lines and provide evidence that the reduced invasive properties of these cells are because of a disruption of the interaction between the EGFR and $\alpha 6\beta 4$, decreased signalling leading to $\beta 4$ phosphorylation and lower PI3K activation in response to EGF. The mechanisms of AR-EGFR interaction as well as of reduced function of EGFR in PC3-AR cells are not clear at the moment but it is possible that AR-EGFR interaction may lead to a sequestration of the EGFR. In fact our results on EGFR internalization pattern indicate a disruption of the endocytotic process in PC3-AR cells compared with parental cells. Recent data have demonstrated that AR is able to interact with plasma membrane-associated proteins such as src kinase family members (Migliaccio *et al.*, 2000) and caveolin-1 (Lu *et al.*, 2001) that is also involved in EGFR signalling by inhibiting EGFR tyrosine kinase activity. In agreement with these results caveolin-1 might interact with AR and EGFR mediating a molecular complex that might negatively regulate invasion by inhibiting EGFR redistribution, signalling and internalization (see Fig. 1: working hypothesis). In conclusion, in androgen-sensitive prostate carcinoma cells, androgens and AR contribute to confer a less malignant phenotype of these cells both by reducing the expression of $\alpha 6\beta 4$ (Bonaccorsi *et al.*, 2000) and by interfering with EGFR- $\alpha 6\beta 4$ interaction




Figure 1. Working hypothesis.

and signalling leading to invasion through an interaction between AR and EGFR (L. Bonaccorsi, V. Carloni,

M. Muratori, L. Formigli, S. Zecchi, C. Forti & E. Baldi, unpublished data).

References

- Amanatullah, D. F., Reutens, A. T., Zafonte, B. T., Fu, M., Mani, S. & Pestell, R. J. (2000) Cell-cycle dysregulation and the molecular mechanisms of prostate cancer. *Frontiers in Bioscience* **5**, d372–d390.
- Bonaccorsi, L., Carloni, V., Muratori, M., Salvatori, A., Giannini, A., Carini, M., Serio, M., Forti, G. & Baldi, E. (2000) Androgen receptor expression in prostate carcinoma cells suppresses $\alpha 6 \beta 4$ integrin-mediated invasive phenotype. *Endocrinology* **141**, 3172–3182.
- Burke, P., Schooler, K. & Wiley, H. S. (2001) Regulation of epidermal growth factor receptor signaling by endocytosis and intracellular trafficking. *Molecular Biology of the Cell* **12**, 1897–1910.
- Cinar, B., Koenenman, K. S., Edlund, M., Prins, G. S., Zhau, H. E. & Chung, L. W. (2001) Androgen receptor mediates the reduced tumor growth, enhanced androgen responsiveness, and selected target gene transactivation in a human prostate cancer cell line. *Cancer Research* **61**, 7310–7317.
- Dans, M., Gagnoux-Palacios, L., Blaikie, P., Klein, S., Mariotti, A. & Giaccotti, F. G. (2001) Tyrosine phosphorylation of the beta 4 integrin cytoplasmic domain mediates Shc signaling to extracellular signal-regulated kinase and antagonizes formation of hemidesmosomes. *Journal of Biological Chemistry* **276**, 1494–1502.
- Gambaletta, D., Marchetti, A., Benedetti, L., Mercurio, A. M., Sacchi, A. & Falcioni, R. (2000) Cooperative signaling between alpha (6) beta (4) integrin and ErbB-2 receptor is required to promote phosphatidylinositol 3-kinase-dependent invasion. *Journal of Biological Chemistry* **275**, 10604–10610.
- Juliano, R. L. & Verner, J. A. (1993) Adhesion molecules in cancer: the role of integrins. *Current Opinions in Cell Biology* **5**, 812–818.
- Lu, M. L., Schneider, M. C., Zheng, Y., Zhang, X. & Richie, J. P. (2001) Caveolin-1 interacts with androgen receptor. A positive modulator of androgen receptor mediated transactivation. *Journal of Biological Chemistry* **276**, 13442–13451.
- Mainiero, F., Pepe, A., Yeon, M., Ren, Y. & Giaccotti, F. G. (1996) The intracellular functions of $\alpha 6 \beta 4$ integrin are regulated by EGF. *Journal of Cell Biology* **134**, 241–253.
- Manos, E. J., Kim, M. L., Kassis, J., Chang, P. Y., Wells, A. & Jones, D. A. (2001) Dolichol-phosphate-mannose (DPM3)/prost-1 is a novel phospholipase C-gamma regulated gene negatively associated with prostate tumor invasion. *Oncogene* **20**, 2781–2790.
- Meredith, J. E., Winitz, S., McArthur Lewis, J., Hess, S., Ren, X.-D., Renshaw, M. W. & Schwartz, M. A. (1996) The regulation of growth and intracellular signalling by integrins. *Endocrine Reviews* **17**, 207–220.
- Migliaccio, A., Castoria, G., Di Domenico, M., de Falco, A., Bilancio, A., Lombardi, M. et al. (2000) Steroid-induced androgen receptor-oestradiol receptor beta-*Src* complex triggers prostate cancer cell proliferation. *EMBO Journal* **19**, 5406–5417.
- Moasser, M. M., Basso, A., Averbuch, S. D. & Rosen, N. (2001) The tyrosine kinase inhibitor ZD1839 ('Iressa') inhibits HER2-driven signaling and suppresses the growth of HER2-overexpressing tumor cells. *Cancer Research* **61**, 7184–7188.
- Shaw, L. M. (2001) Identification of insulin receptor substrate 1 (IRS-1) and IRS-2 as signaling intermediates in the $\alpha 6 \beta 4$ integrin-dependent activation of phosphoinositide 3-OH kinase and promotion of invasion. *Molecular Cell Biology* **15**, 5082–5093.
- Simoncini, T., Hafezi-Moghadam, A., Brazil, D. P., Ley, K., Chin, W. W. & Liao, J. K. (2000) Interaction of oestrogen receptor with the regulatory subunit of phosphatidylinositol-3-OH kinase. *Nature* **407**, 538–541.

- Sumitomo, M., Shen, R., Walburg, M., Dai, J., Geng, Y., Navarro, D., Boileau, G., Papandreou, C. N., Giancotti, F. G., Knudsen, B. & Nanus, D. M. (2000) Neutral endopeptidase inhibits prostate cancer cell migration by blocking focal adhesion kinase signalling. *Journal of Clinical Investigations* **106**, 1399–1407.
- Trusolino, L., Bertotti, A. & Comoglio, P. M. (2001) A signalling adapter function for $\alpha 6 \beta 4$ integrin in the control of HGF-dependent invasive growth. *Cell* **107**, 643–654.
- Van der Kwast, T. H., Schalken, J., Ruizeveld de Winter, J. A., Van Vroonhoven, C. C., Muller, E. & Boersma, W. (1993) Androgen receptors in endocrine therapy-resistant human prostate cancer. *International Journal of Cancer* **48**, 189–193.
- Wiley, H. S. & Burke, P. M. (2001) Regulation of receptor tyrosine kinase signaling by endocytic trafficking. *Traffic* **2**, 12–18.
- Witkowski, C. M., Rabinovitz, I., Nagle, R. B., Affinito, K. S. & Cress, A. E. (1993) Characterization of integrin subunits, cellular adhesion and tumorigenicity of four human prostate cell lines. *Journal of Cancer Research Clinical Oncology* **119**, 637–644.

Received 15 May 2002; revised 12 July 2002; accepted 12 July 2002