UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Biological invasions in inland waters: an overview.

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Biological invasions in inland waters: an overview / F. GHERARDI. - STAMPA. - (2007), pp. 3-25.

Availability:

This version is available at: 2158/258021 since:

Publisher:

Springer

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

Biological invasions in inland waters: an overview

Francesca Gherardi

INTRODUCTION

The value of inland waters to humankind is obviously infinite and the impacts of changes in the goods and services they provide have a strong impact on human welfare. Lakes/ivers and wetlands currently contribute 20% of the estimated annual global value of the entire biosphere amounting to US\$100 trillion per year (Costanza *et al.* 1997). These elevated numbers may have increased the present general concern about the increasing degradation of freshwater systems, associated with the rapid extinction rate of their biodiversity, in some cases even matching that of tropical forests (Ricciardi and Rasmussen 1999).

Together with other anthropogenic sources of disturbance, such as the impoundment of rivers (e.g. dams and weirs, water removal), water quality deterioration (e.g. pollution, eutrophication, acidification), habitat degradation and fragmentation (e.g. channelization and land use change), and over-exploitation, the introduction of non-indigenous species (NIS) into freshwater systems is today regarded as the main driver of biodiversity change (Millennium Ecosystem Assessment 2005). The effects of such a driver have been estimated to be greater in freshwater than in terrestrial ecosystems (Sala *et al.* 2000). This is particularly apparent in lakes where biological invaders have recently been recognized as one of the greatest causes of species extinctions (Ricciardi 2001).

Inland waters have been the theatres of spectacular biological invasions. Well-known cases are the introduction of the Nile perch *Lates niloticus* (Linnaeus) to Lake Victoria followed by the elimination of about 200 species of haplochromine cichlids (Craig 1992), the alteration of the Laurentian Great Lakes communities and ecosystems by sea lamprey *Petromyzon marinus* Linnaeus, zebra mussels *Dreissena polymorpha* (Pallas), and other invaders (MacIsaac *et al.* 2001), and the complete domination of lowland rivers in the western USA by non-indigenous fish and invertebrates (Moyle and Light 1996a). In several freshwater systems, other less celebrated dramas are however ongoing with the intervention of several, previously unsuspected actors, such as *Lepomis gibbosus* (Linnaeus) (Chapter 15) and *Carassius auratus* (Linnaeus) (Chapter 13) among fish, *Asellus bogammarus villosus* (Sowinsky) (Chapters 12 and 27), *Gmelinoides fasciatus* (Sowinsky) (Chapter 26), and *Pontogammarus robustoides* (Sars) (Chapter 25) among crustaceans, and *Rana catesbeiana* Shaw among amphibians (Chapters 12 and 38). This confirms that invasions by NIS are pervasive and highly diverse phenomena in fresh waters but also that our predictive ability may be limited. Meanwhile, other apparently harmless NIS are spreading (see potamid crabs in southern France, Chapter 3).

The reasons that freshwater systems are vulnerable to NIS are several, including the higher intrinsic dispersal ability of freshwater species compared with terrestrial organisms (Beisel 2001). Lakes and some streams are comparable to islands in that their geographic isolation has led to local adaptation with the evolution of many endemisms and sometimes to a low biodiversity (Lodge 1993). The extensive introduction of organisms in inland waters, whether inadvertent (e.g. via ship ballast, artificial/natural canals, or estuarine salt bridges, Chapters 17, 21, and 22; as parasites of other introduced species such as the oomycete *Aphanomyces astaci*, Chapter 6) or deliberate (e.g. stocked fish and crayfish, Chapters 20 and 31; intentional releases of pets or ornamental organisms, Chapters 8 and 9), is a direct consequence of the intensity with which humans utilize these systems for recreation, food sources, and commerce (Rahel 2000, Ricciardi 2001). Human-mediated dispersal of crustacean plankton, for instance, might exceed the natural rate by up to 50,000% (Hebert and Cristescu 2002). And the frequency of species invasions in inland water systems is likely to continue to grow commensurate with enhanced international commerce and human exploitation of these communities.

Finally, freshwater systems are subject, especially at higher latitudes, to altered seasonal temperature regimes due to global climatic warming, and especially in developed countries, to strong human disturbance. In fact, NIS are migrating to new areas where the climate has warmed, such as introduced warm-water fish [e.g. *Micropterus salmoides* (Lacepède), *Lucioperca macrochirus* Rafinesque, and *Lepomis cyanellus* (Rafinesque)] that are spreading in North America into higher latitudes and altitudes (Eaton and Scheller

more than pristine systems, disturbance results in the reshaping of space and energy resources that are available to indigenous and non-indigenous species and may open new vacant niches for the most adaptable and tolerant invaders (Ross *et al.* 2001).

The vulnerability of inland waters to biological invasions is a cause of the complete domination of vast waterscapes in certain regions by NIS, such as the water hyacinth [*Eichornia crassipes* (Martius) Solms] in many tropical lakes and rivers (Chapter 10) and the red swamp crayfish [*Procambarus clarkii* (Günther)] in several waterbodies of southern Europe (Gherardi 2006, Chapter 2). Species diversity may be extraordinarily high in, for instance, large rivers of developed countries that usually host dozens to hundreds of NIS (Mills *et al.* 2003, Chapter 12). The Hudson River, for instance, contains more than 100 species of non-indigenous fish, vascular plants, and large invertebrates, a considerable fraction of which are ecologically important – such as Eurasian watermilfoil (*Myriophyllum spicatum* Linnaeus), purple loosestrife (*Lythrum salicaria* Linnaeus), zebra mussel (*D. polymorpha*), Atlantic rangia (*Rangia cuneata* Gmelin) and common carp (*Cyprinus carpio* Linnaeus) (Strayer *et al.* 2005). Some regions are particularly affected by species introductions: the New Zealand fish fauna contains 30 NIS (53% of the total) (Vitousek *et al.* 1997); isolated islands have more non-indigenous than indigenous fish species (Hawaii: 19 v. 10) but also continental areas have relatively large numbers of non-indigenous fish species (California: 42 v. 76, Brazil: 76 v. 517; references in Vitousek *et al.* 1996); at least 76 fish species belonging to 21 families have been introduced into European fresh waters (Lehtonen 2002), of which 51 have been established.

Species originating from diverse biogeographical areas now coexist in some river basins; in the Rhine, indigenous crustaceans [*Gammarus pulex* (Linnaeus)] coexist with North American species [*Gammarus tigrinus* Sexton and *Orconectes limosus* (Rafinesque)], Mediterranean species (the freshwater shrimp *Atyaephyra desmaresti* Millet), and Ponto-Caspian species [*Gammarus roeseli* Gervais and *Dreissena polymorpha* (*Gammarus villosus* (Sowinsky)] (Beisel 2001). Biotic homogenization is constantly increasing; freshwater fish similarity among the States of the United States amounts today to 7% (Rahel 2000) and some of them, such as Arizona and Montana, which previously had no fish species in common, now share more than 30 species. National borders are obviously irrelevant and they provide no barrier to the natural dispersal of NIS: the Nile perch released in Tanzania spread to other countries bordering Lake Victoria (Welcomme 1988) and *P. clarkii* introduced into Spain invaded Portugal via the common hydrographic basins (Gherardi 2006).

Some freshwater systems function as “hotspots” where NIS accumulate. The Great Lakes system contains over 145 non-indigenous invertebrates, pathogens, algae, fish, and plants, with approximately 75% originated from Eurasia of which 57% are native to the Ponto-Caspian region (Mills *et al.* 2003).

a wide array of taxa, such as mussels [*D. polymorpha*, *Dreissena* sp. (Andrusov)], amphipods (*Echinogammarus ischnus* Stebbing), cladocera [*Cercopagis pengoi* (Ostroumov)], harpacticoid copepods [*Nitocra incerta* (Richardson) and *Schizopera borutzkyi* (Monchenko)], and fish [*Neogobius melanostomus* (Pallas)], *Proterorhinus marmoratus* (Pallas), *Gymnocephalus cernuus* (Linnaeus)]. It has been ascertained that Ponto-Caspian species reach the Great Lakes via ballast along five shipping “corridors” (MacIsaac *et al.* 2001). Four of these corridors require the first transfer of species via rivers and canals to ports on the North and Baltic seas that, in their turn, function as “hubs”, acting as the starting point for other ships that transport these species in secondary invasions to ports in North America and, potentially, in East Asia, San Francisco Bay, and Australia (e.g. Cohen and Carlton 1998, Ruiz *et al.* 2000).

Finally, many freshwater invaders are moved among biogeographic realms within continents and are transported among continents in association with human economic activity and trade globalization that benefit millions worldwide (MacIsaac and Shrader-Frechette 2003). The inevitable tension between two often competing goals – increasing economic activity and protecting the environment from invasive species – make it difficult to justify the need for decision-makers to contain the spread of these species and to mitigate the environmental risk they pose. For instance, a number of issues has been raised in favor of the introduction of introducing crayfish (Gherardi 2006, Chapter 28). First, in the absence of indigenous species, invasive crayfish were claimed to occupy vacant niches, constituting the unique large macro-consumer within polluted or eutrophic waters, where the native fauna has already been severely decimated (Gherardi *et al.* 2000). The second claim is that they constitute abundant prey for native threatened birds and mammals, like several Ardeidae and the otter (Barbaresi and Gherardi 2000, Rodríguez *et al.* 2005). Third, from a socio-economic perspective, introduced crayfish have contributed to: (1) the restoration of traditional habits, e.g. by crayfishing in Sweden and Finland (Kirjavainen and Sipponen 2004); (2) economic benefits for local crayfishermen, e.g. the Spanish crayfishermen; (3) diversification of agriculture to include astaciculture, e.g. by crayfish farmers in Britain and in Spain; and (4) increased trade between European countries inside Europe as well as between European and extra-European countries (Ackefors 1999).

THE CURRENT STATE OF RESEARCH

Studies on the identity, distribution, and impact of freshwater NIS and the dynamics of their invasion have increased exponentially since the 1980s, resulting in a flood of publications particularly abundant in the last decade (Fig. 1). This pattern of growth in the literature is a reflection of the increasing popularity that invasion biology has gained as an appealing area of research.

Fig. 1 Cumulative number of publications dealing with non-indigenous species (total number = 502). Papers were identified via keywords from Biosis analyzed between January 1977 and December 2005.

among ecologists (Kolar and Lodge 2001). The overall number of published articles is, however, significantly biased towards terrestrial invaders (Fig. 2). This is not surprising: terrestrial systems are the most visible and accessible habitats for humans and, as such, have received the preponderance of ecological attention.

The majority of studies of freshwater invaders has been conducted in North America, and mostly in the Great Lakes (Fig. 3A), and centered on animals (74%) more than on plants (20%), whereas a small fraction (6%) analyzed both kingdoms simultaneously. Of all the animal taxa, fish, particularly salmonids, have received the greatest scientific attention (Fig. 3B), as the result of

Fig. 2 Frequency distribution of research articles published in the journal *BioInvasions* (Springer) since 1999 distinguished among habitats (total number = 33).

Fig. 3 Frequency distributions of research articles published in the journal *Biological Invasions* (Springer) since 1999 dealing with inland waters distinguished among continents (A) and animal taxa (B) (total number = 69).

perceived ecological role in aquatic food webs and their economic importance to humans. In the other taxa, dreissenids among mollusks and crayfish among crustaceans had been most often studied. Only recently has been the taxonomic coverage of freshwater invaders broadened. New privileged study animals have been Eleotridae and Poeciliidae among fish (e.g. Bedarf *et al.* 2001, Lahetähti *et al.* 2006, Pusey *et al.* 2006); the Asian clam [*Corbicula fluminea* (Müller)], the golden mussel [*Limnoperna fortunei* (Dunker)], and the golden snail [*Pomacea canaliculata* (Lamarck)] among mollusks (e.g. Darrigran *et al.* 2005, Carlsson and Lacoursière 2005, Boltovskoy *et al.* 2006, Oliveira *et al.* 2006, Yusa *et al.* 2006); and the spiny waterflea (*Bythotrephes longimanus* Leydig) among cladocerans among crustaceans (e.g. Çelik *et al.* 2002, Shurin and Havel 2002, Branstrator *et al.* 2006).

ARE GENERALIZATIONS POSSIBLE?

A consequence of the concentrated interest on terrestrial biomes and limited geographic and taxonomic breadth in fresh waters is that tradi-

have been rarely tested in aquatic organisms (Becker 2001). However, a number of generalizations about freshwater invasion is emerging today. These can be derived from two main approaches (Moyle and Light 1996b): the first approach is based on case studies of invaders and their distribution (e.g. *P. clarkii*, Chapter 4; mollusks, Chapter 5; amphibians and reptiles, Chapter 7; plants, Chapter 19), whereas the second approach aims to extend recent developments in ecological theory to freshwater invaders (e.g. Chapters 19 and 23), in which the focus has been mainly directed to the interaction between the invader and the receiving community and to the biological characteristics of both the invading species and the ecosystem being invaded. Based on the examples provided by the recent literature, a list of 15 general statements characterizing some of the known events of biological invasion in inland waters can be drawn. The list follows.

(1) The establishment and spread success of freshwater NIS often exceeds the 10% value predicted by the “tens rule” (Williamson 1996).

This is seen in the results obtained by Jeschke and Strayer (2005), who analyzed the introductions of vertebrates between Europe and North America (including Canada). Using corrected data for unrecorded introductions, the authors showed that, of the 220 and 713 fish species native to Europe and North America respectively, 11% and 6% have been introduced into Europe and North America respectively, and 36% and 49% have become established after slightly less than a decade, while 56% and 63% of the established fish had spread and become invasive. On a more global scale, Ruesink (2005) used a database of intentional international transfers of freshwater fish and found that up to 30% of the introduced fish became established and 22% of the established cases exerted a documented impact (i.e. changes in food availability, habitat structure, nutrient dynamics, or top-down trophodynamics).

(2) Propagule pressure is often a major predictor of the establishment of freshwater organisms.

Recent findings showed that the large number of propagules present in the inoculating population, such as the thousands of zebra and quagga mussels carried in the ballast of cargo ships, and the frequencies of sequential introductions, such as multiple introductions of “desired” species, are positively correlated with invasion success (Lonsdale 1999, Kolar and Lodge 2001, MacIsaac 2000, Ricciardi 2001). For instance, as showed by Ruesink (2005), introduced fish species were more likely to establish when humans intended their introduction (76%) rather than when fish were cultivated or used with no explicit desire for naturalization (57%).

several instances in which introduced populations appear to be immune to bottlenecks – usually depleting genetic variation (the “genetic paradox”)

Aquatic organisms introduced in large numbers via ballast or subject to multiple introductions can carry a large fraction of the genetic variability of source populations or bring genetic races from different parts of their native range (Stepien *et al.* 2002). Hence, many colonizers arrive with a high phenotypic and genetic diversity. This adds to other features that may favor their establishment in the recipient areas, such as the fast acquisition of genetic variability after their arrival that results from such sources as hybridization with closely related organisms, epistasis (i.e. an interaction in which one gene influences the expression of another), or the potential for chromosomal restructuring (e.g. inversion, translocation, or duplication (Cox 2004).

(4) Failures of NIS to establish derive most often from their inability to meet the “environmental resistance” on the part of the recipient community – the different regimes of temperature, current, water chemistry, or abiotic resources.

Several examples from different taxa support this statement. Moyle and Light (1996a), for instance, showed that freshwater fish invading North American basins are likely to become established when abiotic conditions are appropriate, regardless of the biota already present. The narrow thermal tolerance of *C. fluminea* may explain its absence from most of the Great Lakes systems (Ricciardi 2001), whereas salinity of that system was too low to allow for successful reproduction of the Chinese mitten crab *Eriocheir sinensis* (Edwards), notwithstanding the frequent introductions of this latter species over the past decades in ship ballast (MacIsaac 1999). The general harshness of the environment may reduce the ability of non-indigenous fish to invade. For example is Eagle Lake, California, which is a highly alkaline (pH: 8–9) temperate lake containing only four indigenous fish species (Moyle and Light 1996b). An attempt to introduce fish failed in the long term, including the introduction of the early 20th century of the largemouth bass, *M. salmoides*.

(5) As a consequence of (4), success in the establishment of freshwater invaders may depend on a close match between their physiological requirements and the environmental characteristics of the system being invaded.

Species from nearby areas are more likely to be successful invaders than those from more distant locations, as found for fish species in North America (Light 1996). Species are most likely to be successful if they are adapted to the local, highly seasonal hydrological regime of the recipient environment (Moyle and Light 1996b).

from nearby areas became established because the artificial flow regime of the recipient river closely resembled that of their native streams (Bestgen 1989). Convergent salinity conditions in donor and recipient ecosystems play a key role in the success of invaders in the Great Lakes (MacIsaac *et al.* 2001). An additional prerequisite for successful invasion that allows a species to survive transportation (e.g. in ballast; Bailey *et al.* 2004) and to become established in a recipient area is its euryoeciousness, i.e. its ability to tolerate a wide range of environmental conditions (Ricciardi and Rasmussen 1998). For instance, the range of salinity in which a species can live and reproduce provides a ready basis for discrimination between invasive and non-invasive North American and East European gammarid amphipods (Devin and Beisel 2007). In general, areas with a wide salinity gradient, like the Baltic Sea, may offer a range of hospitable conditions for invaders, functioning as hot spots of xenodiversity (Leppäkoski *et al.* 2002). Finally, the increased ionic content of large European rivers as the result of pollution has allowed salt tolerant species to spread in river basins in recent decades (Ketelaars *et al.* 1999).

(6) Demographic factors (sexual precocity, fecundity, and number of generations per year; Lodge 1993, Chapter 12), biological plasticity (Chapters 14 and 16), and/or the ability to overcome the biotic resistance posed by the recipient community (the complex of native predators, parasites, pathogens, and competitors, and previously introduced species) may be neither essential nor sufficient for freshwater species to become invasive.

Usually, *r*-selected crayfish (e.g. *P. clarkii*) rather than *K*-selected species [*Austropotamobius pallipes* (Lereboullet)] have a high probability of spreading. Fecundity and number of generations per year often are not sufficient to explain why the amphipod *G. pulex* has invaded the streams in Northern Ireland and it outcompetes *G. duebeni* Liljeborg (Devin and Beisel 2007). Often, the magnitude of an invader's impact may be predicted by its "taxonomic distinctiveness" within the recipient community (Ricciardi and Atkinson 2004). Indeed, the lack of evolutionary experience with the invader – meaning the absence of competitors, predators, or parasites and the inability to respond to them with appropriate behavior – may predispose communities to be altered by invaders (e.g. Diamond and Case 1986). For instance, eliminations of indigenous species by *D. polymorpha* have rarely been reported from the invaded European rivers whose native fauna was previously exposed to *Dreissena* during the Pleistocene era (Ricciardi *et al.* 1998). Similarly, the introduced tilapiine species, *Oreochromis niloticus* (Linnaeus), is one of the few fish species to persist in Lake Victoria in large numbers in face of the Nile perch invasion as the result of its evolutionary experience with similar predators (Moyle and Light 1996b). In some times, the interaction between two species that do not share an evolutionary

tionary release" (Schlaepfer *et al.* 2003). For instance, indigenous pre-
 anuran tadpoles and metamorphs) may be unable to recognize intro-
 predators (e.g. the introduced *R. catesbeiana* in the western USA; Rose
 Schwalbe 2002) and their style of preying. As a result, the naïve predator
 released from the difficulties of finding a prey (Kiesecker and Blaustein 1999)
 they may dispose of a high availability of food during the establishment
 of their invasion, which is a precondition of their fast spread. The phenom-
 of evolutionary release might explain the paradox of why invasive species
 sometimes enjoy a competitive advantage over locally adapted species, although
 there would be a priority effect for residents (e.g. Shea and Chesson 2002).
 Schlaepfer *et al.* 2005). Also a reduced attack from natural enemies (pre-
 and parasites) encountered outside their natural range gives some species
 ability to spread and to become invasive, as predicted by the "enemy release
 hypothesis" (ERH) (e.g. Torchin *et al.* 2003). However, generalizations about
 the role of the naïveté of introduced species and of their potential lack of enemies
 in the recipient community may not be possible in freshwater systems. Some
 examples, in fact, provide contrasting evidence. The sea lamprey, *P. marinus*,
 eliminated large fish from Lake Michigan even if some of these species [e.g.
 lake trout *Salvelinus namaycush* (Walbaum)] coexisted with the lamprey in
 lakes where the species have been together for thousands of years (Walbaum
 1986). In addition, the ERH has been verified in relatively few organisms
 and subject to limited criticism (Colautti *et al.* 2004), being only one of
 several hypotheses that can explain the abundance and/or the impact of
 given invader (Enemy Inversion Hypothesis, climatic variables, selective
 "invasive" genotypes, human disturbance, etc.).

**(7) In fresh waters, species-rich communities may be as vulnerable
 to invasion as less speciose, less biologically
 "sophisticated" communities.**

This statement contrasts with one of the most well-established generalizations
 in the invasion literature since Elton (1958), i.e. that communities with
 diversity and complexity are the least susceptible to invasion because of the
 strength of the community interactions (e.g. Lodge 1993, Levine and D'Antonio
 1999, Kennedy *et al.* 2002, Shea and Chesson 2002). Numerous examples at
 both the global and local levels demonstrate that often the opposite occurs.
 freshwater systems (Moyle 1986, Ross *et al.* 2001). Jeschke and Strayer (2003)
 for instance, showed that there is no clear difference in the probability of
 species becoming established and spreading once introduced from Europe to
 North America and *vice versa*, suggesting that, at the global level, the biota of
 North America do not offer more resistance to invaders, notwithstanding the fact
 is less disturbed by humans and species-richer than Europe. At a local level,
 Lake Victoria, which once contained the richest endemic fish community

because it encountered neither predation nor competition from indigenous fish (Goldschmidt *et al.* 1993). Mississippi River, once the most speciose temperate rivers, has been invaded by several non-indigenous fish, including common carp *C. carpio*, goldfish *C. auratus*, grass carp *Ctenopharyngodon* (Valenciennes), striped bass *Morone saxatilis* (Walbaum), rainbow smelt *Osmerus mordax* (Mitchill), rainbow trout *Oncorhynchus mykiss* Walbaum, and white catfish *Ictalurus catus* (Linnaeus) (Burr and Page 1986). Similarly, the zebra mussel has become established at high densities throughout the Mississippi River basin, which contains the world's richest endemic assemblage of freshwater mussels (Ricciardi *et al.* 1998).

The above examples, however, contrast with the results obtained by experimental disturbance treatments in pond zooplankton communities (Smith and Shurin 2006). Shurin (2000) found that reducing the abundance of native species allowed four times as many invaders to establish and to reach 16 times greater total abundance, therefore showing that in some instances local interactions may be strong enough to exclude a large fraction of potential invaders.

(8) Often, freshwater NIS, instead of interfering with one another, facilitate each other's establishment and/or continued existence, and therefore increase the likelihood and the magnitude of their ecological impacts as predicted by the phenomenon of "invasional meltdown" (Simberloff and Von Holle 1999, Simberloff 2006).

Invasive species may facilitate further invasions by direct effects – providing benefits to another invader – and indirect effects – reducing an invader's enemies or enhancing its prey (Ricciardi 2001). Strong experimental evidence was provided by Adams *et al.* (2003), who showed that introduced fish facilitated invasion by the bullfrog (*R. catesbeiana*) in western North America by preying on native macroinvertebrates (such as dragonfly nymphs) that otherwise precluded establishment or severely limited the numbers of frogs. In the Great Lakes, mutualistic, commensal, and asymmetric exploitative interactions facilitated the survival and population growth of many invaders (Ricciardi 2001). *Dreissena polymorpha* facilitated colonization by several invertebrate NIS (Ricciardi *et al.* 1998) because it increased the surface area and structural heterogeneity, creating settling sites, providing refuge, and trapping sediments and biodeposits. Additionally, *Dreissena* generates filtration currents that are exploited by other invertebrates (Stewart and Haynes 1994) and serves as a food source for several introduced fish, i.e. white bass [*Morone chrysops* (Rafinesque)] and round goby [*Neogobius melanostomus* (Pallas)] (Fausch 1993). Its invasion also increased the spread of Eurasian watermilfoil (MacIsaac 1996). The reduction of piscivores by the parasite sea lamprey had an indirect effect that paved the way for invasion by the planktivore alewife.

Finally, sequential invasions by Ponto-Caspian species complicated the invasion of parasitic organisms, such as the trematode *Bucephalus polymorphus* of western European origin. The introduction of the first intermediate host (the zebra mussel) of the trematode and its definitive host [the pike *Stizostedion lucioperca* (Linnaeus)] allowed it to spread, causing high mortality in its secondary intermediate hosts, the indigenous cyprinids (Comber and Le Brun 1990).

(9) There is still a poor state of knowledge of whether invasive species are the “drivers” of the extinction of indigenous populations or species, or merely the “passengers” along for the environmental ride (MacDougall and Turkington 2005).

Local and global extinctions frequently overlap invasions in space and time. For instance, the loss of genetically distinct populations of unionids in North America has been accelerated by a factor of 10 after the invasion of zebra mussels (Ricciardi *et al.* 1998). Of the 40 fish species known to have become extinct since 1890 in North America, 27 were negatively affected by the introduction of NIS (Wilcove and Bean 1994). However, the dominance of NIS might be an indirect consequence of habitat modifications that by themselves lead to both indigenous species loss and NIS invasion (Gurevitch and Padilla 2004, Didham *et al.* 2005). For instance, the unionid declines began before the introduction in the mid-1980s of zebra mussels and were caused by several stressors, such as habitat destruction and deterioration resulting from water diversion, erosion, an increase in eutrophication (which causes periods of anoxia), pesticides, loss of host fish for parasitic unionid larvae, harvest for the button industry and harvest for the pearl industry (reviewed in Gurevitch and Padilla 2004). In some instances, successful water invaders have been integrated without eliminations of species from communities being invaded. In Lake Malawi, East Africa, 12 species of benthic feeding haplochromine cichlids from one part of the lake were introduced to another part of the lake in which they were absent without any apparent changes in the abundance of the local species (Trendall 1988). After the completion of the Panama Canal in 1914 and the consequent creation of a freshwater corridor between the Rio Chagres on the Caribbean slope and the Rio Grande on the Pacific slope of the Isthmus of Panama, the freshwater fish assemblages of previously isolated drainage basins were enabled to change. However, no cases of local extinctions were recorded but species richness increased by 10% in the Rio Chagres and 22% in the Rio Grande (Smith *et al.* 2004). And there are documented cases (see statements of Gurevitch and Padilla 2004) of indigenous species that, given enough time, learn or evolve the ability to escape the “evolutionary trap” caused by an invasive species (Schlaepfer *et al.* 2005).

existence of indigenous species (e.g. predation, parasitism, vectoring of pathogens, and competition; Mack *et al.* 2000), the most underestimated is hybridization with indigenous species (Olden *et al.* 2004).

Hybridization is thought to alter the integrity of the endemic gene pool of unionids, crayfish, and fish; it produces hybrid swarms that eliminate indigenous taxa often in a very short time frame (Perry *et al.* 2002). Examples are several, especially in fish. Within a 4-year period following its introduction of non-indigenous pupfish, *Cyprinodon variegatus* Lacépède, was involved in a large-scale introgressive hybridization event with the endemic *Cyprinodon tularumensis* Echelle and Echelle in New Mexico, USA (Echelle and Connor 1999). Similarly, anadromous populations of wild brown trout (*Salmo trutta* Linnaeus) were highly introgressed by stocking with hatchery fish and eventually reduced their fitness (Hansen 2002).

(11) Introduced species have effects at multiple ecological levels in freshwater systems (Simon and Townsend 2003, Chapter 24), but a few studies, mostly focused on salmonids (Simon and Townsend 2003), have analyzed this multifaceted impact (Parker *et al.* 1999).

At the level of individual organisms, invaders may alter the behavior of indigenous species, influencing habitat use and foraging. At the population and community levels, they may induce changes in the abundance or distribution of indigenous species and affect both direct and indirect interactions among populations, respectively. Finally, at the ecosystem level, invaders may change the pattern and magnitude of movements of energy and nutrients.

(12) A first strong danger posed by freshwater invaders to native biodiversity arises if they are either macro-enemies (predators or grazers) or micro-enemies (pathogens or parasites) (Williamson 1996).

Specifically, the NIS posed on the top of the food web or those that are generalized predators (Williamson 1996) are likely to produce marked effects on ecosystem processes (see the case of *P. clarkii*; Chapters 29 and 30) because their impact can “cascade” through the entire food web, altering both ecosystem processes and the behavior of the indigenous species (Townsend 1996). This is especially true in aquatic systems, in which trophic cascades appear to be more common than in terrestrial biomes (Strong 1992). For instance, in California, the most successful fish invaders have been piscivores and omnivores (Lorenzen and Light 1996a), while detritivorous fish seemed to have little effect on indigenous fish assemblages (Power 1990).

native biota is their role of "ecosystem engineers", i.e. species that "directly or indirectly control the availability of resources to other organisms by causing physical state changes in biotic or abiotic materials" (Jones *et al.* 1994).

Engineering organisms may cause physical modifications to the environment and influence the maintenance or creation of habitats. Their ecological effects on other species occur because of physical state changes caused, either directly or indirectly, by the engineer, and because, as engineers, they affect the distribution and use of resources by other species. Zebra mussels have all the properties of ecosystem engineers. They change the characteristics of biotic and abiotic environments by their presence and activities, especially their feeding and filtering. These changes are system-wide, affecting species composition, species interactions, community structure, and ecosystem properties (Karatayev *et al.* 2002, Chapters 32 and 33).

(14) In freshwater systems, as in other biomes, invaders may be subject to rapid evolutionary changes that influence several life history characteristics (Cox 2004).

Once established, NIS are freed from the constraints of the gene flow from their parent population and from the biotic pressures of former enemies, they are subject to altered selection pressures, and they impose strong new evolutionary pressures on the indigenous species. Substantial evolution may take place on relatively short timescales (Carroll and Dingle 1996). For instance, following introduction to Pacific rivers in North America, the American shad, *Alosa sapidissima* (Wilson), evolved geographic changes in its life history pattern in less than a century (summarized in Dingle 1980). In its native rivers in the eastern North America, fecundity and the yearly number of spawns vary as a function of latitude, with reduced clutch size and increased repeat spawning being more prevalent in northern versus southern rivers. Fish from Pacific rivers are 32–77% (vs. 20–40% from Atlantic rivers) repeat spawners; their age at maturity varies from 3.3 to 3.8 years for males (vs. 4 years) and from 4 to 4.5 years for females (vs. 4.6 years), and their mean lifetime fecundities range from 321,000 to 500,000 eggs (vs. 300,000–350,000). These variations are a function of latitude and water temperature, reflecting rapid post-invasion evolution under selection by local environmental conditions. The western mosquitofish [*Gambusia affinis holbrooki*] (Baird and Girard) native to North America has shown rapid genetic change at several locations into which it has been introduced (references in Cox 2004). A period of about 70 generations was sufficient to induce adaptive changes in the life history of the populations introduced into Hawaii, whereas, in the populations introduced into thermal springs in Nevada, changes in body composition and size at maturity required about 110–165 generations.

Declines in native populations may be ephemeral if indigenous species are genetically variable in their susceptibility to NIS and can evolve in response to invasion. Alternatively, lack of the ability to evolve in the face of strong selection from invaders can cause extinction (see statement 9). For instance, in 70% of *Rana aurora* Baird and Girard has acquired the ability to recognize the chemical cues emitted by its new predator, the introduced bullfrog *R. catesbeiana*, and behaves accordingly, by reducing their foraging activity and increasing refuge use, whereas frogs from uninvaded ponds do not change their behavior when presented with bullfrogs (Kiesecker and Blaustein 1997). Evolutionary adaptation may also involve habitat and resource use, leading to the phenomenon of character displacement (i.e. increased difference in quantitative characters of two or more species in areas of syntopy compared to areas of allopatry) (Strauss *et al.* 2006). In the Great Lakes, the bloater [*Coregonus hoyi* (Milne-Edwards)] is one of the indigenous fish that survived competition with the introduced alewife for zooplankton. Following the explosion of the alewife populations in the 1960s, the bloater shifted its diet from small zooplankton to larger benthic organisms (Crowder and Binkowski 1983, Crowder and Crawford 1984). In less than 20 years, this shift was accompanied by an adaptive change of its feeding apparatus that showed a decrease by about 15% in the number of gill rakers (Crowder 1984). Similarly, in North American lakes where bluegill sunfish (*L. macrochirus*) have been absent, the pumpkinseed (*L. gibbosus*) exhibits pelagic and littoral ecotypes as adaptations to the diet of zooplankton (for pelagic forms) and of benthic arthropods and mollusks (for littoral forms). In the lakes where bluegills have been introduced, the pumpkinseed populations exhibited exclusively littoral ecotypes, thus restricting its feeding activity to littoral areas in response to the competition with the dominant newcomer (Robinson *et al.* 2002).

CONCLUSIONS

In recent years, a rising awareness of the economic and ecologic costs caused by invasions in fresh waters has encouraged more proactive research and thus has increased our understanding of invasive processes in aquatic systems. Notwithstanding some obvious limitations derived from the relatively small taxonomic coverage of invasion studies and the prevailing focus on certain systems, some general issues regarding freshwater invaders can be raised. First and foremost, predicting the likelihood of the success of a freshwater invader or predicting the invasibility of an aquatic system depends on a detailed understanding of the characteristics of the invader and of the system that is being invaded (Moy and Light 1996b). But both are likely to be idiosyncratic and complex at the

biology. As observed by Simberloff (2006), this is part of the larger problem: “ecology is fundamentally an idiographic science” (p. 917): we will need a catalogue of case studies in order to generate the level of understanding required to deal with many of the environmental problems (Simberloff 2004).

The “tens rule” does not hold for invasion processes in fresh waters. Introduction is a critical step, so the most effective means of minimizing the adverse impact of freshwater invaders is to prevent species transport in the first place. Once introduced, several species have a high potential to establish. Once established, eradication is often impossible and mitigation and control are difficult and expensive, if possible at all (Chapters 34, 36, 37, and 38). The successful establishment of a species is positively related to propagule pressure (Chapter 18). A consequence is that the probability of establishment might be lessened by reducing both the number of individuals accidentally released via commerce-related activities and the frequency of such releases. The importance of propagule pressure also alerts us about the need to construct effective legislative barriers against the introduction of “desirable” species that might turn out to be “Frankensteins” (Moyle *et al.* 1986).

The most likely “monsters” in fresh waters are those species whose physiological requirements closely match with the environmental characteristics of the recipient system or those species able to tolerate a wide range of environmental conditions. With some exceptions, the biotic resistance exerted by the recipient community, including its richness in species and complexity, seems to be less effective in countering the establishment of freshwater invaders, while an *r*-selected strategy is only in some cases a prerequisite for a species to become invasive. Similarly, the naïveté of introduced species and the assumed lack of enemies in the recipient community cannot explain *per se* freshwater invasions. Most freshwater communities are not saturated with species (Cornell and Lawton 1992), but instead are capable of supporting greater numbers of species if the pool of potential colonists and the rate of colonization from the pool are increased (Gido and Brown 1999). Introduced species, in fact, often facilitate each other’s establishment and/or their continued existence, therefore increasing the likelihood and the magnitude of the global ecological impact inflicted by biological invasions.

NIS exert multiform effects on the recipient community, most often acting simultaneously at multiple ecological levels. They may pose threats to individual organisms, species, populations, and genes, and may induce changes to individual populations, communities, and ecosystems. The most dangerous species are parasites but also predators or omnivores that may produce trophic cascades in the recipient community, and ecosystem engineers that may cause physical modifications of the environment and may influence the maintenance and creation of habitats. All these recognized impacts of invaders represent, however, only the “tip” of an ecological and evolutionary iceberg (Palumbi 2001). In inland waters, as in the other biomes, the introduction of species may in-

invasions, and global climate change to create an "evolutionary revolution" (2004). And empirical data and theories are urgently needed to enable prediction, understanding, and management of the acute and chronic effects of species invasions (Strayer *et al.* 2006).

REFERENCES

- Ackefors, H. 1999. The positive effects of established crayfish. Pages 49–61 in F. Gherardi and D. M. Holdich, editors. *Crayfish in Europe as alien species: How to make the most of a bad situation?* A. A. Balkema, Rotterdam, The Netherlands.
- Adams, M. J., C. A. Pearl, and R. B. Bury. 2003. Indirect facilitation of an alien species invasion by non-native fishes. *Ecology Letters* **6**, 343–351.
- Barbaresi, S. and F. Gherardi. 2000. The invasion of the alien crayfish *Procambarus clarkii* in Europe, with particular reference to Italy. *Biological Invasions* **2**, 259–264.
- Bailey, S. A., I. C. Duggan, C. D. A. Van Overdijk, T. H. Johengen, D. F. Reimold, and H. J. MacIsaac. 2004. Salinity tolerance of diapausing eggs of freshwater zooplankton. *Freshwater Biology* **49**, 286–295.
- Bedarf, A. T., K. R. McKaye, E. P. Van Den Berghe, L. J. L. Perez, and D. H. Secor. 2003. Initial six-year expansion of an introduced piscivorous fish in a tropical Central American lake. *Biological Invasions* **3**, 391–404.
- Beisel, J.-N. 2001. The elusive model of a biological invasion process: time to take differences among aquatic and terrestrial ecosystems into account? *Ethology, Ecology & Evolution* **13**, 193–195.
- Bestgen, K. R., S. P. Platania, J. E. Brooks, and D. L. Probst. 1989. Dispersal and life history traits of *Notropis girardi* (Cypriniformes, Cyprinidae) introduced into the Rio Grande, New Mexico. *American Midland Naturalist* **122**, 228–235.
- Boltovskoy, D., N. Correa, D. Cataldo, and F. Sylvester. 2006. Dispersion and ecological impact of the invasive freshwater bivalve *Limnoperna fortunei* in the Río de la Plata watershed and beyond. *Biological Invasions* **8**, 947–963.
- Branstrator, D. K., M. E. Brown, L. J. Shannon, M. Thabes, and K. Heimgartner. 2005. Range expansion of *Bythotrephes longimanus* in North America: evaluating habitat characteristics in the spread of an exotic zooplankton. *Biological Invasions* **8**, 1367–1378.
- Burr, B. M. and L. M. Page. 1986. Zoogeography of fishes in the lower Ohio River and Mississippi Basin. Pages 287–324 in C. H. Hocutt and E. O. Wiley, editors. *Zoogeography of North American freshwater fishes*. Wiley, New York, NY.
- Carlsson, N. O. L. and J. O. Lacoursière. 2005. Herbivory on aquatic vascular plants by the introduced golden apple snail (*Pomacea canaliculata*) in Lao PDR. *Biological Invasions* **7**, 233–241.
- Carroll, S. P. and H. Dingle. 1996. The biology of post-invasion events. *BioScience and Conservation* **78**, 207–214.
- Çelik, K., J. E. Schindler, W. J. Foris, and J. C. Knight. 2002. Predator-mediated coexistence of exotic and native crustaceans in a freshwater lake? *Biological Invasions* **4**, 451–454.
- Cohen, A. N. and J. T. Carlton. 1998. Accelerating invasion rate in a highly invaded estuary. *Science* **279**, 555–558.

- Combes, C. and N. Le Brun. 1990. Invasions by parasites in continental Europe. 285–296 in F. Di Castri, A. J. Hansen and M. Debussche, editors. *Biological Invasions in Europe and the Mediterranean Basin*. Kluwer Academic Publishers, Dordrecht, Netherlands.
- Cornell, H. V. and J. H. Lawton. 1992. Species interactions, local and regional processes and limits to the richness of ecological communities: a theoretical perspective. *Journal of Animal Ecology* **61**, 1–12.
- Costanza, R., R. d'Arge, R. de Groot, S. Farber, M. Grasso, B. Hannon, K. Limburg, S. Naeem, R. V. O'Neill, J. Paruelo, R. G. Raskin, P. Sutton, and M. van den Belt. 1997. The value of the world's ecosystem services and natural capital. *Nature* **387**, 253–260.
- Cox, G. W. 2004. *Alien species and evolution*. Island Press, Washington, DC.
- Craig, J. F. 1992. Human induced changes in the composition of fish communities in the African Great Lakes. *Reviews in Fish Biology and Fisheries* **2**, 93–124.
- Crowder, L. B. 1984. Character displacement and habitat shift in a native cisco in southeastern Lake Michigan: evidence for competition? *Copeia* **1984**, 878–888.
- Crowder, L. B. and F. P. Binkowski. 1983. Foraging behaviors and the interactions of alewife, *Alosa pseudoharengus*, and bloater, *Coregonus hoyi*. *Environmental Biology of Fishes* **8**, 105–113.
- Crowder, L. B. and H. L. Crawford. 1984. Ecological shifts in resource use by bloater in Lake Michigan. *Transactions of the American Fisheries Society* **113**, 694–700.
- Darrigran, G. 2002. Potential impact of filter-feeding invaders on temperate freshwater environments. *Biological Invasions* **4**, 145–156.
- Devin, S. and J.-N. Beisel. 2007. Biological and ecological characteristics of introduced species: a gammarid study. *Biological Invasions* **9**, 13–24.
- Diamond, J. and T. J. Case. 1986. Overview: introductions, extinctions, exterminations and invasions. Pages 65–79 in J. Diamond and T. J. Case, editors. *Community Ecology*. Harper & Row, New York, NY.
- Didham, R. K., J. M. Tylianakis, M. A. Hutchison, R. M. Ewers, and N. J. Gemmell. 2005. Are invasive species the drivers of ecological change? *Trends in Ecology and Evolution* **20**, 470–474.
- Dingle, H. 1980. The ecology and evolution of migration. Pages 1–101 in S. A. Stearns Jr., editor. *Animal migration, orientation, and navigation*. Academic Press, New York, NY.
- Eaton, J. G. and R. M. Scheller. 1996. Effects of climatic warming on fish thermal habitat in streams of the United States. *Limnology and Oceanography* **41**, 1110–1115.
- Echelle, A. A. and P. J. Connor. 1989. Rapid, geographically extensive genetic divergence after secondary contact between two pupfish species (Cyprinodon, Cyprinodontidae). *Evolution* **43**, 717–727.
- Elton, C. S. 1958. *The Ecology of invasions by animals and plants*. The University of Chicago Press, Chicago, IL, Edition 2000.
- French, J. R. P. I. 1993. How well can fishes prey on zebra mussels in eastern North America? *Fisheries* **18**, 13–19.
- Gherardi, F. 2006. Crayfish invading Europe: the case study of *Procambarus clarkii*. *Marine and Freshwater Behaviour and Physiology* **39**, 175–191.

- Swamp sparrowgrass, *Panicum capillare*, in an irrigation ditch in Piacenza, Italy. *Wetland Issues* **12**, 99–108.
- Gido, K. B. and J. H. Brown. 1999. Invasion of North American drainage by alien species. *Freshwater Biology* **42**, 387–399.
- Goldschmidt, T., F. Witte, and J. Wanink. 1993. Cascading effects of the introduced perch on the detritivorous/phytoplanktivorous species in the sublittoral areas of Victoria. *Conservation Biology* **7**, 686–700.
- Gurevitch, J. and D. K. Padilla. 2004. Are invasive species a major cause of extinction? *Trends in Ecology and Evolution* **19**, 470–474.
- Hansen, M. H. 2002. Estimating the long-term effects of stocking domesticated into wild brown trout (*Salmo trutta*) populations: an approach using microsatellite DNA analysis of historical and contemporary samples. *Molecular Ecology* **11**, 1003–1015.
- Hebert, P. D. N. and M. Cristescu. 2002. Genetic perspective on invasions: the case of Cladocera. *Canadian Journal of Fisheries and Aquatic Sciences* **59**, 1229–1234.
- Jeschke, J. M. and D. L. Strayer. 2005. Invasion success of vertebrates in Europe and North America. *Proceedings of the National Academy of Sciences USA* **102**, 7202.
- Jones, C. G., J. H. Lawton, and M. Shachak. 1994. Organisms as ecosystem engineers. *Oikos* **69**, 373–386.
- Karatayev, A. Y., L. E. Burlakova, and D. K. Padilla. 2002. Impacts of zebra mussels on aquatic communities and their role as ecosystem engineers. Pages 433–444 in E. Leppäkoski, S. Gollasch and S. Olenin, editors. *Invasive aquatic species of Europe: distribution, impacts and management*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Kennedy, T. A., S. Naeem, K. M. Howe, J. M. H. Knops, D. Tilman, and P. Relch. 2000. Biodiversity as a barrier to ecological invasion. *Nature* **417**, 636–638.
- Ketelaars, H. A. M., F. E. Lambregts-van de Clundert, C. J. Carpentier, A. J. Wager, and W. Hoogenboezem. 1999. Ecological effects of the mass occurrence of the Caspian invader, *Hemimysis anomala* G.O. Sars, 1907 (Crustacea: Mysidacea) in a freshwater storage reservoir in the Netherlands, with notes on autoecology and records. *Hydrobiologia* **394**, 233–248.
- Kiesecker, J. M. and A. R. Blaustein. 1997. Population differences in responses of long-legged frogs (*Rana aurora*) to introduced bullfrogs. *Ecology* **78**, 1752–1760.
- Kirjavainen, J. and M. Sipponen. 2004. Environmental benefit of different control management strategies in Finland. *Fisheries Management and Ecology* **11**, 213–220.
- Kolar, C. S. and D. M. Lodge. 2001. Progress in invasion biology: predicting invasions. *Trends in Ecology and Evolution* **16**, 199–204.
- Laha, M. and H. T. Mattingly. 2006. Identifying environmental conditions to promote species coexistence: an example with the native Barrens topminnow and introduced western mosquitofish. *Biological Invasions* **8**, 719–725.
- Lambrinos, J. G. 2004. How interactions between ecology and evolution influence contemporary invasion dynamics? *Ecology* **85**, 2061–2070.
- Lehtonen, H. 2002. Alien freshwater fishes of Europe. Pages 153–161 in E. Leppäkoski, S. Gollasch and S. Olenin, editors. *Invasive aquatic species of Europe: distribution, impact and management*. Kluwer Academic Publishers, Dordrecht, The Netherlands.

- Pages 1–11 in L. Espinosa, S. G. Shuman, and S. C. Stepien, editors. *Invasive Aquatic Species of Europe: Distribution, impact and management*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Levine, J. M. and C. M. D'Antonio. 1999. Elton revisited: a review of evidence on diversity and invisibility. *Oikos* **87**, 15–26.
- Lodge, D. M. 1993. Biological invasions: lessons for ecology. *Trends in Ecology and Evolution* **8**, 133–137.
- Lodge, D. M. 2001. Lakes. Pages 277–313 in F. S. Chapin III, O. E. Sala and E. S. Sannwald, editors. *Global biodiversity in a changing environment: scenarios for the 21st century*. Springer, New York, NY.
- Lodge, D. M. and K. Shrader-Frechette. 2003. Nonindigenous species: ecological invasions, environmental ethics, and public policy. *Conservation Biology* **17**, 31–40.
- Lonsdale, W. M. 1999. Global patterns of plant invasions and the concept of invasion. *Ecology* **80**, 1522–1536.
- MacDougall, A. S. and R. Turkington. 2005. Are invasive species the drivers or passengers of change in degraded ecosystems? *Ecology* **86**, 42–55.
- MacIsaac, H. J. 1996. Potential abiotic and biotic impacts of zebra mussels on the waters of North America. *American Zoologist* **36**, 287–299.
- MacIsaac, H. J. 1999. Biological invasions in Lake Erie: past, present, and future. State of Lake Erie (SOLE) – past, present, and future. Pages 305–322 in M. Munawar and T. Edsall, editors. *Ecovision World Monograph Series*, Backhuys, Leiden, The Netherlands.
- MacIsaac, H. J., I. A. Grigorovich, and A. Ricciardi. 2001. Reassessment of biological invasion concepts: the Great Lakes basin as a model. *Biological Invasions* **3**, 403–415.
- Mack, R. N., D. Simberloff, W. M. Lonsdale, H. Evans, M. Clout, and F. A. Bazzaz. 2000. Biotic invasions: causes, epidemiology, global consequences, and control. *Ecological Applications* **10**, 689–710.
- Millennium Ecosystem Assessment. 2005. *Ecosystem and human well-being: biodiversity synthesis*. World Resources Institute, Washington, DC.
- Mills, E. L., R. M. Dermott, E. F. Roseman, D. Dustin, E. Mellina, D. B. Conner, and A. P. Spidle. 1993. Colonization, ecology, and population structure of the “quagga” mussel (*Bivalvia: Dreissenidae*) in the lower Great Lakes. *Canadian Journal of Fisheries and Aquatic Sciences* **50**, 2305–2314.
- Mills, E. L., D. L. Strayer, M. D. Scheuerell, and J. T. Carlton. 1996. Exotic species invasions in the Hudson River basin – a history of invasions and introductions. *Estuaries* **19**, 814–824.
- Moyle, P. B. 1986. Fish introductions into North America: patterns and ecological impact. Pages 27–43 in H. A. Mooney and J. A. Drake, editors. *Ecology of biological invasions of North America and Hawaii*. Springer, New York, NY.
- Moyle, P. B., H. W. Li, and B. A. Barton. 1986. The Frankenstein Effect: impacts of introduced fishes on native fishes in North America. Pages 415–426 in R. H. Squires, editor. *Fish culture in fisheries management*. American Fisheries Society, Bethesda, MD.
- Moyle, P. B. and T. Light. 1996a. Fish invasions in California: do abiotic factors limit success? *Ecology* **76**, 1666–1670.
- Moyle, P. B. and T. Light. 1996b. Biological invasions of fresh water: empirical rules and assembly theory. *Biological Conservation* **78**, 149–161.

- Evolution **19**, 18–24.
- Oliveira, M. D. de, A. M. Takeda, L. F. de Barros, D. S. Barbosa, and E. K. de Rêgo. 2006. Invasion by *Limnoperna fortunei* (Dunker, 1857) (Bivalvia, Mytilidae) in a Pantanal wetland, Brazil. *Biological Invasions* **8**, 97–104.
- Palumbi, S. 2001. Humans as the world's greatest evolutionary force. *Science* **291**, 1786–1790.
- Parker, I. M., D. Simberloff, W. M. Lonsdale, K. Goodell, M. Wonham, P. M. Kareiva, M. H. Williamson, B. Von Holle, P. B. Moyle, J. E. Byers, and L. Goldwasser. 2000. Impact: toward a framework for understanding the ecological effects of invasions. *Biological Invasions* **1**, 3–19.
- Perry, W. L., D. M. Lodge, and J. L. Feder. 2002. Importance of hybridization between indigenous and nonindigenous freshwater species: an overlooked threat to American biodiversity. *Systematic Biology* **51**, 255–275.
- Power, M. E. 1990. Effects of fish in river food webs. *Science* **250**, 811–814.
- Pusey, B., D. Burrows, A. Arthington, and M. Kennard. 2006. Translocation and control of piscivorous fishes in the Burdekin River, north-eastern Australia. *Biological Invasions* **8**, 965–977.
- Rahel, F. J. 2000. Homogenization of fish faunas across the United States. *Science* **288**, 854–856.
- Ricciardi, A. 2001. Facilitative interactions among aquatic invaders: is an “invasion meltdown” occurring in the Great Lakes? *Canadian Journal of Fisheries and Aquatic Sciences* **58**, 2513–2525.
- Ricciardi, A. and S. K. Atkinson. 2004. Distinctiveness magnifies the impact of biological invaders in aquatic ecosystems. *Ecology Letters* **7**, 781–784.
- Ricciardi, A., R. J. Neves, and J. B. Rasmussen. 1998. Impending extinctions of American freshwater mussels (Unionida) following the zebra mussel (*Dreissena polymorpha*) invasion. *Journal of Animal Ecology* **67**, 613–619.
- Ricciardi, A. and J. B. Rasmussen. 1998. Predicting the identity and impact of biological invaders: a priority for aquatic resource managers. *Canadian Journal of Fisheries and Aquatic Sciences* **55**, 1759–1765.
- Ricciardi, A. and J. B. Rasmussen. 1999. Extinction rates of North American freshwater fauna. *Conservation Biology* **13**, 1220–1222.
- Robinson, B. W., D. S. Wilson, and A. S. Margosian. 2000. A pluralistic analysis of character release in pumpkinseed sunfish (*Lepomis gibbosus*). *Ecology* **81**, 2799–2809.
- Rodríguez, C. F., E. Bécáres, M. Fernández-Aláez, and C. Fernández-Aláez. 2005. The loss of biodiversity and degradation of wetlands as a result of introducing exotic species. *Biological Invasions* **7**, 75–85.
- Rosen, P. C. and C. R. Schwalbe. 2002. Widespread effects of introduced species on reptiles and amphibians in the Sonoran Desert region. Pages 220–240 in B. T. Baskin, editor. *Invasive exotic species in the Sonoran region*. University of Arizona Press and the Arizona-Sonora Desert Museum, Tucson, AR.
- Ross, R. M., W. A. Lellis, R. M. Bennett, and C. S. Johnson. 2001. Landscape characteristics as determinants of nonindigenous fish invasions. *Biological Invasions* **3**, 347–361.
- Ruesink, J. L. 2005. Global analysis of factors affecting the outcome of freshwater species introductions. *Conservation Biology* **19**, 1883–1893.

- biases. Annual Review of Ecology and Systematics **31**, 481–531.
- Sala, O. E., F. S. Chapin III, J. J. Armesto, R. Berlow, J. Bloomfield, R. E. Huber-Sanwald, L. F. Huenneke, R. B. Jackson, A. Kinzig, R. Leemans, D. H. A. Mooney, M. Oesterheld, N. L. Poff, M. T. Sykes, B. H. Walker, M. Walker, D. H. Wall. 2000. Global biodiversity scenarios for the year 2100. Science **290**, 1770–1774.
- Schlaepfer, M. A., P. W. Sherman, B. Blossey, and M. C. Runge. 2005. Introduced species as evolutionary traps. Ecology Letters **8**, 241–246.
- Shea, K. and P. Chesson. 2002. Community ecology theory as a framework for biological invasions. Trends in Ecology and Evolution **17**, 170–176.
- Shurin, J. B. 2000. Dispersal limitation, invasion resistance, and the structure of zooplankton communities. Ecology **81**, 3074–3086.
- Shurin, J. B. and J. E. Havel. 2002. Hydrologic connections and overland dispersal of an exotic freshwater crustacean. Biological Invasions **4**, 431–439.
- Simberloff, D. 2004. Community ecology: is it time to move on? American Naturalist **163**, 787–799.
- Simberloff, D. 2006. Invasion meltdown 6 years later: important phenomenon, unfortunate metaphor, or both? Ecology Letters **9**, 912–919.
- Simberloff, D. and B. Von Holle. 1999. Positive interactions of nonindigenous species: invasion meltdown? Biological Invasions **1**, 21–32.
- Simon, K. S. and C. R. Townsend. 2003. Impacts of freshwater invaders at different levels of ecological organization, with emphasis on salmonids and ecosystem consequences. Freshwater Biology **48**, 982–994.
- Smith, S. A., G. Bell, and E. Bermingham. 2004. Cross-Cordillera exchange mediated by the Panama Canal increased the species richness of local freshwater fish assemblages. Proceedings of the Royal Society of London B **271**, 1889–1896.
- Smith, S. A. and J. B. Shurin. 2006. Room for one more? Evidence for invasibility saturation in ecological communities. Pages 423–447 in M. W. Cadotte, J. B. Shurin, J. M. McCarthy, and J. H. Lawler, editors. Conceptual ecology invasion biology: recent approaches to nature. Springer, Dordrecht, The Netherlands.
- Stepien, C. A., C. D. Taylor, and K. A. Dabrowska. 2002. Genetic variability and phylogeographical patterns of a nonindigenous species invasion: a comparison of exotic vs. native zebra and quagga mussel populations. Journal of Evolutionary Biology **15**, 3143–328.
- Stewart, T. W. and J. M. Haynes. 1994. Benthic macroinvertebrate community changes in southwestern Lake Ontario following invasion of *Dreissena*. Journal of Great Lakes Research **20**, 479–493.
- Strauss, S. Y., J. A. Lau, and S. P. Carroll. 2006. Evolutionary responses of native species to introduced species: what do introductions tell us about natural communities? Ecology Letters **9**, 357–374.
- Strayer, D. L., E. A. Blair, N. F. Caraco, J. J. Cole, S. Findlay, W. C. Nieder, and M. I. Newbery. 2005. Interactions between alien species and restoration of large-river ecosystems. Archiv für Hydrobiologie, Supplement **155**/1–4, 133–145.
- Strayer, D. L., V. T. Eviner, J. M. Jeschke, and M. L. Pace. 2006. Understanding the long-term effects of species invasions. Trends in Ecology and Evolution **21**, 645–653.

- species-rich ecosystems. *Ecology* **75**, 717–731.
- Torchin, M. E., K. D. Lafferty, A. P. Dobson, V. J. McKenzie, and A. M. Kuris. 2003. Introduced species and their missing parasites. *Nature* **421**, 628–630.
- Townsend, C. R. 1996. Invasion biology and ecological impacts of brown trout *trutta* in New Zealand. *Biological Conservation* **78**, 13–22.
- Trendall, J. 1988. The distribution and dispersal of introduced fish at Thumbi West in Lake Malawi, Africa. *Journal of Fish Biology* **33**, 357–369.
- Vitousek, P. M., C. M. D'Antonio, L. D. Loope, M. Rejmánek, and R. Westbrooks. 2002. Introduced species: A significant component of human-caused global change. *Zealand Journal of Ecology* **21**, 1–16.
- Vitousek, P. M., C. M. D'Antonio, L. D. Loope, and R. Westbrooks. 1996. Bio invasions as global environmental change. *American Scientist* **84**, 468–478.
- Welcomme, R. I. 1988. International introductions of inland aquatic species. FAO technical paper 294. Food and Agriculture Organization, Rome, Italy.
- Wilcove, D. S. and M. J. Bean. 1994. The big kill: declining biodiversity in America's forests and rivers. Environmental Defense Fund, Washington, DC.
- Williamson, M. 1996. Biological invasions. Chapman & Hall, London, UK.
- Yusa, Y., N. Sugiura, and T. Wada. 2006. Predatory potential of freshwater a on an invasive agricultural pest, the apple snail *Pomacea canaliculata* (Gastropoda: Ampullariidae), in southern Japan. *Biological Invasions* **8**, 137–147.