



FLORE Repository istituzionale dell'Università degli Studi di Firenze

Acylphosphatase interferes with SERCA2a-PLN association.

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Acylphosphatase interferes with SERCA2a-PLN association / C. NEDIANI; A. CELLI; C. FIORILLO; V. PONZIANI; L. GIANNINI; P. NASSI. - In: BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS. - ISSN 0006-291X. - STAMPA. - 301:(2003), pp. 948-951. [10.1016/S0006-291X(03)00078-0]

Availability:

The webpage https://hdl.handle.net/2158/311347 of the repository was last updated on 2020-06-30T16:30:43Z

Published version: DOI: 10.1016/S0006-291X(03)00078-0

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The abovementioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)



Available online at www.sciencedirect.com



Biochemical and Biophysical Research Communications 301 (2003) 948-951

www.elsevier.com/locate/ybbrc

Acylphosphatase interferes with SERCA2a–PLN association

Chiara Nediani, Alessandra Celli, Claudia Fiorillo, Vanessa Ponziani, Lara Giannini, and Paolo Nassi*

Dipartimento di Scienze Biochimiche, Università di Firenze, viale Morgagni 50, 50134 Firenze, Italy

Received 2 January 2003

Abstract

We previously reported that acylphosphatase, a cytosolic enzyme present in skeletal and heart muscle, actively hydrolyzes the phosphoenzyme (EP) of cardiac sarcoplasmic reticulum (SR) Ca²⁺-ATPase (SERCA2a), inducing an increased activity of this pump. We hypothesized that acylphosphatase-induced stimulation of SERCA2a, in addition to enhanced EP hydrolysis, may be due to a displacement of phospholamban (PLN), removing its inhibitory effect. To verify this hypothesis co-immunoprecipitation experiments were performed by adding recombinant muscle acylphosphatase to solubilized heart SR vesicles, used as a source of SERCA2a and PLN. With anti-acylphosphatase antibodies only SERCA2a was co-immunoprecipitated in an amount which increased in parallel to the concentrations of our enzyme. Conversely, using anti-SERCA2a antibody, both PLN and acylphosphatase were co-immunoprecipitated with SERCA2a, and the PLN amount in the precipitate decreased with increasing acylphosphatase concentrations. SERCA2a and PLN were co-immunoprecipitated by anti-phospholamban antibodies, but while the amount of precipitated phospholamban increased in the presence of acylphosphatase, the level of SERCA2a decreased. These preliminary results strengthen the supposed displacement of phospholamban by acylphosphatase. © 2003 Elsevier Science (USA). All rights reserved.

Keywords: SERCA2a; Phospholamban; Acylphosphatase; Cardiomyopathy

Sarco(endo)plasmic reticulum Ca²⁺-ATPases (SER-CAs) are 110-kDa transmembrane proteins that transport Ca²⁺ ions from the sarcoplasm to the lumen of the membrane system [1]. The energy for Ca^{2+} transport is provided by ATP; intermediate steps involve the formation and hydrolysis of a phosphoenzyme intermediate (EP) [2]. The cardiac sarcoplasmic reticulum (SR) Ca^{2+} -ATPase (SERCA2a) is regulated by its interactions with the intrinsic membrane protein phospholamban. Phospholamban (PLN) inhibits the activity of SERCA2a, but its phosphorylation by protein kinase A or Ca^{2+} -calmodulin kinase reverses the inhibitory interaction, presumably through dissociation of phosphorylated phospholamban from SERCA2a [3]. An additional regulatory effect on SERCA2a could be exerted by acylphosphatase (EC 3.6.1.7), a cytosolic 11 kDa enzyme well represented in cardiac muscle that catalyzes the hydrolysis of acylphosphates [4–6]. In previous studies we have

shown that acylphosphatase actively hydrolyzes the EP intermediate of this transport system, an effect that results in an enhanced activity of the SR Ca^{2+} pump [7,8]. We have also demonstrated that acylphosphatase, in addition to its hydrolytic activity on EP, stimulated SER-CA2a activity through another mechanism, probably due to its conformational properties. In fact, using an acylphosphatase mutant, Asn41Ser, devoid of catalytic activity but unchanged as for the three-dimensional structure [9], the stimulatory effect on SERCA2a was retained, but disappeared when the conformational structure of the mutant was altered by thermal denaturation. To explain these results we supposed that acylphosphatase could interact with SERCA2a taking the place of unphosphorylated PLN whose inhibitory effect could be removed [8]. In fact, NMR studies showed that acylphosphatase contains a structural motif where 12 residues (from 55 to 66) form an amphipathic α -helix with a prevalence of basic groups [10] that resembles that of the PLN cytoplasmic 1A domain, essential for the association with SERCA2a [11].

^{*} Corresponding author. Fax: +39-055-422-725.

E-mail address: nassi@scibio.unifi.it (P. Nassi).

In the present study, as a first approach to verify the supposed interaction, we performed a series of coimmunoprecipitation experiments. Mixtures of solubilized cardiac SR vesicles, used as a source of SERCA2a and PLN, and recombinant muscle acylphosphatase were incubated with antibodies against each of these proteins and the composition of the obtained immunoprecipitates was determined by Western blot analysis.

Materials and methods

Materials. All reagents were of analytical grade and were obtained from Sigma. Monoclonal (mouse) anti-SERCA2a ATPase antibody (clone 2A7-A1) was obtained from Affinity Bioreagents (Golden, CO), monoclonal (mouse) anti-phospholamban antibody (clone A1) was from Upstate biotechnology (Lake Placid, NY), and acylphosphatase recombinant form was obtained according to Modesti et al. [12]. Anti-acylphosphatase antibodies were raised in rabbit using the recombinant protein and purified by affinity chromatography [13]. Peroxidase coniugated anti-mouse or anti-rabbit secondary antibody was from Amersham. Super Signal enhanced chemiluminescent kit was from Pierce.

Preparation of SR vesicles (SRVs). Cardiac SRVs were isolated from rabbit heart according to Nediani et al. [14] and measured for protein [15].

Co-immunoprecipitation. SRVs were solubilized with 1% SDS in the immunoprecipitation buffer (10 mM Tris–HCl, pH 7, 150 mM NaCl, 1 mM EGTA, and 1 mM EDTA) at room temperature for 1 h and centrifuged for 15' at 16,000g. The supernatants were rotated with 1% bovine serum albumin-treated G-Sepharose for 30' and centrifuged to remove proteins bound non-specifically to G-Sepharose. The supernatants, at a final concentration of 0.5 mg/ml protein, were then incubated with monoclonal anti-SERCA2a or polyclonal anti-acylphosphatase or monoclonal anti-PLN antibody in the absence and in the presence of 5 or 10 µg of acylphosphatase recombinant form on ice for 1 h. The samples were rotated for 1 h with G-Sepharose and centrifuged and the pellets were washed three times in the immunoprecipitation buffer and solubilized for 10' at 90 °C in Laemmli's sample buffer [16].

Western blot analysis of SERCA2a, phospholamban, and acylphosphatase. Samples were resolved by 10% and 15% SDS–PAGE and transferred to PVDF membranes. After blocking, the membrane were incubated with monoclonal anti-SERCA2a, or polyclonal anti-human muscle acylphosphatase or monoclonal anti-PLB antibody, washed three times in T-TBS solution (50 mM Tris–HCl, pH 7.4, 150 mM NaCl, and 0.1% Tween 20), and treated with horseradish peroxidaseconjugated anti-mouse or anti-rabbit secondary antibody. The signals were detected with a chemiluminescence kit and the signal densities corresponding to SERCA2a or PLN immunoprecipitated were quantified by laser densitometry using Quantity One software (Bio-Rad laboratories).

Results and discussion

Taken together, the results of this study agree with the possibility that acylphosphatase, owing to its conformational properties, displaces PLN from SERCA2a. In this connection, we have previously demonstrated a physical interaction between SERCA2a and acylphosphatase since in controlled immunoprecipitation experiments the two proteins were co-precipitated by a monoclonal anti-SERCA2a antibody. In the same study, however, we found that thermally denaturated acylphosphatase was not more precipitated, suggesting that the maintenance of the native conformation of this enzyme was required for its binding to SERCA2a [8]. Now, in addition to confirming a physical association between SERCA2a and acylphosphatase, the present results also indicate that this binding could interfere with SERCA2a-PLN interaction. This, at least, emerges from the experiments that we performed by incubating SERCA2a and PLN (in the form of solubilized cardiac SR vesicles) with monoclonal anti-SERCA2a antibody in the absence and in the presence of increasing amounts of purified recombinant acylphosphatase. As shown in the Fig. 1A, when acylphosphatase was lacking Western blot analysis of the immunoprecipitate revealed two clear bands at 110 kDa and 25-27 kDa, corresponding, respectively, to SERCA2a and to the PLN pentameric form: a result which was expected since the well-known interaction between these proteins. On the other hand, in the presence of acylphosphatase, the analysis of the immunoprecipitate showed another band at 11kDa, suggesting the co-immunoprecipitation of this enzyme together with the other two proteins. Moreover, as



Fig. 1. Effect of acylphosphatase (ACP) on co-immunoprecipitation of SERCA2a with PLN using monoclonal anti-SERCA2a antibody. Solubilized SRVs were incubated with monoclonal anti-SERCA2a antibody as described under "Materials and methods" in the absence (0) and in presence of 5 or 10 µg of ACP. Immunoprecipitates were analyzed by Western blotting (A) and PLN signals were quantified by densitometric analysis (B). Each bar represents the mean value \pm SD of four different blots. *p < 0.05 vs 0 by Student's *t* test.

indicated by densitometric analysis (Fig. 1B), the amount of PLN found in the immunoprecipitated significantly decreased with increasing acylphosphatase concentrations, with a diminution that, compared to the control values, was 17% for the lower (5 µg) and 37% for the higher acylphosphatase levels $(10 \,\mu g)$. Besides these findings that clearly strengthen the supposed displacement of phospholamban by acylphosphatase, also other results of the present study support such an interpretation. In fact, when the above described mixtures of solubilized heart SR vesicles plus acylphosphatase were added with anti-acylphosphatase antibody (Fig. 2A) only this enzyme and SERCA2a were co-immunoprecipitated and the amount of SERCA2a detected in the precipitate increased significantly with the increase in acylphosphatase concentration. While, on one hand, these data corroborate the view of a selective interaction between acylphosphatase and SERCA2a, on the other hand the constant absence of PLN in the immunoprecipitate obtained with anti-acylphosphatase antibody strongly suggests that acylphosphatase, through a sort of competitive binding with SERCA2, can effectively displace PLN from its association with heart SR Ca²⁺







Fig. 2. Co-immunoprecipitation of SERCA2a and acylphosphatase (ACP) with polyclonal anti-ACP antibody. Solubilized SRVs were incubated with polyclonal anti-ACP antibody as described under "Materials and methods" in the presence of 5 and 10µg of ACP. Immunoprecipitates were analyzed by Western blotting (A) and SERCA2a signals were quantified by densitometric analysis (B). Each bar represents the mean value \pm SD of four different blots. *p < 0.05 vs 5µg by Student's *t* test.

Fig. 3. Effect of acylphosphatase (ACP) on co-immunoprecipitation of SERCA2a with PLN using monoclonal anti-PLN antibody. Solubilized SRVs were incubated with monoclonal anti-SERCA2a antibody as described under "Materials and methods" in the absence (0) and in the presence of 10 µg of ACP. Immunoprecipitates were analyzed by Western blotting (A) and SERCA2a and PLN signals were quantified by densitometric analysis (B). Each bar represents the mean value \pm SD of four different blots. \Box , (0); \blacksquare , 10 µg of ACP. *p < 0.05 vs 0 by Student's *t* test.

attenuation of cardiac hypertrophy and enhancement of cardiac contractility in a mouse dilated cardiomyopathy model [20] and in a mice pressure-overload model [21], suggesting that the removal of PLN inhibition on SERCA2a may be a new strategy to treat and to prevent heart failure. In this context it is our opinion that, although preliminary, the present data may be of special interest because they suggest a novel physiological mechanism which could contribute to modulate SER-CA-PLN interactions, all the more so since in a recent study we found that, in a model of volume overload, increased acylphosphatase expression was associated to a recovery of SR function and an improved heart contractility [22].

On the basis of these considerations we think that it would be of interest to probe more deeply the molecular basis of the interaction between SERCA2a and acylphosphatase. More in particular, further studies will be aimed, in our intention, at identifying the residues that are involved in this interaction and to determine, with site-specific mutagenesis, the mutations that could cause a loss of the native conformation of acylphosphatase and a reduction of its hypothesized displacement effect.

Acknowledgments

The financial support of Telethon-Italy (Grant No. 1132) is gratefully acknowledged. We are grateful to Dr. ssa Letizia Niccolai for the assistance.

References

- D.H. MacLennan, W.J. Rice, N.M. Green, The mechanism of Ca²⁺ transport by sarco(endo)plasmic reticulum Ca²⁺-ATPases, J. Biol. Chem. 272 (1997) 28815–28818.
- [2] M. Tada, M. Kadoma, M. Inui, J. Fujii, Regulation of Ca²⁺ pump from cardiac sarcoplasmic reticulum, in: S. Fleischer, B. Fleisher (Eds.), Methods in Enzymology, 157, Academic Press, London, 1988, pp. 107–154.
- [3] K. Simmerman, L.R. Jones, Phospholamban: protein structure, mechanism of action, and role in cardiac function, Physiol. Rev. 78 (1998) 921–947.
- [4] A. Berti, M. Stefani, G. Liguri, G. Camici, G. Manao, G. Ramponi, Acylphosphatase action on dicarboxylic acylphosphates, Ital. J. Biochem. 26 (1977) 377–378.
- [5] G. Ramponi, F. Melani, A. Guerritore, Azione dell' acilfosfatasi di muscolo sul carbamilfosfato, J. Biochim. 10 (1961) 189–196.
- [6] G. Ramponi, C. Treves, A. Guerritore, Hydrolytic activity of muscle acylphosphatase on 3-phosphoglyceroyl-phosphate, Experientia 23 (1967) 1019–1020.
- [7] C. Nediani, C. Fiorillo, E. Marchetti, A. Pacini, G. Liguri, P. Nassi, Stimulation of cardiac sarcoplasmic reticulum calcium pump by acylphosphatase. Relationship to phospholamban phosphorylation, J. Biol. Chem. 271 (1996) 19066–19073.
- [8] C. Nediani, C. Fiorillo, S. Rigacci, F. Magherini, M. Francalanci, G. Liguri, A. Pacini, P. Nassi, A novel interaction mechanism accounting for different acylphosphatase effects on cardiac and fast twitch skeletal muscle sarcoplasmic reticulum calcium pumps, FEBS Lett. 443 (1999) 308–312.

- [9] N. Taddei, M. Stefani, F. Magherini, F. Chiti, A. Modesti, G. Raugei, G. Ramponi, Looking for residues involved in the muscle acylphosphatase catalytic mechanism and structural stabilization: role of Asn41, Thr42, and Thr46, Biochemistry 35 (1996) 7077–7083.
- [10] A. Pastore, V. Saudek, G. Ramponi, R.J.P. Williams, Threedimensional structure of acylphosphatase. Refinement and structure analysis, J. Mol. Biol. 224 (1992) 427–440.
- [11] T. Toyofuku, K. Kurzydlowsky, M. Tada, D.H. MacLennan, Amino acids Glu2 to Ile18 in the cytoplasmic domain of phospholamban are essential for functional association with the Ca²⁺-ATPase of sarcoplasmic reticulum, J. Biol. Chem. 269 (1994) 3088–3094.
- [12] A. Modesti, N. Taddei, F. Chiti, M. Bucciantini, F. Magherini, S. Rigacci, M. Stefani, G. Raugei, G. Ramponi, Expression, purification, and characterization of acylphosphatase muscular isoenzyme as fusion protein with glutathione-S-transferase, J. Prot. Chem. 15 (1996) 27–34.
- [13] A. Berti, G. Liguri, M. Stefani, P. Nassi, G. Ramponi, Purification of horse muscle acylphosphatase antibodies by affinity chromatography, Physiol. Chem. Phys. 14 (1982) 307–311.
- [14] C. Nediani, L. Formigli, A.M. Perna, L. Ibba-Manneschi, S. Zecchi-Orlandini, C. Fiorillo, V. Ponziani, C. Cecchi, P. Liguori, G. Fratini, P. Nass, Early changes induced in the left ventricle by pressure overload. An experimental study on swine heart, J. Mol. Cell Cardiol. 32 (2000) 131–142.
- [15] G. Beisenherz, H.J. Boltze, T.H. Bucher, R. Czok, K.H. Garbade, E. Meyer-Arendt, G. Pfleiderer, Diphosphofructosealdolase, phosphoglyceraldehyd-dehydrogenase, milchsauredehydrogenase, and pyruvat-kinase aus kaninchen-muskulature in einern arbeitsgang, Z. Naturforsch. 8b (1953) 555.
- [16] U.K. Laemmly, Cleavage of structural proteins during the assembly of the head of bacteriophage T4, Nature 227 (1970) 680–685.
- [17] Y. Kimura, M. Inui, Recostitution of the cytoplasmic interaction between phospholamban and Ca²⁺-ATPase of cardiac sarcoplasmic reticulum, Mol. Pharmacol 61 (2002) 667–673.
- [18] J. Zhai, A.G. Schmidt, B.D. Hoit, Y. Kimura, D.H. MacLennan, E.G. Kranias, Cardiac-specific overexpression of a superinhibitory pentameric phospholamban mutant enhances inhibition of cardiac function in vivo, J. Biol. Chem. 275 (2000) 10538–10544.
- [19] E. Zvaritch, P.H. Backx, F. Jirik, Y. Kimura, S. de Leon, A.G. Schmidt, B.D. Hoit, J.W. Lester, E.G. Kranias, D.H. MacLennan, The transgenic expression of highly inhibitory monomeric forms of phospholamban in mouse heart impairs cardiac contractility, J. Biol. Chem. 275 (2000) 14985–14991.
- [20] S. Minamisawa, M. Hoshijima, G. Chu, C.A. Ward, K. Frank, Y. Gu, M.E. Martone, Y. Wang, J. Ross Jr., E.G. Kranias, W.R. Giles, K.R. Chien, Chronic phospholamban-sarcoplasmic reticulum calcium-ATPase interaction is the critical calcium cycling defect in dilated cardiomyopathy, Cell 99 (1999) 313– 322.
- [21] H. Nakayama, K. Otsu, O. Yamaguchi, K. Nishida, M. Date, K. Hongo, Y. Kusakari, T. Toyofuku, S. Hikoso, K. Kashiwase, T. Takeda, Y. Matsumura, S. Kurihara, M. Hori, M. Tada, Cardiacspecific overexpression of high Ca²⁺ affinity mutant of SERCA2a attenuates in vivo pressure overload cardiac hypertrophy, FASEB J., (2002) published online Nov 1.
- [22] C. Nediani, L. Formigli, A.M. Perna, A. Pacini, V. Ponziani, P.A. Modesti, L. Ibba-Manneschi, S. Zecchi-Orlandini, C. Fiorillo, C. Cecchi, P. Liguori, G. Fratini, S. Vanni, P. Nassi, Biochemical changes and their relationship with morphological and functional findings in pig heart subjected to lasting volume overload: a possibile role of acylphosphatase in the regulation of sarcoplasmic reticulum calcium pump, Basic Res. Cardiol. 97 (2002) 469– 478.