



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Microbial activity and hydrolase synthesis in long-term Cd-contaminated soils

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Microbial activity and hydrolase synthesis in long-term Cd-contaminated soils / G. RENELLA; M. MENCH; L. LANDI; P. NANNIPIERI. - In: SOIL BIOLOGY & BIOCHEMISTRY. - ISSN 0038-0717. - STAMPA. - 37:(2005), pp. 133-139. [10.1016/j.soilbio.2004.06.015]

Availability:

The webpage <https://hdl.handle.net/2158/311361> of the repository was last updated on

Published version:

DOI: 10.1016/j.soilbio.2004.06.015

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

Microbial activity and hydrolase synthesis in long-term Cd-contaminated soils

Giancarlo Renella^{a,*}, Michel Mench^b, Loretta Landi^a, Paolo Nannipieri^a

^aDepartment of Soil Science and Plant Nutrition, University of Florence, Piazzale delle Cascine 28, Florence 50144, Italy

^bINRA Centre Bordeaux-Aquitaine, 71, Av. E. Bourlaux, BP81, F-33883, Villenave d'Ornon, France

Received 8 July 2003; received in revised form 19 May 2004; accepted 3 June 2004

Abstract

Alkaline and acid phosphomonoesterase, β -glucosidase, arylsulfatase, protease and urease activities, CO_2 -C evolution and ATP content were monitored in long-term Cd-contaminated (0–40 mg Cd kg⁻¹ dry weight soil) sandy soils, kept under maize or ‘set aside’ regimes, amended with plant residues. The organic matter input increased soil respiration, ATP contents and hydrolase activities in all soils. However, the Cd-contaminated soils had significantly higher metabolic quotients ($q\text{CO}_2$), as calculated by the CO_2 -to-ATP ratio, and significantly lower hydrolase activities and hydrolase activity-to-ATP ratios for alkaline phosphomonoesterase, arylsulfatase and protease activities, compared with the respective uncontaminated soils. The ratios between acid phosphomonoesterase, β -glucosidase and urease activities and ATP were unaffected. A significantly higher $q\text{CO}_2/\mu$ ratio, an expression of maintenance energy, was observed in most of the contaminated soils, indicating that more energy was required for microbial synthesis in the presence of high Cd concentrations. It was concluded that exposure to high Cd concentrations led to a less efficient metabolism, which was responsible for lower enzyme activity and synthesis and lower hydrolase activity-to-ATP ratios observed in these Cd-contaminated soils.

© 2004 Elsevier Ltd. All rights reserved.

Keywords: Cadmium; Soil microbial biomass; Hydrolase synthesis

1. Introduction

The energy flow through soil microflora is regulated by biosynthesis and maintenance processes. Both may be affected by environmental factors such as the availability of nutrients and the presence of toxicants. Soil hydrolase activities are key factors controlling nutrient availability in soil (Nannipieri et al., 2002) and adverse effects of heavy metals on soil enzyme activity have long been recognised under both long-term field experiments and laboratory incubations (Tyler et al., 1989). Such effects have been quantified mainly by assaying soil enzyme activity after fresh additions of heavy metals (Tabatabai, 1977; Doelman and Haanstra, 1986; Eivazi and Tabatabai, 1990; Landi et al., 2000), or by calculation of the ecological dose (ED_{50})

value (Haanstra and Doelman, 1991; Speir et al., 1995; Moreno et al., 2001; Renella et al., 2003).

The microflora of heavy metal contaminated soils generally have altered microbial activities, manifested by higher metabolic quotients ($q\text{CO}_2$) (Brookes, 1995) and less microbial biomass synthesis per unit of organic substrate added (Chander and Brookes, 1991, 1992), compared with values in the respective unpolluted soils. Both of these features indicate a reduced metabolic efficiency in the conversion of substrates into biomass (Pirt, 1975).

Among heavy metals, Cd is considered to be particularly toxic and responsible for serious biological degradation of soils (Reber, 1992; Smith, 1996; Giller et al., 1998). However, severe adverse effects of Cd on soil microbial biomass, microbial growth and microbial activities such as enzyme activities have been generally detected only when Cd concentrations were particularly high (Dar, 1996; Moreno et al., 1998, 1999) and its effects on enzymes generally depend on the enzyme activity studied.

* Corresponding author. Tel.: +39-55-3288-219; fax: +39-55-333273.
E-mail address: giancarlo.renella@unifi.it (G. Renella).

Little information is available on the effects of heavy metals such as Cd on the synthesis of hydrolases induced by the availability of organic substrates. The rationale for this work was that stress due to high Cd concentrations in soil should also be reflected in a measurable effect on hydrolase production.

2. Materials and methods

2.1. Soil characteristics and treatments

Soils (Arenic Udifluent) from the long-term AGIR (Table 1) Experiment (Unité d'Agronomie, INRA, Bordeaux, France) were contaminated between 1988 and 1990 with $\text{Cd}(\text{NO}_3)_2$ to give Cd concentrations of 0 (plots 16 and 17), 10 (plots 11 and 12), 20 (plots 6 and 7), and 40 mg kg^{-1} soil (plots 1 and 2). Soils from plots 16, 11, 6 and 1 were cultivated in maize from period the 1988 to 1990. Thereafter, grass (mainly *Festuca* spp.) was cultivated until the year 2000 when soils were cultivated with maize again. Soils from plots 17, 12, 7 and 2 were under set aside (SA) regime, kept barren by tilling (once a year). Soils were surface (0–20 cm) sampled in 2001, sieved (<2 mm) at field moisture, and kept at 50% WHC for 7 days at 25 °C prior to the analyses. After this pre-incubation, 1000 g soil samples were split into two subsamples: one was amended with 10 g kg^{-1} of dry-milled ryegrass (48.1% C, 3.4% N) and mixed thoroughly by hand to ensure homogenous distribution of the organic amendment; the other was mixed in the same way without organic amendment and these subsamples served as controls. Assays of all properties were carried out after 6 h (day 0), 4, 7, and 28 days after the soil amendments.

2.2. Soil respiration and ATP content determinations

Soil samples were placed in 1 l air-tight conical flasks provided with three-way valves and incubated at 25 °C in the dark, with empty flasks used as blank to account

for the background CO_2 concentration. Respiration of both amended and unamended soils was measured by sampling the head-space gas, using an air-tight syringe and injection into a gas-chromatograph (Hewlett-Packard 6890), equipped with a gas-sampling valve, a packed column (Poropak Q) and a thermal conductivity detector (Blackmer and Bremner, 1977). After soil respiration measurements, samples were assayed for the remaining biochemical parameters.

The ATP content of both amended and unamended soils was measured by the phosphoric acid method of Webster et al. (1984) as modified by Ciardi and Nannipieri (1990).

2.3. Soil hydrolase activities measurements

At each selected incubation time, soil hydrolase activities were measured on both amended and unamended soils. Acid and alkaline phosphomonoesterase activities were assayed according to Tabatabai and Bremner (1969), arylsulphatase activity as reported by Tabatabai and Bremner (1970) and β -glucosidase activity according to Tabatabai (1982). Urease activity was measured by the method of Nannipieri et al. (1978), and protease activity by hydrolysis of *N*-benzoylargininamide (*N*-BAA) according to Ladd and Butler (1972). All enzyme assays were carried out at 37 °C, using a 1 h incubation, followed by centrifugation of soil slurries (6000g) at 4 °C. The concentration of *p*-nitrophenol (*p*-NP) produced in the assays of β -glucosidase, arylsulphatase, and acid and alkaline phosphomonoesterase activities was calculated from a *p*-NP calibration curve after subtraction of the absorbance of the controls at 400 nm wavelength. The NH_4^+ produced by urease and *N*-BAA hydrolysing activities was determined by a Flow Injection Analyzer (FIAS 300-Perkin Elmer) coupled to a spectrophotometer Lambda 2 (Perkin Elmer).

2.4. Microbial physiological indices and statistics

The metabolic quotient ($q\text{CO}_2$) values were calculated by the CO_2 -C-to-ATP ratio. The $q\text{CO}_2/\mu$ ratio, the slope of

Table 1
Main physico-chemical characteristics of the AGIR soils

Soil	pH (H_2O)	Clay (%)	Silt (%)	Sand (%)	CEC (cmol kg^{-1})	TOC (%)	N tot (%)	Total Cd (mg kg^{-1})
<i>Maize-cropped soils</i>								
Plot 16	7.4	17.5	15.0	67.5	14.2	0.65	0.10	0.57
Plot 11	7.5	17.5	15.0	67.5	14.5	0.55	0.03	10.9
Plot 6	7.1	17.5	15.0	67.5	15.1	0.47	0.04	17.4
Plot 1	7.1	17.5	15.0	67.5	14.8	0.49	0.04	38.2
<i>Set aside soils</i>								
Plot 17	7.5	17.5	15.0	67.5	14.6	0.64	0.08	1.07
Plot 12	7.0	17.5	15.0	67.5	14.7	0.43	0.02	8.33
Plot 7	7.1	17.5	15.0	67.5	14.9	0.43	0.03	17.2
Plot 2	7.1	17.5	15.0	67.5	14.5	0.49	0.04	39.1

the regression line between the metabolic quotient ($q\text{CO}_2$) and the growth rate (μ), is an expression of the microbial maintenance energy (Pirt, 1975). The (μ) value was calculated from the following relationship: $[\ln(\text{ATP}_2) - \ln(\text{ATP}_1)]/(\Delta T)$, where ATP_1 and ATP_2 are the ATP values at T_1 (day 0) and T_2 (7 days) and ΔT ($T_2 - T_1$) is the time interval (in hours); thus, dimensionally, μ was expressed as $\mu\text{g ATP h}^{-1}$.

All measurements were carried out in triplicate. The standard error and the least significant differences (LSD) of the means by the Tukey–Kramer test (P level <0.05) were calculated using STATVIEW 5 computer programme (SAS Institute).

3. Results

The addition of plant residues increased the soil respiration, the ATP content and hydrolase activities in all soils. Net cumulative respiration rates, net hydrolase activity values, and net ATP content of the maize-cropped (MC) and SA soils were calculated by subtracting the values of the control soils from those of the respective ryegrass-amended soils.

3.1. Soil cumulative respiration and ATP content

About 40% of the organic C added with plant material evolved as $\text{CO}_2\text{-C}$ during the 28 days of incubation from both the MC and SA soils, with no significant differences due to Cd content or soil management (Fig. 1).

The ATP content measured generally increased throughout after the soil amendment (Fig. 1). At the end of the incubation period, the ATP contents in all of the MC soil treatments were similar, whereas they were significantly lower in SA soils containing 10 mg Cd kg^{-1} soil (Fig. 1).

3.2. Hydrolase activities of soils

All hydrolase activities were stimulated by soil amendment with plant material. The alkaline phosphomonoesterase activity peaked after 7 days of incubation in all soils, whereas acid phosphomonoesterase activity, β -glucosidase, arylsulfatase, protease and urease activities displayed the highest value after 28 days in both the MC and SA soils (Figs. 2 and 3).

After 28 days of incubation, the activity of alkaline phosphomonoesterase was significantly lower in the MC soils containing 20 and 40 mg Cd kg^{-1} soil than the respective soil containing 0 and 10 mg Cd kg^{-1} . Its activity was also significantly lower in all Cd-contaminated SA soils compared with the uncontaminated SA soil (Fig. 2). The activity of acid phosphomonoesterase was significantly reduced in the MC soil containing 40 mg Cd kg^{-1} and in all Cd-contaminated SA soils, compared with the respective control soils (Fig. 2).




Fig. 1. Net cumulative respiration and net ATP content of maize and set aside soils amended with plant residues. The error bars represent the standard errors of the means ($n=3$). The least significant difference values (LSD) were calculated with the Tukey–Kramer test ($P=0.05$) for day 28 only.

The activity of β -glucosidase was inhibited in all Cd-contaminated MC soils and in SA soils containing 20 and 40 mg Cd kg^{-1} (Fig. 2) whereas the activity of arylsulfatase was strongly inhibited in all Cd-contaminated MC soils and in SA soils containing 20 and 40 mg Cd kg^{-1} (Fig. 2).

The activity of urease was not inhibited by Cd in both MC or SA soils whereas the activity of protease was significantly inhibited in MC soils and in SA soils containing 20 and 40 mg Cd kg^{-1} (Fig. 2).

Significantly lower hydrolase activity-to-ATP ratios after 28 days of incubation were found in MC soils for alkaline phosphomonoesterase activity (20 and 40 mg Cd kg^{-1} soil) and for arylsulfatase activity (10, 20 and 40 mg Cd kg^{-1} soil) (Table 2). In SA soils, the hydrolase activity-to-ATP ratios after 28 days of incubation were lower for alkaline phosphomonoesterase and arylsulfatase activities in the presence of 40 mg Cd kg^{-1} soil (Table 2).

3.3. Microbial physiological indices

The metabolic quotient ($q\text{CO}_2$), as calculated by the $\text{CO}_2\text{-C-to-ATP}$ ratio was significantly increased in MC soils with Cd concentrations of 20 and 40 mg Cd kg^{-1} soil and in the SA soil with 40 mg Cd kg^{-1} soil (Table 3).

The $q\text{CO}_2/\mu$ ratio, the slope of the regression line between the metabolic quotient ($q\text{CO}_2$) and growth rate (μ) which represents the maintenance energy of microorganisms (Pirt, 1975), was higher in both MC and SA soils containing 20 and 40 mg Cd kg^{-1} (Fig. 4).




Fig. 2. Net alkaline and acid phosphomonoesterase, β -glucosidase and arylsulphatase activities in maize and set aside soils amended with plant residues. The error bars represent the standard errors of the means ($n=3$). The least significant difference values (LSD) were calculated with the Tukey–Kramer test ($P=0.05$) for day 28 only.

4. Discussion

Previous studies have shown that when soil microflora are exposed for a long time to high metal concentrations, its $q\text{CO}_2$ increases, indicating a greater energy requirement for maintenance (Brookes, 1995). Our results confirm these findings, because the $q\text{CO}_2/\mu$ ratio values, expressing the maintenance energy (Pirt, 1975), showed that more $\text{CO}_2\text{--C}$ was respired per unit of ATP formed as the amount of Cd in soil increased (Fig. 4). A lower metabolic efficiency of soil microflora in the presence of high Cd concentrations might be due to energy diversion into physiological adaptations necessary to tolerate heavy metals, such as synthesis of




Fig. 3. Net urease and protease activities in maize and set aside soils amended with plant residues. The error bars represent the standard errors of the means ($n=3$). The least significant difference values (LSD) were calculated with the Tukey–Kramer test ($P=0.05$) for day 28 only.

intra- and extracellular metal-sequestering proteins or saccharides, and biochemical reactions to precipitate or trap metals onto microbial surfaces (Hughes and Poole, 1991; Gadd, 1993). In fact, any adaptation mechanism is energy-demanding, thus enhancing the energy requirements for growth and/or reducing the conversion of substrates into new microbial biomass and its use for synthesis of enzymes, which represent a large portion of total cellular proteins. This might explain the lower synthesis of some hydrolases following the addition of readily mineralizable organic matter in Cd-contaminated soils (Table 2). Indications of greater energy requirements for biomass synthesis in heavy metal contaminated than in uncontaminated soils, have been reported previously (Killham, 1985; Chander and Brookes, 1991, 1992; Chander and Joergensen, 2001).

Another mechanism that could account for the lower enzyme activity during the microbial growth is the synthesis of different enzyme forms with different kinetic parameters. From in vitro studies, it has been showed that, in some microorganisms, genes encoding sulfur-rich isozymes are specifically repressed during exposure to Cd, whereas genes encoding sulfur-poor isozymes are specifically induced (Fauchon et al., 2002).

Variable effects of heavy metal contamination on soil enzyme activities due to the metal type, the contamination mode and the contact time have been reported previously. For example, Dar (1996) found that Cd as sole contaminant decreased the soil enzyme activities when its concentration was as high as 50 mg kg^{-1} . Moreno et al. (1999) reported that high Cd concentrations ($> 800 \text{ mg Cd kg}^{-1}$) affected dehydrogenase activity, but not β -glucosidase urease

Table 2

Hydrolase activity-to-ATP ratios in control (incubation time 0 days) and ryegrass-amended soils (incubation times 7 and 28 days) during the incubation

Cd content (mg kg ⁻¹)	Hydrolase activity-to-ATP ratios (U/mg ATP ⁻¹) ^a					
	Maize-cropped soils			Set aside soils		
	Incubation time (days)					
	0	7	28	0	7	28
<i>Alkaline phosphomonoesterase</i>						
0	1.22	0.42	0.37	1.08	0.26	0.20
10	1.58	0.43	0.36	0.60	0.23	0.17
20	1.57	0.38	0.21	1.10	0.34	0.11
40	2.26	0.18	0.16	1.38	0.09	0.07
LSD	0.42	0.16	0.15	0.56	0.12	0.12
<i>Acid phosphomonoesterase</i>						
0	0.51	0.15	0.14	0.46	0.13	0.11
10	0.64	0.14	0.12	0.48	0.23	0.18
20	0.91	0.24	0.13	0.72	0.33	0.12
40	1.58	0.13	0.14	1.57	0.16	0.13
LSD	0.46	0.09	0.04	0.67	0.14	0.07
<i>Arylsulphatase</i>						
0	2.24	0.57	0.59	0.54	0.16	0.16
10	1.51	0.33	0.33	0.75	0.35	0.30
20	1.95	0.43	0.29	0.88	0.30	0.12
40	2.61	0.26	0.29	1.52	0.13	0.10
LSD	0.53	0.27	0.18	0.41	0.25	0.05
<i>β-glucosidase</i>						
0	0.10	0.13	0.13	0.11	0.10	0.09
10	0.16	0.05	0.07	0.16	0.08	0.09
20	0.31	0.10	0.08	0.16	0.17	0.07
40	0.40	0.07	0.08	0.29	0.11	0.10
LSD	0.37	0.09	0.06	0.14	0.07	0.01
<i>Urease</i>						
0	0.02	0.01	0.01	0.02	0.00	0.00
10	0.03	0.00	0.01	0.02	0.01	0.01
20	0.02	0.00	0.01	0.03	0.01	0.00
40	0.07	0.01	0.01	0.11	0.01	0.01
LSD	0.012	0.003	0.001	0.021	0.001	0.001
<i>Protease</i>						
0	0.08	0.02	0.02	0.11	0.02	0.01
10	0.12	0.02	0.02	0.07	0.02	0.02
20	0.11	0.02	0.01	0.09	0.02	0.01
40	0.21	0.02	0.02	0.19	0.01	0.01
LSD	0.018	0.001	0.002	0.016	0.003	0.002

The LSD value is least significant difference calculated for each incubation time with the Tukey–Kramer test ($P=0.05$).^a The ratios were calculated by dividing the enzyme units by the ATP content of soils. One enzyme unit will release 1 μM/min of *p*-nitrophenol for alkaline and acid phosphomonoesterase, arylsulphatase and β-glucosidase activities and μM/min of NH₄⁺–N for urease and protease activities at 37 °C and optimal pH.

Table 3

Metabolic quotient ($q\text{CO}_2$) of the soils with increasing Cd concentrations

$q\text{CO}_2$ (mg CO ₂ –C μg ATP ⁻¹ day ⁻¹)									
Maize-cropped soils					Set aside soils				
Plot	Unamended	Ryegrass-amended incubation time (days)			Plot	Unamended	Ryegrass-amended incubation time (days)		
		4	7	28			4	7	28
16	0.09	0.70	0.51	0.75	17	0.12	1.16	0.85	0.95
11	0.09	2.62	0.89	0.85	12	0.11	1.92	1.17	0.88
6	0.13	1.25	0.70	0.92	7	0.13	1.04	0.85	0.97
1	0.18	0.79	0.53	0.79	2	0.21	0.79	0.62	0.84
LSD	0.03	0.32	0.26	0.19	LSD	0.05	0.48	0.32	0.22

The LSD value is least significant difference calculated with the Tukey–Kramer test ($P=0.05$) and it is referred to each incubation time.




Fig. 4. Linear regression between $q\text{CO}_2/\mu$ ratio values and soil total Cd concentration. The labels of the points represent the plots number. For calculations of $q\text{CO}_2$ and μ values, see Section 2.

and phosphomonoesterase activities. Karaca et al. (2002) reported significant reductions of several hydrolase activities by Cd-enriched (50 mg kg^{-1}) sewage sludge, but urease activity was not inhibited. Most of the above mentioned studies investigated the effects of incorporation of multi-metal contaminated sewage sludge into soil and the possibility that the results obtained with this approach could be due to additive toxicity of different heavy metals cannot be excluded (Renella, 2003). The approach we have adopted in this study is different because it involves the addition of plant residues to soils where Cd is the only contaminant rather than the incorporation of metal-contaminated organic residues to soils. However, plant material can bind heavy metals due to the presence of reactive functional groups, thereby reducing their availability (Tiemann et al., 1999) and this effect should be taken into account in assessing the adverse effects of Cd in the contaminated soils, where the addition of plant material might have reduced the heavy metal availability.

A change in the soil functionality might also be caused by changes in composition of the microbial community induced by heavy metals (Sandaa et al., 2001; Kozdroj and van Elsas, 2001). However, Renella et al. (2004) reported no shifts in the dominant members of the bacterial community in the same soils. Reductions of soil enzyme activities in heavy metal contaminated soils without changes in the soil microbial community structure were previously reported by Kandeler et al. (2000).

Our results suggest that negative effects on nutrient turnover in Cd-contaminated soils could be due to reductions in enzyme activity. However, such effects are difficult to predict because, although it has been frequently postulated that the regulation of nutrient cycling in soils should involve both enzyme synthesis and enzyme activity, there is a lack of data on the relationship between enzyme synthesis and organic matter mineralization in heavy metal contaminated soils.

In conclusion our results show that long term exposure of soil microflora to high Cd concentrations leads to lower

hydrolase synthesis and reduced metabolic efficiency, possibly as the result of a physiological adaptation of soil microflora to high cadmium concentrations.

References

- Blackmer, A.M., Bremner, J.M., 1977. Gas chromatographic analysis of soil atmosphere. *Soil Science Society of America Journal* 41, 908–912.
- Brookes, P.C., 1995. Use of microbial parameters in monitoring soil pollution by heavy metals. *Biology and Fertility of Soils* 19, 269–279.
- Chander, K., Brookes, P.C., 1991. Microbial biomass dynamics during the decomposition of glucose and maize in metal-contaminated and non-contaminated soils. *Soil Biology & Biochemistry* 23, 917–925.
- Chander, K., Brookes, P.C., 1992. Synthesis of microbial biomass from added glucose in metal-contaminated and non-contaminated soils following repeated fumigation. *Soil Biology & Biochemistry* 24, 613–614.
- Chander, K., Joergensen, R.G., 2001. Decomposition of ^{14}C glucose in two soils with different amounts of heavy metal contamination. *Soil Biology & Biochemistry* 33, 1811–1816.
- Ciardi, C., Nannipieri, P., 1990. A comparison of methods for measuring ATP in soil. *Soil Biology & Biochemistry* 22, 725–727.
- Dar, G.H., 1996. Effects of cadmium and sewage sludge on soil microbial biomass and enzyme activities. *Bioresource Technology* 56, 141–145.
- Doelman, P., Haanstra, L., 1986. Short- and long-term effects of heavy metals on urease activity in soils. *Biology and Fertility of Soils* 2, 213–218.
- Eivazi, F., Tabatabai, M.A., 1990. Factors affecting glucosidase and galactosidase activities in soils. *Soil Biology & Biochemistry* 22, 891–897.
- Fauchon, M., Lagniel, G., Aude, J.-C., Lombardia, L., Soularue, P., Petat, C., Marguerie, G., Sentenac, A., Werner, M., Labarre, J., 2002. Sulfur sparing in the yeast proteome in response to sulfur demand. *Molecular Cell* 9, 713–723.
- Gadd, G.M., 1993. Interactions of fungi with toxic metals. *New Phytologist* 124, 25–60.
- Giller, K., Witter, E., McGrath, S.P., 1998. Toxicity of heavy metals to microorganisms and microbial processes in agricultural soils: a review. *Soil Biology & Biochemistry* 30, 1389–1414.
- Haanstra, L., Doelman, P., 1991. An ecological dose-response model approach to short- and long-term effects of heavy metals on arylsulfatase activity of soil. *Biology and Fertility of Soils* 11, 18–23.
- Hughes, M.N., Poole, R.K., 1991. Metal speciation and microbial growth, the hard and soft facts. *Journal of General Microbiology* 137, 725–734.
- Kandeler, E., Tcherko, D., Bruce, K.D., Stemmer, M., Hobbs, P.J., Bardgett, R.D., Amelung, W., 2000. Structure and function of the soil microbial community in microhabitats of a heavy metal polluted soil. *Biology and Fertility of Soils* 30, 390–400.
- Karaca, A., Naseby, D.C., Lynch, J.M., 2002. Effect of cadmium contamination with sewage sludge and phosphate fertiliser amendments on soil enzyme activities, microbial structure and available cadmium. *Biology and Fertility of Soils* 35, 428–434.
- Killham, K., 1985. A physiological determination of the impact of environmental stress on the activity of soil microbial biomass. *Environmental Pollution* 38, 283–294.
- Kozdroj, J., van Elsas, J.D., 2001. Structural diversity of microorganisms in chemically perturbed soils assessed by molecular and cytochemical approaches. *Journal of Microbiological Methods* 43, 197–212.
- Ladd, J.N., Butler, J.H.A., 1972. Short-term assays of soil proteolytic enzyme activities using proteins and dipeptide derivatives as substrates. *Soil Biology & Biochemistry* 4, 19–30.
- Landi, L., Renella, G., Moreno, J.L., Falchini, L., Nannipieri, P., 2000. Influence of Cd on the metabolic quotient, L:D-glutamic

- acid respiration ratio and enzyme activity: microbial biomass ratio under laboratory conditions. *Biology and Fertility of Soils* 32, 8–16.
- Moreno, J.L., Garcia, C., Hernandez, T., 1998. Changes in organic matter and enzyme activity of an agricultural soil amended with metal-contaminated sewage sludge compost. *Communications in Soil Science and Plant Analysis* 29, 2247–2262.
- Moreno, J.L., Hernandez, T., Garcia, C., 1999. Effects of cadmium-contaminated sewage sludge compost on dynamics of organic matter and microbial activity in an arid soil. *Biology and Fertility of Soils* 28, 230–237.
- Moreno, J.L., Garcia, C., Landi, L., Falchini, L., Pietramellara, G., Nannipieri, P., 2001. The ecological dose value (ED_{50}) for assessing Cd toxicity on ATP content and dehydrogenase and urease activities of soil. *Soil Biology & Biochemistry* 33, 483–489.
- Nannipieri, P., Johnson, R.L., Paul, E.A., 1978. Criteria for measurement of microbial growth and activity in soil. *Soil Biology & Biochemistry* 10, 223–229.
- Nannipieri, P., Kandeler, E., Ruggiero, P., 2002. Enzyme activity as monitors of soil microbial functional diversity, in: Burns, R.G., Dick, R. (Eds.), *Enzymes in the Environment: Activity, Ecology and Applications*. Marcel Dekker, New York, pp. 234–251.
- Pirt, S.J., 1975. *Principles of Microbe and Cell Cultivation*. Blackwell Scientific Publication, Oxford.
- Reber, H.H., 1992. Simultaneous estimates of the diversity and the degradative capability of heavy-metal-affected soil bacterial communities. *Biology and Fertility of Soils* 13, 181–186.
- Renella, G., Reyes Ortigoza, A.L., Landi, L., Nannipieri, P., 2003. Additive effects of copper and zinc on cadmium toxicity to phosphomonoesterase activities and ATP content of soil as estimated by the ecological dose (ED_{50}). *Soil Biology & Biochemistry* 35, 1203–1210.
- Renella, G., Mench, M., van der Lelie, D., Pietramellara, G., Ascher, J., Ceccherini, M.T., Landi, L., Nannipieri, P., 2004. Hydrolase activity, microbial biomass and community structure in long-term Cd-contaminated soils. *Soil Biology & Biochemistry* 36, 443–451.
- Sandaa, R.-A., Torsvik, V., Enger, E., 2001. Influence of long-term heavy-metal contamination on microbial communities in soil. *Soil Biology & Biochemistry* 33, 287–295.
- Smith, S.R., 1996. *Agricultural Recycling of Sewage Sludge and the Environment*. CABI, Wallingford pp. 382.
- Speir, T.W., Kettles, H.A., Parshotam, A., Searle, P.L., Vlaar, L.N.C., 1995. Simple kinetic approach to derive the ecological dose values, ED_{50} , for the assessment of Cr (VI) toxicity to soil biological properties. *Soil Biology & Biochemistry* 27, 801–810.
- Tabatabai, M.A., 1977. Effects of trace elements on urease activity. *Soil Biology & Biochemistry* 9, 9–13.
- Tabatabai, M.A., 1982. Soil enzymes, in: Page, A.L., Miller, R.H., Keeney, D.R. (Eds.), *Methods of Soil Analysis, Part 2, Chemical and Microbiological Properties*. Soil Science, of America, Madison, WI, pp. 907–943.
- Tabatabai, M.A., Bremner, J.M., 1969. Use of *p*-nitrophenyl phosphate for assay of soil phosphomonoesterase activity. *Soil Biology & Biochemistry* 1, 301–307.
- Tabatabai, M.A., Bremner, J.M., 1970. Arylsulfatase activity of soils. *Soil Science Society of America Proceedings* 34, 225–229.
- Tiemann, K.J., Gardea Torresdey, J., Gamez, G., Dokken, K., Sias, S., 1999. Use of X-ray absorption spectroscopy and esterification to investigate Cr(III) and Ni(II) ligands in alfalfa biomass. *Environmental Science & Technology* 33, 150–154.
- Tyler, G., Balsberg-Pahlson, A.M., Bengtsson, G., Bååth, E., Tranvik, L., 1989. Heavy-metal ecology of terrestrial plants, micro-organisms and invertebrates. *Water Air and Soil Pollution* 47, 189–215.
- Webster, J., Hampton, G., Leach, F., 1984. ATP in soil: a new extractant and extraction procedure. *Soil Biology & Biochemistry* 16, 335–342.