



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Fluoride ion promoted reactivity of thiocarbonyl containing molecules with silyl derivatives: a novel access to allyl sulfides

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Fluoride ion promoted reactivity of thiocarbonyl containing molecules with silyl derivatives: a novel access to allyl sulfides / A. Degl'Innocenti; B.F. Bonini; G. Mazzanti; A. Capperucci; M.C. Ferrara; A. Ricci; P. Zani. - In: PHOSPHORUS SULFUR AND SILICON AND THE RELATED ELEMENTS. - ISSN 1042-6507. - ELETTRONICO. - 74:(1993), pp. 365-366.

Availability:

The webpage <https://hdl.handle.net/2158/338197> of the repository was last updated on 2020-06-30T12:56:00Z

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

FLUORIDE ION PROMOTED REACTIVITY OF THIOCARBONYL CONTAINING MOLECULES WITH SILYL DERIVATIVES : A NOVEL ACCESS TO ALLYL SULFIDES

**ALESSANDRO DEGL'INNOCENTI^a, BIANCA BONINI^b, GERMANA MAZZANTI^b,
ANTONELLA CAPPERUCCI^{*c}, M. CRISTIANA FERRARA^c, ALFREDO RICCI^{a,b} AND
PAOLO ZANI^b**

^a Centro CNR Composti Eterociclici, Via Gino Capponi, 9 - 50121 Firenze
(Italy)

^b Dipartimento di Chimica Organica "A. Mangini", Viale Risorgimento, 4 - 40100
Bologna (Italy)

^c Dipartimento di Chimica Organica "U. Schiff", Via Gino Capponi, 9 - 50121
Firenze (Italy)

Abstract: Fluoride ion induced reaction of allyl- and benzyl-silanes with
thiocarbonyls leads, contrary to the corresponding lithium or Grignard
reagents, to products of S-functionalization.


The chemistry of thiocarbonyl containing molecules has seen an increasing interest in the recent years ¹. Moreover, reaction of such compounds with various organometallic species, like lithium, sodium or magnesium halogenides have been extensively studied². Although thiophilic addition is frequently reported, other various reactions can be observed simultaneously, like carbophilic addition, reduction, double addition and formation of enesulfides.

Our recent interest in the development of organosilicon based procedures for the synthesis of sulfur containing compounds³ led us to investigate the reactivity of thiocarbonyl containing molecules, such as thioketones, dithioesters and trithiocarbonates toward different organometallic species, like derivatives of group 14 elements, namely organosilanes and stannanes.

We report in this communication our findings on the fluoride ion induced reactivity of silylated nucleophiles, such as allylsilanes and benzylsilane toward different thiocarbonyl containing molecules.

Thus, for instance, when reacting different aromatic thioketones either with

benzylsilane or allylsilane, a clean reaction occurs, leading in good yields to the corresponding sulfides outlining a clean inversion of the regioselectivity of addition with respect to allyl- and benzyldervatives of lithium and magnesium^{2,4}.



This reactivity can be conveniently extended to different thiocarbonyl containing molecules, such as dithioesters and trithiocarbonates, in which case again an inversion of regioselectivity is observed, leading to the isolation of S-functionalized products.

Different fluoride ion sources have been tested, like tetrabutylammonium fluoride (TBAF) and tris(dimethylamino)sulfonium difluorotrimethylsilicate (TASF), the former giving generally better yields of the reaction product.

The results obtained then show that the fluoride ion induced reactivity of allylsilanes or benzylsilanes with thioketones and thioesters affords an easy and high yielding access to a wide variety of allylsulfides, otherwise not easily obtainable from the common literature procedures and opens new perspectives in the control of the regiochemistry of such reaction, in strict dependence to the organometallic derivative used.

REFERENCES:

1. E.Vedejs, *Acc. Chem. Res.*, **1984**, 17, 358.
2. P.Beak, I.Yamamoto, C.J.Upton, *J. Org. Chem.*, **1975**, 40, 3052; D.Paquer, *Bull. Soc. Chim. Fr.*, **1975**, 1439.
3. A.Capperucci, A.Degl'Innocenti, A.Ricci, A.Mordini, G.Reginato, *J. Org. Chem.*, **1991**, 56, 7323; G.Capozzi, A.Capperucci, A.Degl'Innocenti, R.Del Duce, S.Menichetti, *Gazz. Chim. Ital.*, **1990**, 120, 421.
4. V.Rautenstrauch, *Helv. Chim. Acta*, **1974**, 57, 496; P.Metzener, J.Vialle, A.Vibet, *Tetrahedron*, **1978**, 34, 2289 and refs. cited therein; S.Masson, M.Saquet, A.Thuillier, *Tetrahedron*, **1977**, 33, 2949 and refs. cited therein.