



UNIVERSITÀ  
DEGLI STUDI  
FIRENZE

## FLORE

# Repository istituzionale dell'Università degli Studi di Firenze

### **Supraspinal inactivation of mitochondrial superoxide dismutase is a source of peroxynitrite in the development of morphine**

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

*Original Citation:*

Supraspinal inactivation of mitochondrial superoxide dismutase is a source of peroxynitrite in the development of morphine antinociceptive tolerance / T. Doyle;L. Bryant;l. Batinic-Haberle;j. Little;S. Cuzzocrea;E. Masini;l. Spasojevic;D. Salvemini. - In: BMC NEUROSCIENCE. - ISSN 1471-2202. - ELETTRONICO. - 164:(2009), pp. 702-710. [10.1016/j.neuroscience.2009.07.019]

*Availability:*

The webpage <https://hdl.handle.net/2158/364936> of the repository was last updated on

*Published version:*

DOI: 10.1016/j.neuroscience.2009.07.019

*Terms of use:*

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

*Publisher copyright claim:*

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

Published in final edited form as:

*Neuroscience*. 2009 December 1; 164(2): 702–710. doi:10.1016/j.neuroscience.2009.07.019.

## Supraspinal Inactivation of Mitochondrial Superoxide Dismutase is a Source of Peroxynitrite in the Development of Morphine Antinociceptive Tolerance

Timothy Doyle<sup>1</sup>, Leesa Bryant<sup>1</sup>, Ines Batinic-Haberle<sup>2</sup>, Joshua Little<sup>3</sup>, Salvatore Cuzzocrea<sup>4,5</sup>, Emanuela Masini<sup>6</sup>, Ivan Spasojevic<sup>7</sup>, and Daniela Salvemini<sup>1,§</sup>

<sup>1</sup>Department of Pharmacological and Physiological Science, Saint Louis University School of Medicine, 1402 South Grand Blvd, St. Louis, MO 63104, USA

<sup>2</sup>Department of Radiation Oncology, Duke University School of Medicine, Durham NC 27710, USA

<sup>3</sup>Department of Surgery, Center for Anatomical Science and Education, 1402 S. Grand Blvd., Saint Louis University School of Medicine, St Louis, MO 63104, USA

<sup>4</sup>Department of Clinical and Experimental Medicine and Pharmacology, School of Medicine, University of Messina, Italy

<sup>5</sup>IRCCS Centro Neurolesi "Bonino-Pulejo" Messina, Italy

<sup>6</sup>Department of Preclinical and Clinical Pharmacology, University of Florence, Italy

<sup>7</sup>Department of Medicine, Duke University School of Medicine, Durham NC 27710, USA

### Abstract

Effective treatment of chronic pain with morphine is limited by decreases in the drug's analgesic action with chronic administration (antinociceptive tolerance). Because opioids are mainstays of pain management, restoring their efficacy has great clinical importance. We have recently reported that formation of peroxynitrite (ONOO<sup>-</sup>, PN) in the dorsal horn of the spinal cord plays a critical role in the development of morphine antinociceptive tolerance and have further documented that nitration and enzymatic inactivation of mitochondrial superoxide dismutase (MnSOD) at that site provides a source for this nitroxidative species. We now report for the first time that antinociceptive tolerance is also associated with the inactivation of MnSOD at supraspinal sites. Inactivation of MnSOD led to nitroxidative stress as evidenced by increased levels of products of oxidative DNA damage and activation of the nuclear factor poly (ADP-ribose) polymerase in whole brain homogenates. Co-administration of morphine with potent Mn porphyrin-based peroxynitrite scavengers, (MnTE-2-PyP<sup>5+</sup> and MnTnHex-2-PyP<sup>5+</sup>) (1) restored the enzymatic activity of MnSOD, (2) attenuated PN derived nitroxidative stress, and (3) blocked the development of morphine induced antinociceptive tolerance. The more lipophilic analogue,

© 2009 IBRO. Published by Elsevier Ltd. All rights reserved.

§**Address correspondence to:** Daniela Salvemini, Ph.D Department of Pharmacological and Physiological Science, Saint Louis University School of Medicine, 1402 South Grand Blvd, St. Louis, MO 63104, USA; Phone: (1) 314. 577. 8856; Fax (1) 314. 577. 8859; salvemd@slu.edu.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Conflict of interest:** The authors have declared that no conflict of interest exists.

MnTnHex-2-PyP<sup>5+</sup> was able to cross the blood brain barrier at higher levels than its lipophilic counterpart MnTE-2-PyP<sup>5+</sup> and was about 30 fold more efficacious. Collectively, these data suggest that peroxynitrite mediated enzymatic inactivation of supraspinal MnSOD provides a source of nitroxidative stress, which in turn contributes to central sensitization associated with the development of morphine antinociceptive tolerance. These results support our general contention that PN-targeted therapeutics may have potential as adjuncts to opiates in pain management.

## Keywords

morphine; mitochondrial superoxide dismutase; superoxide; peroxynitrite; peroxynitrite decomposition catalysts; nitroxidative stress

## Introduction

Opiate/narcotic analgesics, typified by morphine sulfate, are the most effective treatments for acute and chronic severe pain; but their clinical utility is often hampered by the development of analgesic tolerance as well as by *de novo* painful hypersensitivity to innocuous and noxious stimuli, phenomena observed in both animal and human studies (Arner et al., 1988, Mao et al., 1995). With respect to morphine in particular, tolerance necessitates escalating doses to achieve equivalent pain relief (Foley, 1995). This complex pathophysiological cycle contributes to decreased quality of life in the growing population of subjects with chronic pain because of oversedation, reduced physical activity, respiratory depression, constipation, potential for addiction, and other side-effects (Foley, 1995). Accordingly, there is major interest in new approaches to maintain opiate efficacy during repetitive dosing for chronic pain, without engendering tolerance or unacceptable side-effects. Our studies to date have shown that targeting peroxynitrite (ONOO<sup>-</sup>, PN) is an effective therapeutic strategy in blocking the development of antinociceptive tolerance (Salvemini, 2009, Salvemini and Neumann, 2009).

Spinal formation of PN, the reaction product between superoxide (O<sub>2</sub><sup>·-</sup>) and nitric oxide (·NO) (Beckman et al., 1990), is a potent proinflammatory reactive nitroxidative species (Salvemini et al., 1998, Jagtap and Szabo, 2005). Since the rate of interaction between ·NO and O<sub>2</sub><sup>·-</sup> to form PN is faster than the dismutation of O<sub>2</sub><sup>·-</sup> by SOD, the most critical roles of O<sub>2</sub><sup>·-</sup> and ·NO in pain and inflammation may be their formation of PN (Salvemini, 2009). Indeed, PN has been recently implicated in the development of thermal hyperalgesia associated with acute and chronic inflammation (Wang et al., 2004, Khattab, 2006, Bezerra et al., 2007, Ndengele et al., 2008), in response to spinal activation of the N-methyl-D-aspartate receptor (NMDAR) (Muscoli et al., 2004) and in the development of opiate induced antinociceptive tolerance (Muscoli et al., 2007, Batinic-Haberle et al., 2009b, Ndengele et al., 2009). Furthermore, the use of non-selective pharmacological probes (i.e. these agents react not only with O<sub>2</sub><sup>·-</sup> but also with several nitroxidative species and derivatives thereof) (Muscoli et al., 2003) such as PBN [phenyl *N-tert*-butylnitron (PBN)] and TEMPOL [4-hydroxy-2,2,6,6-tetramethylpiperidine 1-oxyl] has supported the role of nitroxidative stress (defined as stress caused by the presence of O<sub>2</sub><sup>·-</sup>, ·NO and/or PN and related derivatives) in both neurogenic nociception (Gao et al., 2007, Lee et al., 2007, Schwartz et al., 2008, Schwartz et al., 2009), visceral pain (Wang et al., 2008) and neuropathic pain (Tal, 1996, Park et al., 2006, Gao et al., 2007, Siniscalco et al., 2007). Ultimately, selective PN decomposition catalysts should confirm the contribution of PN in such settings.

It is well documented that inactivation of manganese superoxide dismutase (MnSOD), the enzyme that normally keeps concentrations of O<sub>2</sub><sup>·-</sup> under tight control (McCord and

Fridovich, 1969) is a central source for  $O_2^{\cdot-}$ -derived PN in many diseases driven by overt production of PN (MacMillan-Crow and Thompson, 1999, MacMillan-Crow et al., 2001). Such enzymatic inactivation results from nitration on Tyr-34 by PN in a Mn-catalysed process (MacMillan-Crow and Thompson, 1999). We have recently extended these findings to provide evidence that spinal inactivation of MnSOD is also a central source for the formation of PN in the development of morphine antinociceptive tolerance (Muscoli et al., 2007, Ndengele et al., 2009). Under these circumstances, increased spinal PN contributes to antinociceptive tolerance through three well defined biochemical pathways within the dorsal horn of the spinal cord: (1) post-translational nitration of proteins involved in glutamate homeostasis such as glutamate transporters and glutamine synthase (2) neuroimmune activation and (3) neuronal apoptosis (Muscoli et al., 2007, Batinic-Haberle et al., 2009b). Whether PN-mediated nitroxidative stress at supraspinal sites contributes to the development of morphine antinociceptive tolerance is not known. In this study we show that the development of antinociceptive tolerance following chronic administration of morphine was associated with supraspinal inactivation of MnSOD which provided a source for PN and PN-mediated nitroxidative stress. The PN decomposition catalysts MnTE-2-PyP<sup>5+</sup> and MnTnHex-2-PyP<sup>5+</sup> (Batinic-Haberle, 2002) potently blocked MnSOD nitration at supraspinal sites and the development of antinociceptive tolerance with MnTnHex-2-PyP<sup>5+</sup> being more potent than MnTE-2-PyP<sup>5+</sup>. These results suggest that in addition to the documented spinal role of PN in tolerance (Muscoli et al., 2007, Batinic-Haberle et al., 2009a, Ndengele et al., 2009), supraspinal PN has an important role as well.

## Methods

### Mn porphyrins

MnTE-2-PyP<sup>5+</sup> and MnTnHex-2-PyP<sup>5+</sup> were synthesized and characterized as previously described (Batinic-Haberle, 2002).

### Thin-layer chromatography

Thin-layer chromatography of the purified compounds was performed on plastic-backed silica gel TLC plates with 1:1:8 KNO<sub>3</sub>-saturated H<sub>2</sub>O: H<sub>2</sub>O: acetonitrile mixture as mobile phase. Typically, 1  $\mu$ L of ~1 mM samples were applied at ~1 cm of the strip border and the solvent front was allowed to run ~12 cm. The absolute R<sub>f</sub> values and their ratio thereof are sensitive to the degree of the saturation of vapor phase in the TLC chamber and may differ slightly from one to another experiment. Thus, the internal standardization of TLC is required for comparison purposes [Kos et al Free Radic Biol Med 2009]. For the purpose of this study and given high levels of ascorbate in brain Mn(III) porphyrins and ascorbate-reduced Mn(II) porphyrins (MnTE-2-PyP<sup>4+</sup> and MnTnHex-2-PyP<sup>4+</sup>) were applied on silica plates. Mn(II) porphyrins are less stable and may lose Mn to some extent. Consequently in solution there may be some amount of metal-free porphyrins that bear same 4+ total charge and may have similar adsorption on the silica as reduced Mn(II) porphyrins. Thus we applied on TLC plates metal-free porphyrins, H<sub>2</sub>TE-2-PyP<sup>4+</sup> and H<sub>2</sub>TnHex-2-PyP<sup>4+</sup> as controls also. The thin-layer chromatographic behavior/lipophilicity of Mn porphyrins was described by retention factor, R<sub>f</sub> which presents the ratio of Mn porphyrin path and solvent path. Based on R<sub>f</sub> values, the P<sub>OW</sub> values were then calculated (see Results).

### Induction of morphine-induced antinociceptive tolerance in mice

The development of morphine induced tolerance was performed as described (Muscoli et al., 2007, Batinic-Haberle et al., 2009b, Ndengele et al., 2009). Male CD-1 mice (24–30g, Charles River) were housed and cared for in accordance with the guidelines of the Institutional Animal Care and Use Committee of the Saint Louis University Health Science Center and in accordance with the National Institute of Health *Guidelines on Laboratory*

*Animal* and the University of Messina, in compliance with Italian regulations on protection of animals used for experimental and other scientific purposes (D.M. 116192) as well as with the EEC regulations (O.J. of E.C. L 358/1 12/18/1986). Mice were housed 4–5 per cage, maintained under identical conditions of temperature ( $21\pm 1^\circ\text{C}$ ), humidity ( $60\pm 5\%$ ) with a 12-hr light-dark cycle, and allowed food *ad libitum*. Nociceptive/pain thresholds were determined by measuring latencies (in seconds, s) of mice placed in a transparent glass cylinder on a hot plate (Ugo Basile, Italy) maintained at  $52^\circ\text{C}$ . Responses indicative of nociception included intermittent lifting and/or licking of the hindpaws or escape behavior. Determination of antinociception/pain relief effects was assessed between 7:00 and 10:00 AM. All injections were given by intraperitoneal (ip) or subcutaneous (sc) means in a 0.1 ml volume at approximately 7 AM and 4 PM. Drug or vehicle (saline) was given before each dose of morphine. Hot plate latencies were taken in mice from all groups on day five before (baseline latency) and 40 min after an acute dose of morphine (3 mg/kg) or its vehicle (saline) (response latency). Baseline values from all groups as measured on day five before injection of the acute dose of morphine or its vehicle, similarly ranged between 6–8 s. Results are expressed as percent maximal possible antinociceptive effect (% MPE) calculated as follows:  $(\text{response latency} - \text{baseline latency}) / (\text{cut off latency} - \text{baseline latency}) \times 100$  (Muscoli et al., 2007). A cut-off latency of 20 s was employed to prevent tissue damage. Twelve mice per group were used and all experiments were conducted with the experimenters blinded to treatment conditions. Unless specified, all drugs were purchased from Sigma. The following experimental groups were used.

**Naïve group (N)**—Mice were injected twice a day with an ip injection of saline and a sc injection of saline. On day five, mice received an ip injection of saline followed fifteen minutes later by a sc injection of saline.

**Vehicle group (V)**—Mice were injected twice a day with an ip injection of saline and a sc injection of saline. On day five, mice received an ip injection of saline followed fifteen minutes later by a sc injection of acute morphine eliciting near-to-maximal antinociception (3 mg/kg) at 40 minutes after injection.

**Tolerant group (T)**—Mice were injected twice a day with an ip injection of saline and sc injection of morphine (20 mg/kg/day). On day five, mice received an ip injection of saline followed fifteen minutes later by a sc injection of acute morphine (3 mg/kg). Hot plate latencies were taken 40 minutes after morphine injection.

**Tolerant + drugs group**—Mice were injected twice a day with an ip injection of varying doses of MnTE-2-PyP<sup>5+</sup> (0.3–3 mg/kg/day) or MnTnHex-2-PyP<sup>5+</sup> (0.01–0.1 mg/kg/day) followed by a sc injection of morphine (20 mg/kg/day). On day five, mice received an ip injection of MnTE-2-PyP<sup>5+</sup> (1.5 mg/kg/day) or MnTnHex-2-PyP<sup>5+</sup> (0.05 mg/kg/day) followed fifteen minutes later by the sc injection of acute morphine (3 mg/kg). Hot plate latencies were taken 40 minutes after morphine injection.

After the behavioral tests on day five, whole brains were removed and tissues processed for Western blot and biochemical analysis.

## Western Blot

Whole brain lysates ( $n = 3$  per group) were prepared as previously described (Wang et al., 2004). Briefly, whole frozen brain (395–500 mg) were pulverized in liquid nitrogen-chilled mortar and pestle prior to homogenization in 1.2 – 1.5 mL of lysis buffer [20 mM Tris-Cl (pH 7.4), 150 mM NaCl, 16.2 mM CHAPS, 12.5 mM EGTA, 1% protease cocktail (Sigma, St. Louis MO) (final concentration: 1 mM 4-(2-aminoethyl)benzenesulfonyl fluoride

(AEBSF), 15  $\mu$ M pepstatinA, 14  $\mu$ M E-64, 40  $\mu$ M bestatin, 20  $\mu$ M leupeptin, and 850 nM aprotinin), 10% glycerol]. The homogenates were sonicated for 10 min on ice in an ultrasonic bath (VWR, Buffalo Grove IL), incubated an additional 10 min on ice, and clarified by centrifugation for 10 min at 12500g, 4°C. The supernatants were frozen at -80°C and the total protein content of an aliquot was measured by bicinchoninic acid assay (Pierce, Rockford IL). The total lysates (40  $\mu$ g) were denatured with an equal volume of loading buffer (2X: 125 mM Tris-Cl (pH 6.8), 15 % glycerol, 3% SDS, 1.1 mM  $\beta$ -mercaptoethanol, and 0.075% bromophenol blue) and heat denatured for 5 min at 95°C. The total lysates were resolved by SDS-PAGE (170 V, ~55 min) with a 4–20% Tris-glycine minigel (Lonza, Basile Switzerland) and electrophoretically transferred (100V, 1.5 h) to 0.2  $\mu$ m nitrocellulose membrane (Whatman, Fordham Park NJ) in Towbin transfer buffer. Additional transfers of total lysates serially diluted (2-fold) were prepared. The relative levels of total MnSOD were determined by enhanced chemiluminescence (Wang et al., 2004). Each membrane was blocked for 2 hr at RT (10% low-fat milk/PBS-T: 1X PBS, 0.05% Tween-20, and 0.1% thimerasol) and incubated for 1 hr at RT with a rabbit anti-MnSOD monoclonal antibody (Millipore) diluted 1:1000 in 5% low-fat milk/PBS-T. The membrane was washed 5 times in PBS-T (15 min and 4  $\times$  5 min), then incubated with a goat anti-rabbit IgG-HRP antibody diluted 1:10000 in 5% low-fat milk/PBS-T. After 5 washes in PBS-T, the bands were visualized with Supersignal<sup>®</sup> West Femto chemiluminescence substrate (Pierce) and captured on film (GE Healthcare, Piscataway NJ). The membranes were subsequently washed twice with PBS-T for 10 min and equal loading was similarly verified with a mouse anti- $\beta$ -actin monoclonal antibody (Sigma, St. Louis MO) diluted 1:10000 and goat anti-mouse IgG-HRP antibody diluted 1:8000 in 5% low-fat milk/PBS-T. The lack of species cross-reactivity of HRP-conjugated antibodies to the anti-MnSOD and anti- $\beta$ -actin was also confirmed. The densities of the MnSOD and  $\beta$ -actin bands were quantified using ImageQuant 5.2 (GE Healthcare) and expressed as relative MnSOD: $\beta$ -actin densities.

### Measurement of MnSOD and Cu,ZnSOD activities

Whole brain tissue was homogenized in a Polytron homogenizer in 4 – 5 ml of 10 mM phosphate buffered saline, pH 7.2; then sonicated twice on ice for 20 s. The sonicated samples were subsequently centrifuged at 1,100 g for 10–15 min and SOD activity was measured in the supernatants (Wang et al., 2004). In brief, a competitive inhibition assay was performed using xanthine-xanthine oxidase-generated superoxide to reduce nitroblue tetrazolium (NBT) to blue tetrazolium salt. The reaction was performed in 50 mM carbonate buffer, pH 10.1, containing 0.1 mM EDTA, 25  $\mu$ M nitroblue tetrazolium, 0.1 mM xanthine and 2 nM xanthine oxidase (Boehringer, Germany). The rate of NTB reduction was monitored spectrophotometrically (Perkin Elmer, Lambda 5 Spectrophotometer, Milan, Italy) at 560 nm. The amount of protein required to inhibit the rate of NBT reduction by 50% was defined as one unit of enzyme activity. Cu,ZnSOD activity was inhibited by performing the assay in the presence of 2 mM NaCN after preincubation for 30 min. Enzymatic activity was expressed in mU per  $\mu$ g of protein as previously described (Wang et al., 2004).

### Determination of 8-hydroxy-2'-deoxyguanosine (8-OHdG)

The isolation of DNA was performed as described previously by our group (Masini et al., 2005). In brief, whole brain tissue was homogenized in 4 – 5 ml of 10 mM PBS, pH 7.4, sonicated on ice 20 s twice, then 1 ml of 10 mM Tris-HCl buffer, pH 8, containing 10 mM EDTA, 10 mM NaCl, and 0.5% sodium dodecyl sulphate was added. The lysates were incubated for 1 h at 37°C with 20  $\mu$ g/ml RNase I (Sigma-Aldrich, Milan, Italy) and then overnight at 37 °C under argon in the presence of 100  $\mu$ g/ml proteinase K (Sigma-Aldrich). The DNA was extracted with chloroform/isoamyl alcohol (10/2 v/v), precipitated from the

aqueous phase with 0.2 volumes of 10 M ammonium acetate, solubilized in 200  $\mu$ l of 20 mM acetate buffer, pH 5.3, and denatured at 90°C for 3 min. The extract was then supplemented with 10 IU of P1 nuclease (Sigma-Aldrich) in 10  $\mu$ l and incubated for 1 h at 37 °C with 5 IU of alkaline phosphatase (Sigma-Aldrich) in 0.4 M phosphate buffer, pH 8.8. All of the procedures were performed in the dark under argon. The mixture was filtered by an Amicon Micropure-EZ filter (Millipore Corporation, Billerica, MA), and 50  $\mu$ l of each sample was used for 8-hydroxy-2'-deoxyguanosine (8-OHdG) determination using a Bioxytech enzyme immunoassay kit (Oxis, Portland, OR), per manufacturer's protocol. The values are expressed as nanogram of 8-OHdG per milligram of protein (Bradford, 1976).

### Measurement of PARP Activity

PARP-1 activity was measured as described previously (Suzuki et al., 2004). Whole brain tissue was gently homogenized at 4 °C, in 4 – 5 mL of 50 mM Tris HCl, pH 8, containing 0.1% tergitol-type NP-40 (NP-40), 200 mM KCl, 2 mM MgCl<sub>2</sub>, 50  $\mu$ M ZnCl<sub>2</sub>, 2 mM dithiothreitol (DTT) and protease inhibitors: 1 mM phenylmethylsulfonyl fluoride (PMSF), 5  $\mu$ l/ml leupeptin, and antipain. Samples were then centrifuged and 10  $\mu$ l of each supernatant were incubated for 5 min at 25 °C with 2  $\mu$ l of [<sup>3</sup>H]NAD<sup>+</sup> (specific activity 25 Ci/mmol, G.E., Health Care, U.K.) in 50 mM Tris HCl, pH 8, containing 20 mM MgCl<sub>2</sub>, 1 mM DTT, and 20  $\mu$ M NAD<sup>+</sup> in the absence or presence of activated calf thymus DNA, in the final volume of 100  $\mu$ l. The reaction was stopped by the addition of 5% trichloroacetic acid. Samples were filtered and the radioactivity in the acid-insoluble fraction was counted by a Beckman LS1801 liquid scintillation spectrometer. PARP activity estimated without activated DNA in the mixture was assigned as "endogenous" activity. Activity estimated in the presence of activated DNA in the assay mixture was assigned as "total" activity of PARP. Ratio between endogenous and total activity was considered as the measure of PARP activity in the tissue. The values are expressed as pmol of NAD<sup>+</sup> per  $\mu$ g of protein/h (Bradford, 1976).

### Statistics

For paired group analysis Students t-test were performed. For paired multiple groups, analysis of variance followed by Student-Newman-Keuls test were employed to analyze the data. Results are expressed as mean $\pm$ s.e.m for n animals. A statistical significant difference was defined as a *P* value <0.05.

## Results

### Lipophilicity of Mn porphyrins

The thin-layer chromatographic behavior of reduced Mn(II) porphyrins, MnTE-2-PyP<sup>4+</sup> and MnTnHEX-2-PyP<sup>4+</sup>, as described by retention factor  $R_f$  is given in Table 1. It is obvious that reduced porphyrins are significantly more lipophilic than their oxidized Mn(III) analogues. In brain due to the high level of ascorbate and their easy reducibility with ascorbate (Ferrer-Sueta et al Chem Res Toxicol 1999, Batinic-Haberle et al, Free Radic Biol Med 2004), Mn porphyrins will at least in part exist in reduced state which would further enhance their accumulation. Utilizing  $P_{OW}$  vs  $R_f$  relationship (Kos et al., 2009) as given by equation,  $\log P_{OW} = 12.18 \times R_f - 7.43$ , we calculated the partition of both porphyrins between n-octanol and water,  $P_{OW}$ .  $P_{OW}$  is a more common indicator of drug lipophilicity. Both  $R_f$  and  $P_{OW}$  values are listed in Table 1.

### **Inactivation of brain mitochondrial superoxide dismutase (MnSOD) as a source of PN in the development of antinociceptive tolerance**

When compared to mice treated over 4 days with saline, chronic administration of morphine over the same time course led to the inactivation of MnSOD in whole brain homogenates as evidenced by loss of the catalytic activity of this enzyme to dismutate superoxide measured spectrophotometrically on day 5 (n=12) (Fig. 2A). Chronic administration of morphine did not change the total amount of MnSOD protein in brain homogenates as measured by Western blotting analysis (Fig. 2B, C). The enzymatic activity of cytosolic SOD (Cu,ZnSOD) was not affected (n=12) (Fig. 2D). Inactivation of brain MnSOD was associated with the development of antinociceptive tolerance (Fig. 3A, B). When compared to animals receiving an equivalent injection of its vehicle (Naïve group), acute injection of morphine (3 mg/kg) in animals that received saline over four days (Vehicle group) produced a significant and near-to-maximal antinociceptive response [percent maximal possible antinociceptive effect (%MPE) ranging between 90–95%] (Fig. 3A, B). On the other hand, when compared to the antinociceptive response to acute morphine in animals that received saline over four days, repeated administration of morphine over the same time course (Tolerant group) led to the development of antinociceptive tolerance as evidenced by a significant loss of its antinociceptive response (Fig. 3A, B). Baseline latencies in groups treated with saline or morphine over four days before acute administration of morphine on day five similarly ranged between 6–8 seconds (n=12).

Co-administration of morphine with varying doses of the PN scavengers MnTE-2-PyP<sup>5+</sup> (0.3–3 mg/kg/day) or MnTnHex-2-PyP<sup>5+</sup> (0.01–0.1 mg/kg/day) prevented the supraspinal inactivation of MnSOD and restored its ability to enzymatically dismutate superoxide (n=12) (Fig. 2A) and blocked in a dose dependent fashion morphine antinociceptive tolerance [0.3–3 mg/kg/day for MnTE-2-PyP<sup>5+</sup> and 0.01–0.1 mg/kg/day for MnTnHex-2-PyP<sup>5+</sup>, n=12] (Fig. 3A, B). We have previously demonstrated that MnTE-2-PyP<sup>5+</sup> or MnTnHex-2-PyP<sup>5+</sup> alone do not increase baseline latencies (Batinic-Haberle et al., 2009b).

### **The development of morphine-induced antinociceptive tolerance is associated with supraspinal oxidative DNA damage and PARP activation: inhibition by MnTE-2-PyP<sup>5+</sup> or MnTnHex-2-PyP<sup>5+</sup>**

It is now well established that inactivation of MnSOD increases superoxide levels thereby favoring PN formation and thus increased nitroxidative stress (Yamakura et al., 1998, MacMillan-Crow and Thompson, 1999, Macmillan-Crow and Cruthirds, 2001, MacMillan-Crow et al., 2001, Yamakura et al., 2001). Here we show that the development of morphine antinociceptive tolerance was associated with increased formation of nitroxidative stress markers as measured in whole brain homogenates. Thus and as can be seen in Fig. 4, when compared to mice treated over 4 days with saline, chronic administration of morphine over the same time course increased 8OHdG (n=12, Fig. 4A), a marker of oxidative DNA damage and increased the activation of PARP a nuclear enzyme known to be activated following oxidative DNA damage (Fig. 4B). Co-administration of morphine with MnTE-2-PyP<sup>5+</sup> (3 mg/kg/day) or MnTnHex-2-PyP<sup>5+</sup> (0.1 mg/kg/day) significantly reduced the increased production of 8-OHdG (n=12) and significantly decreased PARP activity (Fig. 4A, B).

## **Discussion**

Mitochondrial superoxide dismutase (MnSOD), the enzyme that normally keeps concentrations of superoxide under tight control (McCord and Fridovich, 1969) is an exquisite and sensitive target for PN-mediated nitration and enzymatic inactivation (MacMillan-Crow and Thompson, 1999). This process favors the accumulation of PN which

in turn, nitrates and alters additional proteins and receptors, thereby perpetuating and extending the initial damage (Radi et al., 2002). Over the years a great body of experimental evidence has been gathered to link MnSOD inactivation as a source for PN and PN-driven nitroxidative stress to several disease states (Macmillan-Crow and Cruthirds, 2001). More recently, spinal inactivation of MnSOD has been linked to the development of peripheral and central sensitization associated with inflammation and in the development of opiate-induced antinociceptive tolerance (Muscoli et al., 2004, Wang et al., 2004, Muscoli et al., 2007, Schwartz et al., 2008, Batinic-Haberle et al., 2009b, Ndengele et al., 2009, Schwartz et al., 2009). The main objective of the study presented herein was to address whether in addition to its spinal role (Muscoli et al., 2007, Batinic-Haberle et al., 2009b, Ndengele et al., 2009), supraspinal inactivation of MnSOD, as a source of PN, contributes to the development of morphine antinociceptive tolerance. Our results clearly show that the development of antinociceptive tolerance was associated with enzymatic inactivation of MnSOD in whole brain homogenates which in turn contributed to PN formation and PN-mediated nitroxidative stress as evidenced by increased formation of 8-OHdG and increase in PARP activity well known markers of the presence of severe nitroxidative stress. Inhibition of PN with two PN scavengers (MnTE-2-PyP<sup>5+</sup>, MnTnHex-2-PyP<sup>5+</sup>) prevented MnSOD inactivation and attenuated antinociceptive tolerance. The more lipophilic MnTnHex-2-PyP<sup>5+</sup> was found to be approximately 30-fold more potent a finding that can be explained at least in part by the fact that MnTnHex-2-PyP<sup>5+</sup> has better organ uptake (i.e brain). The finding that the development of morphine-induced antinociceptive tolerance was not associated with enzymatic inactivation of the cytosolic isoform of SOD, namely Cu,ZnSOD, is consistent with previous studies that have shown that the interaction of Cu,ZnSOD with PN does not affect the catalytic activity of the protein (Smith et al., 1992). Although little is known about supraspinal locations and mechanisms of nitroxidative species in hyperalgesic states, the rostral ventromedial medulla (RVM) may provide a point of origin for further characterization of the roles of peroxynitrite during central sensitization.

Glutamate neurotransmission, in particular that mediated via activation of NMDAR and glial cell activation in the RVM, contributes to central sensitization (see below). The  $\mu$ -opioid receptors, NMDA receptors, and glial cell activation in the RVM are all potential sources for PN formation since each of these can release its precursors: O<sub>2</sub><sup>-</sup> and ·NO [see (Watkins et al., 2007, Salvemini and Neumann, 2009) for reviews]. Furthermore, PN contributes to NMDA-mediated central sensitization (Muscoli et al., 2004). Investigations using intracerebroventricular injections of non-selective NOS inhibitor (Moore et al., 1991, Kawabata et al., 1993, Salter et al., 1996, Zhao and Bhargava, 1996) or free radical scavengers (Kim et al., 2004, Lee et al., 2007) attenuate central sensitization in various animal models of hypersensitivity including opiate tolerance, implicating roles for nitroxidative species at supraspinal sites. More specifically, RVM microinjections of NOS inhibitors and NMDAR antagonists were all able to significantly reduce secondary hyperalgesia following peripheral insult (Coutinho et al., 1998, 2001). Conversely, RVM microinjections of ·NO donors or NMDA enhanced hyperalgesia (Urban et al., 1999, Guan et al., 2002). Collectively these results suggest that supraspinal formation of nitroxidative species formed upon activation of the NMDAR modulates central sensitization.

The homeostasis of extracellular glutamate, a primary endogenous ligand for the NMDAR, is tightly regulated by sodium-dependent high-affinity glutamate transporters (GTs) in the plasma membranes of both neurons and glial cells (Danbolt, 2001). Loss of the transport function of these proteins leads to increased glutamate levels in the synaptic cleft, overstimulation of NMDAR and neurotoxicity (Mennerick et al., 1999, Lievens et al., 2000). In contradistinction to the key role of GTs, glutamine synthetase (GS) plays a pivotal role in its intracellular metabolic fate by converting glutamate into nontoxic glutamine. Pertinent to the context of our studies, is that PN nitrates and inactivates the transport activity of GTs

and the enzymatic activity of GS (Trotti et al., 1996, Minana et al., 1997, Trotti et al., 1999, Gorg et al., 2005). Furthermore, Zanelli and co-workers demonstrated that PN nitrates tyrosine residues on NMDA receptor subunits, an event associated with constant potentiation of synaptic currents, calcium influx, and ultimately increased NMDA-receptor excitability and neuronal excitotoxicity (Zanelli et al., 2000, Zanelli et al., 2002). We have recently shown that the development of opiate antinociceptive tolerance was associated with nitration of GTs and GS (Muscoli et al., 2007) and of NMDAR (Salvemini, unpublished observations) in dorsal horn tissues. Taken collectively and in an attempt to provide a mechanism whereby PN formation at supraspinal sites modulates opiate tolerance, it is possible that PN fosters tolerance via a pathway involving nitroxidative alterations of glutamate homeostasis.

Spinal neuroimmune activation contributes to opiate antinociceptive tolerance [(Watkins et al., 2007) (Salvemini and Neumann, 2009)for reviews] and inhibition of PN blocks spinal glial cell activation and cytokine formation during antinociceptive tolerance (Muscoli et al., 2007, Batinic-Haberle et al., 2009b, Ndengele et al., 2009, Salvemini and Neumann, 2009). Since neuroimmune activation in the RVM has been associated with central sensitization (Wei et al., 2008) we hypothesize that a second mechanism by which supraspinal PN fosters tolerance is neuroimmune activation.

In summary, we have shown for the first time that supraspinal inactivation of MnSOD and PN mediated nitroxidative stress plays a role in antinociceptive tolerance. We have further shown that potent peroxynitrite scavengers, MnTE-2-PyP<sup>5+</sup> and MnTnHex-2-PyP<sup>5+</sup> were able to prevent MnSOD inactivation and nitroxidative stress at that level thereby inhibiting morphine tolerance. Investigations to determine the region of and cellular sources of supraspinal PN will be important to determine neuroanatomical site of action during antinociceptive tolerance. The RVM appears to provide a plausible site of origin to begin these investigations.

## Abbreviations

8-OHdG	8-hydroxy-2'-deoxyguanosine
%MPE	percent maximal possible antinociceptive effect
Cu,ZnSOD	copper/zinc superoxide dismutase
GS	glutamine synthase
GT	glutamate transporters
$k_{\text{cat}}$ and $k_{\text{red}}$	catalytic and reduction rate constants
MnSOD	manganese superoxide dismutase
MnTE-2-PyP <sup>5+</sup>	Mn(III) 5,10,15,20-tetrakis( <i>N</i> -ethylpyridinium-2-yl)porphyrin
MnTnHex-2-PyP <sup>5+</sup>	Mn(III) 5,10,15,20-tetrakis( <i>N</i> -n-hexylpyridinium-2-yl)porphyrin
NMDA	N-methyl-D-aspartate
NMDAR	N-methyl-D-aspartate receptor
·NO	nitric oxide
O <sub>2</sub> <sup>·-</sup>	superoxide
ONOO <sup>-</sup> and PN	peroxynitrite

PARP	poly(ADP-ribose) polymerase; PMSF
P <sub>OW</sub>	n-octanol/water partitioning constant
RVM	rostral ventromedial medulla
SOD	superoxide dismutase
E <sub>1/2</sub>	metal-centered reduction potential for Mn(III)P/Mn(II)P redox couple
NHE	normal hydrogen electrode
R <sub>f</sub> thin-layer chromatographic retention factor	ration of Mn porphyrin path and solvent path on silica plates
P <sub>OW</sub>	partition coefficient between n-octanol and water

## Acknowledgments

This work was supported by IRCCS Centro Neurolesi “Bonino-Pulejo” grant (SC), by 1R01DA024074-01A1 (DS, IBH), R21DA023056-01A2 (DS), NIH U19AI67798-01 (IBH), William H. Coulter Translation Partners Grant Program (IBH), and NIH/NCI Duke Comprehensive Cancer Center Core Grant (5-P30-CA014236-33) (IS).


## References

- Arner S, Rawal N, Gustafsson LL. Clinical experience of long-term treatment with epidural and intrathecal opioids--a nationwide survey. *Acta Anaesthesiol Scand* 1988;32:253–259. [PubMed: 3364150]
- Batinic-Haberle I, Cuzzocrea S, Reboucas JS, Ferrer-Sueta G, Mazzon E, Di Paola R, Radi R, Spasojevic I, Benov L, Salvemini D. Pure MnTBAP selectively scavenges peroxynitrite over superoxide: Comparison of pure and commercial MnTBAP samples to MnTE-2-PyP in two models of oxidative stress injury, an SOD-specific *Escherichia coli* model and carrageenan-induced pleurisy. *Free Radic Biol Med* 2009a;46:192–201. [PubMed: 19007878]
- Batinic-Haberle I, Ndengele MM, Cuzzocrea S, Reboucas JS, Spasojevic I, Salvemini D. Lipophilicity is a critical parameter that dominates the efficacy of metalloporphyrins in blocking the development of morphine antinociceptive tolerance through peroxynitrite-mediated pathways. *Free Radic Biol Med* 2009b;46:212–219. [PubMed: 18983908]
- Batinic-Haberle ISI, Stevens RD, Hambright P, Fridovich I. Manganese(III) Meso Tetrakis Ortho N-alkylpyridylporphyrins. Synthesis, Characterization and Catalysis of O<sub>2</sub>- Dismutation. *J Chem Soc, Dalton Trans* 2002:2689–2696.
- Beckman JS, Beckman TW, Chen J, Marshall PA, Freeman BA. Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide. *Proceedings of the National Academy of Sciences of the United States of America* 1990;87:1620–1624. [PubMed: 2154753]
- Bezerra MM, Brain SD, Girao VC, Greenacre S, Keeble J, Rocha FA. Neutrophils-derived peroxynitrite contributes to acute hyperalgesia and cell influx in zymosan arthritis. *Naunyn Schmiedebergs Arch Pharmacol* 2007;374:265–273. [PubMed: 17171392]
- Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976;72:248–254. [PubMed: 942051]
- Coutinho SV, Urban MO, Gebhart GF. Role of glutamate receptors and nitric oxide in the rostral ventromedial medulla in visceral hyperalgesia. *Pain* 1998;78:59–69. [PubMed: 9822212]
- Coutinho SV, Urban MO, Gebhart GF. The role of CNS NMDA receptors and nitric oxide in visceral hyperalgesia. *European journal of pharmacology* 2001;429:319–325. [PubMed: 11698052]
- Danbolt NC. Glutamate uptake. *Prog Neurobiol* 2001;65:1–105. [PubMed: 11369436]


- Ferrer-Sueta G, Hannibal L, Batinic-Haberle I, Radi R. Reduction of manganese porphyrins by flavoenzymes and submitochondrial particles: a catalytic cycle for the reduction of peroxynitrite. *Free Radic Biol Med* 2006;41:503–512. [PubMed: 16843831]
- Ferrer-Sueta G, Vitturi D, Batinic-Haberle I, Fridovich I, Goldstein S, Czapski G, Radi R. Reactions of manganese porphyrins with peroxynitrite and carbonate radical anion. *J Biol Chem* 2003;278:27432–27438. [PubMed: 12700236]
- Foley KM. Misconceptions and controversies regarding the use of opioids in cancer pain. *Anticancer Drugs* 1995;6:4–13. [PubMed: 7606036]
- Gao X, Kim HK, Chung JM, Chung K. Reactive oxygen species (ROS) are involved in enhancement of NMDA-receptor phosphorylation in animal models of pain. *Pain* 2007;131:262–271. [PubMed: 17317010]
- Gorg B, Wettstein M, Metzger S, Schliess F, Haussinger D. Lipopolysaccharide-induced tyrosine nitration and inactivation of hepatic glutamine synthetase in the rat. *Hepatology* 2005;41:1065–1073. [PubMed: 15830392]
- Guan Y, Terayama R, Dubner R, Ren K. Plasticity in excitatory amino acid receptor-mediated descending pain modulation after inflammation. *The Journal of pharmacology and experimental therapeutics* 2002;300:513–520. [PubMed: 11805211]
- Jagtap P, Szabo C. Poly(ADP-ribose) polymerase and the therapeutic effects of its inhibitors. *Nat Rev Drug Discov* 2005;4:421–440. [PubMed: 15864271]
- Kawabata A, Umeda N, Takagi H. L-arginine exerts a dual role in nociceptive processing in the brain: involvement of the kyotorphin-Met-enkephalin pathway and NO-cyclic GMP pathway. *British journal of pharmacology* 1993;109:73–79. [PubMed: 8388303]
- Khattab MM. TEMPOL, a membrane-permeable radical scavenger, attenuates peroxynitrite- and superoxide anion-enhanced carrageenan-induced paw edema and hyperalgesia: a key role for superoxide anion. *European journal of pharmacology* 2006;548:167–173. [PubMed: 16973155]
- Kim HK, Park SK, Zhou JL, Taglialatela G, Chung K, Coggeshall RE, Chung JM. Reactive oxygen species (ROS) play an important role in a rat model of neuropathic pain. *Pain* 2004;111:116–124. [PubMed: 15327815]
- Kos I, Reboucas JS, DeFreitas-Silva G, Salvemini D, Vujaskovic Z, Dewhirst MW, Spasojevic I, Batinic-Haberle I. Lipophilicity of potent porphyrin-based antioxidants: comparison of ortho and meta isomers of Mn(III) N-alkylpyridylporphyrins. *Free Radic Biol Med* 2009;47:72–78. [PubMed: 19361553]
- Lee I, Kim HK, Kim JH, Chung K, Chung JM. The role of reactive oxygen species in capsaicin-induced mechanical hyperalgesia and in the activities of dorsal horn neurons. *Pain* 2007;133:9–17. [PubMed: 17379413]
- Lievens JC, Bernal F, Forni C, Mahy N, Kerkerian-Le Goff L. Characterization of striatal lesions produced by glutamate uptake alteration: cell death, reactive gliosis, and changes in GLT1 and GADD45 mRNA expression. *Glia* 2000;29:222–232. [PubMed: 10642749]
- Macmillan-Crow LA, Cruthirds DL. Invited review: manganese superoxide dismutase in disease. *Free Radic Res* 2001;34:325–336. [PubMed: 11328670]
- MacMillan-Crow LA, Cruthirds DL, Ahki KM, Sanders PW, Thompson JA. Mitochondrial tyrosine nitration precedes chronic allograft nephropathy. *Free Radic Biol Med* 2001;31:1603–1608. [PubMed: 11744334]
- MacMillan-Crow LA, Thompson JA. Tyrosine modifications and inactivation of active site manganese superoxide dismutase mutant (Y34F) by peroxynitrite. *Archives of biochemistry and biophysics* 1999;366:82–88. [PubMed: 10334867]
- Mao J, Price DD, Mayer DJ. Mechanisms of hyperalgesia and morphine tolerance: a current view of their possible interactions. *Pain* 1995;62:259–274. [PubMed: 8657426]
- Masini E, Bani D, Vannacci A, Pierpaoli S, Mannaioni PF, Comhair SA, Xu W, Muscoli C, Erzurum SC, Salvemini D. Reduction of antigen-induced respiratory abnormalities and airway inflammation in sensitized guinea pigs by a superoxide dismutase mimetic. *Free Radic Biol Med* 2005;39:520–531. [PubMed: 16043023]
- McCord JM, Fridovich I. Superoxide dismutase. An enzymic function for erythrocyte hemocuprein (hemocuprein). *J Biol Chem* 1969;244:6049–6055. [PubMed: 5389100]

- Mennerick S, Shen W, Xu W, Benz A, Tanaka K, Shimamoto K, Isenberg KE, Krause JE, Zorumski CF. Substrate turnover by transporters curtails synaptic glutamate transients. *J Neurosci* 1999;19:9242–9251. [PubMed: 10531428]
- Minana MD, Kosenko E, Marcaida G, Hermenegildo C, Montoliu C, Grisolia S, Felipe V. Modulation of glutamine synthesis in cultured astrocytes by nitric oxide. *Cell Mol Neurobiol* 1997;17:433–445. [PubMed: 9262869]
- Moore PK, Oluyomi AO, Babbedge RC, Wallace P, Hart SL. L-NG-nitro arginine methyl ester exhibits antinociceptive activity in the mouse. *British journal of pharmacology* 1991;102:198–202. [PubMed: 2043923]
- Muscoli C, Cuzzocrea S, Ndengele MM, Mollace V, Porreca F, Fabrizi F, Esposito E, Masini E, Matuschak GM, Salvemini D. Therapeutic manipulation of peroxynitrite attenuates the development of opiate-induced antinociceptive tolerance in mice. *J Clin Invest* 2007;117:3530–3539. [PubMed: 17975673]
- Muscoli C, Cuzzocrea S, Riley DP, Zweier JL, Thiemeermann C, Wang ZQ, Salvemini D. On the selectivity of superoxide dismutase mimetics and its importance in pharmacological studies. *British journal of pharmacology* 2003;140:445–460. [PubMed: 14522841]
- Muscoli C, Mollace V, Wheatley J, Masini E, Ndengele M, Wang ZQ, Salvemini D. Superoxide-mediated nitration of spinal manganese superoxide dismutase: a novel pathway in N-methyl-D-aspartate-mediated hyperalgesia. *Pain* 2004;111:96–103. [PubMed: 15327813]
- Ndengele MM, Cuzzocrea S, Esposito E, Mazzon E, Di Paola R, Matuschak GM, Salvemini D. Cyclooxygenases 1 and 2 contribute to peroxynitrite-mediated inflammatory pain hypersensitivity. *Faseb J* 2008;22:3154–3164. [PubMed: 18497304]
- Ndengele MM, Cuzzocrea S, Masini E, Vinci MC, Esposito E, Muscoli C, Petrusca DN, Mollace V, Mazzon E, Li D, Petrache I, Matuschak GM, Salvemini D. Spinal ceramide modulates the development of morphine antinociceptive tolerance via peroxynitrite-mediated nitroxidative stress and neuroimmune activation. *The Journal of pharmacology and experimental therapeutics* 2009;329:64–75. [PubMed: 19033555]
- Park ES, Gao X, Chung JM, Chung K. Levels of mitochondrial reactive oxygen species increase in rat neuropathic spinal dorsal horn neurons. *Neurosci Lett* 2006;391:108–111. [PubMed: 16183198]
- Radi R, Cassina A, Hodara R. Nitric oxide and peroxynitrite interactions with mitochondria. *Biol Chem* 2002;383:401–409. [PubMed: 12033431]
- Salter M, Strijbos PJ, Neale S, Duffy C, Follenfant RL, Garthwaite J. The nitric oxide-cyclic GMP pathway is required for nociceptive signalling at specific loci within the somatosensory pathway. *Neuroscience* 1996;73:649–655. [PubMed: 8809786]
- Salvemini D. Peroxynitrite and opiate antinociceptive tolerance: a painful reality. *Archives of biochemistry and biophysics* 2009;484:238–244. [PubMed: 19017525]
- Salvemini D, Jensen MP, Riley DP, Misko TP. Therapeutic manipulations of peroxynitrite. *Drug News Perspect* 1998;11:204–214. [PubMed: 15616662]
- Salvemini D, Neumann WL. Peroxynitrite: a strategic linchpin of opioid analgesic tolerance. *Trends in pharmacological sciences* 2009;30:194–202. [PubMed: 19261337]
- Schwartz ES, Kim HY, Wang J, Lee I, Klann E, Chung JM, Chung K. Persistent pain is dependent on spinal mitochondrial antioxidant levels. *J Neurosci* 2009;29:159–168. [PubMed: 19129394]
- Schwartz ES, Lee I, Chung K, Chung JM. Oxidative stress in the spinal cord is an important contributor in capsaicin-induced mechanical secondary hyperalgesia in mice. *Pain* 2008;138:514–524. [PubMed: 18375065]
- Siniscalco D, Fuccio C, Giordano C, Ferraraccio F, Palazzo E, Luongo L, Rossi F, Roth KA, Maione S, de Novellis V. Role of reactive oxygen species and spinal cord apoptotic genes in the development of neuropathic pain. *Pharmacol Res* 2007;55:158–166. [PubMed: 17207636]
- Smith CD, Carson M, van der Woerd M, Chen J, Ischiropoulos H, Beckman JS. Crystal structure of peroxynitrite-modified bovine Cu,Zn superoxide dismutase. *Archives of biochemistry and biophysics* 1992;299:350–355. [PubMed: 1444476]
- Suzuki Y, Masini E, Mazzocca C, Cuzzocrea S, Ciampa A, Suzuki H, Bani D. Inhibition of poly(ADP-ribose) polymerase prevents allergen-induced asthma-like reaction in sensitized Guinea pigs. *The*

- Journal of pharmacology and experimental therapeutics 2004;311:1241–1248. [PubMed: 15254147]
- Tal M. A novel antioxidant alleviates heat hyperalgesia in rats with an experimental painful peripheral neuropathy. *Neuroreport* 1996;7:1382–1384. [PubMed: 8856680]
- Trotti D, Rolfs A, Danbolt NC, Brown RH Jr, Hediger MA. SOD1 mutants linked to amyotrophic lateral sclerosis selectively inactivate a glial glutamate transporter. *Nat Neurosci* 1999;2:848. [PubMed: 10461226]
- Trotti D, Rossi D, Gjesdal O, Levy LM, Racagni G, Danbolt NC, Volterra A. Peroxynitrite inhibits glutamate transporter subtypes. *J Biol Chem* 1996;271:5976–5979. [PubMed: 8626378]
- Urban MO, Zahn PK, Gebhart GF. Descending facilitatory influences from the rostral medial medulla mediate secondary, but not primary hyperalgesia in the rat. *Neuroscience* 1999;90:349–352. [PubMed: 10215139]
- Wang J, Cochran V, Abdi S, Chung JM, Chung K, Kim HK. Phenyl N-t-butylnitron, a reactive oxygen species scavenger, reduces zymosan-induced visceral pain in rats. *Neurosci Lett* 2008;439:216–219. [PubMed: 18514415]
- Wang ZQ, Porreca F, Cuzzocrea S, Galen K, Lightfoot R, Masini E, Muscoli C, Mollace V, Ndengele M, Ischiropoulos H, Salvemini D. A newly identified role for superoxide in inflammatory pain. *The Journal of pharmacology and experimental therapeutics* 2004;309:869–878. [PubMed: 14988418]
- Watkins LR, Hutchinson MR, Ledebor A, Wieseler-Frank J, Milligan ED, Maier SF. Norman Cousins Lecture. Glia as the "bad guys": implications for improving clinical pain control and the clinical utility of opioids. *Brain Behav Immun* 2007;21:131–146. [PubMed: 17175134]
- Wei F, Guo W, Zou S, Ren K, Dubner R. Supraspinal glial-neuronal interactions contribute to descending pain facilitation. *J Neurosci* 2008;28:10482–10495. [PubMed: 18923025]
- Yamakura F, Matsumoto T, Fujimura T, Taka H, Murayama K, Imai T, Uchida K. Modification of a single tryptophan residue in human Cu,Zn-superoxide dismutase by peroxynitrite in the presence of bicarbonate. *Biochimica et biophysica acta* 2001;1548:38–46. [PubMed: 11451436]
- Yamakura F, Taka H, Fujimura T, Murayama K. Inactivation of human manganese-superoxide dismutase by peroxynitrite is caused by exclusive nitration of tyrosine 34 to 3-nitrotyrosine. *J Biol Chem* 1998;273:14085–14089. [PubMed: 9603906]
- Zanelli SA, Ashraf QM, Delivoria-Papadopoulos M, Mishra OP. Peroxynitrite-induced modification of the N-methyl-D-aspartate receptor in the cerebral cortex of the guinea pig fetus at term. *Neurosci Lett* 2000;296:5–8. [PubMed: 11099820]
- Zanelli SA, Ashraf QM, Mishra OP. Nitration is a mechanism of regulation of the NMDA receptor function during hypoxia. *Neuroscience* 2002;112:869–877. [PubMed: 12088746]
- Zhao GM, Bhargava HN. Nitric oxide synthase inhibition attenuates tolerance to morphine but not to [D-Ala2, Glu4] deltorphin II, a delta 2-opioid receptor agonist in mice. *Peptides* 1996;17:619–623. [PubMed: 8804071]




**Fig. 1.** Structures of MnTE-2-PyP<sup>5+</sup> and MnTnHex-2-PyP<sup>5+</sup>. Replacement of shorter ethyl with longer hexyl side chains increases lipophilicity by several orders of magnitude (see Table 1).




**Fig. 2.**

When compared to the vehicle group (V) repeated administration of morphine (tolerant group, T) led to significant functional enzymatic inactivation of MnSOD (A) but not Cu,ZnSOD (D) as evidenced by loss of its catalytic activity to dismutate superoxide as measured spectrophotometrically. Co-administration of morphine with MnTE-2-PyP<sup>5+</sup> (3 mg/kg/day, n=12) or MnTnHex-2-PyP<sup>5+</sup> (0.1 mg/kg/day, n=12) restored the enzymatic activity of MnSOD. When compared to the vehicle group, repeated administration of morphine did not change the total amount of MnSOD in whole brain tissue homogenates as measured by Western blotting analysis. Gels shown in B are representative from gels obtained in 3 animals. A composite of the densitometry data resulting from these experiments is shown in C. \*P<0.001 for Morphine alone vs Vehicle; †P<0.001 for Morphine+drug-treated vs Morphine alone.

**Fig. 3.**

Inhibition of morphine antinociceptive tolerance with MnTE-2-PyP<sup>5+</sup> or MnTnHex-2-PyP<sup>5+</sup>. On day five, acute injection of morphine (3 mg/kg) in animals that received saline over four days produced a significant antinociceptive response when compared to responses observed in animals that received an equivalent volume of saline (**A, B**, Vehicle, V vs Naïve, N). On the other hand, a significant loss to the antinociceptive effect from the acute injection of morphine was observed in animals that received repeated administration of morphine over four days (tolerant group) (**A, B**). Co-administration of morphine over four days with varying doses of MnTE-2-PyP<sup>5+</sup> (0.3–3 mg/kg/day, n=12; **A**) or MnTnHex-2-PyP<sup>5+</sup> (0.01–0.1 mg/kg/day, n=12; **B**) inhibited the development of tolerance in a dose-dependent manner. Results are expressed as mean  $\pm$  s.e.m. for six animals. °P<0.001 for Vehicle, V vs Naïve, N; \*P<0.001 for Morphine alone vs Vehicle; †P<0.001 for Morphine + drug treated vs Morphine alone.



**Fig. 4.**

When compared to the vehicle group (V) repeated administration of morphine (tolerant group, T) led to significant increase in dorsal horn tissue levels of 8OHdG (**A**) and substantially activated PARP (**B**). These biochemical changes were significantly attenuated by co-administration of morphine over four days with MnTE-2-PyP<sup>5+</sup> (3 mg/kg/day, n=12) or MnTnHex-2-PyP<sup>5+</sup> (0.1 mg/kg/day, n=12) (**A, B**). Results are expressed as mean±s.e.m. for twelve animals. \*P<0.001 for Morphine alone vs Vehicle; †P<0.001 for Morphine + drug-treated vs Morphine alone.

Table 1

The catalytic rate constants for  $O_2^-$  dismutation ( $k_{cat}$  in  $M^{-1}s^{-1}$  at 25°C) and  $ONOO^-$  reduction ( $k_{red}$  in  $M^{-1}s^{-1}$  at 37°C); lipophilicity expressed in terms of partition between n-octanol and water,  $POW$  for oxidized,  $Mn^{III}Ps$  and reduced  $Mn^{II}Ps$ ; thermodynamic property, metal-centered reduction potential for the redox couple  $Mn^{III}P/Mn^{II}P$ ,  $E_{1/2}$  in mV vs NHE; and blood to brain ratio for ethyl, MnTE-2-PyP<sup>5+</sup> and hexyl, MnTnHex-2-PyP<sup>5+</sup> analogues.

MnP	$\log k_{cat}(O_2^-)$	$\log k_{red}(ONOO^-)$	$E_{1/2}$	$POW(Mn^{III}P)$	$POW(Mn^{II}P)$	$R_f$ Blood : brain ratio
MnTE-2-PyP <sup>5+</sup>	7.76 <sup>a</sup>	7.53 <sup>b(&gt;7)c</sup>	+228 <sup>a</sup>	-6.89 <sup>d</sup>	-4.69 <sup>e</sup>	0.23
MnTnHex-2-PyP <sup>5+</sup>	7.48 <sup>a</sup>	7.11 <sup>b</sup>	+314 <sup>a</sup>	-2.76 <sup>d</sup>	-1.64 <sup>e</sup>	0.48

<sup>a</sup> (Batinic-Haberle, 2002);

<sup>b</sup> (Ferrer-Sueta et al., 2003);

<sup>c</sup> value in parenthesis relates to  $Mn^{II}$  (Ferrer-Sueta et al., 2006);

<sup>d</sup> (Kos et al., 2009);

<sup>e</sup> Data calculated based on  $R_f$  values for metal-center reduced Mn(II) porphyrins (by ascorbate) on TLC silica plates in  $KNO_3$  - saturated  $H_2O$  :  $H_2O$  : acetonitrile = 1:1:8 and  $POW$  vs  $R_f$  relationships,  $\log POW = 12.18 \times R_f - 7.43$ , from ref (Kos et al., 2009);  $R_f$ , a thin-layer chromatographic retention factor is a ratio of MnP path over solvent path,  $POW$  partition coefficient between n-octanol and water;

<sup>f</sup> Spasojevic et al, unpublished.