UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Matching perceived depth from disparity and from velocity: Modeling and psychophysics

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Matching perceived depth from disparity and from velocity: Modeling and psychophysics / F.Domini; C.Caudek. - In: ACTA PSYCHOLOGICA. - ISSN 0001-6918. - STAMPA. - 133:(2010), pp. 81-89. [10.1016/j.actpsy.2009.10.003]

Availability:

This version is available at: 2158/369832 since:

Published version:

DOI: 10.1016/j.actpsy.2009.10.003

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

Matching perceived depth from disparity and from velocity: Modeling and psychophysics

Fulvio Domini^{a,*}, Corrado Caudek^b

^a Brown University, Cognitive and Linguistic Sciences, Providence, RI 02912, USA

^b Università degli Studi di Firenze, Firenze, Italy

ARTICLE INFO

Article history:

Received 17 July 2009

Received in revised form 6 October 2009

Accepted 8 October 2009

Available online 5 December 2009

PsycINFO classification:

2323

2340

Keywords:

3D shape

Vision

Perception

Disparity

Motion

Depth

Perceptual biases

Scaling

ABSTRACT

We asked observers to match in depth a disparity-only stimulus with a velocity-only stimulus. The observers' responses revealed systematic biases: the two stimuli appeared to be matched in depth when they were produced by the projection of different distal depth extents. We discuss two alternative models of depth recovery that could account for these results. (1) Depth matches could be obtained by scaling the image signals by constants not specified by optical information, and (2) depth matches could be obtained by equating the stimuli in terms of their signal-to-noise ratios (see Domini & Caudek, 2009). We show that the systematic failures of shape constancy revealed by observers' judgments are well accounted for by the hypothesis that the apparent depth of a stimulus is determined by the magnitude of the retinal signals relative to the uncertainty (i.e., internal noise) arising from the measurement of those signals.

© 2009 Elsevier B.V. All rights reserved.

1. Introduction

When a stationary object is viewed binocularly, the horizontal disparities created by the relief of the object can be used by the visual system to recover information about the three-dimensional (3D) shape. Likewise, when the same object is viewed monocularly, information about 3D shape can be gathered from the pattern of projected velocities generated by the relative motion between an observer and the object. In the present investigation, we will try to understand in which conditions these two different sources of depth information support the perception of an equivalent amount of depth.

1.1. Perceptual depth estimation from retinal images

It has long been recognized that disparity and velocity signals by themselves are insufficient to specify the distal depth map that has generated the two-dimensional (2D) image. For a local analysis of the visual field, in fact, the depth difference (z) with respect to

the fixation point is related to the disparity (d) and velocity (v) signals through parameters which are not specified by optical information (see Fig. 1). For small visual angles, the equations become:

$$d \approx \text{IOD} \frac{z}{z_f^2} + \varepsilon_d, \quad (1)$$

$$v \approx \left(\frac{T_x}{z_f} + \omega \right) \frac{z}{z_f} + \varepsilon_v, \quad (2)$$

where z is the depth-difference with respect to the fixation point, z_f is the fixation distance, IOD is the inter-ocular distance, T_x is the x -axis translation of the observer and ω is the rotation of the object. ε_d and ε_v specify additive Gaussian noise with zero mean and standard deviations σ_d and σ_v .

In order to compute the absolute amount of depth, the horizontal disparities must be scaled inversely with the square of the viewing distance, z_f^2 (by assuming that the IOD is known). Likewise, the velocity signals must be scaled so as to take into account the observer's translational component T_x , the object's rotation ω and the inverse of the viewing distance. To simplify the following discussion, we will denote with:

$$k_{d0} = \frac{\text{IOD}}{z_f^2}, \quad (3)$$

* Corresponding author. Brown University, Cognitive and Linguistic Sciences, 190 Thayer Street, Providence, RI 02912, USA. Tel.: +1 401 996 1863.

E-mail addresses: fulvio.domini@brown.edu (F. Domini), corrado.caudek@unifi.it (C. Caudek).

Fig. 1. Schematic illustration of the viewing geometry. z_f is the viewing distance; α_L and α_R are the angles formed by the visual lines connecting a flanking point to the left and right eyes and the two visual axes; Δz is the front-to-back depth of the stimulus configuration along the line of sight. ω represents the angular rotation about a vertical axis centered at the fixation point. The open circles represent the position of the flanking points after the rotation. α_1 and α_2 are the angles between the visual line connecting a flanking point to the cyclopean eye and the visual axis of the cyclopean eye before and after the rotation, respectively.

the scaling factor for veridical recovery of Euclidean depth from disparity information. Likewise, we will denote with:

$$k_{v_0} = \frac{1}{z_f} \left(\frac{T_x}{z_f} + \omega \right), \quad (4)$$

the scaling factor for veridical recovery of Euclidean depth from velocity information.

In summary, the disparity and velocity signals are related to the depth of the projected object through parameters not specified by optical information. Disparity signals are related to depth through the squared reciprocal fixation distance z_f ; the velocity signals are related to depth through the reciprocal of the fixation distance, the angle of rotation ω , and the translation velocity T_x . How can the “missing parameters” z_f , ω , and T_x be recovered?

1.2. Perceived depth from disparity signals

The parameter z_f may be specified by not-retinal information, such as the vergence angle of the eyes and the state of the accommodation (e.g., Proffitt & Caudek, 2002). Some investigations have provided evidence that observers can estimate fixation distance from vergence (Frisby, Buckley, & Duke, 1996; Tresilian & Mon-Williams, 2000; Tresilian, Mon-Williams, & Kelly, 1999) and accommodation (Fisher & Ciuffreda, 1988). In general, however, extra-retinal information does not guarantee a veridical recovery of viewing distance. The vast majority of experiments on the perception of depth from binocular disparity, in fact, has shown systematic distortions of depth-from-stereo. In the context of the inverse-geometry models of depth perception, these distortions of depth have been attributed to a mis-estimation of the viewing distance (e.g., Johnston, 1991; for a discussion, see Todd & Norman, 2003).

Some of these experiments have been conducted in the laboratory with computer-generated displays (Bradshaw, Glennerster, &

Rogers, 1996; Brenner & Landy, 1999; Brenner & van Damme, 1999; Collett, Schwarz, & Sobel, 1991; Glennerster, Rogers, & Bradshaw, 1996, 1998; Johnston, 1991; Johnston, Cumming, & Landy, 1994; Norman & Todd, 1998; Todd, Oomes, Koenderink, & Kappers, 2001; Tittle, Todd, Perotti, & Norman, 1995), whereas other studies have been carried out by using real objects in fully illuminated natural environments (Baird & Biersdorf, 1967; Battro, Netto, & Rozestraten, 1976; Bradshaw, Parton, & Glennerster, 2000; Cuijpers, Kappers, & Koenderink, 2000a, Cuijpers, Kappers, & Koenderink, 2000b; Gilinsky, 1951; Harway, 1963; Koenderink, van Doorn, Kappers, & Todd, 2002; Koenderink, van Doorn, & Lappin, 2000; Loomis, Da Silva, Fujita, & Fukusima, 1992; Loomis & Philbeck, 1999; Norman, Crabtree, Clayton, & Norman, 2005; Norman, Lappin, & Norman, 2000; Norman, Todd, Perotti, & Tittle, 1996 – see their Experiment 4). In most of the cases, however, the psychophysical literature suggests that human observers do not estimate the viewing distance correctly.

1.3. Perceived depth from velocity signals

The recovery of depth from velocity information, besides being undermined by the indeterminacy of z_f , also suffers from the indeterminacy of the parameters ω and T_x . A veridical description of the Euclidean structure of an object can be derived from second-order optic flow, if appropriate assumptions are introduced in the interpretation process (Longuet-Higgins & Prazdny, 1980; Ullman, 1979). It has been shown, however, that human observers have a very limited sensitivity for the second-order temporal properties of the optic flow and thus rely mainly on velocities to recover 3D information (Hogervorst & Eagle, 2000; Todd & Bressan, 1990). If only the first-order information is used, however, it is not possible, in principle, to derive a veridical estimate of depth.

In our own research, we have showed that observers recover metric information from the optic flow in a heuristic and patchway fashion. We provided evidence that this is achieved by a probabilistic process which assigns the most likely 3D interpretation to (ambiguous) local first-order properties of the optic flow. As a consequence, the perceptual interpretation of velocity information is, in general, neither veridical nor internally consistent (Caudek & Domini, 1998; Caudek & Proffitt, 1993; Caudek & Rubin, 2001; Di Luca, Domini, & Caudek, 2004, 2007; Domini & Braunstein, 1998; Domini & Caudek, 1999, 2003a, 2003b; Domini, Caudek, & Proffitt, 1997; Domini, Caudek, & Richman, 1998; Domini, Caudek, & Skirko, 2003; Domini, Caudek, & Tassinari, 2006; Domini, Caudek, Turner, & Favretto, 1998; Domini, Vuong, & Caudek, 2002; Tassinari, Domini, & Caudek, 2008).

2. Perceptual depth-matching of single-cue stimuli

If the visual system were able to veridically estimate the “missing parameters” z_f , ω and T_x , then two single-cue displays would elicit the perception of the same depth extent when they correspond to the projection of the same distal depth extent z . The literature on the perceptual interpretation of disparity and velocity signals, however, reveals systematic failures of shape constancy (see the previous section). As a consequence, a disparity-only stimulus can be perceived as deeper or shallower than a velocity-only stimulus, even if the two stimuli are generated by the projection of the same distal depth extent, $z_d = z_v$. Likewise, a disparity-only stimulus can be perceived as having the same depth extent of a velocity-only stimulus, even if the two stimuli are generated by the projections of two different depth extents, $z_d \neq z_v$.

The goal of the present investigation is to explain these “depth mis-matches”. Two different hypotheses will be contrasted.

2.1. Hypothesis 1: Perceived depth is recovered by estimating the “missing” scaling constants

According to the first hypothesis, perceived depth is estimated by reversing the following equations:

$$d = k_{d_0} z + \varepsilon_d, \quad (5)$$

$$v = k_{v_0} z + \varepsilon_v, \quad (6)$$

where k_{d_0} and k_{v_0} are defined as indicated by Eqs. (3) and (4). Mis-perceptions of 3D shape are attributed to the mis-estimation of the scaling factors k_{d_0} and k_{v_0} . For example, stereo-processing may fail to appropriately scale horizontal disparities because of the noise in the measurement of eye positions, or because of a conflict between the focus cues and the vergence angle, or because vertical disparities are not sufficiently reliable. Similarly, motion-processing may not accurately measure the head translation T_x or, in the case of a stationary observer, may not accurately estimate the amount ω of object rotation.

The limitation of this account is that we cannot determine *a priori* whether two single-cue stimuli will be perceived to have the same depth or not. Before running the experiment, in fact, we have no way of establishing whether the estimates of the scaling factors k_{d_0} and k_{v_0} are accurate or biased.

2.2. Hypothesis 2: Perceived depth depends on the magnitude of the image signal relative to the uncertainty

Depth perception from disparity or velocity signals does not necessarily require the estimation of the scaling factors k_{d_0} and k_{v_0} as defined by Eqs. (3) and (4), respectively. Recently, we have advanced a different proposal, according to which the apparent depth of a stimulus is determined by the magnitude of motion and/or disparity differences in a display relative to the uncertainty (i.e., internal noise) arising from the measurement of those signals (Domini et al., 2006; Tassinari et al., 2008). Our proposal is formulated in terms of a normative model of depth-cue integration (the *intrinsic constraint* model), but it can also be applied to depth recovery from single cues and to depth-matching tasks. According to the *intrinsic constraint* model, depth estimates are recovered through a weighted combination of image signals. This combination is optimal in the sense that it maximizes the signal-to-noise ratio (SNR) of the combined estimate. A thorough discussion of the *intrinsic constraint* model is provided by Domini and Caudek (2009). What is important for the present discussion is that, according to the *intrinsic constraint* model, two stimuli elicit the same amount of perceived depth when they exhibit the same ratio between signal intensity and discrimination threshold. According to such proposal, therefore, a disparity-only stimulus will be perceived with the same depth as a velocity-only stimulus when they both provide the same signal-to-noise ratio. Note that, in such “retinal model”, the properties of the distal scene are not considered. In the following section, we will illustrate how such proposal can be applied to the stimulus conditions of the present investigation.

2.2.1. Relation between discrimination thresholds and perceived depth magnitudes

The disparity and velocity signals are proportional to the relative depth z and, thus, directly specify the *affine structure* of the distal 3D shape (see Eqs. (5) and (6)). We hypothesize that a perceptual depth match occurs whenever two image signals specify the same affine structure with the same degree of precision, that is, whenever two image signals have the same SNR:

$$\frac{\mathcal{E}(d)}{\sigma_d} = \frac{\mathcal{E}(v)}{\sigma_v}. \quad (7)$$

This hypothesis allows us to predict the disparity and velocity values of a perceptual depth match. In the present experiment, a staircase procedure was used to match the perceived depth of a varying disparity-only stimulus to the perceived depth of a fixed motion-only stimulus. According to Eq. (7), a depth match requires the expected value of the disparity signal to be

$$\mathcal{E}(d) = \frac{\sigma_d}{\sigma_v} v_0, \quad (8)$$

where v_0 is the front-to-back relative-velocity of the velocity-only stimulus.

An estimate of $\mathcal{E}(d)$ is provided by the point of subjective equality (PSE) of the psychometric function estimated from the observers' judgments. Such point of subjective equality (termed PSE_d) was obtained in the first part of the experiment, where the disparity-only stimuli were varied within a staircase procedure and the velocity-only stimuli were kept fixed at the value corresponding to the signal intensity v_0 .

In the second and third part of the experiment, we measured the discrimination thresholds JND_d and JND_v (expressed in terms of the signal's intensities) at the pedestal values PSE_d and v_0 . Such discrimination thresholds are interpreted as estimates of σ_d and σ_v , respectively. We can therefore re-write Eq. (8) as

$$PSE_d = \frac{JND_d}{JND_v} v_0. \quad (9)$$

Eq. (9) can be re-written in terms of the depth of the distal 3D shapes used to generate the stimulus displays. According to Eqs. (5) and (6), $PSE_d = k_{d_0} PSE_{z_d}$, $v_0 = k_{v_0} z_v$, $JND_d = k_{d_0} JND_{z_d}$, and $JND_v = k_{v_0} JND_{z_v}$. By substituting these values in Eq. (9), after simplifying for k_{d_0} and k_{v_0} , we obtain:

$$PSE_{z_d} = \frac{JND_{z_d}}{JND_{z_v}} z_v. \quad (10)$$

In conclusion, the systematic errors made by observers in matching the perceived depth of two single-cue stimuli can be predicted by Eq. (10) in terms of the uncertainty (i.e., the JND) of the perceptual responses. According to Hypothesis 1, conversely, the biases in the perceptual matching tasks cannot be predicted, since there is no way to determine by how much the scaling constants k_{d_0} and k_{v_0} are mis-estimated.

3. Experiment

The experiment had a twofold purpose: (1) to determine the stimulus conditions in which a disparity-only stimulus is perceived to be matched in depth to a velocity-only stimulus, and (2) to measure the discrimination thresholds of the depth-matched stimuli.

3.1. Apparatus

Stereoscopic stimuli were displayed on a haploscope consisting of two CRT monitors (0.22 mm dot pitch) located on swing arms pivoting directly beneath the observer's eyes. Anti-aliasing and spatial calibrating procedures allowed spatial precision of dot location greater than hyper-acuity levels. Each monitor was seen in a mirror by one eye. Head position was fixed with a chin-and-forehead locating apparatus. The distance from each eye to the corresponding monitor was 95 cm. The eyes' vergence was directly manipulated by physically moving the monitors on their swing arms. Since the monitors and mirrors pivot rigidly about the eye's axis of rotation, the retinal images always remain the same for all

Fig. 2. Schematic illustration of the stereo viewing apparatus (haploscope). CRT monitors project through mirrors to each eye individually and are moved about pivots which coincide with the axis of rotation of each eye. The virtual fixation point is specified by vergence angle, which was adjusted properly for each observer's IOD and viewing distance.

positions of the two CRT monitors (see Fig. 2). Thus, the changes in eye position were dissociated from the changes in retinal images.

3.2. Stimuli

The stimuli were 800 high-luminance anti-aliased dots displayed against a low luminance background. 400 dots were randomly placed on three vertical lines 50 mm long (see Fig. 4). The other points were randomly positioned within a volume 50 mm wide, 50 mm high and 25 mm deep.

The actual physical distance to the CRT monitors (suggested by accommodation cues) was always fixed at 95 cm from the observer. In the experiment, we manipulated the eyes' vergence so as to specify two possible viewing distances: 50 and 100 cm.

In different trials, the center line and the two flanking lines were spaced in depth by either 2.5 or 5.0 mm. One of the three vertical lines was positioned at the center of the stimulus display. The other two vertical lines were positioned at 12.5 mm to the left and to the right of the central line. The overall stimulus subtended about 2.86° of visual angle when viewed at 0.5 m; when viewed at 1.0 m, the visual angle was 1.43°.

Depth information was provided by either disparity or velocity cues. Stimuli were drawn in polar projection. Disparities were calculated so as to simulate a 3D structure viewed at 50 or 100 cm from the observer. The vergence angle was computed for each observer, by taking into account her or his inter-ocular distance. For both stereo and motion stimuli, the simulated depth position of the two flanking lines was at fixation.

In the motion-only condition, observers viewed the stimuli with both eyes. This allowed us to equate the vergence information provided in both motion-only and disparity-only conditions. The same image was provided to both eyes, thus producing a nil disparity field. In our previous research, we found that a nil disparity field does not flatten depth from motion for small simulated depth magnitudes.

The 2D motion of the dots in the display was computed by simulating a rotation of the simulated 3D structure about a horizontal axis positioned at fixation. The 3D structure rotated back and forth by 14°. The duration of an entire rotation cycle was 2 s. The rotation increment was 0.23° per frame. The update rate was 60 frames per second.

3.3. Procedure

Observers were asked to determine which of two successively presented stimuli evoked a larger depth separation. The experiment comprised three parts.

1. A constant velocity-only stimulus was shown in one of the two intervals; the other interval showed a disparity-only stimulus which was varied according to a staircase procedure so as to find the PSE.
2. A depth-discrimination threshold was found for the disparity-only stimulus at the pedestal value determined in the first part of the experiment.
3. A depth-discrimination threshold was found for the velocity-only stimulus.

Observers judged the depth separation between the two flanking lines and the central line (see Fig. 4). A two-interval forced-choice task was used in which an observer was two successively presented stimuli and was asked to indicate which stimulus was perceived to be "deeper". No feedback was provided. The stimuli were displayed until the observers' response with a 0.5-s inter-stimulus interval.

It is important to note that the motion stimuli were seen binocularly and, therefore, they could have provided stereo information specifying a flat surface. We chose binocular viewing for the motion stimuli so as to have the same extra-retinal signals for both disparity-only and velocity-only stimuli. According to our hypothesis, in fact, both retinal velocities and binocular disparities are modulated in the same fashion by the extra-retinal signals (Domini & Caudek, 2009; Domini et al., 2006; Tassinari et al., 2008). If the motion stimuli had been seen monocularly, then no vergence angle information would have been provided to scale the optic flow information.

In a pilot study, we checked whether, in the specific stimulus conditions of the present experiment, binocular vision causes a flattening of the 3D structure perceived from the velocity-only stimuli. For the present stimuli, we found no difference in perceived depth from motion, when the stimuli were viewed binocularly or monocularly. In the present experiment, in fact, the fixation distance is either 0.5 or 1.0 m, and the visual angle subtended by the stimulus is very small. In these conditions, the horizontal disparities are negligible and, according to our model, they will not affect the perceptual solution recovered from the velocity signals (e.g., Domini et al., 2006). Most importantly, even if binocular vision of the velocity-only stimuli did produce a decrease of the perceived depth extent, such effect could not explain the results obtained in the depth-matching task.

The stimulus shown in one of the two intervals simulated a constant depth separation (*comparison stimulus*), whereas the depth separation of the other stimulus (*test stimulus*) was varied.

On each trial, the observer indicated which of the two stimuli – one with a fixed depth D and the other at the depth $D \pm \Delta d$ – appeared to be "deeper". We used staircases to control the value of Δd and four reversal rules – 3-down/1-up, 1-down/3-up, 2-down/1-up, and 1-down/2-up – to sample points along the entire psychometric function. Four staircases were employed for each psychometric function, with approximately 350–450 trials per psychometric function. Each staircase was terminated after 6 reversals. The staircase algorithm was the same as that used by Hillis, Watt, Landy, and Banks (2004).

In a first part of the experiment, the *comparison* was a motion stimulus and the *test* was a stereo stimulus which was varied according to a staircase procedure. The purpose was to determine the PSE at which the *test* stereo stimulus appears to be perceptually matched to the *comparison* motion stimulus. Two simulated depth magnitudes were used for the *comparison* stimulus, and each was viewed at two different viewing distances (50 and 100 cm). For each observer, the PSE was estimated from the fitted psychometric function. In a second part of the experiment, the JNDs for the disparity-only stimuli were estimated by means of a depth-discrimination task at the pedestal values determined in the first part

Fig. 3. Schematic representation of the experimental design. The first row represents the first part of the experiment: depth-matching task. The last two rows represent the second and third parts of the experiment: depth-discrimination task for the disparity-only (second part) and the velocity-only stimuli (third part). The two “v” lines represent a bird-view of the imaginary lines connecting the three vertical bars; each of them represents the stimulus shown in one of the two intervals of each trial. The double arrow near to the token figure indicates which of the two stimuli was varied within the staircase procedure. The third column specifies which quantity was estimated from the psychometric function.

of the experiment. In a third part of the experiment, the JNDs for the velocity-only stimuli were estimated by means of a depth-discrimination task at the simulated values of 2.5 and 5.0 mm used in the first part of the experiment (see Fig. 3).

3.4. Participants

A total of three undergraduate and three graduate students from Brown University participated in the experiment. They all had normal or corrected-to-normal vision. All participants were naïve to the purpose of the study, and three of them were experienced psychophysical observers. Participation was voluntary; all participants provided written consent and the undergraduate students were paid for their participation.

3.5. Results and discussion

For each of the three parts of the experiment, the data of each observer in each experimental condition were fitted with a psycho-

Fig. 5. Results of observer R.F. in the four experimental conditions of part 1 of the experiment. Psychometric functions were fitted using psignifit version 2.5.6 (see <http://bootstrap-software.org/psignifit/>), a software package which implements the maximum-likelihood method described by Wichmann and Hill (2001a). The horizontal bars represent 95% confidence intervals for the PSEs and were found by the BC_a bootstrap method implemented by psignifit, based on 1000 simulations (see Wichmann and Hill, 2001b). The points in the figure represent the binned proportion of times the observer indicated the stereo-only stimulus was deeper than the motion-only stimulus (the psychometric functions were estimated from the raw data, not the binned data). Blue: simulated depth of 2.5 mm at a viewing distance of 0.5 m; red: simulated depth of 2.5 mm at a viewing distance of 1.0 m; green: simulated depth of 5.0 mm at a viewing distance of 0.5 m; black: simulated depth of 5.0 mm at a viewing distance of 1.0 m.

metric function. Fig. 5 shows the data of one representative observer for the first part of the experiment.

The means of the fits are points of subjective equality, the values of the velocity-only stimuli that on average had the same apparent depth as the stereo stimuli. If perceived depth were veridical, the PSEs of the psychometric functions would be the following: 123.759 arcsec for the blue function; 30.940 arcsec for the red function; 247.518 arcsec for the green function; 61.880 arcsec for the black function. For this particular observer, judgments are close to veridical at the viewing distance of 1.0 m; at the viewing distance of 0.5 m, instead, the responses of R.F. over-estimated depth from disparity.

Fig. 4. Stereogram representing the stimulus display.

Fig. 6. *Left panel.* Mean depth (in millimeters) of the disparity-only stimuli as a function of the depth (in millimeters) of the velocity-only stimuli, when target and comparison displays appeared to be perceptually matched in depth. Circles (red) and triangles (blue) denote the viewing distances of 0.5 and 1.0 m, respectively. Vertical bars represent one standard error of the mean. Symbol size represents the magnitude of simulated depth: small for 2.5 mm; large for 5.0 mm. *Right panel.* The same data of the left panel have been replotted by recoding the values on the horizontal axis according to Eq. (10).

This pattern of results is confirmed by Fig. 6, which shows the average results of six observers expressed in terms of depth magnitudes (not in terms of signal intensities, as in Fig. 5). In the left panel of Fig. 6, the average depth magnitudes of the disparity-only stimuli at the PSE are shown as a function of the simulated depth of the velocity-only stimuli. The viewing distances of 0.5 and 1.0 m are as coded red circles and blue triangles, respectively. Note that, if the depth matches were veridical, all points should lay on a straight line with zero intercept and unitary slope (the gray line in the Fig. 6).

The effect of viewing distance was confirmed by a linear mixed-effects (LMEs) model¹ applied on the PSEs with two factors as independent variables: *simulated depth* (2.5, 5.0 mm) and *viewing distance* (0.5, 1.0 m). The interaction between simulated depth and viewing distance was not significant ($\chi^2_1 = 2.488$, n.s.). A no-interaction model revealed significant effects for both simulated depth ($B = 0.611$, SE $B = 0.105$, MCMC $p = 0.0002$) and viewing distance ($B = 0.003$, SE $B = 0.0005$, MCMC $p = 0.0001$).

Separate analyses were then performed at each of the two viewing distances. For the viewing distance of 1.0 m, we found that the depth of the disparity-only stimuli at the PSE did not differ from the simulated depth of the velocity-only stimuli ($B = -0.688$, SE $B = 0.520$, MCMC $p = \text{n.s.}$). At the viewing distance of 0.5 m, the depth of the disparity-only stimuli at the PSE was significantly lower than the simulated depth of the velocity-only stimuli ($B = -1.968$, SE $B = 0.383$, MCMC $p = 0.0002$). In summary, the present data suggest a systematic bias: a perceptual depth match at the viewing distance of 0.5 m requires about half of the simulated depth for a disparity-only stimulus than for a velocity-only stimulus.

Having established that, within the present stimulus conditions, the perceptual depth matches are biased, the purpose of the present investigation was to determine whether such biases can be accounted for by the hypothesis described in Section 2.2. According

Fig. 7. Depth (in millimeters) of disparity-only stimuli as a function of the depth (in millimeters) of the velocity-only stimuli, when the two stimuli are perceived as perceptually matched in depth. Circles and triangles denote the viewing distances of 0.5 and 1.0 m, respectively. Symbol size represents the magnitude of simulated depth: small symbols for 2.5 mm; large symbols for 5.0 mm. Different colors indicate different observers.

to Hypothesis 2, a perceptual depth match requires the same signal-to-noise ratio for the two single-cue stimuli, regardless of their simulated depth magnitudes. At the PSE the depth magnitudes of the two stimuli should be related to each other as indicated by Eq. (10). The mean PSE of the disparity-only stimuli as a function of the predictor of Eq. (10) is shown in the right panel of Fig. 6. The data of the individual observers are shown in Figs. 7 and 8². Hypothesis 2 can be tested by estimating the 95% confidence interval for the slope of this empirical regression line and to gauge it against the model's predictions.

² One outlier data point was deleted from Fig. 8. The data analysis was performed on the entire data set.

¹ Inferential statistics are based on a linear mixed-effects (LMEs) model specifying participants as random effects (Pinheiro & Bates, 2000; Quené & Van den Bergh, 2004, for simulations). We used the *lmer* program (*lme4* package; Bates & Sarkar, 2007) in the R system for statistical computing (R Development Core Team, 2009). As indicated by Baayen (2008), p values and confidence intervals are generated from the posterior distribution of parameter estimates with Markov Chain Monte Carlo methods, using the *mcmc* program in the *lme4* package with default specifications (e.g., $n = 1,000$ samples; locally uniform priors for fixed effects; locally non-informative priors for random effects).

Fig. 8. The same data of Fig. 7. The values on the horizontal axis have been recoded according to Eq. (10). Circles and triangles denote the viewing distances of 0.5 and 1.0 m, respectively. Symbol size represents the magnitude of simulated depth: small symbols for 2.5 mm; large symbols for 5.0 mm. Different colors indicate different observers.

Consistently with the predictions of **Hypothesis 2**, an LME analysis found a slope equal to 1.095 (95% HPD CI: 0.911, 1.282, MCMC $p = 0.0001$). Since this CI includes the value of 1.0, we can conclude that the data are consistent with the model's prediction. This test does not prove that **Hypothesis 2** is correct, but rather it provides a measure of its precision: the empirical data are not too far from the expected ones. The small discrepancy between the predictions of **Hypothesis 2** and the empirical data (not statistically significant) should be contrasted with the limitations of **Hypothesis 1**. Since **Hypothesis 1** cannot determine *a priori* the estimation biases for the scaling constants k_{d_0} and k_{v_0} (see Section 2.1), it has no way to predict whether PSE_{z_d} will overestimate or underestimate z_v .

4. General discussion

In the present investigation, observers were asked to discriminate the amounts of depth perceived from disparity-only and from velocity-only stimuli. Given the psychophysical evidence discussed in Sections 1.2 and 1.3, it would be naïve to expect that, when they are both generated by the same distal depth extent, observers judge a velocity-only stimulus to be as deep as a disparity-only stimulus. In general, perceived depth from single-cue stimuli is distorted. The purpose of the present investigation was to understand in which conditions the couplings of *different* distal depth extents support the perception of a depth match.

According to the first hypothesis presented in Section 2, a disparity-only display may appear to be matched in depth to a velocity-only display simulating a different depth magnitude because the scaling constants k_{d_0} and k_{v_0} are mis-estimated (see Eqs. (3) and (4)). Since we do not know the amount of departure of the k_{d_0} and k_{v_0} estimates from their true values, we cannot predict *a priori* when disparity-only and velocity-only stimuli will be perceived as having the same depth.

According to the second hypothesis, two single-cue stimuli are perceived to be equally deep when they both provide the same SNR. This hypothesis stems from the idea that the amount of perceived depth depends on the *retinal* SNR. Consistent with this hypothesis are, for example, the findings of Todd (1985) who stud-

ied structure-from-motion by manipulating the amount of visual noise in the displays. In his study, observers were required to estimate the apparent slant of a random-dot planar surface rotating in depth, on which a proportion of the dots were repositioned randomly on each frame. The results revealed that the apparent slants of the surfaces decreased systematically as the proportion of the displaced dots was increased. If external noise can reduce the magnitude of apparent depth or slant, then it seems reasonable to suppose that internal noise may have the same effect.

More specifically, we proposed that a perceptual depth match is affected by the uncertainty deriving from the measurement of the image signals, as indicated by Eq. (10). It follows that, in general, two single-cue stimuli should be perceived as matched in depth when they are the projection of *specific* different distal depth extents. The results of the present investigation are consistent with this hypothesis.

At the viewing distance of 0.5 m, a perceptual match was obtained when the distal 3D shape of which the disparity-only stimulus is the projection was 49% shallower than the distal shape for the velocity-only stimulus (see Fig. 6). At the viewing distance of 1.0 m, the distal depth for the disparity-only stimulus was 16% shallower than the distal depth of the velocity-only stimulus (a not significant difference). In other words, a perceptual depth match was obtained, at the viewing distance of 0.5 m, when the two stimuli were generated by the projection of two very different distal depths magnitudes; at the viewing distance of 1.0 m, conversely, the distal depths for the two stimuli were quite similar.

As indicated in the right panel of Fig. 6, these data are consistent with **Hypothesis 2**. In all cases, in fact, the two stimuli had to provide the same SNR in order to appear to be matched in depth. Note that the predictions of Eq. (10) hold both when the perceptual matches are “veridical” and when they are not.

Alternative explanations are also important to consider. It may be argued, for example, that observers can interpret correctly disparity information at the “effective viewing distance” of 100 cm, but they over-estimate depth from disparity at the distance of 50 cm (see Johnston, 1991). Disparity information not coupled with a vergence angle specifying the abathic distance, together with a specific mis-estimation of the angle of rotation, thus, could be held responsible for the results of the present experiment.

This alternative explanation, however, misses an important aspect of the present phenomenon. In fact, even if depth from disparity were always over-estimated at small viewing distances and under-estimated at large viewing distances, what about depth from velocity? From our previous research on the perceptual interpretation of the optic flow, we know that apparent depth from motion depends not only on the distal depth magnitude, but also on the amount of angular rotation used to generate the velocity field (for a review, see Domini & Caudek, 2003a). Any explanation of the present results *only* in terms of the role of the abathic distance in disparity processing, therefore, is incomplete.

Another consideration can be made about the fact that our data set is too small to convincingly show that the slope of the relationship shown in Fig. 6 is 1.0. This point can be addressed by applying Eq. (10) to a larger data set provided by the investigation of Domini and Caudek (2009). In that paper, we investigated the relation between two Fechnerian depth scales. One psychophysical depth scale for velocity-only stimuli, the other for disparity-only stimuli. In the third part of that experiment, observers were asked to perceptually match a disparity-only stimulus with a velocity-only stimulus. In a similar manner as in the present experiment, observers judged which of two successively presented stimuli evoked a larger depth separation. The simulated depth separation of the velocity-only stimulus was kept constant, whereas the depth separation of the disparity-only stimulus was varied according to a staircase procedure. The PSE for the disparity-only stimulus was found, together with the JNDs of both stimuli at the PSE.

Fig. 9. *Left panel.* Depth (in millimeters) of disparity-only stimuli as a function of the depth (in millimeters) of the velocity-only stimuli, when the two stimuli are perceived as perceptually matched in depth. The data have been replotted from Domini and Caudek (2009). Vertical bars represent one standard error. *Right panel.* The same data of the left panel. The values on the horizontal axis have been recoded according to Eq. (10).

In Fig. 9 we have replotted the data of Domini and Caudek (2009) in the format of Fig. 6. Each point in the plot represents the distal depth magnitudes used to generate a disparity-only and velocity-only stimuli at the PSE. Note that systematic failures of shape constancy occur also in the experiment of Domini and Caudek (2009). The left panel of Fig. 9 shows that the amount of distal depth for the disparity-only stimuli is consistently smaller than the distal depth for the velocity-only stimuli.

At the PSE, the distal depth magnitudes of the disparity-only stimuli are approximately linearly related to the distal depth magnitudes of the velocity-only stimuli (Fig. 9, left panel). The slope of this linear relation, however, is different from one (gray line, with a nil intercept) and represents the systematic depth misperceptions in depth-matched stimuli. In the right panel of Fig. 9, the PSEs of the disparity-only stimuli are shown as a function of the simulated depths of the velocity-only stimuli weighted by the ratio between the JNDs of the two stimuli. The gray line in the right panel of Fig. 9 has nil intercept and slope of 1.0. It is easy to see, therefore, that Eq. (10) provides an accurate account also for the data of Domini and Caudek (2009). Note an important difference between the stimulus conditions of Domini and Caudek (2009) and those of the present investigation. Differently from the present experiment, in fact, Domini and Caudek (2009) always kept the viewing distance fixed at 1.0 m. The depth distortions found in that investigation, therefore, cannot be attributed to a mis-estimation of the viewing distance.

Another interesting result of the present investigation is the effect of the vergence angle on the perceptual interpretation of velocity information. At the viewing distance of 0.5 m, the simulated depth of 2.5 mm generated a maximum displacement of 50.4 arcsec; at the PSE, the matched disparity-only display exhibited a maximum disparity of 66.7 arcsec. At the viewing distance of 1.0 m, the simulated depth of 5.0 mm generated a maximum velocity of 50.4 arcsec; at the PSE, the matched disparity-only display exhibited a maximum disparity of 53.8 arcsec. The observers' responses in these two conditions were not statistically different ($t_5 = 1.255$, n.s.). This means that the same retinal displacements (50.4 arcsec), when coupled with two different fixation distances (0.5 and 1.0 m), were perceptually matched with (almost) the same disparity (60.25 arcsec, on average) viewed at two different distances. Even though there may be a real difference between these two conditions, which was not detected by the statistical power of the present analysis, this result suggests a potential interesting issue. We know that the same disparity signal coupled with a larger viewing distance produces an increase in the amount of perceived depth (Johnston, 1991). The present data, therefore, are compatible with the hypothesis that the vergence angle may affect the percep-

tual interpretation of velocity information: perceived depth may increase when the same velocity field is associated with a larger fixation distance.³ Even though it is consistent with Eq. (2), the possible effect of the vergence angle on the perceptual interpretation of the velocity field has not been previously addressed in the structure-from-motion literature.

A final point concerns the conflict between vergence and accommodation. Even though the conflict between these two depth-cues may hinder "veridical perception", it is not a problem for the present investigation. In fact, it affects the visual processing of disparity and velocity information in the same manner.

5. Conclusion

Two single-cue stimuli are not necessarily perceived to be matched in depth when they are generated by the projection of the same distal depth magnitude. In the present investigation, we showed that a disparity-only stimulus elicits the same amount of depth as a velocity-only stimulus when they both provide the same signal-to-noise ratio, regardless of the distal depth magnitudes that have generated the two images. Such empirical results are consistent with the hypothesis that the apparent depth of a stimulus is determined by the magnitude of the retinal signals relative to the uncertainty (i.e., internal noise) arising from the measurement of those signals (Domini & Caudek, 2009; Domini et al., 2006).

References

- Baird, J. C., & Biersdorf, W. R. (1967). Quantitative functions for size and distance judgments. *Perception & Psychophysics*, 2, 161–166.
- Bates, D. M., & Sarkar, D. (2007). *lme4: Linear mixed-effects models using Eigen and R package version 0.99875-6*.
- Battro, A. M., Netto, S. P., & Rozestraten, R. J. A. (1976). Riemannian geometries of variable curvature in visual space: Visual alleys, horopters, and triangles in big open fields. *Perception*, 5, 9–23.
- Baayen, R. H. (2008). *Analyzing linguistic data: A practical introduction to statistics using R*. Cambridge: Cambridge University Press.

³ For the other two points of Fig. 6 the 2D signals are the following. For the smallest simulated depth of the motion displays at the distance of 1.0 m, the displacement signal is 25 arcsec with a matched disparity of 28 arcsec; for the largest simulated depth of the motion displays at the distance of 0.5 m, the displacement signal is 100.8 arcsec with a matched disparity of 128 arcsec. In this case it is difficult to interpret the effect of the vergence angle on the processing of velocity information because a "small" disparity paired with a "large" fixation distance must be compared with a "large" disparity paired with a small fixation distance. In these circumstances, any prediction requires the knowledge of how the vergence angle scales disparity information. Such a function would be known if the fixation distance were estimated correctly by the visual system. But this is not so.

- Bradshaw, M. F., Glennerster, A., & Rogers, B. J. (1996). The effect of display size on disparity scaling from differential perspective and vergence cues. *Vision Research*, 36, 1255–1264.
- Bradshaw, M. F., Parton, A. D., & Glennerster, A. (2000). The task-dependent use of binocular disparity and motion parallax information. *Vision Research*, 40, 3725–3734.
- Brenner, E., & Landy, M. S. (1999). Interaction between the perceived shape of two objects. *Vision Research*, 39, 3834–3848.
- Brenner, E., & van Damme, W. J. (1999). Perceived distance, shape and size. *Vision Research*, 39, 975–986.
- Caudek, C., & Domini, F. (1998). Perceived orientation of axis of rotation in structure-from-motion. *Journal of Experimental Psychology: Human Perception and Performance*, 24, 609–621.
- Caudek, C., & Proffitt, D. R. (1993). Depth perception in motion parallax and stereokinesis. *Journal of Experimental Psychology: Human Perception and Performance*, 19, 32–47.
- Caudek, C., & Rubin, N. (2001). Segmentation in structure from motion: Modeling and psychophysics. *Vision Research*, 41, 2715–2732.
- Collett, T. S., Schwarz, U., & Sobel, E. C. (1991). The interaction of oculomotor cues and stimulus size in stereoscopic depth constancy. *Perception*, 20, 733–754.
- Cuijpers, R. H., Kappers, A. M. L., & Koenderink, J. J. (2000a). Investigation of visual space using an exocentric pointing task. *Perception & Psychophysics*, 62, 1556–1571.
- Cuijpers, R. H., Kappers, A. M. L., & Koenderink, J. J. (2000b). Large systematic deviations in visual parallelism. *Perception*, 29, 1467–1482.
- Di Luca, M., Domini, F., & Caudek, C. (2004). Spatial integration in structure from motion. *Vision Research*, 44, 3001–3013.
- Di Luca, M., Domini, F., & Caudek, C. (2007). The relation between disparity and velocity signals of rigidly moving objects constrains depth order perception. *Vision Research*, 25, 1335–1349.
- Domini, F., & Braunstein, M. L. (1998). Recovery of 3-D structure from motion is neither Euclidean nor affine. *Journal of Experimental Psychology: Human Perception and Performance*, 24, 1273–1295.
- Domini, F., & Caudek, C. (1999). Perceiving surface slant from deformation of optic flow. *Journal of Experimental Psychology: Human Perception and Performance*, 25, 426–444.
- Domini, F., & Caudek, C. (2003a). 3D structure perceived from dynamic information: A new theory. *Trends in Cognitive Sciences*, 7, 444–449.
- Domini, F., & Caudek, C. (2003b). Recovering slant and angular velocity from a linear velocity field: Modeling and psychophysics. *Vision Research*, 43, 1753–1764.
- Domini, F., & Caudek, C. (2009). The intrinsic constraint model and Fechnerian sensory scaling. *Journal of Vision*, 9(2), pp. 25, 1–15.
- Domini, F., Caudek, C., & Proffitt, D. R. (1997). Misperceptions of angular velocities influence the perception of rigidity in the kinetic depth effect. *Journal of Experimental Psychology: Human Perception and Performance*, 23, 1111–1129.
- Domini, F., Caudek, C., & Richman, S. (1998). Distortions of depth-order relations and parallelism in structure from motion. *Perception & Psychophysics*, 60, 1164–1174.
- Domini, F., Caudek, C., & Skirko, P. (2003). Temporal integration of motion and stereo cues to depth. *Perception & Psychophysics*, 65, 48–57.
- Domini, F., Caudek, C., & Tassinari, H. (2006). Stereo and motion information are not independently processed by the visual system. *Vision Research*, 46, 1707–1723.
- Domini, F., Caudek, C., Turner, J., & Favretto, A. (1998). Discriminating constant from variable angular velocities in structure from motion. *Perception & Psychophysics*, 60, 747–760.
- Domini, F., Vuong, Q. C., & Caudek, C. (2002). Temporal integration in structure from motion. *Journal of Experimental Psychology: Human Perception and Performance*, 28, 816–838.
- Fisher, K. S., & Ciuffreda, K. J. (1988). Accommodation and apparent distance. *Perception*, 17, 609–621.
- Frisby, J. P., Buckley, D., & Duke, P. A. (1996). Evidence for good recovery of lengths of real objects seen with natural stereo viewing. *Perception*, 25, 129–154.
- Gilinsky, A. S. (1951). Perceived size and distance in visual space. *Psychological Review*, 58, 460–482.
- Glennerster, A., Rogers, B. J., & Bradshaw, M. F. (1996). Stereoscopic depth constancy depends on the subject's task. *Vision Research*, 36, 3441–3456.
- Glennerster, A., Rogers, B. J., & Bradshaw, M. F. (1998). Cues to viewing distance for stereoscopic depth constancy. *Perception*, 27, 1357–1365.
- Harway, N. I. (1963). Judgment of distance in children and adults. *Journal of Experimental Psychology*, 65, 385–390.
- Hillis, J. M., Watt, S. J., Landy, M. S., & Banks, M. S. (2004). Slant from texture and disparity cues: Optimal cue combination. *Journal of Vision*, 4(12), pp. 1, 967–992.
- Hogervorst, M. A., & Eagle, R. A. (2000). The role of perspective effects and accelerations in perceived three-dimensional structure-from-motion. *Journal of Experimental Psychology: Human Perception and Performance*, 26, 934–955.
- Johnston, E. B. (1991). Systematic distortions of shape from stereopsis. *Vision Research*, 31, 1351–1360.
- Johnston, E. B., Cumming, B. G., & Landy, M. S. (1994). Integration of motion and stereopsis cues. *Vision Research*, 34, 2259–2275.
- Koenderink, J. J., van Doorn, A. J., Kappers, A. M. L., & Todd, J. T. (2002). Pappus in optical space. *Perception & Psychophysics*, 64, 380–391.
- Koenderink, J. J., van Doorn, A. J., & Lappin, J. S. (2000). Direct measurement of the curvature of visual space. *Perception*, 29, 69–79.
- Longuet-Higgins, H. C., & Prazdny, K. (1980). The Interpretation of a Moving Retinal Image. *Proceedings of the Royal Society of London B*, 208, 385–397.
- Loomis, J. M., Da Silva, J. A., Fujita, N., & Fukusima, S. S. (1992). Visual space perception and visually directed action. *Journal of Experimental Psychology: Human Perception and Performance*, 18, 906–921.
- Loomis, J. M., & Philbeck, J. W. (1999). Is the anisotropy of perceived 3-D shape invariant across scale? *Perception & Psychophysics*, 61, 397–402.
- Norman, J. F., Crabtree, C. E., Clayton, A. M., & Norman, H. F. (2005). The perception of distances and spatial relationships in natural outdoor environments. *Perception*, 34, 1315–1324.
- Norman, J. F., Lappin, J. S., & Norman, H. F. (2000). The perception of length on curved and flat surfaces. *Perception & Psychophysics*, 62, 1133–1145.
- Norman, J. F., & Todd, J. T. (1998). Stereoscopic discrimination of interval and ordinal depth relations on smooth surfaces and in empty space. *Perception*, 27, 257–272.
- Norman, J. F., Todd, J. T., Perotti, V. J., & Tittle, J. S. (1996). The visual perception of three-dimensional length. *Journal of Experimental Psychology: Human Perception and Performance*, 22, 173–186.
- Pinheiro, J. C., & Bates, D. M. (2000). *Mixed-effects models in s and s-plus*. New York: Springer-Verlag.
- Proffitt, D. R., & Caudek, C. (2002). Depth perception and the perception of events. In A. F. Healy & R. W. Proctor (Eds.), *Comprehensive handbook of psychology. Experimental psychology* (Vol. 4, pp. 213–236). New York: Wiley.
- Quené, H., & Van den Bergh, H. (2004). On multi-level modeling of data from repeated measures designs: A tutorial. *Speech Communication*, 43, 103–121.
- R Development Core Team (2009). *R: A language and environment for statistical computing*. 3-900051-07-0. Vienna, Austria: R Foundation for Statistical Computing <<http://www.R-project.org>>.
- Tassinari, H., Domini, F., & Caudek, C. (2008). The intrinsic constraint model for stereo-motion integration. *Perception*, 37, 79–95.
- Tittle, J. S., Todd, J. T., Perotti, V. J., & Norman, J. F. (1995). Systematic distortion of perceived three-dimensional structure from motion and binocular stereopsis. *Journal of Experimental Psychology: Human Perception and Performance*, 21, 663–678.
- Todd, J. T. (1985). Perception of structure from motion: Is projective correspondence of moving elements a necessary condition? *Journal of Experimental Psychology: Human Perception and Performance*, 11, 689–710.
- Todd, J. T., & Bressan, P. (1990). The perception of 3-dimensional affine structure from minimal apparent motion sequences. *Perception & Psychophysics*, 48, 419–430.
- Todd, J. T., & Norman, J. F. (2003). The visual perception of 3-D shape from multiple cues: Are observers capable of perceiving metric structure? *Perception & Psychophysics*, 65, 31–47.
- Todd, J. T., Oomes, A. H. J., Koenderink, J. J., & Kappers, A. M. L. (2001). On the affine structure of perceptual space. *Psychological Science*, 12, 191–196.
- Tresilian, J. R., & Mon-Williams, M. (2000). Getting the measure of vergence weight in nearness perception. *Experimental Brain Research*, 132, 362–368.
- Tresilian, J. R., Mon-Williams, M., & Kelly, B. M. (1999). Increasing confidence in vergence as a distance cue. *Proceedings of the Royal Society*, B266, 39–44.
- Ullman, S. (1979). *The interpretation of visual motion*. Cambridge, MA: MIT Press.
- Wichmann, F. A., & Hill, N. J. (2001a). The psychometric function: I. Fitting, sampling and goodness-of-fit. *Perception & Psychophysics*, 63(8), 1293–1313.
- Wichmann, F. A., & Hill, N. J. (2001b). The psychometric function: II. Bootstrap-based confidence intervals and sampling. *Perception & Psychophysics*, 63(8), 1314–1329.