



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

On moduli of instanton bundles on P^{2n+1}

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

On moduli of instanton bundles on P^{2n+1} / Ancona, Vincenzo; Ottaviani, GIORGIO MARIA. - In: PACIFIC JOURNAL OF MATHEMATICS. - ISSN 0030-8730. - STAMPA. - 171:(1995), pp. 343-351.

Availability:

The webpage <https://hdl.handle.net/2158/392909> of the repository was last updated on

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

ON MODULI OF INSTANTON BUNDLES ON \mathbb{P}^{2n+1}

VINCENZO ANCONA AND G. OTTAVIANI

Let $MI_{\mathbb{P}^{2n+1}}(k)$ be the moduli space of stable instanton bundles on \mathbb{P}^{2n+1} with $c_2 = k$. We prove that $MI_{\mathbb{P}^{2n+1}}(2)$ is smooth, irreducible, unirational and has zero Euler-Poincaré characteristic, as it happens for \mathbb{P}^3 . We find instead that $MI_{\mathbb{P}^5}(3)$ and $MI_{\mathbb{P}^5}(4)$ are singular.

1. Definition and preliminaries.

Instanton bundles on a projective space $\mathbb{P}^{2n+1}(\mathbb{C})$ were introduced in [OS] and [ST]. In [AO] we studied their stability, proving in particular that special symplectic instanton bundles on \mathbb{P}^{2n+1} are stable, and that on \mathbb{P}^5 every instanton bundle is stable.

In this paper we study some moduli spaces $MI_{\mathbb{P}^{2n+1}}(k)$ of stable instanton bundles on \mathbb{P}^{2n+1} with $c_2 = k$. For $k = 2$ we prove that $MI_{\mathbb{P}^{2n+1}}(2)$ is *smooth, irreducible, unirational and has zero Euler-Poincaré characteristic* (Theor. 3.2), just as in the case of \mathbb{P}^3 [Har].

We find instead that $MI_{\mathbb{P}^5}(k)$ is *singular* for $k = 3, 4$ (theor. 3.3), which is not analogous with the case of \mathbb{P}^3 [ES], [P]. To be more precise, *all points corresponding to symplectic instanton bundles are singular*. Theor. 3.3 gives, to the best of our knowledge, the first example of a singular moduli space of stable bundles on a projective space. The proof of Theorem 3.3 needs help from a personal computer in order to calculate the dimensions of some cohomology group [BaS].

We recall from [OS], [ST] and [AO] the definition of instanton bundle on $\mathbb{P}^{2n+1}(\mathbb{C})$.

Definition 1.1. A vector bundle E of rank $2n$ on \mathbb{P}^{2n+1} is called an instanton bundle of quantum number k if

- (i) The Chern polynomial is $c_i(E) = (1 - t^2)^{-k} = 1 + kt^2 + \binom{k+1}{2}t^4 + \dots$
- (ii) $E(q)$ has natural cohomology in the range $-2n - 1 \leq q \leq 0$ (that is $h^i(E(q)) \neq 0$ for at most one $i = i(q)$)
- (iii) $E|_r \simeq \mathcal{O}_r^{2n}$ for a general line r .

Every instanton bundle is simple [AO]. There is the following characterization:

Proof. See [AO] Theorem 3.13 and Remark 2.22. □

Remark 1.9. If $E \simeq E^*$, then

$$H^2(E \otimes E^*) = H^2[(\text{Ker } A^t) \otimes (\text{Ker } A^t)] = H^2[(\text{Ker } B) \otimes (\text{Ker } B)].$$

Remark 1.10. The single complex associated with the double complex obtained by tensoring the two sequences

$$\begin{aligned} 0 \rightarrow \text{Ker } A^t \rightarrow \mathcal{O}^{2n+2k} \xrightarrow{A^t} \mathcal{O}(1)^k \rightarrow 0 \\ 0 \rightarrow \text{Ker } B^t \rightarrow \mathcal{O}^{2n+2k} \xrightarrow{B^t} \mathcal{O}(1)^k \rightarrow 0 \end{aligned}$$

gives the resolution

$$\begin{aligned} 0 \rightarrow (\text{Ker } A^t) \otimes (\text{Ker } B) \rightarrow \mathcal{O}^{2n+2k} \otimes \mathcal{O}^{2n+2k} \\ \rightarrow \mathcal{O}^{2n+2k} \otimes \mathcal{O}(1)^k \oplus \mathcal{O}(1)^k \otimes \mathcal{O}^{2n+2k} \xrightarrow{\alpha} \mathcal{O}(1)^k \otimes \mathcal{O}(1)^k \rightarrow 0 \end{aligned}$$

where $\alpha = (A^t \otimes \text{id}, \text{id} \otimes B)$.

Hence

$$H^2(E \otimes E^*) = \text{Coker } H^0(\alpha)$$

and its dimension can be computed using [BaS]. For the convenience of the reader we sketch the steps needed in the computations.

A, B^t are given by $k \times (2n + 2k)$ matrices whose entries are linear homogeneous polynomials.

$$A \otimes \text{Id}_k = (a_1, \dots, a_{k(2n+2k)})$$

and

$$\text{Id}_k \otimes B^t = (b_1, \dots, b_{k(2n+2k)})$$

are both $k^2 \times (2n + 2k)k$ matrices. Let

$$C = (a_1, \dots, a_{k(2n+2k)}, b_1, \dots, b_{k(2n+2k)}).$$

We will denote by $\text{syz}_m C$ the dimension of the space of the syzygies of C of degree m . Then

$$\begin{aligned} h^2(E \otimes E^*) &= h^0(\mathcal{O}(2)^{k^2}) - (4n + 4k)h^0(\mathcal{O}(1)^k) + \text{syz}_1 C \\ &= k(n + 1)[k(2n - 5) - 8n] + \text{syz}_1 C \\ h^1(E \otimes E^*) &= h^2(E \otimes E^*) + 1 - k^2 + 8n^2k - 4n^2 + 3nk^2 - 2n^2k^2 \\ &= 1 - 6k^2 - 8kn - 4n^2 + \text{syz}_1 C. \end{aligned}$$

Note also that $h^0(E(1)) = \text{syz}_1 B^t - k$ and $h^0(E^*(1)) = \text{syz}_1 A - k$.

Remark 1.11. In the same way we obtain

$$\begin{aligned} h^1(E \otimes E^*(-1)) &= \text{syz}_0 C \\ h^2(E \otimes E^*(-1)) &= 2k(nk - 2n - k) + \text{syz}_0 C. \end{aligned}$$

2. Example on \mathbb{P}^5 .

Let (a, b, c, d, e, f) be homogeneous coordinates in \mathbb{P}^5 .

Example 2.1. ($k = 3$) Let

$$\begin{aligned} B^t &= \begin{bmatrix} a & b & c & & d & e & f \\ & a & b & c & & d & e & f \\ & & a & b & c & & d & e & f \end{bmatrix} \\ A &= \begin{bmatrix} & f & e & d & & -c & -b & -a \\ & f & e & d & & -c & -b & -a \\ f & e & d & & -c & -b & -a \end{bmatrix}. \end{aligned}$$

The corresponding monad gives a special symplectic instanton bundle on \mathbb{P}^5 with $k = 3$. With the notation of remark 1.10, using [BaS] we can compute $\text{syz}_0 C = 14, \text{syz}_1 C = 174$. Hence $h^2(E \otimes E^*) = 3$ from the formulas of Remark 1.10. Moreover $h^0(E(1)) = 4$.

Example 2.2. ($k = 3$) Let B^t as in the Example 2.1 and

$$A = \begin{bmatrix} & f & e & d & & -c & -b & -a \\ e & d & & 2f & -b & -a & & -2c \\ d & & f & e & -a & & -c & -b \end{bmatrix}.$$

We have $\text{syz}_0 C = 10, \text{syz}_1 C = 171$. Hence $h^2(E \otimes E^*) = 0$. We can compute also the syzygies of B^t and A and we get $h^0(E(1)) = 4, h^0(E^*(1)) = 3$, hence E is not self-dual.

Example 2.3. ($k = 4$) Let

$$\begin{aligned} B^t &= \begin{bmatrix} a & b & c & & d & e & f \\ & a & b & c & & d & e & f \\ & & a & b & c & & d & e & f \\ & & & a & b & c & & d & e & f \end{bmatrix} \\ A &= \begin{bmatrix} & f & e & d & & -c & -b & -a \\ & f & e & d & & -c & -b & -a \\ f & e & d & & -c & -b & -a \\ f & e & d & & -c & -b & -a \end{bmatrix} \end{aligned}$$

E is a special symplectic instanton bundle with $k = 4$. We compute

$$h^2(E \otimes E^*) = 12.$$

Example 2.4. ($k = 4$) Let B^t as in the Example 2.3. Let

$$A = \begin{bmatrix} & f & e & d & & -c & -b & -a \\ e & d & & 2f & -b & -a & & -2c \\ 3d & & f & e & -3a & & -c & -b \\ & f & e & d & & -c & -b & -a \end{bmatrix}.$$

In this case $h^2(E \otimes E^*) = 6$, $h^0(E(1)) = 4$, $h^0(E^*(1)) = 3$.

Example 2.5. ($k = 4$) Let B^t as in the Example 2.3. Let

$$A = \begin{bmatrix} & f & e & d & & -c & -b & -a \\ e & d & & 2f & -b & -a & & -2c \\ 3d & & f & e & -3a & & -c & -b \\ 5d & f & e & d + f & e & -5a & -c & -b - a - c - b \end{bmatrix}.$$

Now $H^2(E \otimes E^*) = 0$, $h^0(E(1)) = 4$, $h^0(E^*(1)) = 2$.

3. On the singularities of moduli spaces.

The stable Schwarzenberger type bundles on \mathbb{P}^m (see (1.2)) form a Zariski open subset of the moduli space of stable bundles. Let $N_{\mathbb{P}^m}(k, q)$ be the moduli space of stable STB whose first Chern class is k and whose rank is q . The following proposition is easy and well known:

Proposition 3.1. *The space $N_{\mathbb{P}^m}(k, q)$ is smooth, irreducible of dimension $1 - k^2 - (q + k)^2 + k(q + k)(m + 1)$.*

We denote by $MI_{\mathbb{P}^{2n+1}}(k)$ the moduli space of stable instanton bundles with quantum number k . It is an open subset of the moduli space of stable $2n$ -bundles on \mathbb{P}^{2n+1} with Chern polynomial $(1 - t^2)^{-k}$.

On \mathbb{P}^5 (as on \mathbb{P}^3) all instanton bundles are stable by [AO], Theorem 3.6. $MI_{\mathbb{P}^{2n+1}}(2)$ is smooth ([AO] Theorem 3.14), unirational of dimension $4n^2 + 12n - 3$ and has zero Euler-Poincaré characteristic ([BE], [K]).

Theorem 3.2. *The space $MI_{\mathbb{P}^{2n+1}}(2)$ is irreducible.*

Proof. The moduli space $N = N_{\mathbb{P}^{2n+1}}(2, n + 2)$ of stable STB of rank $2n + 2$ and $c_1 = 2$ is irreducible of dimension $4n^2 + 8n - 3$ by Prop. 3.1

For a given instanton bundle E there is a STB S associated with E , which is stable ([AO], Theorem 2.8) and unique (ibid., Prop. 2.17). It is easy to prove that the map $\pi : M \rightarrow N$ defined by $\pi([E]) = [S]$ is algebraic, moreover π is dominant by [ST]. If $m = [E] \in M$, the fiber $\pi^{-1}(\pi(m))$ is a Zariski open subset of the grassmannian of planes in the vector space $H^0(\mathbb{P}^{2n+1}, S^*(1))$, where $\pi(m) = [S]$; by the Theorem 3.14 of [AO], $h^0(\mathbb{P}^{2n+1}, S^*(1)) = 2n + 2$, hence $\dim \pi^{-1}(\pi(m)) = 4n$.

In order to prove that M is irreducible, we suppose by contradiction that there are at least two irreducible components M_0 and M_1 of M . Then $M_0 \cap M_1 = \emptyset$ (M is smooth), $\pi(M_0)$ and $\pi(M_1)$ are constructible subset of N by Chevalley's theorem. Looking at the dimensions of M_0, M_1, N and the fibers of π we conclude that both $\pi(M_0)$ and $\pi(M_1)$ must contain an open subset of N , which implies $\pi(M_0) \cap \pi(M_1) \neq \emptyset$ by the irreducibility of N . This is a contradiction because the fibers of π are connected. \square

For $n \geq 2$ and $k \geq 3$, it is no longer true that $\text{MI}_{\mathbb{P}^{2n+1}}(k)$ is smooth. In fact on \mathbb{P}^5 we have:

Theorem 3.3. *The space $\text{MI}_{\mathbb{P}^5}(k)$ is singular for $k = 3, 4$. To be more precise, the irreducible component $M_0(k)$ of $\text{MI}_{\mathbb{P}^5}(k)$ containing the special instanton bundles is generically reduced of dimension $54(k = 3)$ or $65(k = 4)$, and $\text{MI}_{\mathbb{P}^5}(k)$ is singular at the points corresponding to special symplectic instanton bundles.*

Proof. Let E_0 be the special instanton bundle on \mathbb{P}^5 of the Example 2.2 ($k = 3$) or of the Example 2.5 ($k = 4$). Then $h^2(E_0 \otimes E_0^*) = 0$ and $M_0(k)$ is smooth at the point corresponding to E_0 , of dimension $h^1(E_0 \otimes E_0^*) = 54(k = 3)$ or $65(k = 4)$. In particular, $M_0(k)$ is generically reduced. If E_1 is a special symplectic instanton bundle on \mathbb{P}^5 , the computations in 2.1 and 2.3 show that $h^2(E_1 \otimes E_1^*) = 3(k = 3)$ or $12(k = 4)$, and $h^1(E_1 \otimes E_1^*) = 57$ or 77 respectively. Hence $\text{MI}_{\mathbb{P}^5}(k)$ is singular at E_1 for $k = 3$ and 4 . \square

Remark 3.4. It is natural to conjecture that $\text{MI}_{\mathbb{P}^{2n+1}}(k)$ is singular for all $n \geq 2$ and $k \geq 3$.

Theorem 3.5. *Let E be an instanton bundle on \mathbb{P}^{2n+1} with $c_2(E) = k$. Then*

$$h^1(E(t)) = 0 \text{ for } t \leq -2 \text{ and } k - 1 \leq t.$$

Proof. The result is obvious for $t \leq -2$. It is sufficient to prove $h^1(S^*(t)) = 0$ for $t \geq k - 1$. We have

$$S^*(t) = \bigwedge^{2n+k-1} S(t - k).$$

Taking wedge products of (1.2) we have the exact sequence

$$0 \rightarrow \mathcal{O}(t + 1 - 2n - 2k)^{\alpha_0} \rightarrow \dots \rightarrow \mathcal{O}(t - k - 1)^{\alpha_{2n+k-2}} \rightarrow \mathcal{O}(t - k)^{\alpha_{2n+k-1}} \rightarrow \bigwedge^{2n+k-1} S(t - k) \rightarrow 0$$

for suitable $\alpha_i \in \mathbb{N}$ and from this sequence we can conclude.

Ellia proves Theorem 3.5 in the case of \mathbb{P}^3 ([E], Prop. IV.1). He also remarks that the given bound is sharp. This holds on \mathbb{P}^{2n+1} as it is shown by the following theorem, which points out that the special symplectic instanton bundles are the “furthest” from having natural cohomology. \square

Theorem 3.6. *Let E be a special symplectic instanton bundle on \mathbb{P}^{2n+1} with $c_2 = k$. Then*

$$h^1(E(t)) \neq 0 \text{ for } -1 \leq t \leq k - 2.$$

Proof. For $n = 1$ the thesis is immediate from the exact sequence

$$0 \rightarrow \mathcal{O}(t - 1) \rightarrow E(t) \rightarrow \mathcal{J}_C(t + 1) \rightarrow 0$$

where C is the union of $k + 1$ disjoint lines in a smooth quadric surface. Then the result follows by induction on n by considering the sequence

$$0 \rightarrow E(t - 2) \rightarrow E(t - 1)^2 \rightarrow E(t) \rightarrow E(t)|_{\mathbb{P}^{2n-1}} \rightarrow 0$$

and the fact that, for a particular choice of the subspace \mathbb{P}^{2n-1} , the restriction $E|_{\mathbb{P}^{2n-1}}$ splits as the direct sum of a rank-2 trivial bundle and a special symplectic instanton bundle on \mathbb{P}^{2n-1} ([ST] 5.9). \square

Remark 3.7. In [OT] it is proved that if E_k is a special symplectic instanton bundle on \mathbb{P}^5 with $c_2 = k$ then $h^1(\text{End } E_k) = 20k - 3$.

In the following table we summarize what we know about the component $M_0(k) \subset \text{MI}_{\mathbb{P}^5}(k)$ containing E_k .

Table 3.10

	$h^1(E_k \otimes E_k^*)$	$h^2(E_k \otimes E_k^*)$	$\dim M_0(k)$	$\text{MI}_{\mathbb{P}^5}(k)$
$k = 1$	14	0	14	open subset of \mathbb{P}^{14}
$k = 2$	37	0	37	smooth, irreduc., unirat.
$k = 3$	57	3	54	singular
$k = 4$	77	12	65	singular
$k \geq 2$	$20k - 3$	$3(k - 2)^2$?	?

References

- [AO] V. Ancona and G. Ottaviani, *On the stability of special instanton bundles on \mathbb{P}^{2n+1}* , Trans. Amer. Math. Soc., **341** (1994), 677-693.
- [BE] J. Bertin and G. Elencwajg, *Symétries des fibrés vectoriels sur \mathbb{P}^n et nombre d'Euler*, Duke Math. J., **49** (1982), 807-831.
- [BaS] D. Bayer and M. Stillman, *Macaulay, a computer algebra system for algebraic geometry*.
- [BoS] G. Bohnhorst and H. Spindler, *The stability of certain vector bundles on \mathbb{P}^n* , Proc. Bayreuth Conference "Complex Algebraic Varieties", LNM **1507**, Springer Berlin (1992), 39-50.
- [E] Ph. Ellia, *Some vanishings for the cohomology of stable rank two vector bundles on \mathbb{P}^3* , J. reine angew. Math., **451** (1994), 1-14.
- [ES] G. Ellingsrud and S.A. Stromme, *Stable rank-2 bundles on \mathbb{P}^3 with $c_1 = 0$ and $c_2 = 3$* , Math. Ann., **255** (1981), 123-137.
- [Har] R. Hartshorne, *Stable vector bundles of rank 2 on \mathbb{P}^3* , Math. Ann., **238** (1978), 229-280.
- [K] T. Kaneyama, *Torus-equivariant vector bundles on projective spaces*, Nagoya Math. J., **111** (1988), 25-40.
- [M] M. Maruyama, *Moduli of stable sheaves, II*, J. Math. Kyoto Univ., **18** (1978), 557-614.
- [OS] C. Okonek and H. Spindler, *Mathematical instanton bundles on \mathbb{P}^{2n+1}* , Journal reine angew. Math., **364** (1986), 35-50.
- [OT] G. Ottaviani and G. Trautmann, *The tangent space at a special symplectic instanton bundle on \mathbb{P}^{2n+1}* , Manuscr. Math., **85** (1994), 97-107.
- [P] J. LePotier, *Sur l'espace de modules des fibrés de Yang et Mills*, in Mathématique et Physique, Sémin. E.N.S. 1979-1982, Basel-Stuttgart-Boston 1983.
- [S] R.L.E. Schwarzenberger, *Vector bundles on the projective plane*, Proc. London Math. Soc., **11** (1961), 623-640.
- [ST] H. Spindler and G. Trautmann, *Special instanton bundles on \mathbb{P}^{2n+1} , their geometry and their moduli*, Math. Ann., **286** (1990), 559-592.

Received January 5, 1993. Both authors were supported by MURST and by GNSAGA of CNR.

DIPARTIMENTO DI MATEMATICA
 VIALE MORGAGNI 67 A
 I-50134 FIRENZE
E-mail address: ancona@udininw.math.unifi.it

AND

DIPARTIMENTO DI MATEMATICA
 VIA VETOIO, COPPITO
 I-67010 L'AQUILA
E-mail address: ottaviani@vxscqaquila.infn.it

Added in proof. After this paper has been written we received a preprint of R. Miró-Roig and J. Orus-Lacort where they prove that the conjecture stated in the Remark 3.4 is true.

