



UNIVERSITÀ  
DEGLI STUDI  
FIRENZE

# FLORE

## Repository istituzionale dell'Università degli Studi di Firenze

### Some properties of the solution set for integral differential equations

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

*Original Citation:*

Some properties of the solution set for integral differential equations / G.Anichini; G.Conti. - In: FAR EAST JOURNAL OF MATHEMATICAL SCIENCES: FJMS. - ISSN 0972-0871. - STAMPA. - 24:(2007), pp. 415-423.

*Availability:*

This version is available at: 2158/403257 since:

*Terms of use:*

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

*Publisher copyright claim:*

(Article begins on next page)

# SOME PROPERTIES OF THE SOLUTION SET FOR INTEGRAL DIFFERENTIAL EQUATIONS

## 1 Introduction and Notations

In this paper we are concerned with the solution sets for Volterra integral equation and integrodifferential equations like:

$$\begin{cases} x(t) &= h(t) + \int_0^t k(t, s)g(s, x(s))ds \\ x(0) &= x_0, \end{cases} \quad (1)$$

or

$$\begin{cases} x(t) &= f(t, x(t), \int_0^t k(t, s)g(s, x(s))ds) \\ x(0) &= x_0, \end{cases} \quad (2)$$

where  $h : I = [0, T) \longrightarrow \mathbb{R}^n$ ,  $k : I \times \mathbb{R}^n \longrightarrow \mathbb{R}^n$  are continuous functions,  $x_0$  is a given vector of  $\mathbb{R}^n$ ,  $I$  a (possible unbounded) interval of  $\mathbb{R}$ .

In the following  $B(x_0, r)$  will denote an  $r$ -ball (in the metric space  $(X, d)$ ) i.e. the set  $\{x \in X : d(x, x_0) < r\}$  where  $x_0$  is any point in  $X$ ;  $\bar{B}(x_0, r)$  will denote the closed ball centered in  $x_0 = 0$ .

Let now consider the (Hilbert) space  $L^2(I, \mathbb{R}^n)$  normed, as usually, by  $\|x\|_2 = (\int_I x^2(t)dt)^{\frac{1}{2}}$  and its (affine) subspace  $E = \{x \in L^2(I, \mathbb{R}^n) : x(0) = x_0\}$ . Let  $X$  be some Banach space; if  $V \subset X$  is some subset then  $\bar{V}$  will denote its (topological) closure and  $V^c$  will denote the complement of  $V$ . Finally  $\mathcal{B}(\mathcal{X})$  will denote the set of all nonempty and bounded subsets of  $X$ .

**Definition 1 :** Let  $X$  be a Banach space and  $A \subset X$  a subset. A measure  $\mu : \mathcal{B}(X) \longrightarrow \mathbb{R}^+$  defined by  $\mu(V) = \inf\{\epsilon > 0 : V \in \mathcal{B}(\mathcal{X}) \text{ admits a finite cover by sets of diameter } \leq \epsilon\}$  where diameter of  $V$  is the  $\sup\{\|x - y\| : x \in V, y \in V\}$ , is called the (Kuratowski) measure of noncompactness.

A measure like  $\mu$  has interesting properties, some of which are listed in the sequel:

a)  $\mu(V) = 0$  if and only if  $\overline{V}$  is compact;

b)  $\mu(V) = \mu(\overline{V})$ ;  $\mu(\text{conv}(V)) = \mu(V)$ ; ( $\text{conv}(V)$  = convex hull of  $V$ );

c)  $\mu(\alpha(V_1) + (1 - \alpha)V_2) \leq \alpha\mu(V_1) + (1 - \alpha)\mu(V_2)$ ,  $\alpha \in [0, 1]$ ;

d) if  $V_1 \subset V_2$  then  $\mu(V_1) \leq \mu(V_2)$ ;

e) if  $\{V_n\}$  is a nested sequence of closed sets of  $B_d(X)$

and if  $\lim_{n \rightarrow +\infty} \mu(V_n) = 0$  then  $\bigcap_{n=1}^{\infty} V_n \neq \emptyset$ .

The analogous measure of noncompactness for an operator is defined by  $\mu(F(V)) = \inf\{k > 0 : \mu(F(V)) \leq k\mu(V)\}$  for all bounded subsets  $V \subset X$ .

When  $X$  is a complete metric space and  $f : X \rightarrow X$  is a continuous mapping  $f$  is called an *mu*-set contraction if there exists  $k \in [0, 1)$  such that, for all bounded non-compact subsets  $V$  of  $X$ , the following relation holds:  $\mu(f(V)) \leq k\mu(V)$  ([?], pag 160).

A continuous operator  $F : X \rightarrow X$  such that  $\mu(F(V)) < \mu(V)$ , for any bounded  $V \subset X$ , is called *condensing* or *densifying*.

(The concept of measure of noncompactness is considerably dealt with in the references [?], [?] or [?].)

Let  $S$  and  $S_1$  be topological spaces and let  $f : S \rightarrow S_1$ . Then  $f$  is said to be *proper* if, whenever  $K_1$  is a compact subset of  $S_1$ ,  $f^{-1}(K_1)$  is a compact set in  $S$ . It is also known ([?], pag 160) that if  $X$  is a Banach space and  $f : X \rightarrow X$  is a continuous  $k$ -set contraction, then  $I - f$  is a proper mapping.

The following result, due to R.K. Juberg ([?]), will be useful in the proof of our main result:

**Proposition 1 :** *Let  $(a, b)$  be any real (possible unbounded) interval and let  $L^p(a, c)$ ,  $1 \leq p \leq +\infty$  be the Lebesgue's space of (the power  $p$ ) summable*

functions over  $(a, c)$  for every  $c \in (a, b)$ . For  $u \in L^p(c, b)$ ,  $v \in L^q(a, c)$ , where  $\frac{1}{p} + \frac{1}{q} = 1$ , we set

$$\begin{aligned} \rho = & \limsup_{\epsilon \rightarrow 0} \{ [\int_x^{a+\epsilon} |u(y)|^p dy]^{\frac{1}{p}} [\int_a^x |v(y)|^q dy]^{\frac{1}{q}}, a < x \leq a + \epsilon \} + \\ & + \limsup_{\delta \rightarrow 0} \{ [\int_x^b |u(y)|^p dy]^{\frac{1}{p}} [\int_{b-\delta}^x |v(y)|^q dy]^{\frac{1}{q}}, b - \delta \leq x < b \}. \end{aligned}$$

Let  $D$  be the linear operator defined by:  $D(f(y))(x) = \int_0^x u(x)v(y)f(y)dy$ ; in the sequel we shall assume that  $D$  is a bounded operator in the space  $L^p(0, T)$ . We want to recall that the operator  $D$  is bounded (in the  $L^p(a, b)$  space) if and only if the function

$\psi(x) = [\int_x^b |u(y)|^p dy]^{\frac{1}{p}} [\int_a^x |v(y)|^q dy]^{\frac{1}{q}}$  is bounded on  $(a, b)$ . This operator is not necessarily a compact operator; as matter of fact it is well known (see [?], for instance), that  $D$  is a compact operator if the functions  $u(\cdot)$  e  $v(\cdot)$  belongs to  $L^2(a, b)$ .

Furthermore the measure of noncompactness of  $D$ , i.e.  $\mu(D)$  satisfies  $(\frac{1}{2})^{1+\frac{1}{p}} \leq \mu(D) \leq p^{\frac{1}{q}} q^{\frac{1}{p}} \rho$ ; in the special case when  $p = q = 2$ , i.e. when the (Lebesgue) space  $L^p$  is a Hilbert space  $L^2$ , we obtain  $\rho\sqrt{\frac{1}{8}} \leq \mu(D) \leq 2\rho$ .

**Definition 2 :** An  $R_\delta$ -set is the intersection of a decreasing sequence  $\{A_n\}$  of compact AR (metric absolute retracts; see [?] or [?], for a reference.) Moreover it is known (see [?] for instance) that an  $R_\delta$ -set is an acyclic set in the Čech homology.

The following result also will be crucially used in the sequel:

**Proposition 2 :** ([?], pag 159). Let  $X$  be a space and let  $Y, \|\cdot\|$  be a Banach space and  $f : X \rightarrow X$  be a proper mapping. Assume further that for each  $\epsilon_n > 0$ ,  $n > 0 \in \mathbb{N}$  a proper mapping  $f_n : X \rightarrow X$  is given and the couple of conditions is satisfied:

- $\|f_n(x) - f(x)\| < \epsilon_n, \forall x \in X$ ;
- for any  $\epsilon_n > 0$  and  $y \in E$  such that  $\|y\| \leq \epsilon_n$ , the equation  $f_{\epsilon_n}(x) = y$  has exactly one solution.

Then the set  $S = f^{-1}(0)$  is an  $R_\delta$ -set.

*Remark:* a sequence  $f_{\epsilon_n}$  is called an  $\epsilon_n$  approximation (of the function  $f$ ).

**Proposition 3 :** ([?], pag ???). Let  $F, F_n : \overline{B}(0, r) \longrightarrow Y$  be condensing operators such that

- $\delta_n = \sup\{ \|F_n(x) - F(x)\|, x \in \overline{B}(0, r) \} \rightarrow 0$ , as  $n \rightarrow +\infty$ ;
- the equation  $x = F_n(x) + y$  has at most one solution if  $\|y\| \leq \delta_n$ .

Then the set of fixed points of  $F$  is an  $R_\delta$ -set.

### Main result

We are ready to establish out (main) existence result for the (initial value problems for) integral equations of the type here introduced.

First of all let  $F : B(0, r) \rightarrow E$  be defined as follows:

$$F(y) = h(t) + \int_0^t k(t, s)g(s, y(s))ds$$

where  $r$  is a real number (suitably defined below) and put  $m_0 = \|F(0)\|_2$ .

**Theorem 1 :** Let  $\rho$  the number defined in Proposition 1; then we assume that:

1. i) there are functions  $\alpha, \phi, : I \rightarrow \mathbb{R}^n$  belonging to  $L^2(I)$  such that  $k(t, s) = \alpha\phi(s)$  for every  $(t, s) \in I \times I$ ; moreover we assume that  $\|k\|_2 < 2\rho$ ;
2. ii)  $\|g(t, x)\| \leq \frac{1}{2\rho}\|x\| + b(t)$ , for  $(t, x) \in I \times \mathbb{R}^n$ ,  $b \in L^2(I)$ ,  $b(t) \geq 0$ ;
3. iii) there is a ball  $B(0, r)$  such that  $r > \frac{2m_0\rho}{2\rho - \|k\|_2}$ .

Then the set of solution of the integral problem (??) is an  $R_\delta$ -set.

*Remark:* The first part of the assumption i) is satisfied in many cases: for instance when  $k(t, s)$  is a Green function; see, for instance, [?] for similar cases.

*Proof:* Clearly the above operator  $F$  is a single value mapping and a possible fixed point of  $F$  is a solution of the integral problem (??).

In order to prove the theorem the following steps in the proof have to be established:

- a )  $F$  has a closed graph;
- b)  $F$  is a condensing mapping;
- c) The set of fixed point of  $F$  is  $R_\delta$ -set.

*Proof of Step a):* in fact, let  $y_n \rightarrow y_0$  and put  $G(y)(t) = g(t, y(t))$ . Now, from assumption ii), it follows that the superposition operator  $G$  mapping the space  $L^2$  into  $L^2$  is condensing (see [?]); thus we have  $\lim_n \|G(y_n) - G(y_0)\|_2 = 0$ . By using the Holder inequality, we get:

$$\begin{aligned} \|F(y_n) - F(y_0)\|_2 &= [\int_I |F(y_n)(s) - F(y_0)(s)|^2 ds]^{\frac{1}{2}} = \\ &= [\int_I [\int_0^t (k(t, s)g(s, y_n(s)) - k(t, s)g(s, y_0(s))) ds]^2 dt]^{\frac{1}{2}} \leq \|k\|_2 \|G(y_n) - G(y_0)\|_2 \end{aligned}$$

and this quantity is going to zero whenever  $n \rightarrow +\infty$ .

*Proof of Step b):* Always working from  $B(0, r)$  into  $E$ , we have  $F(y) = (H \circ G)(y)$ , where

$$H(y)(t) = \int_0^t \phi(s) \alpha(s) y(s) ds + h(t).$$

Now, by assumptions i) and ii), we have (see [?])  $\mu(G(V)) \leq \frac{1}{2\rho} \mu(V)$ , for any bounded set  $V \subset L^2(I \times \mathbb{R}^n)$  and also  $\mu(H) < 2\rho$ ; so (see [?])  $\mu(F) = \mu(H \circ G)(y) \leq \mu(H) \mu(G) < 1$ .

*Proof of Step c):* Finally we have to prove that the set of fixed points of the operator  $F$  is an  $R_\delta$ -set (in the sequel we assume that  $(a, b) = (0, T)$ .)

Let us consider the mappings  $F_n : L^2(0, T) \rightarrow L^2(0, T)$  defined as:

$$F_n(x)(t) = \begin{cases} h(t) & \text{if } 0 \leq t \leq \frac{T}{n}; \\ h(t) + \int_0^{t-\frac{T}{n}} \phi(s) \alpha(s) g(s, y(s)) ds & \text{if } \frac{T}{n} \leq t \leq T. \end{cases} \quad (3)$$

The mappings  $F_n$  are continuous mappings; by assumption i) and ii) we have that they are also condensing. The intervals  $[0, \frac{T}{n}]$ ,  $[\frac{T}{n}, \frac{2T}{n}]$ ,  $\dots$ ,  $[\frac{kT}{n}, \frac{(k+1)T}{n}]$ ,  $\dots$ ,  $[\frac{(n-1)T}{n}, T]$  are now coming in one after the other: each time the mappings  $F_n$  are bijective and their inverses  $F_n^{-1}$  are continuous. Moreover we have  $\|F_n - F\|_2 \rightarrow 0$  as  $n \rightarrow +\infty$ . The latter fact allows us to say that the mappings  $I - F_n$  and  $I - F$  are proper maps. Finally we can conclude that the set of fixed points of  $F$  is an  $R_\delta$ -set.

## Riferimenti bibliografici

- [1] G. Anichini - G. Conti *Existence of Solutions of a Boundary Value Problem through the solution mapping of a linearized type problem*, Rendiconti del Seminario Mate. Univ. Torino, Fascicolo speciale dedicato a *Mathematical theory of dynamical systems and ordinary differential equations*, 1990, vol 48 (2), p. 149 – 160,
- [2] G. Anichini - G. Conti - P. Zecca *Using solution sets for solving boundary value problems for ordinary differential equations*, Nonlinear Analysis Theory Meth.& Appl., 1991, vol 5, p. 465–474,
- [3] G. Anichini - G. Conti *A direct approach to the existence of solutions of a Boundary Value Problem for a second order differential system*, Differential Equations and Dynamical Systems, 1995, vol 3 (1), p. 23 – 34,
- [4] G. Anichini - G. Conti *About the Existence of Solutions of a Boundary Value Problem for a Carathéodory Differential System*, Zeitschrift für Analysis und ihre Anwendungen, 1997, vol 16 (3), p. 621 – 630,
- [5] G. Anichini - G. Conti *Boundary Value Problem for Implicit ODE's in a singular case*, Differential Equations and Dynamical Systems, 1999, vol 7 (4), p. 437 – 459,
- [6] G. Anichini - G. Conti *How to make use of the solutions set to solve Boundary Value Problems*, Progress in Nonlinear Differential Equations and their Applications, Springer Verlag (Basel), 2000, vol 40,
- [7] G. Anichini - G. Conti *Boundary value problems for perturbed differential systems on an unbounded interval*, International Mathematical Journal, 2002, vol 2 (3), p. 221 – 234 (?),
- [8] J. Banas - K. Goebel *Measures of noncompactness in Banach spaces*, Lecture Notes in Pure and Applied Mathematics, Marcel Dekker, New York, 1980,
- [9] F.E. Browder - C.P. Gupta *Topological Degree and Nonlinear Mappings of Analytic Type in Banach spaces*, Journal of Mathematical Analysis and Applications, 1969, vol 26 (4), p. 390 – 402 ?),

- [10] G. Conti - J. Pejsachowicz *Fixed point theorems for multivalued maps*, Annali Matem. Pura Appl., 1980, vol 126 (4), p. 319 – 341
- [11] G. Darbo *Punti uniti in trasformazioni a codominio non compatto*, Rend. Sem. Matem. Univ. Padova, 1955, vol 24, p. 84 – 92
- [12] A. Deimling *Nonlinear Functional Analysis*, Springer Verlag, Berlin, 1984  
 bibitem13 L. Gorniewicz, *Topological Approach to differential inclusions*, NATO-ASI Series, A.Granas – M. Frigon editors, Kluwer, 1990, vol 472, p. 129 – 190,
- [13] H. Hochstadt *Integral Equations*, Pure and Applied Matheamtics, Wiley, New York, 1973,
- [14] V.I. Istrăţescu *Fixed point theory*, D. Reidel Publishing Company, Dordrecht, 1981,
- [15] R.K. Juberg *The measure of noncompactness in  $L^p$  for a Class of Integral Operators*, Indiana Math. Journal, 1973/74, vol 23, p. 925 – 936,
- [16] M.A. Krasnoselkii - P.P. Zabreiko *Geometrical methods of nonlinear analysis*, Springer Verlag, Berlin, 1984
- [17] J. Lasry – R.Robert *Analyse nonlineare multivoque*, U.E.R. Math de la Décision, 1979, vol 249, Paris Dauphine,
- [18] W.V. Petryshyn *Solvability of various boundary value problems for the equation  $x'' = f(t, x, x', x'') - y$* , Pacific Journal of Math. 1986, vol. 122, p. 169 – 195
- [19] E.Spanier *Algebraic Topology*, McGraw Hill, New York, 1966