



UNIVERSITÀ  
DEGLI STUDI  
FIRENZE

## FLORE

# Repository istituzionale dell'Università degli Studi di Firenze

### **Sphingosine 1-phosphate induces Ca<sup>2+</sup> transients and cytoskeletal rearrangement in C2C12 myoblastic cells**

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

*Original Citation:*

Sphingosine 1-phosphate induces Ca<sup>2+</sup> transients and cytoskeletal rearrangement in C2C12 myoblastic cells / L. Formigli; F. Francini; E. Meacci; M. Vassalli; D. Nosi; F. Quercioli; B. Tiribilli; C. Bencini; C. Piperio; P. Bruni; S.Z. Orlandini. - In: AMERICAN JOURNAL OF PHYSIOLOGY. CELL PHYSIOLOGY. - ISSN 0363-6143. - ELETTRONICO. - 282:(2002), pp. 1361-1373. [10.1152/ajpcell.00378.2001]

*Availability:*

The webpage <https://hdl.handle.net/2158/771749> of the repository was last updated on 2016-11-22T12:21:16Z

*Published version:*

DOI: 10.1152/ajpcell.00378.2001

*Terms of use:*

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

*Publisher copyright claim:*

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

# Sphingosine 1-phosphate induces $\text{Ca}^{2+}$ transients and cytoskeletal rearrangement in $\text{C}_2\text{C}_{12}$ myoblastic cells

Lucia Formigli, Fabio Francini, Elisabetta Meacci, Massimo Vassalli, Daniele Nosi, Franco Quercioli, Bruno Tiribilli, Chiara Bencini, Claudia Piperio, Paola Bruni and Sandra Zecchi Orlandini

*Am J Physiol Cell Physiol* 282:C1361-C1373, 2002. First published 30 January 2002;  
doi: 10.1152/ajpcell.00378.2001

---

## You might find this additional info useful...

This article has been cited by 9 other HighWire-hosted articles:

<http://ajpcell.physiology.org/content/282/6/C1361#cited-by>

Updated information and services including high resolution figures, can be found at:

<http://ajpcell.physiology.org/content/282/6/C1361.full>

Additional material and information about *American Journal of Physiology - Cell Physiology* can be found at:

<http://www.the-aps.org/publications/ajpcell>

---

This information is current as of October 23, 2012.

# Sphingosine 1-phosphate induces $\text{Ca}^{2+}$ transients and cytoskeletal rearrangement in $\text{C}_2\text{C}_{12}$ myoblastic cells

LUCIA FORMIGLI,<sup>1</sup> FABIO FRANCINI,<sup>2</sup> ELISABETTA MEACCI,<sup>3</sup> MASSIMO VASSALLI,<sup>4</sup>  
DANIELE NOSI,<sup>1</sup> FRANCO QUERCIOLO,<sup>4</sup> BRUNO TIRIBILLI,<sup>4</sup> CHIARA BENCINI,<sup>2</sup>  
CLAUDIA PIPERIO,<sup>2</sup> PAOLA BRUNI,<sup>3</sup> AND SANDRA ZECCHI ORLANDINI<sup>1</sup>

Departments of <sup>1</sup>Anatomy, Histology, and Forensic Medicine, <sup>2</sup>Physiological Sciences,  
and <sup>3</sup>Biochemical Sciences, University of Florence, 50134 Florence; and <sup>4</sup>Biophotonics  
Lab, National Institute of Applied Optics, 50125 Florence, Italy

Received 6 August 2001; accepted in final form 25 January 2002

**Formigli, Lucia, Fabio Francini, Elisabetta Meacci, Massimo Vassalli, Daniele Nosi, Franco Quercioli, Bruno Tiribilli, Chiara Bencini, Claudia Piperio, Paola Bruni, and Sandra Zecchi Orlandini.** Sphingosine 1-phosphate induces  $\text{Ca}^{2+}$  transients and cytoskeletal rearrangement in  $\text{C}_2\text{C}_{12}$  myoblastic cells. *Am J Physiol Cell Physiol* 282: C1361–C1373, 2002. First published January 30, 2002; 10.1152/ajpcell.00378.2001.—In many cell systems, sphingosine 1-phosphate (SPP) increases cytosolic  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_i$ ) by acting as intracellular mediator and extracellular ligand. We recently demonstrated (Meacci E, Cencetti F, Formigli L, Squecco R, Donati C, Tiribilli B, Quercioli F, Zecchi-Orlandini S, Francini F, and Bruni P. *Biochem J* 362: 349–357, 2002) involvement of endothelial differentiation gene (Edg) receptors (Rs) specific for SPP in agonist-mediated  $\text{Ca}^{2+}$  response of a mouse skeletal myoblastic ( $\text{C}_2\text{C}_{12}$ ) cell line. Here, we investigated the  $\text{Ca}^{2+}$  sources of SPP-mediated  $\text{Ca}^{2+}$  transients in  $\text{C}_2\text{C}_{12}$  cells and the possible correlation of ion response to cytoskeletal rearrangement. Confocal fluorescence imaging of  $\text{C}_2\text{C}_{12}$  cells preloaded with  $\text{Ca}^{2+}$  dye fluo 3 revealed that SPP elicited a transient  $\text{Ca}^{2+}$  increase propagating as a wave throughout the cell. This response required extracellular and intracellular  $\text{Ca}^{2+}$  pool mobilization. Indeed, it was significantly reduced by removal of external  $\text{Ca}^{2+}$ , pretreatment with nifedipine (blocker of L-type plasma membrane  $\text{Ca}^{2+}$  channels), and inositol 1,4,5-trisphosphate [ $\text{Ins}(1,4,5)\text{P}_3$ ]-mediated  $\text{Ca}^{2+}$  pathway inhibitors. Involvement of EdgRs was tested with suramin (specific inhibitor of Edg-3). Fluorescence associated with  $\text{Ins}(1,4,5)\text{P}_3$ Rs and L-type  $\text{Ca}^{2+}$  channels was evident in  $\text{C}_2\text{C}_{12}$  cells. SPP also induced  $\text{C}_2\text{C}_{12}$  cell contraction. This event, however, was unrelated to  $[\text{Ca}^{2+}]_i$  increase, because the two phenomena were temporally shifted. We propose that SPP may promote  $\text{C}_2\text{C}_{12}$  cell contraction through  $\text{Ca}^{2+}$ -independent mechanisms.

calcium ion transients; cytoskeleton; cell contraction; confocal microscopy

SPHINGOSINE 1-PHOSPHATE (SPP) is a bioactive lysophospholipid mediator that is recognized as a highly versatile molecule capable of affecting many cellular processes, including cell proliferation and differentiation,

apoptotic cell death, cell motility, and cytoskeletal organization (19). Some of these biological effects of SPP have long been attributed to its action as second messenger. Indeed, the mitogenic responses to several growth factors, such as platelet-derived growth factor, epidermal growth factor, nerve growth factor, and insulin, as well as the inhibition of apoptosis induced by antiproliferative drugs have been related to the activation of sphingosine kinase and to the subsequent enhanced production of intracellular SPP (35, 36, 38). More recently, the identification of a subset of receptors belonging to the endothelial differentiation gene (Edg) receptor (R) family has suggested that SPP may also act as an extracellular lipid mediator. In agreement with this suggestion, SPP is released on platelet activation and is an important constituent of serum (18). Moreover, extracellular SPP stimulation is required for inhibition of cell motility in vascular smooth muscle and melanoma cells, neurite retraction, and stimulation of DNA synthesis in 3T3 fibroblasts (8, 39, 50, 52). Only very recently has a role for exogenous SPP in the pathogenesis of inflammatory diseases such as asthma also been proposed (2).

It is becoming apparent that both modes of action of SPP may involve  $\text{Ca}^{2+}$  mobilization from intracellular stores and/or from the external pool. However, at present, the mechanisms by which SPP affects cytosolic  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_i$ ) are far from being fully delineated. Several studies have suggested that SPP, when acting as a second messenger, can directly promote  $\text{Ca}^{2+}$  release from the endoplasmic reticulum (ER) through mechanisms independent of the activation of the ryanodine (Ry)R or inositol 1,4,5-trisphosphate [ $\text{Ins}(1,4,5)\text{P}_3$ ]R pathways (5, 54). Consistent with this, a novel sphingolipid-gated  $\text{Ca}^{2+}$ -permeable channel has been discovered on isolated ER vesicles of *Xenopus* oocytes (26). In contrast, dissection of the  $\text{Ca}^{2+}$  signaling pathways triggered by the interaction of SPP with its G protein-coupled receptors is of great difficulty and complexity considering that multiple ef-

Address for reprint requests and other correspondence: S. Zecchi Orlandini, Dept. of Anatomy, Histology, and Forensic Medicine, Univ. of Florence, Viale Morgagni, 85, 50134 Florence, Italy (E-mail: zecchi@unifi.it).

The costs of publication of this article were defrayed in part by the payment of page charges. The article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

effector systems, including phospholipase C (PLC) and protein kinase C (PKC), may be involved (3, 37, 40). In particular, although activation of phospholipases has been reported in several cell types after SPP stimulation, only limited data have provided conclusive evidence for a role of Ins(1,4,5)P<sub>3</sub> in the SPP-induced Ca<sup>2+</sup> signaling pathway. Moreover, we recently found (28) in a myoblastic C<sub>2</sub>C<sub>12</sub> cell line that SPP elicits a Ca<sup>2+</sup> response in the form of Ca<sup>2+</sup> waves that are dependent on extracellular Ca<sup>2+</sup>, thus suggesting a role for the lipid metabolite in mediating the influx of the ion through plasma membrane Ca<sup>2+</sup> channels. Furthermore, in the same study we also showed that inhibition of Edg-3R and Edg-5R by specific antisense oligodeoxyribonucleotides totally abolished SPP-induced Ca<sup>2+</sup> response (28). Voltage-dependent dihydropyridine receptors (L-type Ca<sup>2+</sup> channels), located on the plasma membrane, represent the major Ca<sup>2+</sup> entry pathway in excitable cells. In particular, Ca<sup>2+</sup> influx through these channels is critical for the activation of Ca<sup>2+</sup>-induced Ca<sup>2+</sup> release via RyR channels of the sarcoplasmic reticulum (SR) and for contractility stimulation in cardiac muscle cells (5). In contrast, L-type Ca<sup>2+</sup> channels were shown to couple conformationally with RyRs on depolarization to release Ca<sup>2+</sup> during contraction in mature skeletal muscle cells (33, 41, 49). Nevertheless, even though numerous studies exist on the physiological significance of L-type Ca<sup>2+</sup> channels in striated muscle cells, their possible role in SPP-induced Ca<sup>2+</sup> response remains to be studied.

Because a better understanding of the molecular basis of SPP action may be of crucial importance in understanding the physiological significance and possible pathological implications of this metabolite, it seemed worthwhile, in the present study, to further characterize the Ca<sup>2+</sup> response elicited by exogenous SPP in skeletal muscle cells, particularly in view of the crucial role exerted by Ca<sup>2+</sup> effector molecules in skeletal muscle development and differentiation. Confocal laser scanning microscopy equipped with Time Course software was then used to determine the spatiotemporal distribution of SPP-mediated Ca<sup>2+</sup> transients in a myoblastic C<sub>2</sub>C<sub>12</sub> cell line and the extracellular and intracellular sources of Ca<sup>2+</sup> mobilization and to characterize the pattern of expression of voltage- and ligand-gated plasma membrane and intracellular Ca<sup>2+</sup> channels in these cells. The effects of SPP-induced Ca<sup>2+</sup> transients on the cytoskeletal reorganization were also considered in view of the well-known role that this ion plays in the regulation of cell contractility.

## MATERIALS AND METHODS

### Cell Cultures

Mouse skeletal C<sub>2</sub>C<sub>12</sub> myoblasts (51) were obtained from American Type Culture Collection (Manassas, VA) and maintained in Dulbecco's modified minimum essential medium (DMEM) with 10% fetal calf serum, penicillin (100 U/ml), and streptomycin (100 µg/ml) (Sigma, Milan, Italy) at 37°C and exposed to a humidified atmosphere of 5% CO<sub>2</sub>.

### Confocal Analysis of Calcium Transients

To reveal variations in intracellular concentrations of calcium in C<sub>2</sub>C<sub>12</sub> cells incubated with SPP (Calbiochem, San Diego, CA), ~2 × 10<sup>4</sup> cells were plated on glass coverslips and incubated at room temperature for 10 min in serum-free DMEM with 0.1% bovine serum albumin (BSA) containing fluo 3-acetoxymethyl ester as fluorescent calcium indicator at a final concentration of 10 µM and 0.1% anhydrous dimethyl sulfoxide and Pluronic F-127 (0.01% wt/vol) as dispersing agent (Molecular Probes, Eugene, OR). The cells were then washed and maintained in fresh medium for 10 min to allow complete deesterification of fluo 3. After that, the cells were placed in open slide flow-loading chambers that were mounted on the stage of a confocal Bio-Rad MRC 1024 ES scanning microscope (Bio-Rad, Hercules, CA) equipped with a krypton/argon laser source (15 mW) for fluorescence measurements. The microscope was also equipped with differential interference contrast (DIC) optics. The fluorescence of fluo 3-loaded cells was monitored by using a 488-nm wavelength and collecting the emitted fluorescence with a Nikon Plan Apo ×60 oil-immersion objective through a 510-nm long-wave pass filter. The time course analysis of Ca<sup>2+</sup> waves after SPP stimulation was performed with Time Course Kinetic software (Bio-Rad).

Some experiments were performed in Ca<sup>2+</sup>-free, 2 mM Mg<sup>2+</sup>-containing medium and/or after pretreatment of C<sub>2</sub>C<sub>12</sub> cells with various modulators of known voltage- and ligand-gated calcium channels. In particular, caffeine (100 mM; Sigma), 2-aminoethyl-diphenylborate (2-APB, 100 µM; Alexis, San Diego, CA), heparin (50 mM), and 1-[6-[(17β)-3-methoxyestra-1,3,5(10)-trien-17-yl]amino]hexyl]-1H-pyrrole-2,5-dione (U-73122, 10 µM; Alexis) were used to inhibit any potential Ins(1,4,5)P<sub>3</sub>-mediated release, Ry (100 µM; Sigma) to inhibit RyR-Ca<sup>2+</sup> release channels, and nifedipine (100 nM; Sigma) to inhibit Ca<sup>2+</sup> influx through L-type Ca<sup>2+</sup> channels. To test the involvement of EdgRs in the Ca<sup>2+</sup> response, the cells were treated with suramin (100 µM) before stimulation. SPP was dissolved in the medium by fast perfusion; the small size of the chamber used (0.2 ml) and the perfusion flux of ~0.2 ml/s allowed a complete replacement of the bathing medium in ~1 s.

Usually, cells did not reach confluence on coverslips; a single coverslip with adherent cells was used for only one experiment. For each cell preparation a variable number of cells ranging from 10 to 22 were analyzed. Multiple regions of interest (ROIs) of 25 µm<sup>2</sup> were selected in single cells to monitor the spatiotemporal distribution of Ca<sup>2+</sup> transients. Fluorescence signals are expressed as fractional changes above the resting baseline, ΔF/F, where F is the averaged baseline fluorescence before the application of SPP and ΔF represents the fluorescence changes from the baseline. The latency (T<sub>0</sub>) of the Ca<sup>2+</sup> wave was measured as the lag between the addition of the agonist and the beginning of the fluorescence increase over the basal noise. The time to peak (T<sub>p</sub>) was measured as the time interval between T<sub>0</sub> and the peak level. The time to half-decay (T<sub>0.5</sub>) of fluorescence was measured as the time for the fluorescence to decay from the peak to half its peak value. The temporal delay to peak amplitude between adjacent ROIs was used to calculate the extent of synchronization vs. propagation of Ca<sup>2+</sup> transients. Usually, two ROIs were placed within the nucleus, whereas a variable number ranging from 3 to 10 ROIs were placed inside the cytoplasm. Confocal fluorescence images were also used to evaluate intracellular Ca<sup>2+</sup> spatial distribution with a purpose-developed software running under Interactive Data Language (Research Systems, Boulder, CO).






Fig. 1. Calcium wave images of fluo 3-loaded C<sub>2</sub>C<sub>12</sub> cells by confocal microscopy. The pseudocoloring represents the absolute Ca<sup>2+</sup> increase as indicated by the color bar. Basal fluorescence is shown in A. Next images were taken ~20 s from sphingosine 1-phosphate (SPP; 1 µM) stimulation (B) and as indicated in C–I, L–N. In the top cell in each panel, a burst of fluorescence is initially visible on the left of the myoblast and then propagates onward, invading the nucleus. In the bottom cell, the fluorescence signal invades, at the same time, the cytoplasmic and nuclear regions.

#### Determination of Inositol Phosphate Production

Serum-starved C<sub>2</sub>C<sub>12</sub> myoblasts were incubated for 24 h in inositol-free DME in the presence of 5 µCi/ml *myo*-[2-<sup>3</sup>H(N)]inositol (25 Ci/mmol; NEN, Dreiech, Germany). Two hours before the beginning of the experiment the medium was changed, and 30 min before the addition of the agonist (1 µM SPP) the cells were incubated with 20 mM LiCl. Incubation was stopped by aspirating the medium and washing the monolayer twice with PBS. Inositol phosphate (InsP) accu-

mulated in the cells was extracted with 5% ice-cold perchloric acid for 30 min. Cell extracts were neutralized with K<sub>2</sub>CO<sub>3</sub>, and InsP was separated onto Dowex (Bio-Rad) formate form (1 ml) and quantified essentially as described previously (45).

#### Determination of Diacylglycerol Production

C<sub>2</sub>C<sub>12</sub> cells were labeled with 5 µCi/ml [2-<sup>3</sup>H]glycerol (14.2 Ci/mmol; NEN) for 24 h and then incubated for 30 s with 1 µM SPP. Lipid extraction and [<sup>3</sup>H]diacylglycerol (DAG) sep-

aration by thin-layer chromatography was performed as described previously (31).

### Confocal Immunofluorescence

C<sub>2</sub>C<sub>12</sub> cells grown on coverslips were fixed in 4% buffered paraformaldehyde for 10 min at room temperature. The cells were then washed, permeabilized with 0.2% Triton X-100 in PBS for 5 min, and blocked with a solution containing 0.5% BSA and 3% glycerol in PBS.

**Calcium channel immunodetection.** Cells were incubated with the following primary antibodies diluted in BSA-PBS: rabbit anti- $\alpha_{1c}$  L-type channels (recognizing  $\alpha_{1c}$ -subunit of voltage-gated Ca<sup>2+</sup> channel) and rabbit anti- $\alpha_{1D}$  L-type channels (reacting with all forms of  $\alpha_{1D}$ -subunit of voltage-gated Ca<sup>2+</sup> channel; 1:100; Chemicon International), mouse anti-Ins(1,4,5)P<sub>3</sub>R [recognizing COOH-terminal cytoplasmic domain of Ins(1,4,5)P<sub>3</sub>R types 1, 2, and 3; 1:200; Chemicon], and mouse anti-RyR (reacting with COOH-terminal domain of RyR; 500 kDa, 1:50; Chemicon) for 1 h at room temperature. After incubation with the primary antibodies, the cells were washed to remove unbound antibodies (Abs) and incubated with Alexa 488-conjugated anti-mouse or anti-rabbit secondary Abs (1:200 dilution; Chemicon). Counterstaining for F-actin was performed with rhodamine-phalloidin (Sigma).

**Cytoskeletal protein immunodetection.** C<sub>2</sub>C<sub>12</sub> cells were incubated with a monoclonal anti-myosin Ab (1:50 dilution; Sigma) and a monoclonal anti-vinculin Ab (1:100 dilution; Sigma) for 1 h at room temperature. The cells were subsequently incubated with Alexa 488-labeled anti-mouse IgG (Molecular Probes). Actin filaments were stained with tetramethylrhodamine-isothiocyanate-labeled phalloidin. After a

series of washes the coverslips containing the immunostained cells were mounted with an antifade mounting medium (Biomed; Electron Microscopic Sciences). Negative controls were performed by substituting blocking solution for the primary Abs. The fluorescence signals were revealed by a confocal laser scanning microscope (Bio-Rad). To this purpose, a series of optical sections (512 × 512 pixels) was taken through the depth of the cells with a thickness of 1  $\mu$ m at intervals of 0.8  $\mu$ m. Twenty optical sections for each examined sample were then projected as a single composite image by superimposition.


### Imaging of Dynamic Changes of Actin Cytoskeleton

Alexa 488-labeled G-actin monomers were used as probes for live cell cytoskeleton. The monomers were introduced into C<sub>2</sub>C<sub>12</sub> cells by the scrape-loading technique. Briefly, C<sub>2</sub>C<sub>12</sub> cells grown to confluence on 100-mm petri dishes were washed twice with PBS and 2 ml of scraping buffer (in mM: 114 KCl, 1 NaCl, 5.5 MgCO<sub>3</sub>, and 10 Tris·HCl). Fluorescent G-actin monomers (250  $\mu$ g/ml; Molecular Probes) were added in 0.5 ml of scraping buffer, and the cells were gently scraped, suspended in DMEM-10% FCS, and split among six-well dishes on coverslips. After 1 h of incubation, we estimated that ~70% of the treated cells were viable. Labeled cells were then observed by confocal microscope and used to test the effect of SPP on dynamic changes of actin cytoskeleton.

### Statistics

All data are expressed as means  $\pm$  SD. Data were analyzed by one-way ANOVA with Bonferroni's correction for multiple comparisons. The  $\alpha$ -value was at  $P < 0.05$  for all tests.

Fig. 2. SPP-mediated Ca<sup>2+</sup> waves in C<sub>2</sub>C<sub>12</sub> myoblastic cells. Time course of Ca<sup>2+</sup> response from representative cells in Ca<sup>2+</sup>-containing medium (A), Ca<sup>2+</sup>-free medium (B), and Ca<sup>2+</sup>-containing medium after pretreatment with nifedipine (100 nM; C) or ryanodine (100  $\mu$ M; D). The data are from representative cells. Inset, number of regions of interest (ROIs; cy and n indicate ROIs taken in cytoplasm and nucleus, respectively). Arrows indicate the addition of SPP (time 0). In A and D, the left vertical lines indicate latency ( $T_0$ ) values in the cytoplasmic ROIs (2), the dashed vertical lines indicate  $T_0$  in the nuclear ROIs (9 or 7), and the right vertical lines indicate  $T_0$  in cytoplasmic ROIs (13 or 12) located at the opposite edge of the cells examined. In B and C,  $T_0$  values are doubled and are the same in all ROIs.  $\Delta F/F$ , fractional fluorescence changes above resting baseline.





## RESULTS

*SPP-Induced Ca<sup>2+</sup> Transients in C<sub>2</sub>C<sub>12</sub> Myoblasts*

C<sub>2</sub>C<sub>12</sub> cells cultured on coverslips and observed under light microscopy showed variable morphology, being rounded or spindle-shaped with several cytoplasmic projections emanating from the cell surfaces and anchoring to the substrate. We first verified the responsiveness of the C<sub>2</sub>C<sub>12</sub> myoblasts. To this aim, the cells were loaded with fluorescent Ca<sup>2+</sup> dye fluo 3 and ATP (1 mM) was added to Ca<sup>2+</sup>-containing medium. In accordance with previous reports (24), ATP induced a significant intracellular Ca<sup>2+</sup> elevation in all cells (data not shown). The application of exogenous SPP (1  $\mu$ M) to C<sub>2</sub>C<sub>12</sub> cells also promoted a substantial increase in intracellular Ca<sup>2+</sup> that was evident in both the cytoplasmic and nuclear compartments. This increase was concentration dependent, with an EC<sub>50</sub> of  $\sim$ 50 nM. The SPP-induced Ca<sup>2+</sup> increase was transient and was followed by a return to near resting levels within 1 min (Fig. 1). In particular, a synchronous Ca<sup>2+</sup> increase was observed in some cells, whereas the Ca<sup>2+</sup> response propagated as a wave in others (Figs. 1 and 2A). As shown in Fig. 3A, almost 60% of the examined cells were responsive to SPP, exhibiting relative fluorescence changes significantly ( $P < 0.001$ ) higher in the nuclear than in the cytoplasmic region (Fig. 3B). Moreover, differences in the time course of the fluorescence signal were found between the cytoplasmic and nuclear regions; in fact,  $T_p$  and  $T_{0.5}$  of the nucleus were significantly higher ( $P < 0.001$ ) and lower ( $P < 0.05$ ), respectively, compared with those of the cytoplasm (Fig. 3, C–E). The Ca<sup>2+</sup> response elicited by exogenous SPP could consist of at least two components: Ca<sup>2+</sup> influx across the plasma membrane and Ca<sup>2+</sup> release from the endogenous stores. To better investigate this issue, we stimulated C<sub>2</sub>C<sub>12</sub> cells in Ca<sup>2+</sup>-free, Mg<sup>2+</sup>-containing medium (Fig. 2B). Under these particular conditions, the number of cells responsive to SPP was reduced to  $\sim$ 40% (Fig. 3A), and the cytosolic and nuclear  $\Delta F/F$  increase in response to 1  $\mu$ M SPP was significantly reduced by  $\sim$ 35% ( $P < 0.001$ ) (Fig. 3B). Moreover, the Ca<sup>2+</sup> increase was evident as a synchronous rather than propagated Ca<sup>2+</sup> wave with a  $T_0$  significantly increased with respect to controls (Figs. 2B and 3C). In myoblasts cultured in the absence of external Ca<sup>2+</sup>, a slight increase in  $T_p$  and a remarkable increase in  $T_{0.5}$  ( $P < 0.001$ ) in both the cytoplasmic and nuclear ROIs (Fig. 3, D and E) were found. All these data together suggested that SPP-induced Ca<sup>2+</sup> response was attributable to both extracellular and intracellular Ca<sup>2+</sup> pool mobilization.

A role for extracellular Ca<sup>2+</sup> influx was further confirmed in two-step experimental protocols consisting of addition of SPP to cells cultured in Ca<sup>2+</sup>-free medium and subsequent readministration of Ca<sup>2+</sup> to the medium once the ion transients had occurred (Fig. 4, A and B). Indeed, Ca<sup>2+</sup> readministration caused a faster elevation of intracellular Ca<sup>2+</sup> in all cells in both the cytoplasmic and nuclear ROIs (Fig. 4, A, B, F, and G).




Fig. 3. SPP-mediated Ca<sup>2+</sup> waves in C<sub>2</sub>C<sub>12</sub> myoblastic cells: statistical analysis. In A–E, bars indicated as Ca, Ca free, Ca + nif and Ca + Ry refer to experiments like those reported in Fig. 2, A–D. A: % of SPP-responsive cells. B–E: values of  $\Delta F/F$ ,  $T_0$ , time to peak ( $T_p$ ), and time to half-decay ( $T_{0.5}$ ) from the cytoplasmic (open bars) and nuclear (filled bars) ROIs examined are reported. Data are means  $\pm$  SD from cells cultured on 10 (A), 9 (B), 9 (C), and 6 (D) different coverslips; the total number of the cells analyzed was 120 (A), 102 (B), 106 (C), and 80 (D). Significant differences were evaluated by 1-way ANOVA test. +Data statistically different with respect to controls ( $P < 0.001$ ). In all experimental condition  $\Delta F/F$  values were significantly ( $P < 0.01$ ) larger in nucleus than in corresponding cytoplasmic ROIs. No statistically significant differences were found between Ca free and Ca + nif data or between control (Ca) and Ca + Ry. nif., Nifedipine; Ry, ryanodine.

The Ca<sup>2+</sup> increase was transient but, in contrast to that elicited by SPP, it rapidly decayed (Fig. 4H) to steady-state intracellular Ca<sup>2+</sup> levels that remained elevated above the baseline. This latter response was absent when the two-step procedure was applied to cells not previously stimulated by SPP (data not shown), thus suggesting the existence of a SPP-dependent Ca<sup>2+</sup> influx pathway mediated by the activation of putative plasma membrane Ca<sup>2+</sup> channels.

To verify this latter hypothesis, C<sub>2</sub>C<sub>12</sub> myoblasts were treated with nifedipine (100 nM), a prototypical blocker of plasma membrane L-type Ca<sup>2+</sup> channels, 20 min before stimulation with SPP (Fig. 2C). The presence of nifedipine in Ca<sup>2+</sup>-containing medium significantly reduced the Ca<sup>2+</sup> transients, which became similar to those observed in Ca<sup>2+</sup>-free medium (Figs. 2B and 3), supporting a role for nifedipine-sensitive receptors in SPP-mediated Ca<sup>2+</sup> influx.




Fig. 4. Effects of Ca<sup>2+</sup> readministration on SPP-mediated Ca<sup>2+</sup> waves in C<sub>2</sub>C<sub>12</sub> myoblasts from 2-step experimental protocol. **A** and **B**: cells were stimulated with SPP in Ca<sup>2+</sup>-free medium, and Ca<sup>2+</sup> was readministered to the medium once intracellular Ca<sup>2+</sup> had returned to baseline. An SPP-responsive and an unresponsive cell are shown in **A** and **B**, respectively. Note that both cells respond to the readministration of Ca<sup>2+</sup> (indicated by horizontal line marked Ca<sup>2+</sup>). **C**: experimental protocol as in **A** and **B** except that cells were pretreated with caffeine (100 mM). The cell is unresponsive to SPP but responds to the readministration of Ca<sup>2+</sup> to the medium. In **D–H**, bars labeled A, B, and C refer to the Ca<sup>2+</sup> response observed after readministration of Ca<sup>2+</sup> to the medium as reported in **A**, **B**, and **C**, respectively. The values obtained in cells in control conditions are similar to those in Fig. 3. Other symbols, labeling, and scales as in Fig. 3. Data are means  $\pm$  SD from cells cultured on 8 (**A** and **B**) and 7 (**C**) different coverslips; the total number of cells analyzed was 90 (**A**), 80 (**B**), and 76 (**C**). Significant differences were evaluated by 1-way ANOVA test. Data statistically different with respect to controls: \* $P < 0.05$ , + $P < 0.001$ . In all experimental conditions  $\Delta F/F$  values were significantly ( $P < 0.05$  for **A** and  $P < 0.001$  for **B** and **C**) larger in nucleus than in corresponding cytoplasmic ROIs. In **D** and **E**, which indicate the number of responsive cells,  $\Delta F/F$  values of **A** were significantly reduced ( $P < 0.001$ ) with respect to **B** and **C**. In **F–H**, no statistical differences were found among **A**, **B**, and **C** data.

We next tested C<sub>2</sub>C<sub>12</sub> myoblasts for the presence of voltage-dependent ionic channels in C<sub>2</sub>C<sub>12</sub> cells by adding KCl (100 mM) to the medium. High extracellular K<sup>+</sup> was not able to elicit any Ca<sup>2+</sup> response in undifferentiated C<sub>2</sub>C<sub>12</sub> cells. This finding indicated the absence of voltage-dependent Ca<sup>2+</sup> channels and demonstrated that the observed nifedipine-sensitive Ca<sup>2+</sup> influx in response to SPP was independent from the existence of functional voltage-dependent L-type Ca<sup>2+</sup> channels.

Subsequently, we explored the involvement of intracellular receptors in the Ca<sup>2+</sup> mobilization elicited by SPP by adding caffeine (100 mM), a known agonist of RyRs, to the medium. Caffeine was without effect on induction of Ca<sup>2+</sup> mobilization in all C<sub>2</sub>C<sub>12</sub> cells examined. Moreover, pretreatment with Ry (100  $\mu$ M) did not affect SPP-mediated intracellular Ca<sup>2+</sup> transients (Figs. 2D and 3), suggesting that RyR-mediated Ca<sup>2+</sup> mobilization was absent in C<sub>2</sub>C<sub>12</sub> myoblasts.






Fig. 5. Effects of pretreatment with inhibitors of inositol-1,4,5-trisphosphate [Ins(1,4,5)P<sub>3</sub>]-mediated Ca<sup>2+</sup> pathway and of inhibitor of endothelial differentiation gene (Edg) receptors (Rs) on SPP-mediated intracellular Ca<sup>2+</sup> elevation in C<sub>2</sub>C<sub>12</sub> myoblasts. Time course of the Ca<sup>2+</sup> response from cells in Ca<sup>2+</sup>-containing medium is shown. Data are referred to representative cells. Cells were pretreated with caffeine (100 mM; A), heparin (50 mM; B), 2-aminoethyl-diphenylborate (2-APB, 100  $\mu$ M; C), U-73122 (10  $\mu$ M; D), and suramin (100  $\mu$ M; E).


Interestingly, pretreatment with caffeine 1 min before SPP stimulation (Fig. 5A) significantly reduced the number of responsive cells ( $P < 0.05$ ) and the peak of Ca<sup>2+</sup> transients in both the cytoplasmic and nuclear ROIs by  $\sim 35\%$  ( $P < 0.001$ ; Fig. 6, A and B). Moreover,  $T_0$  and  $T_{0.5}$  of the Ca<sup>2+</sup> response were significantly increased with respect to controls in both the cytoplasm and nuclear ROIs ( $P < 0.001$ ; Fig. 6). Because it has been reported that caffeine, besides activating RyRs, inhibits Ins(1,4,5)P<sub>3</sub>Rs (53), these latter data are suggestive for an involvement of the Ins(1,4,5)P<sub>3</sub> signaling pathway in the cytosolic Ca<sup>2+</sup> response. As it occurred in the cells cultured in Ca<sup>2+</sup>-free medium, the pretreatment with caffeine before stimulation with SPP elicited a Ca<sup>2+</sup> response in the form of a synchronous wave (Fig. 5A). Quite similar results were obtained by pretreating C<sub>2</sub>C<sub>12</sub> cells with other inhibitors of the Ins(1,4,5)P<sub>3</sub> signaling Ca<sup>2+</sup> pathway such as heparin and 2-APB, both known blockers of Ins(1,4,5)

P<sub>3</sub>Rs (Figs. 5, B and C, and 6). In contrast, pretreatment with U-73122, a specific inhibitor of PLC activation, completely blocked the occurrence of Ca<sup>2+</sup> transients elicited by SPP (Figs. 5D and 6). These data are in favor of a role of PLC activation not only in the Ins(1,4,5)P<sub>3</sub> signaling pathway but also in mediating plasma membrane Ca<sup>2+</sup> influx.

The involvement of EdgRs in the Ca<sup>2+</sup> signaling pathway induced by SPP was tested by pretreating C<sub>2</sub>C<sub>12</sub> cells with suramin, an inhibitor of Edg-3R (Ref. 4; Fig. 5E). Pretreatment with suramin, although significantly ( $P < 0.001$ ) reducing the number of responsive cells (by  $\sim 40\%$ ), only slightly affected the time course of the intracellular Ca<sup>2+</sup> transients evoked by SPP in C<sub>2</sub>C<sub>12</sub> myoblasts (Figs. 5E and 6).

The dual origin, extracellular and intracellular, of Ca<sup>2+</sup> transients elicited by SPP was further supported by the finding of a complete inhibition of SPP-mediated Ca<sup>2+</sup> response after pretreatment with caffeine in

Fig. 6. Effects of pretreatment with inhibitors of Ins(1,4,5)P<sub>3</sub>-mediated Ca<sup>2+</sup> pathway on SPP-mediated intracellular Ca<sup>2+</sup> elevation in C<sub>2</sub>C<sub>12</sub> myoblasts: statistical analysis. The values obtained in cells in control conditions, indicated as Ca, are as in Fig. 3. In A–E, Ca + caf, Ca + hep, Ca + 2A, Ca + U7, and Ca + sur refer to experiments like those reported in Fig. 5, A–E, respectively. Other symbols, labeling, and scales as in Fig. 3. Data are means  $\pm$  SD from cells cultured on 10 (A), 9 (B), 9 (C), 9 (D), and 10 (E) different coverslips; the total number of cells analyzed was 99 (A), 80 (B), 86 (C), 102 (D), and 108 (E). Significant differences were evaluated by 1-way ANOVA test. Data statistically different with respect to controls (Ca): \* $P < 0.05$ , +  $P < 0.001$ . In all experimental conditions,  $\Delta F/F$  values were significantly ( $P < 0.001$ ) larger in nucleus than in corresponding cytoplasmic ROIs. No statistical differences for  $\Delta F/F$ ,  $T_0$ , and  $T_D$  were found among Ca + caf, Ca + hep, and Ca + 2-APB. caf, Caffeine; hep, heparin; 2A, 2-APB; U7, U-73122; sur, suramin.



Ca<sup>2+</sup>-free medium (Fig. 4C). The subsequent readministration of Ca<sup>2+</sup> caused fast elevation of intracellular Ca<sup>2+</sup> in all cells examined (Fig. 4D). The Ca<sup>2+</sup> increase was transient and decreased to steady-state intracellular Ca<sup>2+</sup> levels that remained elevated over baseline. A complete inhibition of the SPP-mediated Ca<sup>2+</sup> response in C<sub>2</sub>C<sub>12</sub> cells was also observed after pretreatment with caffeine and nifedipine in Ca<sup>2+</sup>-containing medium (data not shown), strongly suggesting that the plasma membrane Ca<sup>2+</sup> channels implicated in the Ca<sup>2+</sup> influx were sensitive to nifedipine.

#### InsP and DAG Production

To further study the characteristics and mechanisms of the Ca<sup>2+</sup> response elicited by SPP, we measured the production of radiolabeled total InsP as well as [<sup>3</sup>H]DAG after SPP stimulation. Treatment of C<sub>2</sub>C<sub>12</sub> cells with 1  $\mu$ M SPP increased the cellular levels of both InsP and DAG after 30 s by  $\sim 25\% \pm 3$  [10,141  $\pm$  697 (control) vs. 13,747  $\pm$  1,107 disintegrations/min (dpm) InsP/10<sup>6</sup> dpm labeled phospholipid (SPP);  $n = 4$ ,  $P = 0.05$ ] and  $41\% \pm 7$  [18,756  $\pm$  2,014 (control) vs. 27,530  $\pm$  2,950 dpm DAG/10<sup>6</sup> dpm labeled phospholipid (SPP);  $n = 3$ ,  $P < 0.01$ ], respectively.

#### Confocal Immunocytochemistry

To further probe the Ca<sup>2+</sup> sources implicated in SPP-induced Ca<sup>2+</sup> mobilization, we evaluated, by confocal immunofluorescence, the expression of several known plasma membrane as well as intracellular Ca<sup>2+</sup>

channels in C<sub>2</sub>C<sub>12</sub> myoblastic cells. We found that some cells abundantly expressed L-type Ca<sup>2+</sup> channels in the form of small, granular fluorescence bodies, whereas others displayed virtually no reactivity (Fig. 7, A and B). When present, both the  $\alpha_{1c}$ - and  $\alpha_{1d}$ -subunits were distributed in a diffuse pattern within the cytoplasm and also in close association with the plasma membrane. Ins(1,4,5)P<sub>3</sub>Rs were also markedly expressed and were distributed throughout the cytoplasm with a denser staining in the perinuclear regions, whereas the cell surfaces remained virtually unstained (Fig. 7C). However, not all the cells were stained with the same intensity, and cells negative for Ins(1,4,5)P<sub>3</sub>Rs were also present. In agreement with the lack of effects of ryanodine on the Ca<sup>2+</sup> transients stimulated by SPP, RyRs showed a low and often barely detectable intracellular labeling (data not shown). The latter data strongly suggest that these receptors, which are specific markers for SR Ca<sup>2+</sup> release channels of striated muscle fibers, may develop in later phases of myogenic differentiation.

#### Cytoskeletal Modifications Induced by SPP

We finally examined whether the SPP-dependent effects on intracellular Ca<sup>2+</sup> mobilization were associated with variations in cytoskeletal organization. We first analyzed the organization of the cytoskeletal apparatus in fixed C<sub>2</sub>C<sub>12</sub> cells and found that actin myofilaments represented the main cytoskeletal components in myoblastic cells at this stage of differentiation;




Fig. 7. Localization of L-type channels and Ins(1,4,5)P<sub>3</sub>Rs in C<sub>2</sub>C<sub>12</sub> myoblasts. Confocal immunofluorescence microscopy of C<sub>2</sub>C<sub>12</sub> myoblasts that double-stained either with polyclonal antibodies to the  $\alpha_{1c}$ - and  $\alpha_{1D}$ -subunits of L-type channels (A and B) or with monoclonal antibodies to Ins(1,4,5)P<sub>3</sub>Rs (C) and with tetramethylrhodamine isothiocyanate-labeled phalloidin is shown. A and B: both subunits of L-type channels appear as bright granular bodies scattered in the cytoplasm and in close association with the plasma membrane (brighter dots). Note that some cells (arrows) display lower immunolabeling and even absence of immunolabeling. C: Ins(1,4,5)P<sub>3</sub>R staining also shows a spotlike pattern that is diffuse throughout the cytoplasm, tending to concentrate in the perinuclear regions (arrowheads).

actin filaments were arranged in a weblike structure that invaded all the cytoplasm, anchoring to the plasma membrane, and terminated in typical focal adhesion sites containing vinculin immunostaining (Fig. 8). A less defined reaction was, however, observed for myosin filaments, which appeared as scattered fluorescent small cytoplasmic aggregates (data not shown). These results indicated that the cytoskeletal organization of these cells was quite different from the orderly arrays of myofilaments forming the sarcomeric units of mature skeletal muscle cells. Interestingly, in Ca<sup>2+</sup>-containing medium, SPP stimulated contraction and shortening of living C<sub>2</sub>C<sub>12</sub> cells, but, unexpectedly, this phenomenon did not correlate with the onset of Ca<sup>2+</sup> rise in these cells. In fact, by comparing time-lapse video imaging obtained by DIC with the fluorescence Ca<sup>2+</sup> images, we found that 88.4% of the cells underwent cytoskeletal contraction whereas only 64% displayed Ca<sup>2+</sup> transients in response to SPP stimulation. Moreover, whenever the two phenomena coexisted in the same cell, a clear temporal shift existed between SPP-stimulated cell contractility and SPP-stimulated intracellular Ca<sup>2+</sup> increase (Fig. 9). In fact, cell contraction was a very rapid event, occurring within 3–5 s, whereas the rise in intracellular Ca<sup>2+</sup> became evident

within ~14–35 s (mean value  $24 \pm 1.5$  s; Figs. 1, 2A, and 3C) from SPP stimulation. These data were further confirmed by statistical analyses of the time behaviors of the intracellular spatial Ca<sup>2+</sup> distribution (Fig. 10). By plotting together the time course of the area occupied by fluo 3 fluorescence with the temporal behavior of the total fluo 3 fluorescence, it was found that significant modifications in the fluorescence area (i.e., an initial decrease followed by a return to basal levels, indicating cell contraction and cell relaxation, respectively) occurred before the beginning of Ca<sup>2+</sup> transients. The small increase in the fluorescence signal ( $\Delta F/F = 0.05$ – $0.1$ ) corresponding to the time of cell contraction was probably due to a reduction in the cell volume caused by contraction rather than to a small intracellular calcium elevation.

To confirm that SPP was able to affect the cytoskeletal reorganization, C<sub>2</sub>C<sub>12</sub> cells were preloaded with fluorescent probes for actin. With a time interval quite similar to that observed with DIC video imaging (~5 s), a remarkable reorganization of the actin cytoskeleton could be visualized (Fig. 11). Indeed, small fluorescent dots, representing G-actin short polymers, tended to move coordinately and concentrate toward the nucleus in response to SPP.

## DISCUSSION

In the present study we have shown that SPP is capable of producing intracellular Ca<sup>2+</sup> transients in myoblastic C<sub>2</sub>C<sub>12</sub> cells. In light of our previous findings (28), this response was mediated by the interaction of




Fig. 8. Localization of vinculin and actin in C<sub>2</sub>C<sub>12</sub> myoblasts. Confocal fluorescence microscopy of C<sub>2</sub>C<sub>12</sub> myoblasts double-stained with polyclonal antibodies to vinculin and with tetramethylrhodamine isothiocyanate-labeled phalloidin to reveal actin filaments is shown. A myoblast exhibits a well-developed actin cytoskeleton with stress fibers converging toward focal adhesion complexes that contain vinculin (green staining).






Fig. 9. Differential interference contrast and  $\text{Ca}^{2+}$  fluorescence time-lapse images of C<sub>2</sub>C<sub>12</sub> myoblasts. Images were acquired sequentially according to the indicated times. Two consecutive applications of PBS were made before SPP stimulation to avoid misinterpretation of the data. SPP was added at 65 s from the beginning of the experiment. The images were acquired before SPP addition (A) and 10 (B), 56 (C), and 85 (D) s from the agonist addition. Basal distribution of fluorescence is seen in A; note that the  $\text{Ca}^{2+}$  dye is particularly concentrated in some regions corresponding to the nucleus and perinuclear areas and is absent in the cytoplasmic projections anchoring the cell to the substrate. B: after stimulation with SPP remarkable changes in the cell size (arrowheads), strongly indicative for cell contraction, become visible, whereas  $\text{Ca}^{2+}$  signals are absent, apart from the apparent brighter fluorescence staining, probably due to the redistribution of the fluorescence signal during contraction. C: after contraction the cells return to the original size of A. D:  $\text{Ca}^{2+}$  transient is now clearly visible both in the cytoplasmic and nuclear regions of 1 C<sub>2</sub>C<sub>12</sub> cell. Arrowheads point to identical spots of the same cell in parallel images.




Fig. 10. Statistical evaluation of intracellular  $\text{Ca}^{2+}$  distribution. The SPP-responsive cell shown in Fig. 9 is considered. The solid line represents the normalized area occupied by fluo 3 fluorescence, and the lower dot-dashed line indicates the total increase in intracellular  $\text{Ca}^{2+}$ . Addition of PBS within the first 50 s does not modify the characteristics of the fluorescence distribution. In contrast, after the addition of SPP, a substantial decrease in the fluorescence area (solid line), suggestive of cell contraction, is clearly evident. During the relaxation phase, the fluorescence area returns to the basal levels whereas the  $\text{Ca}^{2+}$  transients become visible.

the bioactive lipid with specific EdgRs. Evidence is reported here that the  $\text{Ca}^{2+}$  response elicited by SPP involved both the cytoplasmic and nuclear compartments as propagated or synchronous waves and required contributions from intracellular and extracellular  $\text{Ca}^{2+}$  sources. In trying to dissect the  $\text{Ca}^{2+}$  signaling pathway we also found that intracellular Ins(1,4,5) $\text{P}_3$ -sensitive  $\text{Ca}^{2+}$  release channels [Ins(1,4,5) $\text{P}_3$ R] and nifedipine-sensitive  $\text{Ca}^{2+}$  channels (undifferentiated/non-voltage-dependent L-type channels) of the plasma membrane were likely involved in the SPP action on myoblasts. Our experimental evidence to support this hypothesis includes the  $\leq 40\%$  reduction of the SPP-mediated  $\text{Ca}^{2+}$  response in cells in the absence of external  $\text{Ca}^{2+}$ ; the requirement of functional PLC cascade and Ins(1,4,5) $\text{P}_3$ R for this response, as evidenced by its reduction up to 60% on pretreatment with inhibitors of Ins(1,4,5) $\text{P}_3$ R, such as heparin, caffeine and 2-APB, or with U-73122, an inhibitor of PLC; the ability of nifedipine, a prototypical blocker of L-type  $\text{Ca}^{2+}$  channels, to affect significantly the  $\text{Ca}^{2+}$  transients elicited by SPP; and, finally, the complete inhibition of the  $\text{Ca}^{2+}$  response in cells pretreated with caffeine and nifedipine in  $\text{Ca}^{2+}$ -containing medium or with caffeine in  $\text{Ca}^{2+}$ -free medium.

Both the cytosolic and nuclear  $\text{Ca}^{2+}$  signals elicited by SPP action were dependent on intracellular  $\text{Ca}^{2+}$




Fig. 11. Dynamic changes of actin cytoskeleton in SPP-stimulated C<sub>2</sub>C<sub>12</sub> myoblasts. The cells were previously loaded with fluorescence-conjugated G-actin monomers. The images were acquired before SPP addition (A) and 9.1 (B) and 32.5 (C) s from agonist stimulation. A substantial redistribution of G-actin polymers, suggestive of activation of cell contractility, is clearly evident after the addition of SPP to the culture medium.

increase, regardless of whether it originated from external or endogenous stores. Indeed, removal of external Ca<sup>2+</sup> and pretreatment with nifedipine or caffeine significantly affected the latency time of the SPP-mediated Ca<sup>2+</sup> response. In these particular circumstances, in fact,  $T_0$  was about twofold that of control myoblasts. A possible explanation of these results may be that Ca<sup>2+</sup> influx, namely through nifedipine-sensitive channels, and Ca<sup>2+</sup> release from the endogenous stores in the cytoplasm, perinuclear, and nuclear regions may be a Ca<sup>2+</sup>-sensitive, autocatalytic processes. Moreover, under these particular experimental conditions the decrease in the number of cells responsive to SPP with respect to that of control conditions may be linked to the expression of different levels of Ca<sup>2+</sup> channels in these cells.

The involvement of Ins(1,4,5)P<sub>3</sub>R and L-type Ca<sup>2+</sup> channels in the SPP-mediated Ca<sup>2+</sup> response in both the cytoplasmic and nuclear compartments was further confirmed by confocal immunofluorescence studies. Ins(1,4,5)P<sub>3</sub>R were, in fact, found throughout the cytoplasm and in association with the nuclear envelope, whereas L-type channels were localized both at the plasma membrane and inside the cytoplasm. However, not all the cells revealed the same degree of fluorescence labeling. SPP did not affect Ca<sup>2+</sup> release through RyR channels in the myoblastic cells, in agreement with the absence of any detectable immunostaining associated with anti-RyR antibody in C<sub>2</sub>C<sub>12</sub> cells and with previous findings on the absence of RyRs in proliferating undifferentiated skeletal muscle cells (1, 23, 25, 27).

The role for Ins(1,4,5)P<sub>3</sub>R in the generation of Ca<sup>2+</sup> signals has been well established in several cell types, including immature and developing skeletal muscle cells (12, 20). In particular, C<sub>2</sub>C<sub>12</sub> myoblasts have been shown to express several isoforms of G protein-coupled SPP receptors (Edg-1, Edg-3, and Edg-5; Ref. 30), and numerous studies on signal transduction have demonstrated that these receptors, activated by SPP, stimulate PLC and, in turn, promote Ins(1,4,5)P<sub>3</sub> generation

(34, 37, 40). In contrast, Ca<sup>2+</sup> influx through L-type Ca<sup>2+</sup> channels in myoblastic cells has not been investigated in depth. Voltage-dependent, dihydropyridine-sensitive L-type Ca<sup>2+</sup> channels are multisubunit transmembrane proteins that allow Ca<sup>2+</sup> influx necessary for the excitation-contraction coupling of cardiac fibers and for modulation of RyR Ca<sup>2+</sup>-release channels of skeletal muscle cells (14). Their expression is developmentally regulated in embryos and in muscle cell lines, because their plasma membrane density sharply increases on muscle differentiation (23, 44). In such a view, the presence of the  $\alpha_1$ -subunit of nifedipine-sensitive L-type Ca<sup>2+</sup> channels in the cytoplasm of C<sub>2</sub>C<sub>12</sub> cells probably denotes that they are on course to be transferred to the cell surface during this stage of myogenic differentiation. Moreover, the presence of these receptors at the cell surface, where no voltage-gated Ca<sup>2+</sup> currents were ever seen, further indicates the existence of "immature" L-type channels in undifferentiated C<sub>2</sub>C<sub>12</sub> myoblasts. Nevertheless, a possible effect of SPP on these channels during muscle cell differentiation should be taken into account in view of the findings that L-type Ca<sup>2+</sup> channels of cardiac muscle cells (21) and developing skeletal muscle cells (10, 46) are modulated by PKC activation, which is a critical step in the signal transduction of exogenous SPP in C<sub>2</sub>C<sub>12</sub> cells (30). Moreover, activation of L-type Ca<sup>2+</sup> channels, which involves phosphorylation of both  $\alpha$ - and  $\beta$ -subunits, may occur whether or not the auxiliary subunits are coexpressed (42). Therefore, it is likely that SPP interacting with Edg cell surface receptors expressed on myoblastic cells triggers a signaling pathway that, through DAG production and PKC activation, targets regulation of immature plasma membrane L-type Ca<sup>2+</sup> channels, thus promoting extracellular Ca<sup>2+</sup> influx in C<sub>2</sub>C<sub>12</sub> cells.

In searching for a possible morphological-functional correlation of the SPP-mediated Ca<sup>2+</sup> transients, we examined whether the Ca<sup>2+</sup> response was associated with corresponding changes in the cytoskeletal organization in the myoblastic cells, particularly in view of

the well-known effects that the sphingolipid has on cytoskeletal remodeling in several cell types (2, 6, 39, 48). Interestingly, we found that SPP promoted cell contractility in the myoblastic cells, but this event did not require intracellular Ca<sup>2+</sup> mobilization. Indeed, cell contraction was a rapid event that occurred early after stimulation, whereas the intracellular Ca<sup>2+</sup> changes become evident only after longer times, thus suggesting that binding of SPP to Edg receptors was able to activate multiple signaling pathways. Indeed, differential coupling to G proteins and effector systems have been shown for Edg-3 and Edg-5 receptors, which are involved in Ca<sup>2+</sup> mobilization and cytoskeletal remodeling, respectively (17, 32, 37, 40). In particular, Edg-5 receptors ligated by SPP stimulate Rho proteins and Rho-dependent kinases have been shown to play an important role in the regulation of smooth muscle and nonmuscle cell contractility by modulating the levels of phosphorylation of myosin light chain (16, 22, 43). Interestingly, in a previous study we demonstrated (29) that RhoA activation also occurs in C<sub>2</sub>C<sub>12</sub> myoblasts after SPP stimulation. In view of that finding and in consideration of the findings reported here that C<sub>2</sub>C<sub>12</sub> cytoskeleton has structural similarities to that of nonmuscle cells, being formed by an extensive network of actin filaments rather than a highly ordered array of myofilaments forming the contractile units of striated fibers, it may be speculated that Ca<sup>2+</sup>-independent/Rho-dependent contraction of the actin cytoskeleton may also occur in undifferentiated skeletal muscle cells. Although not directly addressed here, it seems likely, on the basis of these data, that Ca<sup>2+</sup> transients elicited by exogenous SPP in C<sub>2</sub>C<sub>12</sub> cells may play a role in the myogenic differentiation program rather than in the regulation of cell contractility. This hypothesis would be consistent with an involvement of calcineurin in skeletal muscle differentiation (15) and with recent data showing that C<sub>2</sub>C<sub>12</sub> myoblasts subjected to genetic and metabolic mitochondrial stress release high Ca<sup>2+</sup> transients that are capable of enhancing the expression of RyR-1 and of modifying the activity of several Ca<sup>2+</sup>-dependent transcription factor pathways (7). In keeping with all these findings, it is becoming apparent that Ca<sup>2+</sup> released from the muscle Ins(1,4,5)P<sub>3</sub>Rs may not be significantly involved in muscle cell contraction (20, 47).

In conclusion, the present study, taking advantage of updated techniques and instrumentation, substantially contributes to the understanding of the molecular and functional properties of the Ca<sup>2+</sup> signaling pathway triggered by exogenous SPP in myoblastic cells. It appears that SPP exerts profound biological effects on myoblasts that could have physiological and pathophysiological implications. Indeed, very recently, it was shown that cystic fibrosis transmembrane regulator (CFTR), also expressed in skeletal muscle (13), mediates cellular uptake of SPP, thus attenuating SPP signaling (9). In view of this finding, it is tempting to speculate that alterations of the SPP signaling and skeletal muscle cell response may occur in cystic fibro-

sis, accounting for the peripheral muscle weakness observed in this pathology (11).

The technical assistance of Dr. Alessia Tani and Dr. Ferdinando Paternostro is gratefully acknowledged.

This work was supported by Telethon Grants no. 945 to F. Francini and no. 1086 to P. Bruni and by Ministero dell'Università e della Ricerca Scientifica e Tecnologica grants (Cofin-1999) to S. Zecchi Orlandini, F. Francini, and P. Bruni.

## REFERENCES

1. Airey JA, Baring MD, and Sutko JL. Ryanodine receptor protein is expressed during differentiation in the muscle cell lines BC<sub>3</sub>H1 and C<sub>2</sub>C<sub>12</sub>. *Dev Biol* 148: 365–374, 1991.
2. Ammit AJ, Hastie AT, Edsall LC, Hoffman RK, Amrani Y, Krymskaya VP, Kane SA, Peters SP, Penn RB, Spiegel S, and Panettieri RA. Sphingosine-1-phosphate modulates human airway smooth muscle cell functions that promote inflammation and airway remodeling in asthma. *FASEB J* 15: 1212–1214, 2001.
3. An S, Bleu T, and Zheng Y. Transduction of intracellular calcium signals through G protein-mediated activation of phospholipase C by recombinant sphingosine 1-phosphate receptors. *Mol Pharmacol* 55: 787–794, 1999.
4. Ancellin N and Hla T. Differential pharmacological properties and signal transduction of the sphingosine 1-phosphate receptors EDG-1, EDG-3, and EDG-5. *J Biol Chem* 274: 18997–19002, 1999.
5. Berridge MJ, Lipp P, and Bootman MD. The versatility and universality of calcium signalling. *Nature Rev Mol Cell Biol* 1: 11–21, 2000.
6. Bischoff A, Czyborra P, Fetscher C, Meyer Zu Heringdorf D, Jakobs KH, and Michel MC. Sphingosine-1-phosphate and sphingosylphosphorylcholine constrict renal and mesenteric microvessels in vitro. *Br J Pharmacol* 130: 1871–1877, 2000.
7. Biswas G, Adebajo OA, Freedman FB, Anandatheerthavarada HK, Vijayasarathy C, Zaidi M, Kotlikoff M, and Avadhani NG. Retrograde Ca<sup>2+</sup> signaling in C<sub>2</sub>C<sub>12</sub> skeletal myocytes in response to mitochondrial genetic and metabolic stress: a novel mode of inter-organelle crosstalk. *EMBO J* 18: 522–533, 1999.
8. Bornfeldt KE, Graves LM, Raines EW, Igarashi Y, Wayman G, Yamamura S, Yatomi Y, Sidhu JS, Krebs EG, Hakomori S, and Ross R. Sphingosine-1-phosphate inhibits PDGF-induced chemotaxis of human arterial smooth muscle cells: spatial and temporal modulation of PDGF chemotactic signal transduction. *J Cell Biol* 130: 193–206, 1995.
9. Boujaoude LC, Bradshaw-Wilder C, Mao C, Cohn J, Ogretmen B, Hannun YA, and Obeid L. CFTR regulates uptake of sphingoid base phosphates and LPA. *J Biol Chem* 276: 35258–35264, 2001.
10. Capiati DA, Vazquez G, Tellez Inon MT, and Boland RL. Role of protein kinase C in 1,25(OH)<sub>2</sub>-vitamin D<sub>3</sub> modulation of intracellular calcium during development of skeletal muscle cells in culture. *J Cell Biochem* 77: 200–212, 2000.
11. De Meer K, Gultmans VA, and van Der Laag J. Peripheral muscle weakness and exercise capacity in children with cystic fibrosis. *Am J Respir Care Med* 159: 748–745, 1999.
12. Estrada M, Cardenas C, Liberona JL, Carrasco MA, Mignery GA, Allen PD, and Jaimovich E. Calcium transients in 1B5 myotubes lacking ryanodine receptors are related to inositol trisphosphate receptors. *J Biol Chem* 276: 22868–22874, 2001.
13. Fiedler MA, Nemecek ZK, and Shull GE. Cloning and sequence analysis of rat cystic fibrosis transmembrane conductance regulator. *Am J Physiol Lung Cell Mol Physiol* 262: L779–L784, 1992.
14. Francini F, Bencini C, and Squecco R. Activation of L-type calcium channels in twitch skeletal muscle fibers of the frog. *J Physiol* 494: 121–140, 1996.
15. Friday BB, Horsley V, and Pavlath GK. Calcineurin activity is required for the initiation of skeletal muscle differentiation. *J Cell Biol* 149: 657–665, 2000.



16. Fukata Y, Amano M, and Kaibuchi K. Rho-Rho-kinase pathway in smooth muscle contraction and cytoskeletal reorganization of non-muscle cells. *Trends Pharmacol Sci* 22: 32–39, 2001.
17. Goetzl EJ and An S. Diversity of cellular receptors and functions for the lysophospholipid growth factors lysophosphatidic acid and sphingosine 1-phosphate. *FASEB J* 12: 1589–1598, 1998.
18. Hla T. Sphingosine-1-phosphate receptors. *Prostaglandins* 64: 135–142, 2001.
19. Hla T, Lee Ancellin N M, Liu CH, Thangada S, Thompson BD, and Klux M. Sphingosine-1-phosphate: extracellular or intracellular second messenger? *Biochem Pharmacol* 58: 201–207, 1999.
20. Jaimovich E, Reyes R, Liberona JL, and Powell JA. IP<sub>3</sub> receptors, IP<sub>3</sub> transients, and nucleus-associated Ca<sup>2+</sup> signals in cultured skeletal muscle. *Am J Physiol Cell Physiol* 278: C998–C1010, 2000.
21. Kamp TJ and Hell JW. Regulation of cardiac L-type calcium channels by protein kinase A and protein kinase C. *Circ Res* 87: 1095–1102, 2000.
22. Kimura K, Ito M, Amano M, Chihara K, Fukata Y, Nakafuku M, Yamamuri B, Feng J, Nakano K, Okawa K, Iwamatsu A, and Kaibuchi K. Regulation of myosin phosphatase by Rho and Rho-associated kinase (Rho-kinase). *Science* 273: 245–248, 1996.
23. Liberona JL, Caviedes P, Tascon S, Hidalgo J, Giglio JR, Sampaio SV, Caviedes R, and Jaimovich E. Expression of ion channels during differentiation of a human skeletal muscle cell line. *J Muscle Res Cell Motil* 18: 587–598, 1997.
24. Lorenzon P, Giovannelli A, Ragozzino D, Eusebi F, and Ruzzier F. Spontaneous and repetitive calcium transients in C<sub>2</sub>C<sub>12</sub> mouse myotubes during in vitro myogenesis. *Eur J Neurosci* 9: 800–808, 1997.
25. Lorenzon P, Grohovaz F, and Ruzzier F. Voltage- and ligand-gated ryanodine receptors are functionally separated in developing C<sub>2</sub>C<sub>12</sub> mouse myotubes. *J Physiol* 525: 499–507, 2000.
26. Mao C, Kim SH, Almenoff JS, Rudner XL, Kearney DM, and Kindman LA. Molecular cloning and characterization of SCaM-PCR, a sphingolipid Ca<sup>2+</sup> release-mediating protein from endoplasmic reticulum. *Proc Natl Acad Sci USA* 93: 1993–1996, 1996.
27. McMahon DK, Anderson PAW, Nassar R, Bunting JB, Saba Z, Oakeley AE, and Malouf NN. C<sub>2</sub>C<sub>12</sub> cells: biophysical, biochemical, and immunocytochemical properties. *Am J Physiol Cell Physiol* 266: C1795–C1802, 1994.
28. Meacci E, Cencetti F, Formigli L, Squecco R, Donati C, Tiribilli B, Quercioli F, Zecchi-Orlandini S, Francini F, and Bruni P. Sphingosine 1-phosphate evokes calcium signals in C<sub>2</sub>C<sub>12</sub> myoblasts via EDG3 and EDG5 receptors. *Biochem J* 362: 349–357, 2002.
29. Meacci E, Donati C, Cencetti F, Romiti E, and Bruni P. Permissive role of protein kinase C $\alpha$  but not protein kinase C $\delta$  in sphingosine 1-phosphate-induced RhoA activation in C<sub>2</sub>C<sub>12</sub> myoblasts. *FEBS Lett* 482: 97–101, 2000.
30. Meacci E, Vasta V, Donati C, Farnararo M, and Bruni P. Receptor-mediated activation of phospholipase D by sphingosine 1-phosphate in skeletal muscle C<sub>2</sub>C<sub>12</sub> cells. A role for protein kinase C. *FEBS Lett* 457: 184–188, 1999.
31. Meacci E, Vasta V, Faraoni P, Farnararo M, and Bruni P. Potentiated bradykinin-induced increase of 1,2-diacylglycerol generation and phospholipase D activity in human senescent fibroblasts. *Biochem J* 312: 799–803, 1995.
32. Moolenaar WH, Kranenburg O, Postma FR, and Zondag GCM. Lysophosphatidic acid: G-protein signaling and cellular responses. *Curr Opin Cell Biol* 9: 168–173, 1997.
33. Ogawa Y, Kurebayashi N, and Murayama T. Ryanodine receptor isoforms in excitation-contraction coupling. *Adv Biophys* 36: 27–64, 1999.
34. Okamoto H, Takuwa N, Gonda K, Okazaki H, Chang K, Yatomi Y, Shigematsu H, and Takuwa Y. EDG1 is a functional sphingosine-1-phosphate receptor that is linked via a G<sub>i/o</sub> to multiple signaling pathways, including phospholipase C activation, Ca<sup>2+</sup> mobilization, Ras-mitogen-activated protein kinase activation and adenylate cyclase inhibition. *J Biol Chem* 273: 27104–27110, 1998.
35. Olivera A, Kohama T, Edsall L, Nava V, Cu villier O, Poulton S, and Spiegel S. Sphingosine kinase expression increases intracellular sphingosine-1-phosphate and promotes cell growth and survival. *J Cell Biol* 147: 545–557, 1999.
36. Olivera A and Spiegel S. Sphingosine-1-phosphate as second messenger in cell proliferation induced by PDGF and FCS mitogens. *Nature* 365: 557–560, 1993.
37. Panetti TS, Magnusson MK, Peyruchaud O, Zhang Q, Cooke ME, Sakai T, and Mosher DF. Modulation of cell interactions with extracellular matrix by lysophosphatidic acid and sphingosine 1-phosphate. *Prostaglandins* 64: 93–106, 2001.
38. Perez GJ, Knudson CM, Leykin L, Korsmeyer SJ, and Tilly JL. Apoptosis-associated signaling pathways are required for chemotherapy-mediated female germ cell destruction. *Nat Med* 3: 1228–1232, 1997.
39. Postma FR, Jalink K, Hengeveld T, and Moolenaar WH. Sphingosine-1-phosphate rapidly induces Rho-dependent neurite retraction: action through a specific cell surface receptor. *EMBO J* 15: 2388–2395, 1996.
40. Pyne S and Pyne NJ. Sphingosine 1-phosphate signaling in mammalian cells. *Biochem J* 349: 385–402, 2000.
41. Schneider MF. Control of calcium release in functioning skeletal muscle fibers. *Ann Rev Physiol* 56: 463–484, 1994.
42. Singer-Lahat D, Gershon E, Lotan I, Hullin R, Biel M, Flockerzi V, Hofmann F, and Dascal N. Modulation of cardiac Ca<sup>2+</sup> channels in *Xenopus* oocytes by protein kinase C. *FEBS Lett* 306: 113–118, 1992.
43. Somlyo AP and Somlyo AV. Signal transduction by G-proteins, Rho-kinase and protein phosphatase to smooth muscle and non-muscle myosin II. *J Physiol* 522: 177–185, 2000.
44. Varadi G, Orłowski J, and Schwartz A. Developmental regulation of expression of the  $\alpha$ 1 and  $\alpha$ 2 subunits mRNAs of the voltage-dependent calcium channel in a differentiating myogenic cell line. *FEBS Lett* 250: 515–518, 1989.
45. Vasta V, Meacci E, Romiti E, Farnararo M, and Bruni P. A role for phospholipase D activation in the lipid signaling cascade generated by bradykinin and thrombin in C<sub>2</sub>C<sub>12</sub> myoblasts. *Biochim Biophys Acta* 1391: 280–286, 1998.
46. Vazquez G and de Boland AR. Involvement of protein kinase C in the modulation of 1 $\alpha$ ,25-dihydroxy-vitamin D<sub>3</sub>-induced <sup>45</sup>Ca<sup>2+</sup> uptake in rat and chick cultured myoblasts. *Biochim Biophys Acta* 1310: 157–162, 1996.
47. Walker JW, Somlyo AV, Goldman YE, Somlyo AP, and Trentham DR. Kinetics of smooth and skeletal muscle activation by laser pulse photolysis of caged inositol 1,4,5-trisphosphate. *Nature* 327: 249–252, 1987.
48. Wang F, Nobes CD, Hall A, and Spiegel S. Sphingosine 1-phosphate stimulates rho-mediated tyrosine phosphorylation of focal adhesion kinase and paxillin in Swiss 3T3 fibroblasts. *Biochem J* 324: 481–488, 1997.
49. Weigl LW, Hohenegger M, and Kress HG. Dihydropyridine-induced Ca<sup>2+</sup> release from ryanodine-sensitive Ca<sup>2+</sup> pools in human skeletal muscle cells. *J Physiol* 525: 461–469, 2000.
50. Wu J, Spiegel S, and Sturgill TW. Sphingosine 1-phosphate rapidly activates the mitogen-activated protein kinase pathway by a G-protein dependent mechanism. *J Biol Chem* 270: 11484–11488, 1995.
51. Yaffe D and Saxel O. Serial passaging and differentiation of myogenic cells isolated from dystrophic mouse muscle. *Nature* 270: 725–727, 1977.
52. Yamamura S, Yatomi Y, Ruan F, Sweeney EA, Hakomori S, and Igarashi Y. Sphingosine 1-phosphate regulates melanoma cell motility through a receptor-coupled extracellular action and in a pertussis toxin-insensitive manner. *Biochemistry* 36: 10751–10759, 1997.
53. Young KW, Bootman MD, Channing DR, Lipp P, Maycox PR, Meakin J, Challiss CH, and Nahorski SF. Lysophosphatidic acid-induced Ca<sup>2+</sup> mobilization requires intracellular sphingosine 1-phosphate production. *J Biol Chem* 275: 38532–38539, 2000.
54. Young KW, Challiss RA, Nahorski SR, and MacKrell JJ. Lysophosphatidic acid-mediated Ca<sup>2+</sup> mobilization in human SH-SY5Y neuroblastoma cell is independent of phosphoinositide signaling, but dependent on sphingosine kinase activation. *Biochem J* 343: 45–52, 1999.