



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Plant communities of travertine outcrops of the Saturnia area in southern Tuscany (central Italy).

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Plant communities of travertine outcrops of the Saturnia area in southern Tuscany (central Italy) / Viciani D.; Maffei B.; Selvi F.. - In: HACQUETIA. - ISSN 1581-4661. - ELETTRONICO. - 12:(2013), pp. 141-164. [10.2478/HACQ-2013-0005]

Availability:

The webpage <https://hdl.handle.net/2158/800510> of the repository was last updated on 2018-01-06T19:54:26Z

Published version:

DOI: 10.2478/HACQ-2013-0005

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

PLANT COMMUNITIES OF TRAVERTINE OUTCROPS OF THE SATURNIA AREA IN SOUTHERN TUSCANY (CENTRAL ITALY)

Daniele VICIANI¹, Barbara MAFFEI¹ & Federico SELVI²

Abstract

A phytosociological survey was carried out in a poorly known travertine area of southern Tuscany harbouring a rich vegetation mosaic with chamaephytic garrigues, species-rich xerophytic grasslands, chasmophytic coenoses, annual species-dominated communities, shrublands and thermophilous deciduous forests. Field sampling and data analysis allowed to identify and characterize several community types, some of which of significant interest due to their ecological specificity and rarity in peninsular Italy. In particular, our data confirm the associations *Pistacio terebinthi-Paliuretum spinosae* and *Pistacio terebinthi-Quercetum pubescentis*, respectively a shrub and forest community type previously unknown for Tuscany. In addition, a new therophytic association of travertine debris named *Sedetum hispanico-caespitosi* and placed in the *Hypochoerion achyrophori* alliance (*Brachypodietalia distachyi* order, *Tuberarietea* class) is also described. Finally, dynamic relationships between the vegetation types are highlighted and the presence of conservation priority habitats in the area are pointed out.

Key words: phytosociology, vegetation, travertine outcrops, shrublands, annual *Sedum*-rich communities, *Sedetum hispanico-caespitosi*.

Izveleček

Opravili smo fitocenološko raziskavo na malo znanem območju na travertinu v južni Toskani z bogatim mozaikom vegetacije z hamefitskimi garigami, vrstno bogatimi kserofitskimi travniki, hazmofitskimi združbami, združbami prevladujočih enoletnic, grmišči in termofilnimi listantimi gozdovi. S terenskim vzorčenjem in analizo podatkov smo opisali in označili številne vegetacijske tipe. Nekateri so še posebej pomembni, ker so ekološko posebni in redki na italijanskem poltoku. Potrdili smo prisotnost asociacij *Pistacio terebinthi-Paliuretum spinosae* in *Pistacio terebinthi-Quercetum pubescentis*, grmiščnih in gozdnih združb, ki sta bili do sedaj neznan v Toskani. Opisali smo novo terofitsko asociacijo na travertinskem grušču *Sedetum hispanico-caespitosi* in jo uvrstili v zvezo *Hypochoerion achyrophori* (red *Brachypodietalia distachyi*, razred *Tuberarietea*). Prikazali smo dinamične odnose med vegetacijskimi tipi in izpostavili naravovarstveno prednostne habitate na raziskovanem območju.

Ključne besede: fitosociologija, vegetacija, travertin, grmišča, enoletne združbe z vrstami rodu *Sedum*, *Sedetum hispanico-caespitosi*.

1. INTRODUCTION

Although the ecological and landscape value of southern Tuscany is today widely recognized, plant diversity of this large territory is still incompletely unknown at both the species and community level. One of the interesting sites still waiting for a vegetation study is the travertine area

of Poggio Bagno Santo and Pian di Palma (Manciano) in the middle part of the Albegna river valley, in the internal part of the historical region of Maremma (Grosseto province). This site is partially included in the Natura 2000 network, SCI “Medio corso del Fiume Albegna” (Regione Toscana 2010) and is recognized as one of the important biotopes of the Grosseto territory for its

¹ Laboratory of plant Biology, Department of Biology, University of Florence, via G. La Pira 4, I-50121 Florence, Italy; email: daniele.viciani@unifi.it

² Department of Agricultural, Food and Environmental Sciences, Section of Plant and Soil Science, University of Florence, p. le delle Cascine 28, I-50144 Florence, Italy; email: selvi@unifi.it

floristic richness and peculiar vegetation (Selvi & Stefanini 2005). Such a diversity is largely due to the occurrence of a mosaic including shrublands in the upper parts of the hill, alternated with sparse chasmophytic and lithophytic herb communities on the rocky travertine outcrops and

cliffs, and relatively mature forest communities on the lower part of the slopes. Low garrigues and species-rich herbaceous communities, alternatively dominated by hemicryptophytes and chamaephytes or therophytes in the more xeric conditions, colonize the openings of the shrub-




Figure 1: Geographic location of the studied sites, with magnification of the areas of Poggio Bagno Santo and Pian di Palma.
Figure 1: Geografski položaj proučevanih rastišč s povečanim ombočjem Poggio Bagno Santo in Pian di Palma.

lands and the flat areas of Pian di Palma. Along with natural variation of site conditions, grazing by sheep is a major determinant of this mosaic, as typical in many Mediterranean ecosystems.

Since available vegetation data were still largely incomplete, however, additional fieldwork was carried out to better characterize and classify the communities of this area by means of the phytosociological approach. This work allowed to confirm the presence of forest, shrub and herbaceous communities with a distinct eastern-Mediterranean character, some of which were not previously known from Tuscany. Results are described in this paper to provide a contribution to the botanical knowledge of this region and, possibly, a tool for the management of the more valuable habitats in the biotope.

2. METHODS

2.1 STUDY AREA

2.1.1 Geographical location and geomorphological aspects

Poggio Bagno Santo is a low travertine hill (286 m a.s.l.) located at the northwest of Saturnia in southern Maremma, on the right bank of the Albegna river. The northern slopes are gently connected to the flat travertine pavement of Pian di Palma, while the southern slope is characterized by high and rugged cliffs dominating the Albegna river (Figure 1). In this part the river flows with a marked slope in a narrow gorge between Poggio Bagno Santo and Poggio Pancotta (268 m) and Poggio Saturnia (294 m) on the left bank.

The whole area is composed by recent travertine deposits, mainly originated during the Pleistocene period (Servizio Geologico d'Italia 1965). Travertine is a very porous and permeable sedimentary rock with a chemical and organogenic origin, consisting of calcium carbonate crystals (Press & Siever 1985, Uggeri 2003). Travertine quarries are still active in the area of Pian di Palma, while those on Bagno Santo were abandoned several decades ago.

The area is also known since ancient times for its geothermal activity. Hot water springs are located at the foot of the hills, as the one occurring on the southwestern slope of Bagno Santo after which the hill takes its name.

2.1.2 Climate

Thermopluviometric data can be inferred from the Manciano bioclimatic station, which is located at a higher altitude (443 m a.s.l.) and shows a slightly lower mean annual temperature (13.4 °C) and slightly higher mean annual rainfall (867 mm). According to the climate classification by Thornthwaite & Mather (1957), the climate formula of Manciano is B² b⁴ C₂ (Bigi & Rustici (1984), while in the Rivas-Martinez & Rivas-Saenz (1996–2009) bioclimatic system this site falls in the category “Mediterranean pluviseasonal-oceanic” (Figure 2). Summer drought is therefore relatively pronounced especially in the Bagno Santo area, due to the lower altitude with respect of Manciano and especially the shallow rocky substrate consisting of a very porous rock such as the travertine.




Figure 2: Bioclimatic diagram and diagnosis of the Manciano thermopluviometric station, according to Rivas-Martinez & Rivas-Saenz (1996–2009).

Slika 2: Bioklimatski diagram in analiza podatkov termopluviometrične postaje Manciano v skladu z Rivas-Martinez & Rivas-Saenz (1996–2009).

2.2 DATA COLLECTION AND ANALYSIS

This study is based on the classical Zurich-Montpellier approach (Braun-Blanquet, 1932; 1964), including his recent developments (Westhoff & Van Der Maarel, 1973; Géhu & Rivas-Martínez, 1981; Biondi, 2011; Blasi et al., 2011; Pott, 2011). After a preliminary study of the local flora in the years 2005–2008, we carried out 24 phytosociological relevés between late April and early May of the years 2008–2010, in order to sample the most representative community types.

Field data were first grouped in a raw table that was explored using cluster analysis as implemented in PAST software (Hammer et al. 2001). Abundance-dominance values were transformed according to Van Der Maarel (1979) and Noest et al. (1989), before calculating the Euclidean Distance matrix between the relevés and submit it to hierarchical clustering using the UPGMA algorithm. Plant names follow basically Tutin et al. (1964–80, 1993) and Pignatti (1982), with some later modifications as reported in the Checklist of the Italian Vascular Flora (Conti et al. 2005, 2007) and the recent checklist of the flora of Tuscan Maremma (Selvi 2010). Syntaxonomical nomenclature above the association level is in line with the Phytosociological Nomenclature Code (Weber et al. 2000) and largely based on Rivas-Martínez et al. (2002).

3. RESULTS AND DISCUSSION

Cluster analysis resulted in the dendrogram shown in Figure 3. Five physiognomically consistent groups could be recognized: herb communi-

ties of flat travertine pavements (group 1), small herb communities of cliffs and rocks (group 2), garrigues and grasslands (group 3), open woods and shrublands (group 4) and forests (group 5). Each of these groups is described below.

3.1 ANNUAL-DOMINATED COMMUNITIES OF FLAT TRAVERTINE PAVEMENTS

The relevés of group no. 1 in the dendrogram are shown in Table 1. They describe sparse communities dominated by small Mediterranean annuals colonizing the flat surfaces of travertine debris, with a thin layer of organic soil and abundant mosses. Dominating species are the succulent *Sedum caespitosum* and *Sedum hispanicum*, along with *Plantago lagopus*, *Saxifraga tridactylites*, *Arenaria leptoclados*, *Cerastium brachypetalum* s.l., *Minuartia hybrida*, *Trifolium tomentosum*, *Trifolium nigrescens* and the grasses *Poa bulbosa*, *Catapodium rigidum* and *Vulpia ciliata*.

Selvi & Stefanini (2005) provisionally referred these communities to the *Alyso-Sedion albi*, that correspond to the EU priority habitat “Rupicolous calcareous or basophilic grasslands of the *Alyso-Sedion albi*” (Natura 2000 code: 6110; European Commission, 2007). According to the authors who described this alliance (Oberdorfer 1978), however, the vegetation of this *syntaxon* is characterized by abundant *Sedum album* and/or perennial species of *Sedum*, with also other annual species. Several authors have reported vegetation types of the *Alyso-Sedion albi* from central Italy, such as Venanzoni & Gigante (1999) from Umbria, Scoppola & Angiolini (2001) and Angiolini et al. (2009) from southern Tuscany, Di Pietro et al. (2006) from the Prenestini mountains around Rome, Foggi & Venturi (2009) from the Prato province in northern Tuscany, and Lastucci et al. (2009) from eastern Tuscany. These authors describe associations or groupings usually dominated by perennial species of *Sedum*. Recently Fanelli (2007) described the association *Chaenorhino rubrifolii-Saxifragetum tridactylites* from the travertine area of Bagni di Tivoli near Rome and classified it in the *Alyso-Sedion albi* al-




Figure 3: Dendrogram from UPGMA cluster analysis of the vegetation relevés.

Slika 3: Dendrogram UPGMA klusterske analize vegetacijskih popisov.

liance in spite of the large prevalence of therophytes. Even more recently, Fanelli et al. (2010) recognized that several European authors exclude the annual-dominated communities from the alliance *Alyso-Sedion albi* (e.g. De Foucault 1999, Rivas-Martinez et al. 1997, 2002), though finally following the same classification of the *Chaenorhino rubrifolii-Saxifragetum tridactylites* association.

Due to the dominance of annual species and the nearly complete lack of perennial species of *Sedum* in our relevés, we therefore refer the examined communities to the class of the terophytic Mediterranean meadows. This class has been subject to contrasting nomenclatural interpretations (Braun-Blanquet et al. 1952; De Foucault 1999; Brullo et al. 2001; Fanelli et al. 2010) and indicated as *Thero-Brachypodietea*, *Tuberarietea/Helianthemetea*, *Stipo-Trachynietea* and other names. According to the classification followed here (Rivas-Martinez et al. 2002), this syntaxon is called *Tuberarietea*.

Phytosociological works on carbonatic substrates in the temperate-oceanic areas of Latium refer similar annual-dominated vegetation types to the associations *Trifolio scabri-Hypochoeridetum achyrophori* and *Saxifrago-Hypochoeridetum achyrophori* (Scoppola et al. 2001). These associations include several species also found in our relevés but differ substantially in the dominant species, and cannot be used to classify the Bagno Santo community.

In a recent paper, Fanelli et al. (2010) discussed the syntaxonomy of the basophilous nanophanerophytic vegetation in central Italy based on above 100 phytosociological relevés, both published or unpublished. They suggest that this vegetation type, usually included in the single association *Trifolio scabri-Hypochoeridetum achyrophori*, is characterized by wide floristic variations and should be split in at least five distinct associations (*Medicagini rigidulae-Trifolietum scabri*, *Trifolio scabri-Hypochoeridetum achyrophori*, *Hippocrepido siliquosae-Brachypodietum distachyi*, *Trigonello gladiatae-Brachypodietum distachyi*, *Crucianello latifoliae-Hypochoeridetum achyrophori*). In their large general table, however, only three relevés include *Sedum caespitosum* and *S. hispanicum*. These relevés, which are anyway dominated by other species, represent the meadow vegetation on limestone (not travertine) areas of M. Castagneto-La Spera near Rieti and had already been discussed by Scoppola & Angiolini (2001).

None of the nanophanerophytic associations described to date is therefore physiognomically dominated by *Sedum caespitosum* and characterized by *S. hispanicum*. The former is a stenomediterranean species with a discontinuous distribution in Central and Southern Italy (Pignatti, 1982), whereas the latter is a more widely distributed Southeastern European taxon relatively un-frequent in Tuscany. Accordingly, these community can be referred to the new association *Sedetum hispanico-caespitosi* ass. nova hoc loco (typus rel. 20, Table 1). Besides the two *Sedum* species, constant differential species of this association are *Plantago lagopus* and *Saxifraga tridactylites*, as well as several others of higher syntaxonomical rank such as *Arenaria leptoclados*, *Catapodium rigidum*, etc. or typical of annual Mediterranean meadows (*Anthemis arvensis*, *Poa bulbosa*, etc.).

Concerning the higher ranks, Biondi & Guerra (2008) recently analyzed the syntaxonomy of the Italian annual meadows and described the endemic alliance *Hypochoerion achyrophori*. Following this classification, the new association can be placed in the northern and central Italian sub-alliance *Hypochoerion achyrophori*. Alternative placements would have been in the *Sedion caespitosi* suballiance according to Rivas-Martinez et al. (2002), or in the *Trachynion distachyae* alliance according to Fanelli et al. (2010).

3.2 CHASMOPHYTIC AND COMOPHYTIC VEGETATION OF STEEP ROCKS AND CLIFF EDGES

Another small group consists of two relevés (Table 2) representing the community type colonizing the steep and rugged travertine rocks (10–30° inclination) and the edge of cliffs over the Albegna river on Poggio Bagno Santo. This community includes either chasmophytic or comophytic species (plants growing in small soil pockets found in rock fissures), often developed in partial shade by forest and shrub vegetation surrounding such rocky surfaces. Dominant species are the hemicryptophyte *Cymbalaria muralis* (casmophytic) and the geophytes *Umbilicus rupestris*, *Muscari commutatum* and *Allium subhirsutum* (comophytic). Though not recorded in the relevés presented here, *Anthirrhinum latifolium* and *Sedum rupestre* are often found in these phytocoenoses, as well as the xerophytic ferns *Polypodium cambricum* and *Asplenium trichomanes*. Other species appear in

the fissures of the partially shaded rocks containing pockets of nutrient-rich soil from the mineralization of forest litterfall, such as the therophytes *Geranium purpureum*, a species of dry forest edges, *Galium aparine*, *Centranthus calcitrapa*, *Arabis turrita* and *Parietaria officinalis*.

From the floristic point of view it is especially worth of note the presence of a conspicuous population of *Muscari commutatum* (*Liliaceae*) in the narrow fissures of travertine rocks of Bagno Santo, due the rarity of this bulbous geophyte in Italy. As a rare Mediterranean *taxon* it is included in the national list of the threatened plants and as a relevant species for the recognition of the IPAs in the country (Blasi et al. 2010).

Concerning the syntaxonomical classification of this type of vegetation we refer to Brullo & Guarino (2002), who published a synthetic treatment of the chasmophytic synantropic vegetation in Italy. Unlike in previous works (cited in Brullo & Guarino 1999), these authors place the phytocoenoses of natural cliffs and rock ledges and those of walls in two distinct classes, respectively *Asplenetea trichomanis* and *Parietarietea judaicae*. Among the characteristic species of the latter class, our samples include *Cymbalaria muralis* and *Umbilicus rupestris*, along with several other rupicolous plants. The presence of the somewhat nitrophilous species *Parietaria officinalis*, instead of *P. judaica*, indicates the presence of local deposits of nutrient-rich soil connected to forest edges.

Accordingly we provisionally refer the relevés of Table 2 to an informal grouping with *Cymbalaria muralis*, *Muscari commutatum* and *Allium subhirsutum* within the *Parietarietea* class.

3.3 CHAMAEPHYTIC GARRIGUES WITH *EUPHORBIA SPINOSA* AND *SATUREJA* *MONTANA*

The travertine plateau of Pian di Palma is characterized by a low garrigue community dominated by the two Mediterranean chamaephytes *Euphorbia spinosa* and *Satureja montana* (Table 3). Soil cover resulted relatively high also due to the abundance of other shrublets such as *Teucrium capitatum* and *Helichrysum italicum* and robust herbs and grasses such as *Cephalaria leucantha*, *Bromus erectus*, *Convolvulus cantabrica* and *Thymus longicaulis*. On the whole, these species indicate shallow carbonatic lithosoils with little

organic material, under very slow pedogenetic evolution.

Satureja montana is a widespread Mediterranean species of mountain areas. Along the whole Apennine chain it is often dominant in the garrigues developed on various carbonatic substrates, even at relatively high altitudes. *Euphorbia spinosa* is also a basophilous species often abundant in limestone and serpentine garrigues, but more thermophilous and usually at lower altitudes than *S. montana* (Pignatti 1982, Allegrezza et al. 1997). According to Allegrezza et al. (1997), the rocky habitats of the limestone hills in central Italy harbour distinct associations with *Satureja montana*, some of which show floristic and ecological similarity with the Pian di Palma travertine community. In particular, the more thermophilous aspect of the *Cephalario leucanthae-Saturejetum montanae*, described as subass. *euphorbietosum spinosae*, is the community type that better corresponds to our vegetation samples for the cenological conditions and the presence of several characteristic species. The same association was also reported from some travertine areas between Siena and Viterbo (Scoppola & Angiolini 1997), but in a distinct variant characterized by the endemic *Santolina etrusca* that is lacking at Pian di Palma because of the carbonatic lithosoil.

According to Scoppola & Angiolini (1997), the *Cephalario leucanthae-Saturejetum montanae* association belongs to the alliance *Artemisio albae-Saturejion montanae* of the Mediterranean *Rosmarinetea* class, although the number of Mediterranean species decreases going towards the more inland areas and the communities become progressively more atypical for the class.

3.4 MESOXEROPHILOUS *BROMUS ERECTUS* GRASSLANDS

A second community type (Table 4) found on the travertine plateau of Pian di Palma occurs on slightly more evolved soils with a higher content of organic matter. Here, soil cover by xerophytic perennial herbs and grasses reaches 70–80%, while chamaephytes are less abundant than in the *Cephalario-Saturejetum*. Most typical species of this community are *Bromus erectus* and other hemicryptophytes such as *Koeleria splendens*, *Eryngium campestre*, *Convolvulus cantabrica*, *Cephalaria leucantha*, *Urospermum dalechampii*, *Galium corrudifolium*, *Potentilla hirta*, *Tragopogon dubius*,

Carlina corymbosa, *Anthyllis vulneraria* subsp. *rubriflora*, etc. The mainly W Mediterranean geophytes *Allium moschatum* and *Iris lutescens* are also relatively common, as well as the orchids *Orchis morio*, *Serapias vomeracea* and others observed outside the vegetation relevés. Annual species were not numerous, among them *Helianthemum salicifolium* and *Tordylium apulum*.

The phytosociological classification of the perennial grasslands of the *Brometalia erecti* order in the Apennines has been revised by Biondi et al. (1995, 2005), though recent studies suggest to place these communities in a separate *syntaxon* at the order level due to their floristic and ecological distinctness (Di Pietro 2011, Di Pietro et al. 2012). According to the recent study by Biondi & Galdenzi (2012), the order *Brometalia erecti* is restricted to north-western Italy (up to western Tuscany), while most Apennine communities belong to the *Scozonero villosae-Chrysopogonetalia grylli*.

Our vegetation samples include characteristic and differential species of the order *Brometalia erecti*, such as *Bromus erectus*, *Thymus longicaulis*, *Anthyllis vulneraria* ssp. *rubriflora*, *Eryngium campestre*, *Koeleria splendens*. The presence of also *Convolvulus cantabrica* and *Satureja montana* of the suborder *Artemisio albae-Bromenalia erecti* suggest to refer the Pian di Palma community to this *syntaxon*, that also fits in terms of edaphic and bioclimatic features. Classification at the association level is more difficult. Angiolini et al. (2003) referred the *Bromus erectus* grasslands from the calcareous massifs in southern Tuscany to two associations: *i*) the more xeric *Cerastio etrusci-Brometum erecti* to *Phleo-Bromion* alliance; *ii*) *Trifolio incarnati-Brometum erecti*, more mesophilous and rich in species of the *Molinio-Arrhenatheretea* class, attributed to *Bromion* alliance. However, floristic and ecological differences seem too large to refer our relevés to one of these associations, especially to the more mesic one. Also due to their small number, the present relevés can only be provisionally referred to a generic *Bromus erectus* grouping of xerophytic type within the *Artemisio albae-Bromenalia erecti*. These communities are likely to represent a dynamic stage of the edaphoxerophilous series leading to the supramediterranean *Quercus pubescens* forest. This type of series with *Bromus erectus* as an intermediate stage is also reported by Scoppola & Angiolini (1997) for other travertine garrigues between Siena and Viterbo.

3.5 OPEN SCRUB AND SHRUBLANDS

The shrub vegetation colonizing the rocky travertine slopes and summit area of Bagno Santo is dominated by mainly eastern Mediterranean species such as *Pistacia terebinthus* and *Paliurus spina-christi*, mixed with a few evergreen sclerophyllous species, especially *Phillyrea latifolia* and *Rhamnus alaternus*. *Paliurus spina-christi* is a vigorously resprouting spiny shrub favoured by sheep grazing and therefore locally dominant. Together with *Cercis siliquastrum*, *Pistacia terebinthus* and *Carpinus orientalis* this plant participates to distinct *syntaxa* of the Illyrian shrublands found in the coastal and subcoastal sectors of the Balkans, and commonly indicated as “Sibljak” (Blasi & Di Pietro 2002). All of these species are well represented in our relevés (Table 5), with the only exception of *Carpinus orientalis*. Though relatively widespread in the thermophilous woodlands of Latium and most regions of central and southern Italy, this tree is still to be confirmed in the Tuscan Maremma (Selvi 2010). Other abundant tree species in our samples are *Fraxinus ornus*, *Acer monspessulanum* and *Quercus pubescens*, all frequent on especially carbonatic substrates of the supramediterranean hill belt of southern Tuscany. The pubescent oak plays a key role in the dynamics of this vegetation, in most cases becoming increasingly dominant and causing the progressive transformation of the *Paliurus* shrubland into thermophilous oak forests, especially where sheep grazing is not longer present.

Numerous species of openings and meadows occur in the herbaceous layer of this community (*Bromus erectus*, *Allium subhirsutum*, *Carex caryophylla*, etc.), due to the sparse soil cover by the woody species. Well developed herbaceous communities are found in the openings of the *Paliurus* shrubland, where ruderal and somewhat nitrophilous perennials of synanthropic vegetation such as *Calamintha nepeta* and *Marrubium incanum* are often dominant. Annual weeds of the *Stellarietea* class also occur in these phytocoenoses, together with some other therophytes of the travertine pavements.

In Tuscany, *Paliurus spina-christi* is relatively widespread in the shrublands in the warmer and drier sites of the southern parts of the region, especially in the more inland areas. However, the phytosociological affinities of the *Paliurus* vegetation has been scarcely studied. According to Angiolini et al. (2006), a dense shrub vegetation with *Paliurus* and stenomediterranean species such as

Phillyrea latifolia, *Rosa sempervirens* and *Clematis flammula* occurs in the Trasubbie river bed, a site of Regional Importance in the central-eastern part of the Grosseto province. However, this vegetation is only recognized as a generic grouping with no syntaxonomical position. On the other hands it differs from the Bagno Santo community in the lack of *Pistacia terebinthus* and *Cercis siliquastrum*.

In the central Tyrrhenian region (Latium), the shrublands with *Paliurus spina-christi* have been referred to two syntaxa of the *Pruno-Rubenion* suballiance: the *Pistacio terebinthi-Paliuretum* association from southern Latium (Blasi & Di Pietro 2002) and the subassociation *paliuretosum spina-christi* of *Lonicero etruscae-Rosetum sempervirentis*. The latter represents dense and structured shrub formations in the hill belt of northern Latium, mainly on clayey substrates (Cutini et al. 1996, Blasi et al. 2002). Although the *Pistacio terebinthi-Paliuretum spinae-christi* has never been recorded from Tuscany, the vegetation represented by our samples show clear floristic and ecologic affinities to this association. In particular, the frequency of the characteristic shrub species allows to refer the Bagno Santo community to this *syntaxon*, despite the lack of *Carpinus orientalis*.

3.6 THERMOPHILOUS DECIDUOUS FORESTS

The slopes of Bagno Santo are covered by a mature and relatively undisturbed broad-leaf woodland with tall trees and rich understory vegetation (Selvi & Stefanini 2005).

Based on our relevés (Table 6) this thermophilous forest community is dominated by *Quercus pubescens* and *Acer monspessulanum*, with abundant *Fraxinus ornus* and, to lesser extent, the relatively more mesophilous *Ostrya carpinifolia*. Of particular interest is the presence of mainly eastern Mediterranean tree species that are unfrequent in Tuscany, such as *Cercis siliquastrum*, *Celtis australis* and *Pistacia terebinthus*. The latter occurs with aged, single-stem individuals of remarkable size, that reach the upper layers of the forest canopy. The shrub layer is rich in *Paliurus spina-christi*, while the herbaceous understory includes *Ruscus aculeatus*, *Hedera helix*, *Tamus communis* and some species linked to relatively rich and mesic forest soils, such as *Melica uniflora* and *Brachypodium sylvaticum*. A relatively good level of nutrients in the soil is indicated by slightly nitrophilous species that colonize the fissures of

the rocks rich in organic forest soil, such as *Parietaria officinalis* and *Geranium robertianum*.

The syntaxonomy of the thermophilous deciduous forests in the Italian peninsula is still subject to different treatments, because of the difficult interpretation of the effects of the orographic, geopedologic and bioclimatic variations of this area and the strong impact of human activities since millennia (Blasi et al. 2004). Therefore, the Apennine deciduous forests have been classified in more or less contrasting ways by different authors (Ubaldi et al. 1987, Scoppola et al. 1995, Arrigoni 1998, Pignatti 1998, Biondi et al. 2002, 2010, Ubaldi 2003, Blasi et al. 2004, Di Pietro et al. 2010). In this study, we refer to the treatment of the order *Quercetalia pubescenti-petraeae* in the Italian peninsula by Blasi et al. (2004). Among the five alliances proposed in this work, that of *Carpinion orientalis* is the one that best matches with our relevés in terms of floristic composition. This grouping is widely distributed in the suboceanic areas of peninsular Italy (meso-mediterranean to upper submontane thermotypes), mainly on carbonatic soils. Four suballiances have been recognized: *Laburno-Ostryenion*, *Lauro-Quercenion pubescentis*, *Cytiso sessilifolii-Quercenion pubescentis*, *Campanulo mediae-Ostryenion*. The more thermophilous woodlands of the mid-mesomediterranean or lower hill belt are included in the *Lauro-Quercenion pubescentis*, characterized and usually dominated by *Carpinus orientalis*, *Quercus pubescens*, *Acer monspessulanum*, *Fraxinus ornus*, *Pistacia terebinthus* and *Cercis siliquastrum*. All these species are well represented in our relevés from Bagno Santo (Table 6), with the only exception of *Carpinus orientalis* as already observed for the shrubland communities described above. In the synoptic table of the *Lauro-Quercenion pubescentis* by Blasi et al. (2004), the associations floristically closer to our samples are *Clematido flammulae-Quercetum pubescentis* (Ubaldi et al. 1993), *Roso sempervirentis-Quercetum pubescentis carpinetosum orientalis* (Blasi & Di Pietro 1998) and *Lonicero etruscae-Carpinetum orientalis* (Blasi et al. 2001).

Although *Clematido flammulae-Quercetum pubescentis* shares several species with our samples it includes a group of more mesophilous species usually found at higher elevations (*Chamaecytisus hirsutus*, *Cytisus sessilifolius*) and should therefore be excluded. *Lonicero etruscae-Carpinetum orientalis* is dominated by *Carpinus orientalis* and *Ostrya carpinifolia*, the former absent and the latter scarcely represented in our area; *Quercus pubescens* has a rela-

tively secondary role. Furthermore one of the most important characteristic species of this association, *Sesleria autumnalis*, is not present in our study area.

Scoppola & Angiolini (2001) had already reported the occurrence of woodlands close to the *Roso sempervirentis-Quercetum pubescentis* from the travertine outcrops in the western sector of central Italy. In particular, the floristic assemblage of the Bagno Santo forest matches that of the subassociation *carpinetosum orientalis* described from the Ausoni and Aurunci mountains in southern Latium (Blasi & Di Pietro 1998). The classification problem is further complicated by the studies of Allegrezza et al. (2002) and Biondi et al. (2010). Allegrezza et al. (2002) re-analyzed the relevés of Blasi & Di Pietro (1998) and stated that *Roso sempervirentis-Quercetum pubescentis carpinetosum orientalis* has to be treated as a distinct association named *Pistacio terebinthi-Quercetum pubescentis*; this new association was divided in two subassociations, of which the more thermophilous one was named *rosetosum sempervirentis*. Our relevés differ from this *syntaxon* by the lack of *Carpinus orientalis*, the presence of *Celtis australis* and *Paliurus spina-christi*, and the higher cover of *Acer monspesulanum* (Table 6). However, the ecological similarities and the presence of important differential species of subass. *rosetosum sempervirentis* such as *Cercis siliquastrum* and *Rosa sempervirens* allows to refer to the Bagno Santo forest to this *syntaxon*.

Finally, a taxonomical problem with potential consequences on the syntaxonomy of this forest vegetation concerns the identity of the oak species upon which *Roso sempervirentis-Quercetum* and *Lauro-Quercenion* are based. According to Biondi

et al. (2010) the oak originally found in the communities of these *syntaxa* was not *Quercus pubescens* s.s. but *Q. virgiliana* Ten. In this case, the association and subassociation names found in the literature should be used only in a broad sense. *Quercus virgiliana* is a weakly differentiated species still not reported from Tuscany, even in recent floristic checklists focusing on the Tuscan Maremma (Selvi, 2010). At present there are no sufficient elements to confirm its presence in the Saturnia area, but this is possible and more detailed observations are needed to bring more light on this issue.

4. CONCLUSION

This study describes the vegetation types of a poorly known area of southern Tuscany, some of which of significant interest due to their ecological specificity and rarity in peninsular Italy. In particular, our data confirm the presence in Tuscany of shrub communities of the *Pistacio-Paliuretum*, an association previously known only from southern Latium, and allow to characterize a new therophytic association of travertine outcrops, named *Sedetum hispanico-caespitosi*. Besides the ecological and floristic features of the communities described, field relevés have also allowed to interpret their dynamic relationships, as schematically shown in the transect of Figure 4.




Figure 4: Schematic transects of the vegetation dynamic stages on the travertine outcrops of Poggio Bagno Santo and Pian di Palma: 1 = *Sedetum hispanico-caespitosi* ass. nova; 2 = *Bromus erectus*-dominated conenoses; 3 = *Cephalario leucanthae-Saturejetum montanae euphorbietosum spinosae*; 4 = *Pistacio terebinthi-Paliuretum spinae-christi*; 5 = *Calamintha nepeta* and *Marrubium incanum* conenoses; 6 = *Cymbalaria muralis*, *Muscari commutatum* and *Allium subhirsutum* conenoses; 7 = *Pistacio terebinthi-Quercetum pubescentis rosetosum sempervirentis*.

Slika 4: Shema transekta dinamike vegetacijskih tipova na travertinu na območju Poggio Bagno Santo in Pian di Palma: 1 = *Sedetum hispanico-caespitosi* ass. nova; 2 = združba z vrsto *Bromus erectus*; 3 = *Cephalario leucanthae-Saturejetum montanae euphorbietosum spinosae*; 4 = *Pistacio terebinthi-Paliuretum spinae-christi*; 5 = združbi z vrstama *Calamintha nepeta* in *Marrubium incanum*; 6 = združbe z vrstami *Cymbalaria muralis*, *Muscari commutatum* in *Allium subhirsutum*; 7 = *Pistacio terebinthi-Quercetum pubescentis rosetosum sempervirentis*.

Based on the Habitat Directive 92/43/CE, the European Interpretation Manual (European Commission, 2007) and the Italian Manual (Biondi & Blasi 2009), our data do not support the presence of typical aspects of the priority habitat *Alyso-Sedion albi* (Rupicolous calcareous or basophilic grasslands, Natura 2000 code 6110), while they confirm the priority habitats *Festuco-Brometalia* (Semi-natural dry grasslands and scrubland *facies* on calcareous substrates, Natura 2000 code 6210) and *Thero-Brachypodietea* (Pseudo-steppe with grasses and annuals, Natura 2000 code 6220).

Syntaxonomic scheme

Tuberarietea guttatae (Br.-Bl. in Br.-Bl., Roussine & Nègre 1952) Rivas Goday & Rivas-Martínez 1963

Brachypodietalia distachyi Rivas-Martínez 1978

Hypochoerion achyrophori Biondi & Guerra 2008

Hypochoerenion achyrophori Biondi & Guerra 2008

Sedetum hispanico-caespitosi ass. nova hoc loco

Parietarietea judaicae Rivas-Martínez ex Rivas Goday 1964

Gymnalaria muralis, *Muscari commutatum* and *Alilium subhirsutum* coenoses

Rosmarinetea officinalis Rivas-Martínez, T.E. Diaz, F. Prieto, Loidi & Penas 1991

Rosmarinetalia officinalis Br.-Bl. ex Molinier 1934

Artemisio albae-Saturejon montanae Allegrezza, Biondi, Formica & Ballelli 1997

Cephalario leucanthae-Saturejetum montanae Allegrezza, Biondi, Formica & Ballelli 1997

euphorbietosum spinosae Allegrezza, Biondi, Formica & Ballelli 1997

Festuco-Brometea Br.-Bl. & Tx. ex Br.-Bl. 1949

Brometalia erecti Br.-Bl. 1936

Artemisio albae-Bromenalia erecti Biondi, Blasi ex Biondi, Ballelli, Allegrezza, Zuccarello 1995

Bromus erectus coenoses

Rhamno-Prunetea Rivas Goday & Borja ex Tuxen 1962

Prunetalia spinosae Tuxen 1952

Pruno-Rubion ulmifolii Bolòs 1954

Pruno-Rubenion ulmifolii Arnaiz & Loidi 1983

Pistacio terebinthi-Paliuretum spinae-christi Blasi & Di Pietro 2001

Quercus-Fagetea Br.-Bl. & Vlieg. in Vlieg. 1937

Quercetalia pubescenti-petreae Klika 1933 corr. Blasi, Di Pietro & Filesi 2004

Carpinion orientalis Horvat 1958

Lauro nobilis-Quercenion virgiliana Ubaldi 1995 corr. Biondi, Casavecchia & Pesaresi 2010

Pistacio terebinthi-Quercetum pubescentis Allegrezza, Baldoni, Biondi, Taffetani & Zuccarello 2002

rosetosum sempervirentis Allegrezza, Baldoni, Biondi, Taffetani & Zuccarello 2002

Other syntaxa quoted in text and tables

Alyso-Sedion albi Oberdorfer et Müller in Müller 1961

Asplenietea trichomanis (Br.-Bl. in Meier & Br.-Bl. 1934) Oberdorfer 1977

Campanulo mediae-Ostryenion Ubaldi 1995

Cerastio etrusci-Brometum erecti Angiolini, Riccucci & De Dominicis 2003

Chaenorhino rubrifolii-Saxifragetum tridactylites Fanelli 2007

Clematido flammulae-Quercetum pubescentis Ubaldi, Zanotti & Puppi 1993

Crucianello latifoliae-Hypochoeridetum achyrophori Filesi, Blasi & Di Marzio 1996

Cytiso sessilifolii-Quercenion pubescentis Ubaldi 1995

Galio-Urticetea Passarge ex Kopecký 1969

Hippocrepido siliquosae-Brachypodietum distachyi Fanelli, Bianco, De Sanctis & Serafini Sauli 2010

Laburno-Ostryenion (Ubaldi 1995) Blasi, Di Pietro & Filesi 2004

Lonicero etruscae-Carpinetum orientalis Blasi, Di Pietro, Filesi & Fortini 2001

Lonicero etruscae-Rosetum sempervirentis Cutini, Fabozzi, Fortini, Armanini & Blasi 1996 *paliuretsum spina-christi* Blasi, Cutini, Di Pietro & Fortini 2002

Lygeo-Stipetea Rivas-Martínez 1978

Medicagini rigidulae-Trifolietum scabri Fanelli, Bianco, De Sanctis & Serafini Sauli 2010 Tüxen

Phleo-Bromion Biondi & Blasi ex Biondi et al. 1995

Poetea bulbosae Rivas Goday & Rivas-Martínez in Rivas-Martínez 1978

Quercetalia ilicis Br.-Bl. ex Molinier 1934 em. Rivas-Martínez 1975

Quercetea ilicis Br.-Bl. ex A. & O. Bolòs 1950

Roso sempervirentis-Quercetum pubescentis Biondi 1986 *carpinetosum orientalis* Blasi & Di Pietro 1998

Saxifrago-Hypochoeridetum achyrophori Biondi, Izco, Ballelli & Formica 1997

Sedenion caespitosi Rivas-Martínez 1978
Stellarietea mediae (Br.-Bl. 1931) Tüxen, Lohmeyer
 & Preising. in Tüxen 1950
Stipo-Trachynietea Brullo 2001
Thero-Brachypodietea Br.-Bl. ex A. Bolos & Bolos
 in A. Bolos 1950
Trachynion distachyae Rivas-Martínez 1978 *Trifolio-
 Geranietea sanguinei* Th. Müller 1961
Trifolio scabri-Hypochoeridetum achyrophori Bion-
 di, Izco, Ballelli & Formica 1997
Trigonello gladiatae-Brachypodietum distachyi Fa-
 nelli, Bianco, De Sanctis, Serafini Sauli, 2010

5. REFERENCES

- Allegrezza, M., Baldoni, M., Biondi, E., Taffetani, F. & Zuccarello, V. 2002: Studio fitosociologico dei boschi a *Quercus pubescens* s.l. delle Marche e delle zone contigue dell'Appennino centro-settentrionale (Italia centrale). *Fitosociologia* 39 (1): 161–171.
- Allegrezza, M., Biondi, E., Formica, E. & Ballelli, S. 1997: La vegetazione dei settori rupestri calcarei dell'Italia centrale. *Fitosociologia* 32: 91–120.
- Angiolini, C., Gabellini, A., Frignani, F., Giallonardo, T. & De Dominicis, V. 2009: Contributo alla conoscenza della vegetazione della Riserva naturale provinciale "Monte Penna" (Grosseto, Toscana meridionale) *Atti Soc. tosc. Sci. nat., Mem., Ser. B*, 115 (2008): 1–20.
- Angiolini, C., Landi, M., Boddi, M. & Frignani, F. 2006: La vegetazione dell'alveo fluviale del sito d'importanza regionale torrente Trasubbie (Grosseto, Toscana meridionale). *Atti Soc. tosc. Sci. nat., Mem., Ser. B*, 112 (2005): 127–151.
- Angiolini, C., Riccucci, C. & De Dominicis V. 2003: *Brometalia erecti* Br.-Bl. 1936 grasslands from Antiappennine calcareous massifs of central-southern Tuscany (Central Italy). *Lazaroa* 24: 61–85.
- Arrigoni, P. V., 1998: La vegetazione forestale. Boschi e macchie di Toscana. Edizioni Regione Toscana.
- Bigi, L. & Rustici, L. 1984: Regime idrico dei suoli e tipi climatici in Toscana. Dipartimento Agricoltura e Foreste, regione Toscana, 130 pp.
- Biondi, E. 2011: Phytosociology today: methodological and conceptual evolution. *Plant Biosystems* 145: 19–29.
- Biondi, E., Allegrezza, M. & Zuccarello, V. 2005: Syntaxonomic revision of the Apennine grasslands belonging to *Brometalia erecti*, and an analysis of their relationships with the xerophyllous vegetation of *Rosmarinetea officinalis* (Italy). *Phytocoenologia* 35 (1): 129–163.
- Biondi, E., Ballelli, S., Allegrezza, M. & Zuccarello, V. 1995: La vegetazione dell'ordine *Brometalia erecti* Br.-Bl. 1936 nell'Appennino (Italia). *Fitosociologia* 30: 3–45.
- Biondi, E. & Blasi, C. (Coord.) 2009: Manuale italiano di Interpretazione degli Habitat della Direttiva 92/43/CEE. <http://vnr.unipg.it/habitat/>. Accessed 05/07/2012
- Biondi, E., Casavecchia, S. & Pesaresi, S. 2010: Interpretation and management of the forest habitats of the Italian peninsula. *Acta Botanica Gallica* 157: 687–719.
- Biondi, E., Casavecchia, S., Pinzi, M., Allegrezza, M. & Baldoni, M. 2002: The syntaxonomy of the mesophilous woods of the Central and Northern Apennines (Italy). *Fitosociologia* 39 (2): 71–93.
- Biondi, E. & Galdenzi, D. 2012: Phytosociological analysis of the grasslands of Montagna dei Fiori (central Italy) and syntaxonomic review of the class *Festuco-Brometea* in the Apennines. *Plant Sociology* 49(1): 91–112.
- Biondi, E. & Guerra, V. 2008: Vegetazione e paesaggio vegetale delle gravine dell'arco jonico. *Fitosociologia* 45 (1) suppl. 1: 57–125.
- Blasi, C., Biondi, E. & Izco, J. 2011: 100 years of plant sociology: A celebration. *Plant Biosystems* 145: 1–3.
- Blasi, C., Cutini, M., Di Pietro, R. & Fortini, P. 2002: Contributo alla conoscenza della suballeanza *Pruno-Rubionium ulmifolii* in Italia. *Fitosociologia* 39 (1), suppl. 2: 129–143.
- Blasi, C. & Di Pietro, R. 1998: Two new phytosociological types of *Quercus pubescens* s.l. woodland communities in southern Latium. *Plant Biosystems* 132 (3): 207–223.
- Blasi, C. & Di Pietro, R. 2002: Sulla presenza di una nuova associazione a *Paliurus spina-christi* L. nel Lazio meridionale. *Inform. Bot. Ital.* 33(2) (2001): 407–415.
- Blasi, C., Di Pietro, R., Filesi, L. & Fortini, P. 2001: Syntaxonomy, chorology and syndynamics of *Carpinus orientalis* communities in Central Italy. *Phytocoenologia* 31 (1): 33–62.
- Blasi, C., Di Pietro, R. & Filesi, L. 2004: Syntaxonomical revision of *Quercetalia pubescenti-petraeae* in the Italian Peninsula. *Fitosociologia* 41(1): 87–164.
- Blasi, C., Marignani, M., Copiz, R., Fipaldini, M.

- & Del Vico, E. (Eds.) 2010: *Le Aree Importanti per le Piante nelle Regioni d'Italia: il presente e il futuro della conservazione del nostro patrimonio botanico*. Sapienza Università di Roma, Progetto Artiser, Roma, 224 pp.
- Braun-Blanquet, J. 1932: *Plant Sociology*. Mac Graw Hill Book Company, New York.
- Braun-Blanquet, J. 1964: *Pflanzensoziologie*. Springer Verlag, Wien and New York.
- Braun-Blanquet, J., Roussine, N. & Nègre, R. 1952: *Les groupements végétaux de la France méditerranéenne*. CNRS, Paris.
- Brullo, S., Scelsi, F. & Spampinato, G. 2001: *La vegetazione dell'Aspromonte*. Studio fitosociologico. Laruffa Editore, Reggio Calabria
- Brullo, S. & Guarino, R. 1999: Syntaxonomy of the *Parietarietea judaicae* class in Europe. *Annali di Botanica* 56 (1): 109–146.
- Brullo, S. & Guarino, R. 2002: La classe *Parietarietea judaicae* Oberd. 1977 in Italia. *Fitosociologia* 39(1) (suppl. 2): 5–27.
- Conti, F. & Abbate, G., Alessandrini, A. & Blasi, C., 2005: An annotated Checklist of the Italian Vascular Flora. *Min. Ambiente e Tutela Territ., Dip. Biol. Veg. Univ. "La Sapienza", Palombi Ed., Roma*.
- Conti, F., Alessandrini, A., Bacchetta, G., Banfi, E., Barberis, G., Bartolucci, F., Bernardo, L., Bonacquisti, S., Bouvet, D., Bovio, M., Brusa, G., Del Guacchio, E., Foggi, B., Frattini, S., Galasso, G., Gallo, L., Gangale, C., Gottschlich, G., Grünanger, P., Gubellini, L., Iriti, G., Lucarini, D., Marchetti, D., Moraldo, B., Peruzzi, L., Poldini, L., Prosser, F., Raffaelli, M., Santangelo, A., Scassellati, E., Scortegagna, S., Selvi, F., Soldano, A., Tinti, D., Ubaldi, D., Uzunov, D. & Vidali, M., 2007: Integrazioni alla checklist della flora vascolare italiana. *Natura Vicentina* 10 (2006): 5–74.
- Cutini M., Fabozzi F., Fortini P., Armanini E. & Blasi C., 1996. Coenological and phytosociological characterization of Latium. *Arch. Geobot.* 2 (2): 113–122.
- De Foucault, B. 1999: Nouvelle contribution à une synsystème des pelouses sèches à thérophytes. *Doc. Phytosoc.* 19: 47–105.
- Di Pietro, R., 2011: New dry grassland associations from the Ausoni–Aurunci mountains (Central Italy) – syntaxonomical updating and discussion on the higher rank syntaxa. *Hacquetia* 10 (2): 183–231.
- Di Pietro, R., Azzella, M. M. & Facioni, L. 2010: The forest vegetation of the Tolfa-Ceriti mountains (northern Latium – central Italy). *Hacquetia* 9 (1): 91–150.
- Di Pietro, R., Burrascano, S. & Blasi, C., 2006: Le comunità dell'*Alyso–Sedion* sui Monti Prenestini (Lazio). *Allionia* 40: 39–49.
- Di Pietro, R., Terzi, M. & Čarni, A. 2012: Syntaxonomy and phytogeography of the *Festuco-Brometea* dry grasslands from an alpha-Adriatic point of view. *Vegetation Databases and large-scale classification, Biogeographical patterns in vegetation, Vegetation and global change, 21st Workshop of European Vegetation Survey, Vienna (Austria) 24th–27th May 2012*. Book of Abstracts: 16.
- European Commission 2007: *Interpretation Manual of European Union Habitats*, vers. EUR27. European Commission, DG Environment, Brussel.
- Fanelli, G. 2007: Cryptogams-rich nanotherophytic vegetation on travertine outcrops near Tivoli. *Annali di Botanica*, VII: 85–92.
- Fanelli, G., Bianco, P. M., De Sanctis, M. & Serafini Sauli, A. 2010: The alliance *Trachynion distachyae* Rivas-Martínez 1978 in central Italy. *Annali di Botanica* 2010: 1–13.
- Foggi, B. & Venturi, E. 2009: Gli habitat meritevoli di conservazione della Provincia di Prato. In: Fancelli, E. (ed.), *Biodiversità in Provincia di Prato*. 4. Habitat. Provincia di Prato, Le Balze Editrice, Roma.
- Géhu, J. M., & Rivas-Martínez, S. 1981: Notions fondamentales de phytosociologie. *Berich. int. Symp. int. Verein Vegetationsk. Syntaxonomie*: 5–33.
- Hammer, Ø., Harper, D. A. T. & Ryan, P. D. 2001: PAST: Paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4(1): 9pp. http://palaeo-electronica.org/2001_1/past/issue1_01.htm. PAST software available from: <http://folk.uio.no/ohammer/past>.
- Lastrucci, L., Foggi, B., Raffaelli, M., Benesperi, R., Gonnelli, V., Zinetti, F., Principi, G. & Cavazzoni, D. 2009: Contributo alla conoscenza naturalistica dell'Area Naturale Protetta di Interesse Locale (ANPIL) "Serpentine di Pieve S. Stefano" (Arezzo, Toscana). *Boll. Mus. reg. Sci. nat. Torino* 27 (1): 101–151.
- Noest, V., Van Der Maarel, E., Van Der Meulten, F. & Van Der Loan, D. 1989: Optimum–transformation of plant species cover abundance values. *Vegetatio* 83: 167–178.
- Oberdorfer, E. 1978: *Süddeutsche Pflanzenge-*

- sellschaften – Teil II. G. Fischer Verlag. – Stuttgart, New York. pp. 355.
- Pignatti, S. 1982: Flora d'Italia. 1–3. Edagricole, Bologna.
- Pignatti, S. 1998: I boschi d'Italia. Edizioni UTET, Torino.
- Pott, R. 2011: Phytosociology – A modern geobotanical method. *Plant Biosystems* 145: 9–18.
- Press, F & Siever, R. 1985: Introduzione alle scienze della Terra. Zanichelli, Bologna, pp. 127–150.
- Regione Toscana 2010: Rete Natura 2000 – Toscana, Siti di importanza regionale. http://www.regione.toscana.it/regione/multimedia/RT/documents/2010/08/26/cfcd9d65030cd89e0ee-b371acb941093_cartografiasir2010.pdf. Accessed 05.07.2012.
- Rivas-Martínez, S., Fernando-Gonzalez, F. & Loidi, J. 1997: Syntaxonomical check-list of the Iberian Peninsula and Balearic and Canary Islands (Spain and Continental Portugal). Phytosociological Research Center (CIF), Madrid.
- Rivas-Martínez, S., Díaz, T. E., Fernández-González, F., Izco, J., Loidi, J., Lousã, M. & Penas, Á. 2002: Vascular plant communities of Spain and Portugal. Addenda to the Syntaxonomical checklist of 2001. *Itinera Geobotanica* 15 (1–2): 5–922.
- Rivas-Martínez, S., & Rivas-Saenz, S. 1996–2009: Worldwide Bioclimatic Classification System, Phytosociological Research Center, Spain. <http://www.globalbioclimatics.org>. Accessed 05/07/2012
- Scoppola, A. & Angiolini, C. 1997: Considerazioni ecologiche e sintassonomiche su alcune garighe dell'entroterra fra Siena e Viterbo (Italia centrale). *Fitosociologia* 32: 121–134.
- Scoppola, A. & Angiolini, C. 2001: Therophytic vegetation of carbonate soils of central Tyrrhenian Italy: synecology and syntaxonomy. *Fitosociologia* 38 (1): 77–89.
- Scoppola, A., Blasi, C., Abbate, G., Cutini, M., Di Marzio, P., Fabozzi, C. & Fortini, P. 1995: Analisi critica e considerazioni fitogeografiche sugli ordini e le alleanze dei querceti e boschi misti a caducifogliedell'Italia peninsulare. *Ann. Bot. LI, Suppl. 10* (1993): 81–112.
- Selvi, F. & Stefanini, P. 2005: Biotopi naturali e aree protette nella provincia di Grosseto: componenti floristiche e ambienti vegetazionali. *Quaderni delle Aree Protette, Prov. Grosseto. U.O.C. Aree Protette e Biodiversità, Città di Castello*, 143 pp.
- Selvi, F. 2010: A critical checklist of the vascular flora of Tuscan Maremma (Grosseto province, Italy). *Fl. Medit.* 20: 47–139.
- Servizio Geologico d'Italia 1965: Carta Geologica d'Italia, Foglio 129, S. Fiora, scala 1:100.000, Poligrafica & Cartevalori, Napoli.
- Thorntwaite, C. W. & Mather, J. R. 1957: Instruction and tables for computing potential evapotranspiration and the water balance. *Pubbl. Climatol.*, 10 (3): 1–311.
- Tutin, T. G., et al. 1964–1980: *Flora Europaea*. 1–5. Cambridge University Press.
- Tutin, T.G., et al. 1993: *Flora Europaea*. 1, 2nd ed. Cambridge University Press.
- Ubaldi, D. 2003: La vegetazione boschiva d'Italia: Manuale di Fitosociologia forestale. Clueb, Bologna, 368 pp.
- Ubaldi, D., Zanotti, A. L., Puppi, G., Speranza, M. & Corbetta, F. 1987: Sintassonomia dei boschi caducifogli mesofili dell'Italia peninsulare. *Not. Fitosoc.* 23: 31–62.
- Ubaldi, D., Zanotti, A. L. & Puppi, G. 1993: Les paysages forestiers de l'Emilie-Romagne et leur signification bioclimatique. *Colloques phytosociologiques* 21: 269–286.
- Uggeri, A. 2003: Geologia dei travertini. In: AA. VV., *Le Sorgenti Petrificanti: linee guida per lo studio, il monitoraggio e la gestione. Parco Regionale Campo dei Fiori*. http://www.parcocurone.it/files/retenatura/travertino_2003/2%20UGGERI%20geologia%20travertini.pdf. Accessed 05/07/2012
- Van der Maarel, E. 1979: Transformation of cover-abundance values in phytosociology and its effects on community similarity. *Vegetatio* 39 (1979): 97–114.
- Venanzoni, R. & Gigante, D. 1999: Contributo allo studio dei pascoli sommitali del Monte Tezio (Perugia, Italia). *Fitosociologia* 36 (1): 157–174.
- Weber, H. E., Moravec, J. & Theurillat, J. P. 2000: International Code of Phytosociological Nomenclature, 3rd edition. *J. Veg. Sc.* 11: 739–768.
- Westhoff, V. & Van Der Maarel, E. 1973: The Braun-Blanquet approach. In: Whittaker R. H. (ed.), *Ordination and classification of communities. Handbook of Veg. Science. Vol. 5*, The Hague, Junk, pp. 617–626.

Received 29. 9. 2012

Revision received 11. 2. 2013

Accepted 14. 2. 2013

Table 1: Annual-dominated communities of flat travertine pavements, *Sedetum hispanico-caespitosi* ass. nova.**Tabella 1:** Združbe enoletnic na travertinskih ploščah, *Sedetum hispanico-caespitosi* ass. nova.

Relevè nr.	19	20*	14	15	18
Aspect	0	0	0	0	0
Slope (°)	0	0	0	0	0
Altitude (m asl)	235	235	250	257	245
Area (m ²)	2	1,5	1	1	1
Total cover (%) (without bryophytes)	20	25	50	20	35
Site (BS: Poggio Bagno Santo; PP: Pian di Palma)	PP	PP	BS	BS	BS
Number of species per relevè	27	25	25	20	22
<i>Sedetum hispanico-caespitosi</i>					
<i>Sedum caespitosum</i>	1	2	3	1	2
<i>Plantago lagopus</i>	+	+	1	1	.
<i>Saxifraga tridactylites</i>	1	+	.	1	+
<i>Sedum hispanicum</i>	+	+	+	1	.
<i>Hypochoerion</i> / <i>Hypochoerion achyrophori</i>, <i>Brachypodietalia distachyi</i>, <i>Tuberarietea</i>					
<i>Arenaria leptoclados</i>	1	+	1	+	+
<i>Catapodium rigidum</i>	.	r	+	+	1
<i>Vulpia ciliata</i>	+	+	+	.	+
<i>Cerastium brachypetalum</i> s.l.	.	.	1	+	+
<i>Minuartia hybrida</i>	+	.	+	.	1
<i>Crepis neglecta</i>	+	+	.	.	r
<i>Filago pyramidata</i>	r	+	.	.	.
<i>Hedypnois rhagadioloides</i>	r	.	.	+	.
<i>Trifolium campestre</i>	r	.	.	.	r
<i>Hypochoeris achyrophorus</i>	.	.	.	+	.
<i>Campanula erinus</i>	.	.	.	+	.
<i>Helianthemum salicifolium</i>	.	r	.	.	.
<i>Sideritis romana</i>	+
<i>Valerianella eriocarpa</i>	.	.	+	.	.
<i>Trifolium scabrum</i>	+
<i>Medicago minima</i>	r
Other annual-dominated community species					
(<i>Stellarietea</i>, <i>Poetea bulbosae</i>)					
<i>Anthemis arvensis</i>	+	r	1	+	r
<i>Poa bulbosa</i>	+	+	.	1	1
<i>Capsella rubella</i>	.	r	+	+	r
<i>Crepis sancta</i>	+	+	+	+	.
<i>Trifolium tomentosum</i>	.	.	2	+	1
<i>Veronica acinifolia</i>	r	.	+	.	r
<i>Cerastium ligusticum</i>	1	1	.	.	.
<i>Aphanes arvensis</i>	.	.	.	+	1
<i>Trifolium nigrescens</i>	.	+	1	.	.
<i>Erodium cicutarium</i>	+	r	.	.	.
<i>Geranium pusillum</i>	r	.	+	.	.
<i>Parentucellia latifolia</i>	+	+	.	.	.
<i>Poa annua</i>	.	.	+	+	.
<i>Ranunculus flabellatus</i>	+	+	.	.	.
<i>Sherardia arvensis</i>	+	+	.	.	.

Relevè nr.	19	20*	14	15	18
<i>Trisetaria panicea</i>	.	.	.	+	+
<i>Sagina apetala</i>	.	.	1	.	.
<i>Sedum stellatum</i>	.	.	.	1	.
<i>Adonis annua</i>	.	.	+	.	.
<i>Calendula arvensis</i>	.	+	.	.	.
<i>Koeleria cristata</i>	.	.	+	.	.
<i>Papaver rhoeas</i>	.	.	+	.	.
<i>Reichardia picroides</i>	+
<i>Trifolium subterraneum</i>	.	+	.	.	.
<i>Bunias erucago</i>	.	.	+	.	.
<i>Vulpia ligustica</i>	r
Other species					
<i>Salvia clandestina</i>	1	+	.	.	.
<i>Eryngium campestre</i>	+	+	.	.	.
<i>Romulea bulbocodium</i>	+	+	.	.	.
<i>Calamintha nepeta</i>	r
<i>Centaurea solstitialis</i>	.	.	.	r	.
<i>Convolvulus cantabrica</i>	+
<i>Dactylis hispanica</i>	+
<i>Hordeum murinum</i>	.	.	+	.	.
<i>Juncus bufonius</i>	.	.	r	.	.
<i>Salvia verbenaca</i>	.	.	+	.	.
<i>Torilis nodosa</i>	r

* = type relevé

Table 2: Chasmophytic and comophytic vegetation of steep rocks and cliff edges, *Cymbalaria muralis*, *Muscari commutatum* and *Allium subhirsutum* coenoses, *Parietarietea*.

Tabella 2: Hazmofitska in komofitska vegetacija na strmih skalah in robovih klifov, *Cymbalaria muralis*, *Muscari commutatum* and *Allium subhirsutum* coenoses, *Parietarietea*.

Relevè nr.	16	17	Relevè nr.	16	17
Aspect	0	0	<i>Galium aparine</i>	r	.
Slope (°)	30	10	<i>Arabis turrata</i>	r	.
Altitude (m asl)	255	250	Other species		
Area (m ²)	2	1	<i>Aetheorrhiza bulbosa</i>	1	.
Total cover (%)	40	20	<i>Lens ervoides</i>	+	.
Site (BS: Poggio Bagno Santo; PP: Pian di Palma)	BS	BS	<i>Aristolochia lutea</i>	+	.
Number of species per relevé	16	7	<i>Asparagus acutifolius</i>	+	.
<i>Parietarietea</i> and other chasmophytic and comophytic species			<i>Ranunculus millefoliatus</i>	+	.
<i>Cymbalaria muralis</i>	1	1	<i>Teucrium chamaedrys</i>	+	.
<i>Muscari commutatum</i>	+	+			
<i>Asplenium trichomanes</i>	+	+			
<i>Umbilicus rupestris</i>	.	1			
<i>Polypodium cambricum</i>	+	.			
<i>Galio-Urticetea</i> and <i>Trifolio-Geranietea</i>					
<i>Allium subhirsutum</i>	1	1			
<i>Geranium purpureum</i>	+	+			
<i>Parietaria officinalis</i>	.	+			
<i>Centranthus calcitrapa</i>	+	.			

Table 3: Chamaephytic garrigues, *Cephalario leucanthae-Saturejetum montanae euphorbietosum spinosae*, low altitude variant.

Tabela 3: Hamefitske garige, *Cephalario leucanthae-Saturejetum montanae euphorbietosum spinosae*, nižisnka varianta.

Relevè nr.	23	24	Relevè nr.	23	24
Aspect	0	0	<i>Carlina corymbosa</i>	r	r
Slope (°)	0	0	<i>Potentilla hirta</i>	+	+
Altitude (m asl)	235	235	<i>Sanguisorba minor</i>	.	+
Area (m ²)	10	10	<i>Salvia verbenaca</i>	.	+
Total cover (%)	80	70	<i>Koeleria splendens</i>	.	+
Site (BS: Poggio Bagno Santo; PP: Pian di Palma)	PP	PP	Travertine annual-dominated community species		
Number of species per relevè	32	35	<i>Ranunculus flabellatus</i>	r	r
<i>Cephalario leucanthae-Saturejetum montanae</i>			<i>Reichardia picroides</i>	+	+
<i>Satureja montana</i>	2	1	<i>Sherardia arvensis</i>	+	+
<i>Thymus longicaulis</i>	1	1	<i>Geranium pusillum</i>	r	r
<i>Cephalaria leucantha</i>	2	1	<i>Helianthemum salicifolium</i>	+	+
<i>euphorbietosum spinosae</i>			<i>Hypochoeris achyrophorus</i>	+	.
<i>Euphorbia spinosa</i>	2	3	<i>Sideritis romana</i>	.	+
<i>Galium corrudifolium</i>	+	+	<i>Trifolium campestre</i>	.	+
other <i>Rosmarinetea</i> species			<i>Arenaria leptoclados</i>	.	+
<i>Teucrium capitatum</i>	1	2	<i>Biscutella mollis</i>	+	.
<i>Convolvulus cantabrica</i>	1	1	<i>Crepis sancta</i>	r	.
<i>Alyso-Sedion albi</i>			Other species		
<i>Sedum rupestre</i>	+	+	<i>Avena barbata</i>	+	+
<i>Sedum acre</i>	.	+	<i>Dactylis hispanica</i>	+	+
Grassland species			<i>Melica transylvanica</i>	+	+
<i>Bromus erectus</i>	2	2	<i>Marrubium incanum</i>	.	+
<i>Allium moschatum</i>	1	1	<i>Romulea bulbocodium</i>	r	+
<i>Iris lutescens</i>	1	1	<i>Anemone hortensis</i>	+	.
<i>Urospermum dalechampii</i>	+	+	<i>Coronilla scorpioides</i>	r	.
<i>Eryngium campestre</i>	+	+	<i>Crepis neglecta</i>	.	r
<i>Filipendula vulgaris</i>	r	r	<i>Echium italicum</i>	.	r
			<i>Osyris alba</i>	r	.
			<i>Pallenis spinosa</i>	r	.

Table 4: *Bromus erectus* grasslands, *Artemisio albae*-*Bromenalia erecti*.**Tabela 4:** Travišča z vrsto *Bromus erectus*, *Artemisio albae*-*Bromenalia erecti*.

Relevè nr.	21	22	Relevè nr.	21	22
Aspect	0	0	<i>Coronilla scorpioides</i>	r	.
Slope (°)	0	0	<i>Pallenis spinosa</i>	r	.
Altitude (m asl)	235	235	<i>Bromus rubens</i>	.	+
Area (m ²)	40	40	<i>Poa bulbosa</i>	.	+
Total cover (%)	80	70	<i>Sedum hispanicum</i>	.	+
Site (BS: Poggio Bagno Santo; PP: Pian di Palma)	PP	PP	<i>Sideritis romana</i>	.	+
Number of species per relevè	35	36	<i>Trifolium campestre</i>	.	+
<i>Brometalia erecti</i> and <i>Artemisio albae</i>-<i>Bromenalia erecti</i>			<i>Trifolium stellatum</i>	.	+
<i>Bromus erectus</i>	4	3	<i>Vulpia ciliata</i>	.	+
<i>Eryngium campestre</i>	1	1	<i>Crepis setosa</i>	.	r
<i>Convolvulus cantabrica</i>	1	1	<i>Hippocrepis ciliata</i>	.	r
<i>Teucrium capitatum</i>	+	1	Other species		
<i>Urospermum dalechampii</i>	+	1	<i>Euphorbia spinosa</i>	1	2
<i>Galium corrudifolium</i>	+	+	<i>Cephalaria leucantha</i>	1	1
<i>Koeleria splendens</i>	+	+	<i>Romulea bulbocodium</i>	+	1
<i>Potentilla hirta</i>	+	+	<i>Sedum rupestre</i>	+	1
<i>Thymus longicaulis</i>	+	+	<i>Dactylis hispanica</i>	+	+
<i>Tragopogon dubius</i>	+	+	<i>Orchis morio</i>	r	+
<i>Carlina corymbosa</i>	+	+	<i>Serapias vomeracea</i>	r	+
<i>Filipendula vulgaris</i>	r	+	<i>Ornithogalum narbonense</i>	r	r
<i>Satureja montana</i>	1	+	<i>Anemone hortensis</i>	+	.
<i>Anthyllis vulneraria</i> ssp. <i>rubriflora</i>	+	.	<i>Osyris alba</i>	r	.
<i>Cleistogenes serotina</i>	.	+	<i>Sedum acre</i>	.	+
<i>Salvia verbenaca</i>	.	+	<i>Bellis perennis</i>	.	r
<i>Sanguisorba minor</i>	.	+	<i>Echium italicum</i>	.	r
<i>Melica transylvanica</i>	+	.	<i>Marrubium incanum</i>	.	r
<i>Crupina vulgaris</i>	r	.			
<i>Lygeo-Stipetea</i>					
<i>Allium moschatum</i>	1	1			
<i>Iris lutescens</i>	1	1			
Travertine annual-dominated community species					
<i>Reichardia picroides</i>	1	1			
<i>Helianthemum salicifolium</i>	+	1			
<i>Tordylium apulum</i>	+	1			
<i>Crepis neglecta</i>	+	+			
<i>Geranium pusillum</i>	+	+			
<i>Ranunculus flabellatus</i>	+	+			
<i>Sherardia arvensis</i>	+	+			
<i>Anthemis arvensis</i>	r	+			
<i>Arenaria leptocladus</i>	.	1			
<i>Crepis sancta</i>	+	.			
<i>Filago pyramidata</i>	+	.			
<i>Hypochoeris achyrophorus</i>	+	.			
<i>Orlaya grandiflora</i>	+	.			
<i>Veronica acinifolia</i>	+	.			
<i>Biscutella mollis</i>	r	.			
<i>Cerastium ligusticum</i>	r	.			

Table 5: Open scrub and shrublands. a: Pistacio terebinthi-Paliuretum spinae-christi; b: transition aspect between Pistacio-Paliuretum shrublands and Quercus pubescens woods

Tabela 5: Odrpta grmišča. a: Pistacio terebinthi-Paliuretum spinae-christi; b: prehod med grmišči Pistacio-Paliuretum in gozdovi Quercus pubescens woods

	a	a	a	a	a	a	a	b
Relevè nr.	1	2	6	7	4	5	3	13
Aspect	0	SE	W	SE	E	E	E	0
Slope (°)	0	<5	5	7	3	5	2	0
Altitude (m asl)	275	261	245	255	230	245	240	215
Area (m ²)	100	100	50	100	60	70	70	200
Total cover (%) (without bryophytes)	70	75	75	90	85	90	80	90
Site (BS: Poggio Bagno Santo; PP: Pian di Palma)	BS	BS	BS	BS	BS	BS	BS	PP
Number of species per relevè	38	44	23	27	49	50	33	39
<i>Pistacio terebinthi-Paliuretum spinae-christi</i>								
<i>Paliurus spina - christi</i>	3	3	1	1	3	3	3	2
<i>Pistacia terebinthus</i>	3	2	1	2	2	3	1	2
<i>Phyllirea latifolia</i>	2	2	3	3	r	1	+	.
<i>Cercis siliquastrum</i>	.	3	2	1	1	2	r	.
<i>Melica transylvanica</i>	+	r	1	.	+	+	.	.
<i>Pruno-Rubenion, Pruno-Rubion ulmifolii, Prunetalia spinosae, Rhamno-Prunetea</i>								
<i>Asparagus acutifolius</i>	1	1	+	+	r	1	.	1
<i>Ruscus aculeatus</i>	.	.	2	.	.	1	.	.
<i>Clematis flammula</i>	.	.	r	.	.	.	+	.
<i>Pyrus amygdaliformis</i>	1	.
<i>Rubia peregriana</i>	+	.
<i>Rosa sempervirens</i>	+	.
<i>Crataegus monogyna</i>	+
<i>Prunus spinosa</i>	r	.
<i>Tamus communis</i>	.	.	r
<i>Quercetalia pubescenti-petraeae, Quercus-Fagetea</i>								
<i>Fraxinus ornus</i>	1	1	1	1	.	2	1	.
<i>Acer monspessulanum</i>	1	2	.	2	.	1	1	.
<i>Quercus pubescens</i>	.	.	.	+	.	.	2	3
<i>Ulmus minor</i>	+	+	.	.
<i>Celtis australis</i>	.	+
<i>Quercetalia, Quercetea ilicis</i>								
<i>Rhamnus alaternus</i>	+	1	.	.	.	r	.	.
<i>Cyclamen repandum</i>	r	+	r
<i>Olea sylvestris</i>	.	1
<i>Osyris alba</i>	2	.
<i>Carex distachya</i>	.	.	.	1
<i>Smilax aspera</i>	r	.	.
Other species								
<i>Bromus erectus</i>	+	.	.	r	1	1	1	2
<i>Allium subhirsutum</i>	2	2	+	+	.	r	.	.
<i>Calamintha nepeta</i>	1	+	.	r	r	.	.	1
<i>Carex caryophylla</i>	+	.	+	+	.	1	.	1
<i>Dactylis hispanica</i>	1	+	r	.	+	1	.	.
<i>Convolvulus cantabrica</i>	+	+	.	.	+	r	1	.

Relevè nr.	1	2	6	7	4	5	3	13
<i>Arabis hirsuta</i>	+	+	r	.	.	r	.	r
<i>Orlaya grandiflora</i>	.	r	.	.	r	+	+	r
<i>Urospermum dalechampii</i>	+	.	.	r	r	.	+	+
<i>Brachypodium distachyum</i>	.	r	+	.	l	l	.	.
<i>Bromus sterilis</i>	l	l	+	+
<i>Cynosurus echinatus</i>	.	.	+	+	l	l	.	.
<i>Knautia integrifolia</i>	.	.	r	+	+	l	.	.
<i>Teucrium chamaedrys</i>	.	l	.	r	.	+	.	+
<i>Hypochoeris achyrophorus</i>	.	.	.	+	+	+	r	.
<i>Serratula cichoracea</i>	.	r	.	+	.	r	r	.
<i>Sideritis romana</i>	.	.	.	r	r	.	+	+
<i>Thymus longicaulis</i>	+	+	+	+
<i>Aristolochia rotunda</i>	.	.	l	l	.	+	.	.
<i>Eryngium campestre</i>	+	.	l	l
<i>Euphorbia spinosa</i>	l	l	+
<i>Brachypodium ramosum</i>	.	+	.	.	.	+	.	l
<i>Gastridium ventricosum</i>	l	+	+	.
<i>Geranium robertianum</i>	.	.	l	+	.	r	.	.
<i>Crepis zacintha</i>	+	r	.	r
<i>Geranium purpureum</i>	+	+	+
<i>Teucrium capitatum</i>	r	.	+	+
<i>Trifolium angustifolium</i>	.	+	.	.	+	r	.	.
<i>Trifolium stellatum</i>	+	r	.	+
<i>Aegilops geniculata</i>	l	l	.	.
<i>Aristolochia lutea</i>	+	2
<i>Gaudinia fragilis</i>	l	+	.	.
<i>Ranunculus millefoliatus</i>	+	l
<i>Trifolium lappaceum</i>	.	.	.	r	l	.	.	.
<i>Aetheorrhiza bulbosa</i>	+	+
<i>Anemone hortensis</i>	+	+
<i>Anthemis arvensis</i>	+	.	.	.	+	.	.	.
<i>Arenaria leptoclados</i>	.	+	+
<i>Arum italicum</i>	r	+
<i>Bromus hordeaceus</i>	+	+	.	.
<i>Bromus rubens</i>	+	+	.	.
<i>Campanula rapunculus</i>	.	.	.	+	.	r	.	.
<i>Carlina corymbosa</i>	r	.	.	+
<i>Catapodium rigidum</i>	.	+	.	r
<i>Crepis neglecta</i>	+	.	.	+
<i>Euphorbia exigua</i>	r	.	+
<i>Galium corrudifolium</i>	+	+
<i>Geranium dissectum</i>	r	+	.	.
<i>Lathyrus cicera</i>	r	+
<i>Orchis tridentata</i>	+	r
<i>Poa bulbosa</i>	+	+
<i>Poa trivialis</i>	+	+
<i>Salvia clandestina</i>	+	+	.	.
<i>Sanguisorba minor</i>	r	.	+	.
<i>Sedum rupestre</i>	+	+
<i>Torilis japonica</i>	.	+	r
<i>Torilis nodosa</i>	+	+	.	.
<i>Trifolium campestre</i>	+	+	.	.

Relevè nr.	1	2	6	7	4	5	3	13
<i>Trifolium scabrum</i>	+	.	.	+
<i>Vicia sativa ssp. nigra</i>	.	+	.	.	.	r	.	.
<i>Ammoides pusilla</i>	l	.	.	.
<i>Brachypodium rupestre</i>	l	.
<i>Cerastium ligusticum</i>	l
<i>Reichardia picroides</i>	l
<i>Satureja montana</i>	l
<i>Asplenium ceterach</i>	.	r
<i>Avena barbata</i>	+	.	.	.
<i>Biscutella mollis</i>	+
<i>Bombycilaena erecta</i>	+	.	.	.
<i>Briza maxima</i>	+	.	.	.
<i>Bupleurum baldense</i>	+	.	.
<i>Campanula erinus</i>	.	.	.	r
<i>Carex divulsa</i>	+
<i>Centaurea bracteata</i>	+	.
<i>Centaureum erythraea</i>	r	.	.	.
<i>Coronilla scorpioides</i>	r
<i>Crepis sancta</i>	+
<i>Cynoglossum creticum</i>	r
<i>Dorycnium hirsutum</i>	+	.
<i>Filipendula vulgaris</i>	r	.
<i>Fumaria capreolata</i>	.	r
<i>Galium aparine</i>	.	+
<i>Geranium lucidum</i>	.	.	r
<i>Iris lutescens</i>	r
<i>Lathyrus ochrus</i>	r
<i>Linum strictum</i>	r	.	.	.
<i>Lolium perenne</i>	+	.	.
<i>Marrubium incanum</i>	r
<i>Medicago arabica</i>	.	+
<i>Medicago minima</i>	+
<i>Melilotus neapolitana</i>	r	.
<i>Nigella damascena</i>	+	.	.	.
<i>Ophrys sphecodes</i>	r	.	.
<i>Pallenis spinosa</i>	r	.	.	.
<i>Plantago bellardi</i>	+	.	.	.
<i>Plantago lagopus</i>	+	.	.	.
<i>Poa pratensis</i>	+	.	.
<i>Potentilla hirta</i>	+	.
<i>Ruscus hypoglossum</i>	.	.	.	r
<i>Tanacetum corymbosum ssp. achilleae</i>	r	.	.
<i>Theligonium cynocrambe</i>	.	r
<i>Tordylium apulum</i>	+
<i>Tragopogon dubius</i>	+	.
<i>Trifolium cherleri</i>	+	.	.	.
<i>Trifolium striatum</i>	+	.	.
<i>Urospermum picroides</i>	.	.	r
<i>Veronica acinifolia</i>	+
<i>Vulpia bromoides</i>	+

Table 6: Thermophilous deciduous forests dominated by *Quercus pubescens* and *Acer monspessulanum*, *Pistacio terebinthi-Quercetum pubescentis rosetosum sempervirentis*.**Tabella 6:** Termofilni listopadni gozdovi z prevladujočimi vrstami *Quercus pubescens* in *Acer monspessulanum*, *Pistacio terebinthi-Quercetum pubescentis rosetosum sempervirentis*.

Relevè nr.	8	9	10	12	11
Aspect	0	E	0	W	W
Slope (°)	0	8	0	20	15
Altitude (m asl)	220	240	240	210	255
Area (m ²)	200	250	200	300	250
Total cover (%) (without bryophytes)	95	95	95	95	95
Site (BS: Poggio Bagno Santo; PP: Pian di Palma)	PP	BS	BS	PP	BS
Number of species per relevè	28	46	34	31	29
<i>Pistacio terebinthi-Quercetum pubescentis</i>					
<i>Smilax aspera</i>	.	2	.	+	2
<i>Acer campestre</i>	.	1	1	.	.
<i>Pistacia terebinthus</i>	.	+	.	.	.
<i>rosetosum sempervirentis</i>					
<i>Cercis siliquastrum</i>	.	1	3	2	1
<i>Rosa sempervirens</i>	.	+	.	.	+
<i>Brachypodium ramosum</i>	2
<i>Cyclamen repandum</i>	+
local variant differential species					
<i>Paliurus spina - christi</i>	2	1	1	.	2
<i>Celtis australis</i>	.	.	2	1	1
<i>Lauro-Quercenion, Carpinion orientalis</i>					
<i>Acer monspessulanum</i>	3	4	3	4	2
<i>Asparagus acutifolius</i>	.	+	.	r	+
<i>Clematis flammula</i>	1	+	+	.	.
<i>Rubia peregrina</i>	.	1	+	.	r
<i>Phyllirea latifolia</i>	.	+	.	.	1
<i>Ostrya carpinifolia</i>	.	1	.	.	.
<i>Lonicera etrusca</i>	.	+	.	.	.
<i>Anemone hortensis</i>	+
<i>Arabis turrita</i>	.	+	.	.	.
<i>Quercetalia pubescenti-petraeae, Quercus-Fagetea</i>					
<i>Quercus pubescens</i>	4	3	3	2	4
<i>Fraxinus ornus</i>	2	2	1	2	2
<i>Ruscus aculeatus</i>	2	3	2	+	2
<i>Hedera helix</i>	1	1	1	1	1
<i>Tamus communis</i>	.	1	1	r	+
<i>Cornus mas</i>	.	1	.	1	+
<i>Melica uniflora</i>	.	1	+	1	.
<i>Brachypodium sylvaticum</i>	.	+	1	.	+
<i>Melittis melissophyllum</i>	.	1	.	r	+
<i>Viola alba</i> ssp. <i>dehnhardtii</i>	+	+	.	+	.
<i>Lithospermum purpureocaeruleum</i>	.	r	+	.	r
<i>Helleborus foetidus</i>	.	r	r	+	.
<i>Poa sylvicola</i>	.	+	+	+	.
<i>Silene italica</i>	1	.	r	.	.

Relevè nr.	8	9	10	12	11
<i>Quercus cerris</i>	.	1	.	.	.
<i>Stachys officinalis</i>	1
<i>Scutellaria columnae</i>	.	+	.	.	.
<i>Pulicaria odora</i>	+
<i>Lathyrus venetus</i>	.	+	.	.	.
<i>Cyclamen hederifolium</i>	.	.	.	+	.
<i>Geum urbanum</i>	.	.	.	r	.
Other species					
<i>Geranium robertianum</i>	.	2	1	2	2
<i>Allium subhirsutum</i>	.	+	+	+	+
<i>Aristolochia rotunda</i>	.	.	1	+	1
<i>Brachypodium rupestre</i>	2	+	.	.	r
<i>Crataegus monogyna</i>	.	.	1	+	+
<i>Myrrhoides nodosa</i>	.	+	+	r	.
<i>Parietaria officinalis</i>	.	.	r	+	+
<i>Calamintha nepeta</i>	1	.	.	.	r
<i>Quercus ilex</i>	.	1	+	.	.
<i>Serratula cichoracea</i>	1	.	.	.	+
<i>Arabis hirsuta</i>	+	.	r	.	.
<i>Asplenium onopteris</i>	.	r	.	r	.
<i>Cardamine graeca</i>	.	r	.	+	.
<i>Carex distachya</i>	.	.	+	.	+
<i>Geranium lucidum</i>	.	.	+	+	.
<i>Ranunculus velutinus</i>	.	+	.	r	.
<i>Torilis japonica</i>	.	.	+	.	+
<i>Bromus erectus</i>	1
<i>Carex hallerana</i>	1
<i>Filipendula vulgaris</i>	1
<i>Stellaria media</i>	.	.	.	1	.
<i>Alliaria petiolata</i>	r
<i>Asplenium trichomanes</i>	.	r	.	.	.
<i>Bellis perennis</i>	+
<i>Bromus sterilis</i>	.	.	+	.	.
<i>Carex depauperata</i>	.	.	r	.	.
<i>Carlina corymbosa</i>	+
<i>Cruciata glabra</i>	.	r	.	.	.
<i>Cymbalaria muralis</i>	.	r	.	.	.
<i>Cynosurus echinatus</i>	.	.	+	.	.
<i>Ficus carica</i>	.	+	.	.	.
<i>Helianthemum nummularium</i>	+
<i>Lotus hispidus</i>	r
<i>Melica transylvanica</i>	.	.	+	.	.
<i>Moehringia pentandra</i>	.	r	.	.	.
<i>Ornithogalum etruscum</i>	.	r	.	.	.
<i>Prunus spinosa</i>	.	+	.	.	.
<i>Ranunculus bulbosus</i>	+
<i>Rhamnus alaternus</i>	.	.	+	.	.
<i>Sonchus oleraceus</i>	.	.	.	r	.
<i>Theligonium cynocrambe</i>	.	.	.	r	.
<i>Thymus longicaulis</i>	+
<i>Trifolium ochroleucum</i>	+
<i>Umbilicus rupestris</i>	.	r	.	.	.



Pian di Palma *Bromus erectus* grasslands, with cushions of *Euphorbia spinosa*.

Travišča z vrsto *Bromus erectus* in blazinicami *Euphorbia spinosa* na območju Pian di Palma.



Pian di Palma chamaephytic garrigues with *Euphorbia spinosa*, *Satureja montana*, *Helichrysum italicum* and others.

Hamefitska gariga z vrstami *Euphorbia spinosa*, *Satureja montana*, *Helichrysum italicum* in drugimi na območju Pian di Palma.



Shrub community of *Pistacio terebinthi*-*Paliuretum spinae-christi* on the rocky hill slopes of Poggio Bagno Santo.

Grmiščne združbe *Pistacio terebinthi*-*Paliuretum spinae-christi* na gričevju Poggio Bagno Santo.



Aerial view of the mixed thermophilous deciduous forest (*Pistacio terebinthi-Quercetum pubescentis rosetosum sempervirentis*) on the slopes of Poggio Bagno Santo facing the Albegna river.

Pogled iz zraka na mešane termofilne listopadne gozdove (*Pistacio terebinthi-Quercetum pubescentis rosetosum sempervirentis*) na pobočju gričevja Poggio Bagno Santo proti reki Albegna.



The flat travertine pavement of Pian di Palma, with open woodland and species-rich grassland communities.

Ravne plošče travertina na območju Pian di Palma z odprtimi gozdovi in vrstno bogatimi travišči.



Sedum caespitosum, one of the most physiognomically typical species of the annual-dominated communities on flat travertine pavements (*Sedetum hispanico-caespitosi*).

Sedum caespitosum je vrsta, ki daje značilen fiziognomski videz združbam na travnatih ploščah, v katerih prevladujejo anoletnice (*Sedetum hispanico-caespitosi*).