



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Derived algebraic geometry, determinants of perfect complexes, and applications to obstruction theories for maps and complexes

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Derived algebraic geometry, determinants of perfect complexes, and applications to obstruction theories for maps and complexes / T. Schurg; B. Toen; G. Vezzosi. - In: JOURNAL FÜR DIE REINE UND ANGEWANDTE MATHEMATIK. - ISSN 1435-5345. - STAMPA. - Volume 2015:(2015), pp. 1-40. [10.1515/crelle-2013-0037]

Availability:

The webpage <https://hdl.handle.net/2158/800683> of the repository was last updated on 2021-03-08T14:18:12Z

Published version:

DOI: 10.1515/crelle-2013-0037

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

Derived algebraic geometry, determinants of perfect complexes, and applications to obstruction theories for maps and complexes

By *Timo Schürg* at Bonn, *Bertrand Toën* at Montpellier and *Gabriele Vezzosi* at Paris

Abstract. A quasi-smooth derived enhancement of a Deligne–Mumford stack \mathcal{X} naturally endows \mathcal{X} with a functorial perfect obstruction theory in the sense of Behrend–Fantechi. We apply this result to moduli of maps and perfect complexes on a smooth complex projective variety.

For *moduli of maps*, for $X = S$ an algebraic $K3$ -surface, $g \in \mathbb{N}$, and $\beta \neq 0$ in $H_2(S, \mathbb{Z})$ a curve class, we construct a derived stack $\mathbb{R}\overline{\mathbf{M}}_{g,n}^{\text{red}}(S; \beta)$ whose truncation is the usual stack $\overline{\mathbf{M}}_{g,n}(S; \beta)$ of pointed stable maps from curves of genus g to S hitting the class β , and such that the inclusion $\overline{\mathbf{M}}_g(S; \beta) \hookrightarrow \mathbb{R}\overline{\mathbf{M}}_g^{\text{red}}(S; \beta)$ induces on $\overline{\mathbf{M}}_g(S; \beta)$ a perfect obstruction theory whose tangent and obstruction spaces coincide with the corresponding *reduced* spaces of Okounkov–Maulik–Pandharipande–Thomas. The approach we present here uses derived algebraic geometry and yields not only a full rigorous proof of the existence of a reduced obstruction theory – not relying on any result on semiregularity maps – but also a new global geometric interpretation.

We give two further applications to *moduli of complexes*. For a $K3$ -surface S we show that the stack of simple perfect complexes on S is smooth. This result was proved with different methods by Inaba for the corresponding coarse moduli space. Finally, we construct a map from the derived stack of stable embeddings of curves (into a smooth complex projective variety X) to the derived stack of simple perfect complexes on X with vanishing negative Ext's, and show how this map induces a morphism of the corresponding obstruction theories when X is a Calabi–Yau 3-fold.

An important ingredient of our construction is a *perfect determinant map* from the derived stack of perfect complexes to the derived stack of line bundles whose tangent morphism is given by Illusie's trace map for perfect complexes.

Note 1:
removed the cross
references

Note 2:
unified to '3-fold'

Introduction

It is well known in Algebraic Geometry – e.g. in Gromov–Witten and Donaldson–Thomas theories – the importance of endowing a Deligne–Mumford moduli stack with a (perfect) *obstruction theory*, as defined in [3]: such an obstruction theory gives a *virtual fundamental class*

in the Chow group of the stack. If the stack in question is the stack of pointed stable maps to a fixed smooth projective variety ([4]), then integrating appropriate classes against this class produces all versions of Gromov–Witten invariants ([1]).

Now, it is a distinguished feature of *Derived Algebraic Geometry* ([36]) that any quasi-smooth *derived extension* of such a stack F , i.e. a derived stack that is locally of finite presentation whose cotangent complex is of perfect amplitude in $[-1, 0]$, and whose underived part or *truncation* is the given stack F , induces a *canonical* obstruction theory on F : we have collected these results in Section 1 below. A morphism of derived stacks induces naturally a morphism between the induced obstruction theories – so that functoriality results like [3, **Proposition 5.10**] or the so-called virtual pullback result in [17] follow immediately. Moreover the functoriality of obstruction theories induced by morphisms of derived extensions is definitely richer than the usual one in [3], that is restricted to special situations (e.g. [3, **Proposition 5.10**]), and requires the axiomatics of compatible obstruction theories. In other words, a suitable reformulation of a moduli problem in derived algebraic geometry, immediately gives us a canonical obstruction theory, in a completely geometric way, with no need of clever choices.

In this paper we apply this ability of derived algebraic geometry in producing obstruction theories – functorial with respect to maps of derived stacks – to the cases of moduli of maps and moduli of perfect complexes on a complex smooth projective variety X .

Moduli of maps. For moduli of maps, we show how the standard obstruction theory yielding Gromov–Witten invariants comes from a natural derived extension of the stack of pointed stable maps to X . Then we concentrate on a geometrically interesting occurrence of two different obstruction theories on a given stack, namely the stack $\overline{\mathbf{M}}_g(S; \beta)$ of stable maps of type (g, β) to a smooth projective complex $K3$ -surface S . The stack $\overline{\mathbf{M}}_g(S; \beta)$ has a *standard* obstruction theory, yielding trivial Gromov–Witten invariants in the n -pointed case, and a so-called *reduced* obstruction theory, first considered by Okounkov–Maulik–Pandharipande–Thomas (often abbreviated to O-M-P-T in the text), giving interesting – and extremely rich in structure – curve counting invariants in the n -pointed case (see [20, 21, 25], and Section 4.1 below, for a detailed review). In this paper we use derived algebraic geometry to give a construction of a global reduced obstruction theory on $\overline{\mathbf{M}}_g(S; \beta)$, and compare its deformation and obstruction spaces with those of Okounkov–Maulik–Pandharipande–Thomas. More precisely, we use a perfect determinant map from the derived stack of perfect complexes to the derived stack of line bundles, and exploit the peculiarities of the derived stack of line bundles on a $K3$ -surface, to produce a derived extension $\mathbb{R}\overline{\mathbf{M}}_g^{\text{red}}(S; \beta)$ of $\overline{\mathbf{M}}_g(S; \beta)$. The derived stack $\mathbb{R}\overline{\mathbf{M}}_g^{\text{red}}(S; \beta)$ arises as the canonical homotopy fiber over the *unique derived factor* of the derived stack of line bundles on S , so it is, in a very essential way, a purely derived geometrical object. We prove quasi-smoothness of $\mathbb{R}\overline{\mathbf{M}}_g^{\text{red}}(S; \beta)$, and this immediately gives us a global reduced obstruction theory on $\overline{\mathbf{M}}_g(S; \beta)$. Our proof is self-contained (inside derived algebraic geometry), and does not rely on any previous results on semiregularity maps.

Moduli of complexes. We give two applications to moduli of perfect complexes on smooth projective varieties. In the first one we show that the moduli space of simple perfect complexes on a $K3$ -surface is smooth. Inaba gave a direct proof of this result in [13], by generalizing methods of Mukai ([22]). Our proof is different and straightforward. We use the perfect determinant map, and the peculiar structure of the derived Picard stack of a $K3$ -surface, to produce a derived stack of simple perfect complexes. Then we show that this derived stack

Note 3:
Red parts indicate
major changes.
Please check them
carefully.

is actually *underived* (i.e trivial in the derived direction) and *smooth*. The moduli space studied by Inaba is exactly the coarse moduli space of this stack.

In the second application, for X an arbitrary smooth complex projective scheme X , we first construct a map C from the derived stack $\mathbb{R}\mathbf{M}_{g,n}(X)^{\text{emb}}$ consisting of pointed stable maps which are closed immersions, to the derived stack $\mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}^{\text{si}, > 0}$ of simple perfect complexes with no negative Ext's and fixed determinant \mathcal{L} (for arbitrary \mathcal{L}). Then we show that, if X is a *Calabi–Yau 3-fold*, the derived stack $\mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}^{\text{si}, > 0}$ is actually *quasi-smooth*, and use the map C to compare (according to Section 5.2) the canonical obstruction theories induced by the source and target derived stacks on their truncations. Finally, we relate this second applications to a simplified, open version of the Gromov–Witten/Donaldson–Thomas conjectural comparison. In such a comparison, one meets two basic problems. The first, easier, one is in producing a map enabling one to compare the obstruction theories – and derived algebraic geometry, as we show in the open case, is perfectly suited for this (see Sections 1 and 5.2). Such a comparison would induce a comparison (via a virtual pullback construction as in [29, Theorem 7.4]) between the corresponding *virtual fundamental classes*, and thus a comparison between the GW and DT invariants. The second problem, certainly the most difficult one, is to deal with problems arising at the boundary of the compactifications. For this second problem, derived algebraic methods unfortunately do not provide at the moment any new tool or direction.

One of the main ingredients of all the applications given in this paper is the construction of a *perfect determinant map* $\det_{\mathbf{Perf}} : \mathbf{Perf} \rightarrow \mathbf{Pic}$, where \mathbf{Perf} is the stack of perfect complexes, \mathbf{Pic} the stack of line bundles, and both are viewed as derived stacks (see Section 3.1 for details), whose definition requires the use of a bit of Waldhausen K -theory for simplicial commutative rings, and whose tangent map can be identified with Illusie's trace map of perfect complexes ([12, Chapter 5]). We expect that this determinant map might be useful in other moduli contexts as well.

An important remark – especially for applications to Gromov–Witten theory – is that, in order to simplify the exposition, we have chosen to write the proofs only in the non-pointed case, since obviously no substantial differences except for notational ones are involved. The relevant statements are however given in both the unpointed and the n -pointed case.

Finally, let us observe that most of the natural maps of complexes arising in moduli problems can be realized as tangent or cotangent maps associated to morphisms between appropriate derived moduli stacks. This suggestion is confirmed in the present paper for the standard obstruction theories associated to the stack of maps between a fixed algebraic scheme and a smooth projective target, to the stack of stable maps to a smooth projective scheme or to the Picard stack of a smooth projective scheme, for the trace map, the Atiyah class map, and the first Chern class map for perfect complexes ([12, Chapter 5]), and for the map inducing O-M-P-T's reduced obstruction theory.

Organization of the paper. The first three sections and the beginning of the fifth are written for an *arbitrary* smooth complex projective scheme X . We explain how a derived extension induces an obstruction theory on its truncation (Section 1), how to define the standard derived extensions of the Picard stack of X , and of the stack of stable maps to X (Section 2), and finally define the perfect determinant map (Section 3). In Section 4, we *specialize* to the case where $X = S$ is a smooth complex projective $K3$ -surface. We first give a self-contained description of O-M-P-T's pointwise reduced tangent and obstruction spaces (Section 4.1). Then, by exploiting the features of the derived Picard stack of S (Section 4.2), we define in Sec-

Note 4:
unified to $K3$ -
surface (with
hyphen)

tion 4.3 a derived extension $\mathbb{R}\overline{\mathbf{M}}_g^{\text{red}}(S; \beta)$ of the usual stack $\overline{\mathbf{M}}_g(S; \beta)$ of stable maps of type $(g, \beta \neq 0)$ to S , having the property that, for the canonical inclusion

$$j_{\text{red}} : \overline{\mathbf{M}}_g(S; \beta) \hookrightarrow \mathbb{R}\overline{\mathbf{M}}_g^{\text{red}}(S; \beta),$$

the induced map

$$j_{\text{red}}^* \mathbb{L}_{\mathbb{R}\overline{\mathbf{M}}_g^{\text{red}}(S; \beta)} \rightarrow \mathbb{L}_{\overline{\mathbf{M}}_g(S; \beta)}$$

is a perfect obstruction theory with the same tangent and obstruction spaces as the reduced theory introduced by Maulik–Okounkov–Pandharipande–Thomas (Section 4.4, Theorem 4.8).

In Section 5, for a complex smooth projective variety X , we define the derived stack $\mathbb{R}\overline{\mathbf{M}}_{g,n}(X)^{\text{emb}}$ of pointed stable maps to X that are closed embeddings, the derived stack $\mathcal{M}_X \equiv \mathbb{R}\mathbf{Perf}(X)^{\text{si}, > 0}$ of simple perfect complexes on X with vanishing negative Ext's, and the derived stack $\mathcal{M}_{X, \mathcal{L}} \equiv \mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}^{\text{si}, > 0}$ of simple perfect complexes on X with vanishing negative Ext's and fixed determinant \mathcal{L} , and we define a morphism $C : \mathbb{R}\overline{\mathbf{M}}_{g,n}(X)^{\text{emb}} \rightarrow \mathcal{M}_{X, \mathcal{L}}$. When X is a K3-surface, we show that the truncation stack of \mathcal{M}_X is smooth. When X is a Calabi–Yau 3-fold, we prove that $\mathcal{M}_{X, \mathcal{L}}$ is quasi-smooth, and that the map C induces a map between the obstruction theories on the underlying underived stacks.

In an appendix we give a derived geometrical interpretation of the Atiyah class map and the first Chern class map for a perfect complex \mathbf{E} on a scheme Y , by relating them to the tangent of the corresponding map $\varphi_{\mathbf{E}} : Y \rightarrow \mathbb{R}\mathbf{Perf}$; then we follow this reinterpretation to prove some properties used in the main text.

Note 5:
you use ϕ and φ .
Unify?

Acknowledgement. Our initial interest in the possible relationships between reduced obstruction theories and derived algebraic geometry was positively boosted by comments and questions by B. Fantechi, D. Huybrechts and R. Thomas. We are grateful to R. Pandharipande for pointing out a useful classical statement, and to H. Flenner for some important remarks. We especially thanks A. Vistoli for generously sharing his expertise on stable maps with us, and R. Thomas for his interest and further comments on this paper.

The first author was supported by the SFB/TR 45 ‘Periods, Moduli Spaces and Arithmetic of Algebraic Varieties’ of the DFG (German Research Foundation). The second and third authors acknowledge financial support from the French ANR grant HODAG (ANR-09-BLAN-0151).

Frequently used notions: Notations and references. For background and basic notations in derived algebraic geometry we refer the reader to [36, Chapter 2.2] and to the overview [33, Sections 4.2 and 4.3]. In particular, $\mathbf{St}_{\mathbb{C}}$ (respectively, $\mathbf{dSt}_{\mathbb{C}}$) will denote the (homotopy) category of *stacks* (respectively, of *derived stacks*) on $\text{Spec } \mathbb{C}$ with respect to the étale (resp., strongly étale) topology. We will most often omit the inclusion functor $i : \mathbf{St}_{\mathbb{C}} \rightarrow \mathbf{dSt}_{\mathbb{C}}$ from our notations, since it is fully faithful; its left adjoint, the truncation functor, will be denoted $t_0 : \mathbf{dSt}_{\mathbb{C}} \rightarrow \mathbf{St}_{\mathbb{C}}$ ([36, Definition 2.2.4.3]). In particular, we will write $t_0(F) \hookrightarrow F$ for the adjunction morphism $i t_0(F) \hookrightarrow F$. However recall that the inclusion functor i does *not* commute with taking internal HOM (derived) stacks nor with taking homotopy limits. All fibered products of derived stacks will be implicitly derived (i.e. they will be homotopy fibered products in the model category of derived stacks).

When useful, we will freely use Quillen result and switch back and forth between (the model category of) simplicial commutative k -algebras and (the model category of) commuta-

tive differential non-positively graded k -algebras, where k is a field of characteristic 0 (details can be found also in [37, Appendix A]).

To any derived stack F , there is an associated dg-category $L_{\text{qcoh}}(F)$ of quasi-coherent complexes, and for any map $f : F \rightarrow G$ of derived stacks, we have a (left, right) adjunction

$$(\mathbb{L}f^* : L_{\text{qcoh}}(G) \rightarrow L_{\text{qcoh}}(F), \mathbb{R}f_* : L_{\text{qcoh}}(F) \rightarrow L_{\text{qcoh}}(G))$$

(see [33, Section 4.2] or [34, Section 1.1]).

All complexes will be cochain complexes and, for such a complex C^\bullet , either $C_{\leq n}$ or $C^{\leq n}$ (depending on typographical convenience) will denote its good truncation in degrees $\leq n$. Analogously for either $C_{\geq n}$ or $C^{\geq n}$ ([39, Section 1.2.7]).

To ease notation we will often write \otimes for the derived tensor product $\otimes^{\mathbb{L}}$ whenever no confusion is likely to arise.

In what follows, X will denote a smooth complex projective scheme while S a smooth complex projective $K3$ -surface.

As a purely terminological remark, for a given obstruction theory, we will call its *deformation space* what is usually called its *tangent space* (while we keep the terminology *obstruction space*). We do this to avoid confusion with tangent spaces, tangent complexes or tangent cohomologies of related (derived) stacks.

We will often abbreviate the list of authors Okounkov–Maulik–Pandharipande–Thomas to O-M-P-T.

1. Derived extensions, obstruction theories and their functoriality

We briefly recall here the basic observation that a derived extension of a given stack \mathcal{X} induces an obstruction theory (in the sense of [3]) on \mathcal{X} , and deduce a richer functoriality with respect to the one known classically. Everything in this section is true over an arbitrary base ring, though it will be stated for the base field \mathbb{C} .

1.1. Derived extensions induce obstruction theories. Let $t_0 : \mathbf{dSt}_{\mathbb{C}} \rightarrow \mathbf{St}_{\mathbb{C}}$ be the truncation functor between derived and underived stacks over \mathbb{C} for the étale topologies ([36, Def. 2.2.4.3]). It has a left adjoint $i : \mathbf{St}_{\mathbb{C}} \rightarrow \mathbf{dSt}_{\mathbb{C}}$ which is fully faithful (on the homotopy categories), and is therefore usually omitted from our notations.

Definition 1.1. Given a stack $\mathcal{X} \in \text{Ho}(\mathbf{St}_{\mathbb{C}})$, a *derived extension* of \mathcal{X} is a derived stack \mathcal{X}^{der} together with an isomorphism

$$\mathcal{X} \simeq t_0(\mathcal{X}^{\text{der}}).$$

Proposition 1.2. Let \mathcal{X}^{der} be a derived geometric stack which is a derived extension of the (geometric) stack \mathcal{X} . Then, the closed immersion

$$j : \mathcal{X} \simeq t_0(\mathcal{X}^{\text{der}}) \hookrightarrow \mathcal{X}^{\text{der}}$$

induces a morphism

$$j^*(\mathbb{L}\mathcal{X}^{\text{der}}) \rightarrow \mathbb{L}\mathcal{X}$$

which is 2-connective, i.e. its cone has vanishing cohomology in degrees ≥ -1 .

Proof. The proof follows easily from the remark that if A is a simplicial commutative \mathbb{C} -algebra and $A \rightarrow \pi_0(A)$ is the canonical surjection, then the cotangent complex $\mathbb{L}_{\pi_0(A)/A}$ is 2-connective, i.e. has vanishing cohomology in degrees ≥ -1 . \square

The previous proposition shows that a derived extension always induces an obstruction theory (whenever such a notion is defined by [3, Definition 4.4], e.g. when \mathcal{X} is a Deligne–Mumford stack).

Definition 1.3. A derived stack is *quasi-smooth* if it is locally of finite presentation and its cotangent complex is of perfect amplitude contained in $[-1, 0]$.

For quasi-smooth derived stacks we have the following result.

Corollary 1.4. *Let \mathcal{X}^{der} be a quasi-smooth derived Deligne–Mumford stack which is a derived extension of a (Deligne–Mumford) stack \mathcal{X} . Then*

$$j^*(\mathbb{L}_{\mathcal{X}^{\text{der}}}) \rightarrow \mathbb{L}_{\mathcal{X}}$$

is a $[-1, 0]$ -perfect obstruction theory as defined in [3, Definition 5.1].

1.2. Functoriality of deformation theories induced by derived extensions. If the map $f : X \rightarrow Y$ is a morphism of (Deligne–Mumford) stacks, and if $o_X : E_X \rightarrow \mathbb{L}_X$ and $o_Y : E_Y \rightarrow \mathbb{L}_Y$ are $([-1, 0]$ -perfect) obstruction theories, the classical theory of obstructions (see [3]) does not provide in general a map $f^*E_Y \rightarrow E_X$ such that the square

$$\begin{array}{ccc} f^*E_Y & \xrightarrow{f^*(o_Y)} & f^*\mathbb{L}_Y \\ \downarrow & & \downarrow \\ E_X & \xrightarrow{o_X} & \mathbb{L}_X \end{array}$$

is commutative (or commutative up to an isomorphism) in the derived category $D(X)$ of complexes on X , where $f^*\mathbb{L}_Y \rightarrow \mathbb{L}_X$ is the canonical map on cotangent complexes induced by f ([12, Chapter 2, (1.2.3.2)']). On the contrary, if $j_X : X \hookrightarrow \mathbb{R}X$ and $j_Y : Y \hookrightarrow \mathbb{R}Y$ are quasi-smooth derived (Deligne–Mumford) extensions of X and Y , respectively, and $F : \mathbb{R}X \rightarrow \mathbb{R}Y$ is a morphism of derived stacks

$$\begin{array}{ccc} X & \xrightarrow{t_0 F} & Y \\ \downarrow & & \downarrow \\ \mathbb{R}X & \xrightarrow{F} & \mathbb{R}Y, \end{array}$$

then $j_X^*\mathbb{L}_{\mathbb{R}X} \rightarrow \mathbb{L}_X$ and $j_Y^*\mathbb{L}_{\mathbb{R}Y} \rightarrow \mathbb{L}_Y$ are $([-1, 0]$ -perfect) obstruction theories by Corollary 1.4, and moreover there is indeed a canonical morphism of triangles in $D(X)$ (we denote $t_0(F)$ by f)

$$\begin{array}{ccccc} f^*j_Y^*\mathbb{L}_{\mathbb{R}Y} & \longrightarrow & f^*\mathbb{L}_Y & \longrightarrow & f^*\mathbb{L}_{\mathbb{R}Y/Y} \\ \downarrow & & \downarrow & & \downarrow \\ j_X^*\mathbb{L}_{\mathbb{R}X} & \longrightarrow & \mathbb{L}_X & \longrightarrow & \mathbb{L}_{\mathbb{R}X/X} \end{array}$$

(see [36, Proposition 1.2.1.6] or [12, Chapter 2, (2.1.1.5)]). This map relates the two induced obstruction theories and may be used to relate the corresponding virtual fundamental classes, too (when they exist). We will not do this here since we will not need it for the results in this paper. However, the type of result we are referring to is the following

Proposition 1.5 ([29, Theorem 7.4]). *Let $F : \mathbb{R}X \rightarrow \mathbb{R}Y$ be a quasi-smooth morphism between quasi-smooth Deligne–Mumford stacks, and $f : X \rightarrow Y$ be the induced morphism on the truncations. Then, there is an induced virtual pullback (as defined in [17])*

$$f^! : A_*(Y) \rightarrow A_*(X),$$

between the Chow groups of Y and X , such that $f^!([Y]^{\text{vir}}) = [X]^{\text{vir}}$, where $[X]^{\text{vir}}$ (resp., $[Y]^{\text{vir}}$) is the virtual fundamental class ([3]) on X (resp., on Y) induced by the $[-1, 0]$ perfect obstruction theory $j_X^ \mathbb{L}_{\mathbb{R}X} \rightarrow \mathbb{L}_X$ (resp., by $j_Y^* \mathbb{L}_{\mathbb{R}Y} \rightarrow \mathbb{L}_Y$).*

2. Derived stack of stable maps and derived Picard stack

In this section we prove a correspondence between derived open substacks of a derived stack and open substacks of its truncation, and use it to construct the derived Picard stack $\mathbb{R}\text{Pic}(X; \beta)$ of type $\beta \in H^2(X, \mathbb{Z})$, for any complex projective smooth variety X . After recalling the derived version of the stack of (pre-)stable maps to X , possibly pointed, the same correspondence will lead us to defining the derived stack $\mathbb{R}\overline{\mathbf{M}}_g(X; \beta)$ of stable maps of type (g, β) to X and its pointed version.

Throughout the section X will denote a smooth complex projective scheme, g a nonnegative integer, c_1 a class in $H^2(X, \mathbb{Z})$ (which, for our purposes, may be supposed to belong to the image of $\text{Pic}(X) \simeq H^1(X, \mathcal{O}_X^*) \rightarrow H^2(X, \mathbb{Z})$, i.e. belonging to $H^{1,1}(X) \cap H^2(X, \mathbb{Z})$), and $\beta \in H_2(X, \mathbb{Z})$ an effective curve class.

We will frequently use of the following

Proposition 2.1. *Let F be a derived stack and $t_0(F)$ its truncation. There is a bijective correspondence of equivalence classes*

$$\phi_F : \{\text{Zariski open substacks of } t_0(F)\} \rightarrow \{\text{Zariski open derived substacks of } F\}.$$

For any Zariski open substack $U_0 \hookrightarrow t_0(F)$, we have a homotopy cartesian diagram in $\mathbf{dSt}_{\mathbb{C}}$

$$\begin{array}{ccc} U_0 & \hookrightarrow & t_0(F) \\ \downarrow & & \downarrow \\ \phi_F(U_0) & \hookrightarrow & F \end{array}$$

where the vertical maps are the canonical closed immersions.

Proof. The statement is an immediate consequence of the fact that F and $t_0(F)$ have the same topology ([36, Corollary 2.2.2.9]). More precisely, let us define ϕ_F as follows. If $U_0 \hookrightarrow t_0(F)$ is an open substack, $\phi_F(U_0)$ is the functor

$$\mathbf{SAlg}_{\mathbb{C}} \rightarrow \mathbf{SSets} : A \mapsto F(A) \times_{t_0(F)(\pi_0(A))} U_0(\pi_0(A))$$

where $F(A)$ maps to $t_0(F)(\pi_0(A))$ via the morphism (induced by the truncation functor t_0)

$$F(A) \simeq \mathbb{R}\underline{\mathrm{Hom}}_{\mathbf{dSt}_{\mathbb{C}}}(\mathbb{R}\mathrm{Spec}(A), F) \rightarrow \mathbb{R}\underline{\mathrm{Hom}}_{\mathbf{St}_{\mathbb{C}}}(\mathbf{t}_0(\mathbb{R}\mathrm{Spec}(A)), \mathbf{t}_0(F)) \simeq \mathbf{t}_0(F)(\pi_0(A)).$$

The inverse to ϕ_F is simply induced by the truncation functor t_0 . \square

2.1. The derived Picard stack.

Definition 2.2. The *Picard stack* of X/\mathbb{C} is the stack

$$\mathbf{Pic}(X) := \mathbb{R}\mathrm{HOM}_{\mathbf{St}_{\mathbb{C}}}(X, B\mathbb{G}_m).$$

The *derived Picard stack* of X/\mathbb{C} is the derived stack

$$\mathbb{R}\mathbf{Pic}(X) := \mathbb{R}\mathrm{HOM}_{\mathbf{dSt}_{\mathbb{C}}}(X, B\mathbb{G}_m).$$

By definition we have a natural isomorphism

$$\mathbf{t}_0(\mathbb{R}\mathbf{Pic}(X)) \simeq \mathbf{Pic}(X)$$

in $\mathrm{Ho}(\mathbf{dSt}_{\mathbb{C}})$. Note that even though $\mathbf{Pic}(X)$ is smooth, it is *not* true that $\mathbb{R}\mathbf{Pic}(X) \simeq \mathbf{Pic}(X)$ if $\dim(X) > 1$; this can be seen on tangent spaces since

$$\mathbb{T}_L \mathbb{R}\mathbf{Pic}(X) \simeq C^\bullet(X, \mathcal{O}_X)[1] := \mathbb{R}\Gamma(X, \mathcal{O}_X)[1]$$

for any global point $x_L : \mathrm{Spec}(\mathbb{C}) \rightarrow \mathbb{R}\mathbf{Pic}(X)$ corresponding to a line bundle L over X .

Given $c_1 \in H^2(X, \mathbb{Z})$, we denote by $\mathbf{Pic}(X; c_1)$ the open substack of $\mathbf{Pic}(X)$ classifying line bundles with first Chern class c_1 . More precisely, for any $R \in \mathbf{Alg}_{\mathbb{C}}$, let us denote by $\mathrm{Vect}_1(R; c_1)$ the groupoid of line bundles L on $\mathrm{Spec}(R) \times X$ such that, for any point $x : \mathrm{Spec}(\mathbb{C}) \rightarrow \mathrm{Spec}(R)$ the pullback line bundle L_x on X has first Chern class equal to c_1 . Then, $\mathbf{Pic}(X; c_1)$ is the stack

$$\mathbf{Alg}_{\mathbb{C}} \rightarrow \mathbf{SSets} : R \mapsto \mathrm{Nerve}(\mathrm{Vect}_1(R; c_1))$$

where $\mathrm{Nerve}(C)$ is the nerve of the category C .

Note that we have

$$\mathbf{Pic}(X) = \coprod_{c_1 \in H^2(X, \mathbb{Z})} \mathbf{Pic}(X; c_1).$$

Definition 2.3. Let $c_1 \in H^2(X, \mathbb{Z})$. The *derived Picard stack of type c_1* of X/\mathbb{C} is the derived stack

$$\mathbb{R}\mathbf{Pic}(X; c_1) := \phi_{\mathbb{R}\mathbf{Pic}(X)}(\mathbf{Pic}(X; c_1)).$$

In particular, we have a natural isomorphism $\mathbf{t}_0(\mathbb{R}\mathbf{Pic}(X; c_1)) \simeq \mathbf{Pic}(X; c_1)$, and a homotopy cartesian diagram in $\mathbf{dSt}_{\mathbb{C}}$

$$\begin{array}{ccc} \mathbf{Pic}(X; c_1) & \hookrightarrow & \mathbf{Pic}(X) \\ \downarrow & & \downarrow \\ \mathbb{R}\mathbf{Pic}(X; c_1) & \hookrightarrow & \mathbb{R}\mathbf{Pic}(X). \end{array}$$

2.2. The derived stack of stable maps. We recall from [33, Section 4.3 (4.d)] the construction of the derived stack $\mathbb{R}\mathbf{M}_g^{\text{pre}}(X)$ (respectively, $\mathbb{R}\mathbf{M}_{g,n}^{\text{pre}}(X)$) of prestable maps (resp., of n -pointed prestable maps) of genus g to X , and of its open derived substack $\mathbb{R}\overline{\mathbf{M}}_g(X)$ (respectively, $\mathbb{R}\overline{\mathbf{M}}_{g,n}(X)$) of stable maps (resp., of n -pointed stable maps) of genus g to X . Then we move to define the derived version of the stack of (pointed) stable maps of type (g, β) to X .

Let $\mathbf{M}_g^{\text{pre}}$ (respectively, $\mathbf{M}_{g,n}^{\text{pre}}$) be the stack of (resp., n -pointed) pre-stable curves of genus g , and $\mathcal{C}_g^{\text{pre}} \rightarrow \mathbf{M}_g^{\text{pre}}$ (resp., $\mathcal{C}_{g,n}^{\text{pre}} \rightarrow \mathbf{M}_{g,n}^{\text{pre}}$) its universal family (see e.g. [1, 23]).

Definition 2.4. The derived stack $\mathbb{R}\mathbf{M}_g^{\text{pre}}(X)$ of prestable maps of genus g to X is defined as

$$\mathbb{R}\mathbf{M}_g^{\text{pre}}(X) := \mathbb{R}\text{HOM}_{\mathbf{dSt}_{\mathbb{C}}/\mathbf{M}_g^{\text{pre}}}(\mathcal{C}_g^{\text{pre}}, X \times \mathbf{M}_g^{\text{pre}}).$$

Then $\mathbb{R}\mathbf{M}_g^{\text{pre}}(X)$ is canonically a derived stack over $\mathbf{M}_g^{\text{pre}}$, and the corresponding *derived universal family* $\mathbb{R}\mathcal{C}_{g;X}^{\text{pre}}$ is defined by the following homotopy cartesian square:

$$\begin{array}{ccc} \mathbb{R}\mathcal{C}_{g;X}^{\text{pre}} & \longrightarrow & \mathbb{R}\mathbf{M}_g^{\text{pre}}(X) \\ \downarrow & & \downarrow \\ \mathcal{C}_g^{\text{pre}} & \longrightarrow & \mathbf{M}_g^{\text{pre}}. \end{array}$$

The derived stack $\mathbb{R}\mathbf{M}_{g,n}^{\text{pre}}(X)$ of n -pointed prestable maps of genus g to X is defined as

$$\mathbb{R}\mathbf{M}_{g,n}^{\text{pre}}(X) := \mathbb{R}\text{HOM}_{\mathbf{dSt}_{\mathbb{C}}/\mathbf{M}_{g,n}^{\text{pre}}}(\mathcal{C}_{g,n}^{\text{pre}}, X \times \mathbf{M}_{g,n}^{\text{pre}}).$$

Then $\mathbb{R}\mathbf{M}_{g,n}^{\text{pre}}(X)$ is canonically a derived stack over $\mathbf{M}_{g,n}^{\text{pre}}$, and the corresponding *derived universal family* $\mathbb{R}\mathcal{C}_{g,n;X}^{\text{pre}}$ is defined by the following homotopy cartesian square

$$\begin{array}{ccc} \mathbb{R}\mathcal{C}_{g,n;X}^{\text{pre}} & \longrightarrow & \mathbb{R}\mathbf{M}_{g,n}^{\text{pre}}(X) \\ \downarrow & & \downarrow \\ \mathcal{C}_{g,n}^{\text{pre}} & \longrightarrow & \mathbf{M}_{g,n}^{\text{pre}}. \end{array}$$

Remark 2.5. The derived stack $\mathbb{R}\mathbf{M}_{g,n}^{\text{pre}}(X)$ has the following derived-moduli space description. Roughly speaking, it associates to any simplicial commutative or differential non-positively graded \mathbb{C} -algebra A the nerve of equivalences of the category of pairs

$$(g : C \rightarrow \mathbb{R} \text{Spec } A, h : C \rightarrow X)$$

where g is a pointed proper flat pointed curve over $\mathbb{R} \text{Spec } A$, and h is a map that when restricted to the fiber over any complex point of $\mathbb{R} \text{Spec } A$, yields a stable map to X . We will not use this derived-moduli interpretation in the rest of the paper.

Note that, by definition, $\mathbb{R}\mathcal{C}_{g;X}^{\text{pre}}$ comes also equipped with a canonical map

$$\mathbb{R}\mathcal{C}_{g;X}^{\text{pre}} \rightarrow \mathbb{R}\mathbf{M}_g^{\text{pre}}(X) \times X.$$

We also have $t_0(\mathbb{R}\mathbf{M}_g^{\text{pre}}(X)) \simeq \mathbf{M}_g^{\text{pre}}(X)$ (the stack of prestable maps of genus g to X), and $t_0(\mathbb{R}\mathcal{C}_{g;X}^{\text{pre}}) \simeq \mathcal{C}_{g;X}^{\text{pre}}$ (the universal family over the stack of pre-stable maps of genus g to X), since the truncation functor t_0 commutes with homotopy fibered products. The same is true for the pointed version.

We can now use Proposition 2.1 to define the derived stable versions. Let $\overline{\mathbf{M}}_g(X)$ (respectively, $\overline{\mathbf{M}}_{g,n}(X)$) be the open substack of $\mathbf{M}_g^{\text{pre}}(X)$ (resp., of $\mathbf{M}_{g,n}^{\text{pre}}(X)$) consisting of *stable maps of genus g to X* (resp., *n -pointed stable maps of genus g to X*), and $\mathcal{C}_g; X \rightarrow \mathbf{M}_g^{\text{pre}}(X)$ (resp., $\mathcal{C}_{g,n}; X \rightarrow \mathbf{M}_{g,n}^{\text{pre}}(X)$) the (induced) universal family ([1, 23]).

Definition 2.6. The *derived stack $\mathbb{R}\overline{\mathbf{M}}_g(X)$ of stable maps of genus g to X* is defined as

$$\mathbb{R}\overline{\mathbf{M}}_g(X) := \phi_{\mathbb{R}\mathbf{M}_g^{\text{pre}}(X)}(\overline{\mathbf{M}}_g(X)).$$

The *derived stable universal family*

$$\mathbb{R}\mathcal{C}_g; X \rightarrow \mathbb{R}\overline{\mathbf{M}}_g(X)$$

is the derived restriction of $\mathbb{R}\mathcal{C}_g^{\text{pre}}; X \rightarrow \mathbb{R}\mathbf{M}_g^{\text{pre}}(X)$ to $\mathbb{R}\overline{\mathbf{M}}_g(X)$.

The *derived stack $\mathbb{R}\overline{\mathbf{M}}_{g,n}(X)$ of n -pointed stable maps of genus g to X* is defined as

$$\mathbb{R}\overline{\mathbf{M}}_{g,n}(X) := \phi_{\mathbb{R}\mathbf{M}_{g,n}^{\text{pre}}(X)}(\overline{\mathbf{M}}_{g,n}(X)).$$

The *derived stable universal family*

$$\mathbb{R}\mathcal{C}_{g,n}; X \rightarrow \mathbb{R}\overline{\mathbf{M}}_{g,n}(X)$$

is the derived restriction of $\mathbb{R}\mathcal{C}_{g,n}^{\text{pre}}; X \rightarrow \mathbb{R}\mathbf{M}_{g,n}^{\text{pre}}(X)$ to $\mathbb{R}\overline{\mathbf{M}}_{g,n}(X)$.

Recall that

- $t_0(\mathbb{R}\overline{\mathbf{M}}_g(X)) \simeq \overline{\mathbf{M}}_g(X)$;
- $t_0(\mathbb{R}\mathcal{C}_g; X) \simeq \mathcal{C}_g; X$;
- $\mathbb{R}\mathcal{C}_g; X$ comes equipped with a canonical map

$$\pi : \mathbb{R}\mathcal{C}_g; X \rightarrow \mathbb{R}\overline{\mathbf{M}}_g(X) \times X;$$

- we have a homotopy cartesian diagram in $\mathbf{dSt}_{\mathbb{C}}$

$$\begin{array}{ccc} \overline{\mathbf{M}}_g(X) & \hookrightarrow & \mathbf{M}_g^{\text{pre}}(X) \\ \downarrow & & \downarrow \\ \mathbb{R}\overline{\mathbf{M}}_g(X) & \hookrightarrow & \mathbb{R}\mathbf{M}_g^{\text{pre}}(X). \end{array}$$

With the obvious changes, this applies to the pointed version too.

Let g be a non-negative integer, $\beta \in H_2(X, \mathbb{Z})$, and $\overline{\mathbf{M}}_g(X; \beta)$ (resp., $\overline{\mathbf{M}}_{g,n}(X; \beta)$) be the stack of stable maps (resp., of n -pointed stable maps) of type (g, β) to X (see e.g. [1] or [23]); its derived version is given by the following

Definition 2.7. The *derived stack of stable maps of type (g, β) to X* is defined as the open substack of $\mathbb{R}\overline{\mathbf{M}}_g(X)$

$$\mathbb{R}\overline{\mathbf{M}}_g(X; \beta) := \phi_{\mathbb{R}\overline{\mathbf{M}}_g(X)}(\overline{\mathbf{M}}_g(X; \beta)).$$

The *derived stable universal family of type* $(g; \beta)$,

$$\mathbb{R}\mathcal{C}_{g,\beta;X} \rightarrow \mathbb{R}\overline{\mathbf{M}}_g(X; \beta),$$

is the (derived) restriction of $\mathbb{R}\mathcal{C}_g; X \rightarrow \mathbb{R}\overline{\mathbf{M}}_g(X)$ to $\mathbb{R}\overline{\mathbf{M}}_g(X; \beta)$.

The *derived stack of n -pointed stable maps of type* (g, β) *to* X is defined as the open substack of $\mathbb{R}\overline{\mathbf{M}}_{g,n}(X)$

$$\mathbb{R}\overline{\mathbf{M}}_{g,n}(X; \beta) := \phi_{\mathbb{R}\overline{\mathbf{M}}_{g,n}(X)}(\overline{\mathbf{M}}_{g,n}(X; \beta)).$$

The *derived stable universal family of type* $(g; \beta)$,

$$\mathbb{R}\mathcal{C}_{g,n,\beta;X} \rightarrow \mathbb{R}\overline{\mathbf{M}}_{g,n}(X; \beta),$$

is the (derived) restriction of $\mathbb{R}\mathcal{C}_{g,n}; X \rightarrow \mathbb{R}\overline{\mathbf{M}}_{g,n}(X)$ to $\mathbb{R}\overline{\mathbf{M}}_{g,n}(X; \beta)$.

Remark 2.8. A derived-moduli space description similar to the one given in Remark 2.5 is available for $\mathbb{R}\overline{\mathbf{M}}_g(X)$, $\mathbb{R}\overline{\mathbf{M}}_{g,n}(X)$, $\mathbb{R}\overline{\mathbf{M}}_g(X; \beta)$, and $\mathbb{R}\overline{\mathbf{M}}_{g,n}(X; \beta)$. We leave the details to the reader, since we will not need this result in the rest of the paper.

The fiber of the canonical projection map $\mathbb{R}\mathbf{M}_g^{\text{pre}}(X) \rightarrow \mathbf{M}_g^{\text{pre}}$ over the derived point $x_A : \mathbb{R} \operatorname{Spec} A \rightarrow \mathbf{M}_g^{\text{pre}}$ (A being a cdga) is $\mathbb{R}\operatorname{HOM}_{\mathbf{dSt}_{\mathbb{C}}/\mathbb{R} \operatorname{Spec} A}(C, X)$, where C is the curve over $\mathbb{R} \operatorname{Spec} A$ corresponding to x_A . By [36, Corollary 2.2.6.14], this fiber is geometric, and therefore the projection

$$\mathbb{R}\mathbf{M}_g^{\text{pre}}(X) \rightarrow \mathbf{M}_g^{\text{pre}}$$

is representable. Therefore the derived stack $\mathbb{R}\mathbf{M}_g^{\text{pre}}(X)$, as well as $\mathbb{R}\overline{\mathbf{M}}_g(X; \beta)$ are geometric. Moreover, by definition, $t_0(\mathbb{R}\overline{\mathbf{M}}_g(X; \beta)) \simeq \overline{\mathbf{M}}_g(X; \beta)$, thus $\mathbb{R}\overline{\mathbf{M}}_g(X; \beta)$ is a proper derived Deligne–Mumford stack ([36, Section 2.2.4]).

By the transitivity triangle associated to the representable map $\mathbb{R}\mathbf{M}_g^{\text{pre}}(X) \rightarrow \mathbf{M}_g^{\text{pre}}(X)$, the tangent complex $\mathbb{T}_{(f:C \rightarrow X)}$ of the derived stack $\mathbb{R}\overline{\mathbf{M}}_g(X; \beta)$ at a stable map $(f : C \rightarrow X)$ of type (g, β) (corresponding to a classical point $x_f : \operatorname{Spec}(\mathbb{C}) \rightarrow \mathbb{R}\overline{\mathbf{M}}_g(X; \beta)$) is given by¹⁾

$$\mathbb{T}_{(f:C \rightarrow X)} \simeq \mathbb{R}\Gamma(C, \operatorname{Cone}(\mathbb{T}_C \rightarrow f^*\mathbb{T}_X)),$$

where \mathbb{T}_C is the tangent complex of C and \mathbb{T}_X is the tangent sheaf of X .

The canonical map $\mathbb{R}\overline{\mathbf{M}}_g(X; \beta) \rightarrow \mathbf{M}_g^{\text{pre}}$ is quasi-smooth. In fact, the fiber at a geometric point, corresponding to prestable curve C , is the derived stack $\mathbb{R}\operatorname{HOM}_{\beta}(C, X)$ whose tangent complex at a point $f : C \rightarrow X$ is $\mathbb{R}\Gamma(C, f^*\mathbb{T}_X)$ which, obviously, has cohomology only in degrees $[0, 1]$. But $\mathbf{M}_g^{\text{pre}}$ is smooth, and any derived stack quasi-smooth over a smooth base is quasi-smooth (by the corresponding exact triangle of tangent complexes). Therefore the derived stack $\mathbb{R}\overline{\mathbf{M}}_g(X; \beta)$ is *quasi-smooth*.

Proposition 1.2 **above** then recovers the *standard* (absolute) perfect obstruction theory on $\overline{\mathbf{M}}_g(X; \beta)$ via the canonical map

$$j^*(\mathbb{L}_{\mathbb{R}\overline{\mathbf{M}}_g(X; \beta)}) \rightarrow \mathbb{L}_{\overline{\mathbf{M}}_g(X; \beta)}$$

induced by the closed immersion $j : \overline{\mathbf{M}}_g(X; \beta) \hookrightarrow \mathbb{R}\overline{\mathbf{M}}_g(X; \beta)$.

¹⁾ As communicated by the authors, the $[1]$ -shift in [8, Theorem 5.4.8] is just a typo: their proof is correct and yields no shift.

Note 6:
removed ‘By’ for
line break

In the *pointed* case, the same argument used above for the unpointed case shows that the map $\mathbb{R}\mathbf{M}_{g,n}^{\text{pre}}(X) \rightarrow \mathbf{M}_{g,n}^{\text{pre}}$ is representable, and that the tangent complex of $\mathbb{R}\overline{\mathbf{M}}_{g,n}(X; \beta)$ at a pointed stable map $(f : (C; x_1, \dots, x_n) \rightarrow X)$ of type (g, β) (corresponding to a classical point $x_f : \text{Spec}(\mathbb{C}) \rightarrow \mathbb{R}\overline{\mathbf{M}}_{g,n}(X; \beta)$) is likewise given by

$$\mathbb{T}_{(f:(C;x_1,\dots,x_n)\rightarrow X)} \simeq \mathbb{R}\Gamma\left(C, \text{Cone}\left(\mathbb{T}_C\left(-\sum_i x_i\right) \rightarrow f^*\mathbb{T}_X\right)\right).$$

The pointed variant of the argument above, proving quasi-smoothness of $\mathbb{R}\overline{\mathbf{M}}_g(X; \beta) \rightarrow \mathbf{M}_g^{\text{pre}}$, proves that also the canonical map $\mathbb{R}\overline{\mathbf{M}}_{g,n}(X; \beta) \rightarrow \mathbf{M}_{g,n}^{\text{pre}}$ is quasi-smooth, and Proposition 1.2 then recovers the *standard* absolute perfect obstruction theory on $\overline{\mathbf{M}}_{g,n}(X; \beta)$ via the canonical map

$$j^*(\mathbb{L}_{\mathbb{R}\overline{\mathbf{M}}_{g,n}(X;\beta)}) \rightarrow \mathbb{L}_{\overline{\mathbf{M}}_{g,n}(X;\beta)}$$

induced by the closed immersion

$$j : \overline{\mathbf{M}}_{g,n}(X; \beta) \hookrightarrow \mathbb{R}\overline{\mathbf{M}}_{g,n}(X; \beta).$$

Note that, as observed in [23, Section 5.3.5], this obstruction theory yields trivial Gromov–Witten invariants on $\overline{\mathbf{M}}_{g,n}(X; \beta)$ for $X = S$ a $K3$ -surface. Hence the need for another obstruction theory carrying more interesting curves counting invariants on a $K3$ -surface: this will be the so-called *reduced* obstruction theory (see Section 4.1, Section 4.3, and Theorem 4.8).

Finally, the derived stable universal family $\mathbb{R}\mathcal{C}_{g,\beta;X}$ comes, by restriction, equipped with a natural map

$$\pi : \mathbb{R}\mathcal{C}_{g,\beta;X} \rightarrow \mathbb{R}\overline{\mathbf{M}}_g(X; \beta) \times X.$$

We have a homotopy cartesian diagram in $\mathbf{dSt}_{\mathbb{C}}$

$$\begin{array}{ccc} \overline{\mathbf{M}}_g(X; \beta) & \hookrightarrow & \overline{\mathbf{M}}_g(X) \\ \downarrow & & \downarrow \\ \mathbb{R}\overline{\mathbf{M}}_g(X; \beta) & \hookrightarrow & \mathbb{R}\overline{\mathbf{M}}_g(X). \end{array}$$

Analogous remarks hold in the *pointed* case.

3. The derived determinant morphism

We recall from [35, 36] the definition of the derived stack $\mathbb{R}\mathbf{Perf}$ (denoted as \mathcal{M}_1 in loc. cit). The functor $\mathbb{R}\mathbf{Perf}$ sends a differential non-positively graded \mathbb{C} -algebra A to the nerve of the category of perfect (i.e. homotopically finitely presentable, or equivalently, dualizable in the monoidal model category of A -dg-modules) A -dg-modules which are cofibrant in the projective model structure of all A -dg-modules. It is a locally geometric derived stack, that is a union of open substacks which are derived Artin stacks of finite presentation over $\text{Spec } \mathbb{C}$ ([35, Proposition 3.7]).

For a derived stack Y , the derived stack of perfect complexes on Y is

$$\mathbb{R}\mathbf{Perf}(Y) := \mathbb{R}\mathbf{HOM}_{\mathbf{dSt}_{\mathbb{C}}}(Y, \mathbb{R}\mathbf{Perf}),$$

and the space (or simplicial set) of perfect complexes on Y is, by definition, the mapping space $\mathbb{R}\underline{\mathrm{Hom}}_{\mathbf{dSt}_{\mathbb{C}}}(Y, \mathbb{R}\mathbf{Perf})$ in the model category $\mathbf{dSt}_{\mathbb{C}}$; an element in its π_0 is called a perfect complex on Y . Note that

$$\mathbb{R}\underline{\mathrm{Hom}}_{\mathbf{dSt}_{\mathbb{C}}}(Y, \mathbb{R}\mathbf{Perf}) \simeq \mathbb{R}\mathbf{Perf}(X)(\mathbb{C}).$$

In this section we start by defining a quite general *perfect determinant map* of derived stacks

$$\det_{\mathbf{Perf}} : \mathbb{R}\mathbf{Perf} \rightarrow \mathbf{Pic} = B\mathbb{G}_m$$

whose construction requires a small detour into Waldhausen K -theory. We think this perfect determinant might play an important role in other contexts as well, e.g. in a general GW/DT correspondence.

Using the perfect determinant together with a natural perfect complex on $\mathbb{R}\overline{\mathbf{M}}_g(X; \beta)$, we will be able to define a map

$$\delta_1(X) : \mathbb{R}\overline{\mathbf{M}}_g(X) \rightarrow \mathbb{R}\mathbf{Pic}(X)$$

which will be one of the main ingredients in the construction of the *reduced* derived stack of stable maps $\mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta)$, for a $K3$ -surface S , given in the next section.

3.1. The perfect determinant map. The aim of this subsection is to produce a determinant morphism $\det_{\mathbf{Perf}} : \mathbb{R}\mathbf{Perf} \rightarrow \mathbf{Pic}$ in $\mathrm{Ho}(\mathbf{dSt}_{\mathbb{C}})$ extending the natural determinant morphism $\mathbf{Vect} \rightarrow \mathbf{Pic}$. To do this, we will have to pass through Waldhausen K -theory.

By [36, Lemma 2.2.6.1] we do not have to distinguish between the stack and derived stack of vector bundles \mathbf{Vect} : if $i : \mathbf{St}_{\mathbb{C}} \rightarrow \mathbf{dSt}_{\mathbb{C}}$ is the canonical functor viewing a stack as a derived stack, we have a canonical equivalence $i(\mathbf{Vect}) \simeq \mathbb{R}\mathbf{Vect}$. We will then simply write \mathbf{Vect} for either \mathbf{Vect} or $\mathbb{R}\mathbf{Vect}$.

We start with the classical determinant map in $\mathrm{Ho}(\mathbf{St}_{\mathbb{C}})$, $\det : \mathbf{Vect} \rightarrow \mathbf{Pic}$, induced by the map sending a vector bundle to its top exterior power. Consider the following simplicial stacks:

$$B_{\bullet}\mathbf{Pic} : \Delta^{\mathrm{op}} \ni [n] \mapsto (\mathbf{Pic})^n$$

(with the simplicial structure maps given by tensor products of line bundles, or equivalently, induced by the product in the group structure of $B\mathbb{G}_m \simeq \mathbf{Pic}$), and

$$B_{\bullet}\mathbf{Vect} : \Delta^{\mathrm{op}} \ni [n] \mapsto wS_n\mathbf{Vect},$$

where, for any commutative \mathbb{C} -algebra R , $wS_n\mathbf{Vect}(R)$ is the nerve of the category of sequences of split monomorphisms

$$0 \rightarrow M_1 \rightarrow M_2 \rightarrow \cdots \rightarrow M_n \rightarrow 0$$

with morphisms the obvious equivalences, and the simplicial structure maps are the natural ones described in [38, Section 1.3]. Similarly, we define the simplicial object in stacks

$$B_{\bullet}\mathbf{Perf} : \Delta^{\mathrm{op}} \ni [n] \mapsto wS_n\mathbf{Perf}$$

(see [38, Section 1.3] for the definition of wS_n in this case). Now, $B_{\bullet}\mathbf{Pic}$ and $B_{\bullet}\mathbf{Vect}$, and $B_{\bullet}\mathbf{Perf}$ are pre- Δ^{op} -stacks according to [32, Definition 1.4.1], and the map \det extends to

a morphism

$$\det_{\bullet} : B_{\bullet}\mathbf{Vect} \rightarrow B_{\bullet}\mathbf{Pic}$$

in the homotopy category of pre- Δ^{op} -stacks. By applying the functor $i : \text{Ho}(\mathbf{St}_{\mathbb{C}}) \rightarrow \text{Ho}(\mathbf{dSt}_{\mathbb{C}})$ (that will be, according to our conventions, omitted from notations), we get a determinant morphism (denoted in the same way)

$$\det_{\bullet} : B_{\bullet}\mathbf{Vect} \rightarrow B_{\bullet}\mathbf{Pic}$$

in the homotopy category of pre- Δ^{op} -derived stacks. We now pass to Waldhausen K -theory, i.e. apply $K := \Omega \circ |-|$ (see [32, Theorem 1.4.3], where the loop functor Ω is denoted by $\mathbb{R}\Omega_{*}$, and the realization functor $|-|$ by B), and observe that, by [32, Theorem 1.4.3 (2)], there is a canonical isomorphism in $\text{Ho}(\mathbf{dSt}_{\mathbb{C}})$

$$K(B_{\bullet}\mathbf{Pic}) \simeq \mathbf{Pic}$$

since \mathbf{Pic} is group-like (i.e. an H_{∞} -stack in the parlance of [32, Theorem 1.4.3]). This gives us a map in $\text{Ho}(\mathbf{dSt}_{\mathbb{C}})$

$$K(\det_{\bullet}) : K(B_{\bullet}\mathbf{Vect}) \rightarrow \mathbf{Pic}.$$

Now, consider the map

$$u : \mathbf{K}^{\text{Vect}} := K(B_{\bullet}\mathbf{Vect}) \rightarrow K(B_{\bullet}\mathbf{Perf}) := \mathbf{K}^{\text{Perf}}$$

in $\text{Ho}(\mathbf{dSt}_{\mathbb{C}})$, induced by the inclusion $\mathbf{Vect} \hookrightarrow \mathbb{R}\mathbf{Perf}$. By [38, Theorem 1.7.1], u is an isomorphism in $\text{Ho}(\mathbf{dSt}_{\mathbb{C}})$. Therefore, we get a diagram in $\text{Ho}(\mathbf{dSt}_{\mathbb{C}})$

$$\begin{array}{ccc} \mathbf{K}^{\text{Vect}} & \xrightarrow{K(\det_{\bullet})} & \mathbf{Pic} \\ u \downarrow & & \\ \mathbb{R}\mathbf{Perf} & \xrightarrow{\text{1st-level}} & \mathbf{K}^{\text{Perf}} \end{array}$$

where u is an isomorphism. This allows us to give the following

Definition 3.1. The induced map in $\text{Ho}(\mathbf{dSt}_{\mathbb{C}})$

$$\det_{\text{Perf}} : \mathbb{R}\mathbf{Perf} \rightarrow \mathbf{Pic}$$

is called the *perfect determinant* morphism.

For any complex scheme (or derived stack) X , the perfect determinant morphism

$$\det_{\text{Perf}} : \mathbb{R}\mathbf{Perf} \rightarrow \mathbf{Pic}$$

induces a map in $\text{Ho}(\mathbf{dSt}_{\mathbb{C}})$

$$\det_{\text{Perf}}(X) : \mathbb{R}\mathbf{Perf}(X) := \mathbb{R}\text{HOM}_{\mathbf{dSt}_{\mathbb{C}}}(X, \mathbf{Perf}) \rightarrow \mathbb{R}\text{HOM}_{\mathbf{dSt}_{\mathbb{C}}}(X, \mathbf{Pic}) =: \mathbb{R}\mathbf{Pic}(X).$$

As perhaps not totally unexpected (e.g. [12, Remark 5.3.3]), the tangent morphism to the perfect determinant map is given by the *trace for perfect complexes*. We state the result here only for complex points of $\mathbb{R}\mathbf{Perf}(X)$ because we will only need this case in the rest of the paper.

Proposition 3.2. *Let X be a complex quasi-projective scheme, and*

$$\det_{\text{Perf}}(X) : \mathbb{R}\text{Perf}(X) \rightarrow \mathbb{R}\text{Pic}(X)$$

be the induced perfect determinant map. For any complex point $x_E : \text{Spec } \mathbb{C} \rightarrow \mathbb{R}\text{Perf}(X)$, corresponding to a perfect complex E over X , the tangent map

$$\mathbb{T}_{x_E} \det_{\text{Perf}}(X) : \mathbb{T}_{x_E} \mathbb{R}\text{Perf}(X) \simeq \mathbb{R}\text{Hom}(E, E)[1] \rightarrow \mathbb{R}\text{Hom}(\mathcal{O}_S, \mathcal{O}_S)[1] \simeq \mathbb{T}_{x_E} \mathbb{R}\text{Pic}(X)$$

is given by $\text{tr}_E[1]$, where tr_E is the trace map for the perfect complex E of [12, Chapter 5, Section 3.7.3].

Proof. Let $\mathbb{R}\text{Perf}^{\text{strict}}(X) := \mathbb{R}\text{HOM}_{\mathbf{dSt}_{\mathbb{C}}}(X, \mathbb{R}\text{Perf}^{\text{strict}})$ be the derived stack of *strict* ([5, Exposé I, Section 2.1]) perfect complexes on X . Since X is quasi-projective, the canonical map $\mathbb{R}\text{Perf}^{\text{strict}}(X) \rightarrow \mathbb{R}\text{Perf}(X)$ is an isomorphism in $\text{Ho}(\mathbf{dSt}_{\mathbb{C}})$. Therefore (e.g. [5, Exposé I, Section 8.1.2]), the comparison statement is reduced to the case where E is a vector bundle on X , which is a direct computation and is left to the reader. \square

3.2. The map $\mathbb{R}\overline{\mathbf{M}}_g(X) \rightarrow \mathbb{R}\text{Perf}(X)$. A map

$$\mathbb{R}\overline{\mathbf{M}}_g(X) \rightarrow \mathbb{R}\text{Perf}(X) = \mathbb{R}\text{HOM}_{\mathbf{dSt}_{\mathbb{C}}}(X, \mathbb{R}\text{Perf})$$

in $\text{Ho}(\mathbf{dSt}_{\mathbb{C}})$ is, by adjunction, the same thing as a map

$$\mathbb{R}\overline{\mathbf{M}}_g(X) \times X \rightarrow \mathbb{R}\text{Perf}$$

i.e. a perfect complex on $\mathbb{R}\overline{\mathbf{M}}_g(X) \times X$; so, it is enough to find such an appropriate perfect complex.

Let

$$\pi : \mathbb{R}\mathcal{C}_{g;X} \rightarrow \mathbb{R}\overline{\mathbf{M}}_g(X) \times X$$

be the derived stable universal family (Section 2.2), and recall the existence of a derived direct image functor $\mathbb{R}\pi_* : \text{L}_{\text{qcoh}}(\mathbb{R}\mathcal{C}_{g;X}) \rightarrow \text{L}_{\text{qcoh}}(\mathbb{R}\overline{\mathbf{M}}_g(X) \times X)$ (see the section “Frequently used notions: Notations and references” in the Introduction).

Proposition 3.3. *The set $\mathbb{R}\pi_*(\mathcal{O}_{\mathbb{R}\mathcal{C}_{g;X}})$ is a perfect complex on $\mathbb{R}\overline{\mathbf{M}}_g(X) \times X$.*

Proof. First of all, π is representable, and the truncation of π is proper. Moreover π is quasi-smooth. To see this, observe that both $\mathbb{R}\mathcal{C}_{g;X}$ and $\mathbb{R}\overline{\mathbf{M}}_g(X) \times X$ are smooth over $\mathbb{R}\overline{\mathbf{M}}_g(X)$. Then we conclude, since any map between derived stacks smooth over a base is quasi-smooth. So we have that π is representable, proper and quasi-smooth. Since the statement is local on the target, we conclude by [34, Lemma 2.2]. \square

Remark 3.4. If we fix a class $\beta \in H_2(X, \mathbb{Z})$, the corresponding β -decorated version of Proposition 3.3 obviously holds.

We may therefore give the following

Definition 3.5. We will denote by

$$\mathbf{A}_X : \mathbb{R}\overline{\mathbf{M}}_g(X) \rightarrow \mathbb{R}\text{Perf}(X)$$

the map induced by the perfect complex $\mathbb{R}\pi_*(\mathcal{O}_{\mathbb{R}\mathcal{C}_{g;X}})$.

Note that, in particular, A_X sends a complex point of $\mathbb{R}\overline{\mathbf{M}}_g(X)$, corresponding to a stable map $f : C \rightarrow X$ to the perfect complex $\mathbb{R}f_*\mathcal{O}_C$ on X .

The tangent morphism of A_X . The tangent morphism of A_X is related to the Atiyah class of $\mathbb{R}\pi_*(\mathcal{O}_{\mathbb{R}\mathcal{C}_{g;X}})$, and pointwise on $\mathbb{R}\overline{\mathbf{M}}_g(X)$ to the Atiyah class map of the perfect complex $\mathbb{R}f_*\mathcal{O}_C$: this is explained in detail in Appendix A, so we will just recall here the results and the notations we will need in the rest of the main text.

Let us write $\mathcal{E} := \mathbb{R}\pi_*(\mathcal{O}_{\mathbb{R}\mathcal{C}_{g;X}})$; since this is a perfect complex on $\mathbb{R}\overline{\mathbf{M}}_g(X) \times X$, its Atiyah class map (see Appendix A)

$$\mathrm{at}_{\mathcal{E}} : \mathcal{E} \rightarrow \mathbb{L}_{\mathbb{R}\overline{\mathbf{M}}_g(X) \times X} \otimes \mathcal{E}[1]$$

corresponds uniquely, by adjunction, to a map, denoted in the same way,

$$\mathrm{at}_{\mathcal{E}} : \mathbb{T}_{\mathbb{R}\overline{\mathbf{M}}_g(X) \times X} \rightarrow \mathcal{E}^{\vee} \otimes \mathcal{E}[1].$$

Let x be a complex point x of $\mathbb{R}\overline{\mathbf{M}}_g(X)$ corresponding to a stable map $f : C \rightarrow X$, and let $p : C \rightarrow \mathrm{Spec} \mathbb{C}$ and $q : X \rightarrow \mathrm{Spec} \mathbb{C}$ denote the structural morphisms, so that $p = q \circ f$. Correspondingly, we have a ladder of homotopy cartesian diagrams

$$\begin{array}{ccccc} C & \xrightarrow{\iota_f} & \mathbb{R}\mathcal{C}_{g;X} & & \\ f \downarrow & & \downarrow \pi & & \\ X & \xrightarrow{x} & \mathbb{R}\overline{\mathbf{M}}_g(X) \times X & \xrightarrow{\mathrm{pr}_X} & X \\ q \downarrow & & \downarrow \mathrm{pr} & & \downarrow q \\ \mathrm{Spec} \mathbb{C} & \xrightarrow{x} & \mathbb{R}\overline{\mathbf{M}}_g(X) & \longrightarrow & \mathrm{Spec} \mathbb{C}. \end{array}$$

Let us consider the perfect complex $E := \mathbb{R}f_*\mathcal{O}_C$ on X . By [12, Chapter 4, [Section 2.3.7](#)], the complex E has an Atiyah class map

$$\mathrm{at}_E : E \rightarrow E \otimes \Omega_X^1[1]$$

which corresponds uniquely (E being perfect) by adjunction to a map (denoted in the same way)

$$\mathrm{at}_E : T_X \rightarrow \mathbb{R}\underline{\mathrm{End}}_X(\mathbb{R}f_*\mathcal{O}_C)[1].$$

Proposition 3.6. *In the situation and notations above, we have that:*

- *the tangent map of A_X fits into the following commutative diagram:*

$$\begin{array}{ccccc} \mathbb{T}_{\mathbb{R}\overline{\mathbf{M}}_g(X)} & \xrightarrow{\mathbb{T}A_X} & A_X^* \mathbb{T}_{\mathbb{R}\mathrm{Perf}(X)} & \xrightarrow{\sim} & \mathbb{R}\mathrm{pr}_*(\mathcal{E}^{\vee} \otimes \mathcal{E})[1] \\ \mathrm{can} \downarrow & & & & \uparrow \mathbb{R}\mathrm{pr}_*(\mathrm{at}_{\mathcal{E}}) \\ \mathbb{R}\mathrm{pr}_* \mathrm{pr}^* \mathbb{T}_{\mathbb{R}\overline{\mathbf{M}}_g(X)} & \xrightarrow{\mathrm{can}} & \mathbb{R}\mathrm{pr}_*(\mathrm{pr}^* \mathbb{T}_{\mathbb{R}\overline{\mathbf{M}}_g(X)} \oplus \mathrm{pr}_X^* T_X) & \xrightarrow{\sim} & \mathbb{R}\mathrm{pr}_* \mathbb{T}_{\mathbb{R}\overline{\mathbf{M}}_g(X) \times X} \end{array}$$

where can denote obvious canonical maps, and $\mathcal{E} := \mathbb{R}\pi_*(\mathcal{O}_{\mathbb{R}\mathcal{C}_{g;X}})$.

Note 7:
pr upright

- The tangent map to A_X at $x = (f : C \rightarrow X)$, is the composition

$$\begin{aligned} T_x A_X : T_x \mathbb{R}\overline{\mathbf{M}}_g(X) &\simeq \mathbb{R}\Gamma(C, \text{Cone}(T_C \rightarrow f^* T_X)) \rightarrow \mathbb{R}\Gamma(X, x^* T_{\mathbb{R}\overline{\mathbf{M}}_g(X) \times X}) \\ &\xrightarrow{\mathbb{R}\Gamma(X, x^* \text{at}_E)} \mathbb{R}\text{End}_X(\mathbb{R}f_* \mathcal{O}_C)[1] \simeq T_{\mathbb{R}f_* \mathcal{O}_C} \mathbb{R}\mathbf{Perf}(X) \end{aligned}$$

where $E := \mathbb{R}f_* \mathcal{O}_C$.

- The composition

$$\begin{aligned} \mathbb{R}\Gamma(X, T_X) &\xrightarrow{\text{can}} \mathbb{R}\Gamma(X, \mathbb{R}f_* f^* T_X) \\ &\xrightarrow{\text{can}} \mathbb{R}\Gamma(X, \text{Cone}(\mathbb{R}f_* T_C \rightarrow \mathbb{R}f_* f^* T_X)) \simeq T_x \mathbb{R}\overline{\mathbf{M}}_g(X) \\ &\xrightarrow{T_x A_X} x^* A_X^* T \mathbb{R}\mathbf{Perf}(X) \simeq T_{\mathbb{R}f_* \mathcal{O}_C} \mathbb{R}\mathbf{Perf}(X) \simeq \mathbb{R}\text{End}_X(\mathbb{R}f_* \mathcal{O}_C)[1] \end{aligned}$$

coincides with $\mathbb{R}\Gamma(X, \text{at}_E)$, where $E := \mathbb{R}f_* \mathcal{O}_C$.

Proof. See Appendix A. □

Definition 3.7. We denote by $\delta_1(X)$ the composition

$$\mathbb{R}\overline{\mathbf{M}}_g(X) \xrightarrow{A_X} \mathbb{R}\mathbf{Perf}(X) \xrightarrow{\det_{\mathbf{Perf}}(X)} \mathbb{R}\mathbf{Pic}(X),$$

and, for a complex point x of $\mathbb{R}\overline{\mathbf{M}}_g(X)$ corresponding to a stable map $f : C \rightarrow X$, by

$$\begin{aligned} \Theta_f := T_f \delta_1(X) : T_{(f:C \rightarrow X)} \mathbb{R}\overline{\mathbf{M}}_g(X) &\xrightarrow{T_x A_X} T_{\mathbb{R}f_* \mathcal{O}_C} \mathbb{R}\mathbf{Perf}(X) \\ &\xrightarrow{\text{tr}_X} T_{\det(\mathbb{R}f_* \mathcal{O}_C)} \mathbb{R}\mathbf{Pic}(X). \end{aligned}$$

Note that, as a map of explicit complexes, we have

$$\begin{aligned} \Theta_f : \mathbb{R}\Gamma(C, \text{Cone}(T_C \rightarrow f^* T_X)) &\xrightarrow{T_x A_X} \mathbb{R}\text{Hom}_X(\mathbb{R}f_* \mathcal{O}_C, \mathbb{R}f_* \mathcal{O}_C)[1] \\ &\xrightarrow{\text{tr}_X} \mathbb{R}\Gamma(X, \mathcal{O}_X)[1]. \end{aligned}$$

Remark 3.8 (First Chern class of $\mathbb{R}f_* \mathcal{O}_C$ and the map Θ_f). Let x be a complex point x of $\mathbb{R}\overline{\mathbf{M}}_g(X)$ corresponding to a stable map $f : C \rightarrow X$, and let $p : C \rightarrow \text{Spec } \mathbb{C}$ and $q : X \rightarrow \text{Spec } \mathbb{C}$ denote the structural morphisms, so that $p = q \circ f$. Using Proposition 3.6, we can relate the map Θ_f above to the *first Chern class* of the perfect complex $\mathbb{R}f_* \mathcal{O}_C$ (see [12, Chapter V]). With the same notations as in Proposition 3.6, the following diagram is commutative:

$$\begin{array}{ccccc} \mathbb{R}q_* T_X & \xrightarrow{\mathbb{R}q_*(\text{at}_{\mathbb{R}f_* \mathcal{O}_C})} & \mathbb{R}q_* \mathbb{R}\text{End}_X(\mathbb{R}f_* \mathcal{O}_C)[1] & \xrightarrow{\text{tr}} & \mathbb{R}q_* \mathcal{O}_X[1] \\ \downarrow & & \uparrow T_x A_X & & \uparrow \text{id} \\ \mathbb{R}q_* \mathbb{R}f_* f^* T_X \simeq \mathbb{R}p_* f^* T_X & \longrightarrow & \mathbb{R}p_* \text{Cone}(T_C \rightarrow f^* T_X) & \xrightarrow{\Theta_f} & \mathbb{R}q_* \mathcal{O}_X[1]. \end{array}$$

In this diagram, the composite upper row is the image under $\mathbb{R}q_*$ of the first Chern class $c_1(\mathbb{R}f_* \mathcal{O}_C) \in \text{Ext}_X^1(T_X, \mathcal{O}_X) \simeq H^1(X, \Omega_X^1)$.

Pointed case. In the pointed case, if

$$\pi : \mathbb{R}\mathcal{C}_{g,n;X} \rightarrow \mathbb{R}\overline{\mathbf{M}}_{g,n}(X) \times X$$

is the derived stable universal family (Section 2.2), the same argument as in Proposition 3.3 shows that $\mathbb{R}\pi_*(\mathcal{O}_{\mathbb{R}\mathcal{C}_{g,n;X}})$ is a perfect complex on $\mathbb{R}\overline{\mathbf{M}}_{g,n}(X) \times X$. And we give the analogous

Definition 3.9. We denote by

$$A_X^{(n)} : \mathbb{R}\overline{\mathbf{M}}_{g,n}(X) \rightarrow \mathbb{R}\mathbf{Perf}(X)$$

the map induced by the perfect complex $\mathbb{R}\pi_*(\mathcal{O}_{\mathbb{R}\mathcal{C}_{g,n;X}})$.

We denote by $\delta_1^{(n)}(X)$ the composition

$$\mathbb{R}\overline{\mathbf{M}}_{g,n}(X) \xrightarrow{A_X^{(n)}} \mathbb{R}\mathbf{Perf}(X) \xrightarrow{\det_{\mathbf{Perf}}(X)} \mathbb{R}\mathbf{Pic}(X),$$

and, for a complex point x of $\mathbb{R}\overline{\mathbf{M}}_g(X)$ corresponding to a pointed stable map

$$f : (C; x_1, \dots, x_n) \rightarrow X,$$

by

$$\begin{aligned} \Theta_f^{(n)} := \mathbb{T}_f \delta_1^{(n)}(X) : \mathbb{T}_f \mathbb{R}\overline{\mathbf{M}}_{g,n}(X) &\xrightarrow{\mathbb{T}_x A_X^{(n)}} \mathbb{T}_{\mathbb{R}f_* \mathcal{O}_C} \mathbb{R}\mathbf{Perf}(X) \\ &\xrightarrow{\mathrm{tr}_X} \mathbb{T}_{\det(\mathbb{R}f_* \mathcal{O}_C)} \mathbb{R}\mathbf{Pic}(X). \end{aligned}$$

Again, if we fix a class $\beta \in H_2(X, \mathbb{Z})$, we have the corresponding β -decorated version of Definition 3.9.

4. The reduced derived stack of stable maps to a $K3$ -surface

In this section we specialize to the case of an algebraic $K3$ -surface S , with a fixed nonzero curve class $\beta \in H_2(S; \mathbb{Z})$, and a fixed symplectic form $\sigma \in H^0(S, K_S)$. After recalling in some detail the reduced obstruction theory of O-M-P-T, we first identify canonically the derived Picard stack $\mathbb{R}\mathbf{Pic}(S)$ with $\mathbf{Pic}(S) \times \mathbb{R}\mathrm{Spec}(\mathrm{Sym}(H^0(S, K_S)[1]))$ where K_S is the canonical sheaf of S . This result is then used to define the *reduced* version $\mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta)$ of the derived stack of stable maps of type (g, β) to S (and its n -pointed variant $\mathbb{R}\overline{\mathbf{M}}_{g,n}^{\mathrm{red}}(S; \beta)$), and to show that this induces, via the canonical procedure available for any algebraic derived stack, a modified obstruction theory on its truncation $\overline{\mathbf{M}}_g(S; \beta)$ whose deformation and obstruction spaces are then compared with those of the reduced theory of O-M-P-T. As a terminological remark, given an obstruction theory, we will call *deformation space* what is usually called its *tangent space* (while we keep the terminology *obstruction space*). We do this to avoid confusion with tangent spaces, tangent complexes or tangent cohomologies of possibly related (derived) stacks.

4.1. Reduced obstruction theory. For a $K3$ -surface S , the moduli of stable maps of genus g curves to S with non-zero effective class $\beta \in H^{1,1}(S, \mathbb{C}) \cap H^2(S, \mathbb{Z})$ (note that

Poincaré duality yields a canonical isomorphism

$$H_2(S; \mathbb{Z}) \simeq H^2(S; \mathbb{Z})$$

between singular (co)homologies) carries a relative perfect obstruction theory. This obstruction theory is given by

$$(R\pi_* F^* T_S)^\vee \rightarrow \mathbb{L}_{\overline{\mathbf{M}}_g(S; \beta) / \mathbf{M}_g^{\text{pre}}}.$$

Here $\pi: \mathcal{C}_{g, \beta; S} \rightarrow \overline{\mathbf{M}}_g(S; \beta)$ is the universal curve, $F: \mathcal{C}_{g, \beta; S} \rightarrow S$ is the universal morphism from the universal curve to S , and $\mathbf{M}_g^{\text{pre}}$ denotes the Artin stack of prestable curves. A Riemann–Roch argument along with the fact that a $K3$ -surface has trivial canonical bundle yields the expected dimension of $\overline{\mathbf{M}}_g(S; \beta)$:

$$\exp \dim \overline{\mathbf{M}}_g(S; \beta) = g - 1.$$

We thus expect no rational curves on a $K3$ -surface. This result stems from the deformation invariance of Gromov–Witten invariants. A $K3$ -surface admits deformations such that the homology class β is no longer of type $(1, 1)$, and thus cannot be the class of a curve.

This is unfortunate, given the rich literature on enumerative geometry of $K3$ -surfaces, and is in stark contrast to the well-known conjecture that a projective $K3$ -surface over an algebraically closed field contains infinitely many rational curves. Further evidence that there should be an interesting Gromov–Witten theory of $K3$ -surfaces are the results of Bloch, Ran and Voisin that rational curves deform in a family of $K3$ -surfaces provided their homology classes remain of type $(1, 1)$. The key ingredients in the proof is the semi-regularity map. We thus seek a new kind of obstruction theory for $\overline{\mathbf{M}}_g(S; \beta)$ which is deformation invariant only for such deformations of S which keep β of type $(1, 1)$.

Such a new obstruction theory, called the *reduced obstruction theory*, was introduced in [20, 21, 24]. Sticking to the case of moduli of morphisms from a *fixed* curve C to S , the obstruction space at a fixed morphism f is $H^1(C, f^* T_S)$.

This obstruction space admits a map

$$H^1(C, f^* T_S) \xrightarrow{\sim} H^1(C, f^* \Omega_S) \xrightarrow{H^1(df)} H^1(C, \Omega_C^1) \longrightarrow H^1(C, \omega_C) \simeq \mathbb{C},$$

where the first isomorphism is induced by the choice of a holomorphic symplectic form on S . The difficult part is to prove that all obstructions for *all* types of deformations of f lie in the kernel of this map, called the semi-regularity map for morphisms. Recall that using classical methods as in [6, 7] it has only been possible to show that the semi-regularity map annihilates obstructions to deformations over a base of the form $\mathbb{C}[x]/(x^n)$. For the construction of a reduced virtual fundamental class this is not nearly enough, as this requires annihilation of obstructions over square-zero extensions of arbitrary bases, which are not even assumed to be Artinian [3, Theorem 4.5]. Once this is proven, $\overline{\mathbf{M}}_g(S; \beta)$ carries a reduced obstruction theory which yields a virtual class, called the *reduced class*. This reduced class is one dimension larger than the one obtained from the standard perfect obstruction theory and leads to many interesting enumerative results (see [20, 21, 25]).

We will give below the construction of the reduced deformation and obstruction spaces giving all the details that will be needed in our comparison result (Theorem 4.8).

4.1.1. Deformation and obstruction spaces of the reduced theory according to Okounkov–Maulik–Pandharipande–Thomas. For further reference, we give here a self-contained treatment of the reduced *deformation* and reduced *obstruction* spaces on $\overline{\mathbf{M}}_g(S; \beta)$ according to Okounkov–Maulik–Pandharipande–Thomas.

Let us fix a stable map $f : C \rightarrow S$ of class $\beta \neq 0$ and genus g ; $p : C \rightarrow \text{Spec } \mathbb{C}$ and $q : S \rightarrow \text{Spec } \mathbb{C}$ will denote the structural morphisms. Let $\underline{\omega}_C \simeq p^! \mathcal{O}_{\text{Spec } \mathbb{C}}$ be the dualizing complex of C , and $\omega_C = \underline{\omega}_C[-1]$ the corresponding dualizing sheaf.

First of all, the *deformation spaces* of the standard (i.e. unreduced) and reduced theory, at the stable map f , coincide with $H^0(C, \text{Cone}(\mathbb{T}_C \rightarrow f^* T_S))$ where \mathbb{T}_C is the tangent complex of the curve C .

Let us recall now ([25, Section 3.1]) the construction of the reduced *obstruction space*. We give here version that is independent of the choice of a holomorphic symplectic form σ on S .

Consider the isomorphism²⁾

$$\varphi : T_S \otimes H^0(S, K_S) \xrightarrow{\sim} \Omega_S^1.$$

By tensoring this by $H^0(S, K_S)^\vee \simeq H^2(S, \mathcal{O}_S)$ (this isomorphism is given by Grothendieck–Serre duality that includes the Grothendieck trace map isomorphism $H^2(S, K_S) \rightarrow \mathbb{C}$, see e.g. [9, Section 3.4]) which is of dimension 1 over \mathbb{C} , we get a sequence of isomorphisms of \mathcal{O}_S -Modules

$$T_S \xleftarrow{\sim} T_S \otimes H^0(S, K_S) \otimes H^2(S, \mathcal{O}_S) \xrightarrow{\sim} \Omega_S^1 \otimes H^2(S, \mathcal{O}_S).$$

We denote by

$$\psi : T_S \rightarrow \Omega_S^1 \otimes H^2(S, \mathcal{O}_S)$$

the induced isomorphism. From this, we get an isomorphism of \mathcal{O}_C -Modules

$$f^* \psi : f^* T_S \xrightarrow{\sim} f^*(\Omega_S^1) \otimes H^2(S, \mathcal{O}_S).$$

Now consider the canonical maps

$$f^* \Omega_S^1 \xrightarrow{s} \Omega_C^1 \xrightarrow{t} \omega_C \simeq p^! \mathcal{O}_{\text{Spec } \mathbb{C}}[-1]$$

where $\omega_C \simeq \underline{\omega}_C[-1]$ is the dualizing sheaf of C and $\underline{\omega}_C = p^! \mathcal{O}_{\text{Spec } \mathbb{C}}$ the dualizing complex of C (see [10, Chapter V]). We thus obtain a map

$$\widetilde{v} : f^* T_S \rightarrow \underline{\omega}_C \otimes H^2(S, \mathcal{O}_S)[-1] \simeq \omega_C \otimes H^2(S, \mathcal{O}_S).$$

By the properties of dualizing complexes, we have

$$\underline{\omega}_C \otimes H^2(S, \mathcal{O}_S)[-1] = \underline{\omega}_C \otimes p^*(H^2(S, \mathcal{O}_S))[-1] \simeq p^!(H^2(S, \mathcal{O}_S)[-1]),$$

so we get a morphism

$$f^* T_S \rightarrow p^!(H^2(S, \mathcal{O}_S)[-1])$$

²⁾ The map φ is canonical, and the fact that it is an isomorphism depend on the existence (though not on the choice) of a symplectic form σ on S . Also note that we use throughout the standard abuse of writing $\mathcal{F} \otimes V$ for $\mathcal{F} \otimes_{\mathcal{O}_X} p^* V$, for any scheme $p : X \rightarrow \text{Spec } \mathbb{C}$, any \mathcal{O}_X -Module \mathcal{F} , and any \mathbb{C} -vector space V .

Note 8:
displayed as in-line
formula

which induces, by applying $\mathbb{R}p_*$ and composing with the adjunction map $\mathbb{R}p_*p^! \rightarrow \text{Id}$, a map

$$\begin{aligned} \widetilde{\alpha} : \mathbb{R}\Gamma(C, f^*T_S) &\simeq \mathbb{R}p_*(f^*T_S) \\ &\xrightarrow{\mathbb{R}p_*(\widetilde{v})} \mathbb{R}p_*(\omega_C \otimes H^2(S, \mathcal{O}_S)) \simeq \mathbb{R}p_*p^!(H^2(S, \mathcal{O}_S)[-1]) \\ &\longrightarrow H^2(S, \mathcal{O}_S)[-1]. \end{aligned}$$

Since $\mathbb{R}\Gamma$ is a triangulated functor, to get a unique induced map

$$\alpha : \mathbb{R}\Gamma(C, \text{Cone}(\mathbb{T}_C \rightarrow f^*T_S)) \rightarrow H^2(S, \mathcal{O}_S)[-1]$$

it will be enough to observe that

$$\text{Hom}_{D(\mathbb{C})}(\mathbb{R}p_*\mathbb{T}_C[1], H^2(S, \mathcal{O}_S)[-1]) = 0$$

(which is obvious since $\mathbb{R}p_*\mathbb{T}_C[1]$ lives in degrees $[-1, 0]$, while $H^2(S, \mathcal{O}_S)[-1]$ in degree 1), and to prove the following

Lemma 4.1. *The composition*

$$\mathbb{R}p_*\mathbb{T}_C \longrightarrow \mathbb{R}p_*f^*T_S \xrightarrow{\mathbb{R}p_*(\widetilde{v})} \mathbb{R}p_*(\omega_C \otimes H^2(S, \mathcal{O}_S))$$

vanishes in the derived category $D(\mathbb{C})$.

Proof. If C is smooth, the composition

$$\mathbb{T}_C \longrightarrow f^*T_S \xrightarrow{f^*\psi} f^*\Omega_S^1 \otimes H^2(S, \mathcal{O}_S) \xrightarrow{s \otimes \text{id}} \Omega_C^1 \otimes H^2(S, \mathcal{O}_S)$$

is obviously zero, since $\mathbb{T}_C \simeq T_C$ in this case, and a curve has no 2-forms. For a general prestable C , we proceed as follows. Let us consider the composition

$$\begin{aligned} \theta : \mathbb{T}_C &\longrightarrow f^*T_S \xrightarrow{f^*\psi} f^*\Omega_S^1 \otimes H^2(S, \mathcal{O}_S) \\ &\xrightarrow{s \otimes \text{id}} \Omega_C^1 \otimes H^2(S, \mathcal{O}_S) \xrightarrow{t \otimes \text{id}} \omega_C \otimes H^2(S, \mathcal{O}_S) := \mathcal{L}. \end{aligned}$$

On the smooth locus of C , $\mathcal{H}^0(\theta)$ is zero (by the same argument used in the case C smooth), hence the image of $\mathcal{H}^0(\theta) : \mathcal{H}^0(\mathbb{T}_C) \simeq T_C \rightarrow \mathcal{L}$ is a torsion subsheaf of the line bundle \mathcal{L} . But C is Cohen–Macaulay, therefore this image is 0, i.e. $\mathcal{H}^0(\theta) = 0$; and, obviously, **we have** $\mathcal{H}^i(\theta) = 0$ for any i (i.e. for $i = 1$). Now we use the hypercohomology spectral sequences

$$\begin{aligned} H^p(C, \mathcal{H}^q(\mathbb{T}_C)) &\Rightarrow \mathbb{H}^{p+q}(C, \mathbb{T}_C) \simeq H^{p+q}(\mathbb{R}\Gamma(C, \mathbb{T}_C)), \\ H^p(C, \mathcal{H}^q(\mathcal{L}[0])) &\Rightarrow \mathbb{H}^{p+q}(C, \mathcal{L}[0]) \simeq H^{p+q}(\mathbb{R}\Gamma(C, \mathcal{L}[0])) \simeq H^{p+q}(C, \mathcal{L}), \end{aligned}$$

to conclude that the induced maps

$$H^i(\mathbb{R}\Gamma(\theta)) : H^i(\mathbb{R}\Gamma(C, \mathbb{T}_C)) \rightarrow H^i(\mathbb{R}\Gamma(C, \mathcal{L})) \simeq H^i(C, \mathcal{L})$$

are zero for all i . Since \mathbb{C} is a field, we deduce that the map $\mathbb{R}\Gamma(\theta) = \mathbb{R}p_*(\theta)$ is zero in $D(\mathbb{C})$ as well. \square

By the lemma above, we have therefore obtained an induced map

$$\alpha : \mathbb{R}\Gamma(C, \text{Cone}(\mathbb{T}_C \rightarrow f^*T_S)) \rightarrow H^2(S, \mathcal{O}_S)[-1].$$

Now, O-M-P-T reduced obstruction space is defined as $\ker H^1(\alpha)$.

Moreover, again by Lemma 4.1, we have an induced map

$$v : \mathbb{R}p_*\text{Cone}(\mathbb{T}_C \rightarrow f^*T_S) \rightarrow \mathbb{R}p_*(\omega_C \otimes H^2(S, \mathcal{O}_S)),$$

and, since the map

$$\mathbb{R}p_*(\omega_C \otimes H^2(S, \mathcal{O}_S)) \rightarrow H^2(S, \mathcal{O}_S)[-1]$$

obviously induces an isomorphism on H^1 , we have that O-M-P-T reduced obstruction space is *also* the kernel of the map

$$H^1(v) : H^1(\mathbb{R}\Gamma(C, \text{Cone}(\mathbb{T}_C \rightarrow f^*T_S)) \rightarrow H^1(C, \omega_C \otimes H^2(S, \mathcal{O}_S)).$$

The following result proves the non-triviality of O-M-P-T reduced obstruction space.

Proposition 4.2. *If, as we are supposing, $\beta \neq 0$, the maps $H^1(v)$, $H^1(\alpha)$, $H^1(\tilde{\alpha})$, and $H^1(\mathbb{R}p_*(\tilde{v}))$ are all nontrivial, hence surjective.*

Proof. The non-vanishing of $H^1(\mathbb{R}p_*(\tilde{v}))$ obviously implies all other non-vanishing statements, and the non-vanishing of $H^1(\mathbb{R}p_*(\tilde{v}))$ is an immediate consequence of the following³⁾. □

Lemma 4.3. *Since the curve class $\beta \neq 0$, the map*

$$H^1(t \circ s) : H^1(C, f^*\Omega_S^1) \rightarrow H^1(C, \omega_C)$$

is nonzero (hence surjective).

Proof. By [4, Corollary 2.3], $\beta \neq 0$ implies non-triviality of $df : f^*\Omega_S^1 \rightarrow \Omega_C^1$. But S is a smooth surface and C a prestable curve, hence in the short exact sequence

$$f^*\Omega_S^1 \xrightarrow{s} \Omega_C^1 \rightarrow \Omega_{C/S}^1 \rightarrow 0$$

the sheaf of relative differentials $\Omega_{C/S}^1$ is concentrated at the (isolated, closed) singular points and thus its H^1 vanishes. Therefore the map

$$H^1(s) : H^1(C, f^*\Omega_S^1) \rightarrow H^1(C, \Omega_C^1)$$

is surjective. The same argument yields surjectivity, hence non-triviality (since $H^1(C, \omega_C)$ has dimension 1 over \mathbb{C}), of the map $H^1(t) : H^1(C, \Omega_C^1) \rightarrow H^1(C, \omega_C)$, by observing that, on the smooth locus of C , $\Omega_C^1 \simeq \omega_C$ and $H^1(t)$ is the induced isomorphism. In particular, $H^1(C, \Omega_C^1) \neq 0$. Therefore both $H^1(s)$ and $H^1(t)$ are non-zero and surjective, so the same is true of their composition. □

³⁾ We thank R. Pandharipande for pointing out this statement, of which we give here our proof.

Note 9:
removed 'the map'
for line break

4.2. The canonical projection $\mathbb{R}\mathbf{Pic}(S) \rightarrow \mathbb{R}\mathbf{Spec}(\mathrm{Sym}(H^0(S, K_S)[1]))$. In this subsection we identify canonically the derived Picard stack $\mathbb{R}\mathbf{Pic}(S)$ of a $K3$ -surface with $\mathbf{Pic}(S) \times \mathbb{R}\mathbf{Spec}(\mathrm{Sym}(H^0(S, K_S)[1]))$, where $K_S := \Omega_S^2$ is the canonical sheaf of S ; this allows us to define the canonical map

$$\mathrm{pr}_{\mathrm{der}} : \mathbb{R}\mathbf{Pic}(S) \rightarrow \mathbb{R}\mathbf{Spec}(\mathrm{Sym}(H^0(S, K_S)[1]))$$

which is the last ingredient we will need to define the *reduced* derived stack $\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta)$ of stable maps of genus g and class β to S in the next subsection.

In the proof of the next proposition, we will need the following elementary result (which holds true for k replaced by any semisimple ring, or k replaced by a hereditary commutative ring and E by a bounded above complex of free modules).

Lemma 4.4. *Let k be a field and E be a bounded above complex of k -vector spaces. Then there is a canonical map $E \rightarrow E_{<0}$ in the derived category $\mathbf{D}(k)$, such that the obvious composition*

$$E_{<0} \rightarrow E \rightarrow E_{<0}$$

is the identity.

Proof. Any splitting p of the map of k -vector spaces

$$\ker(d_0 : E_{-1} \rightarrow E_0) \hookrightarrow E_{-1}$$

yields a map $\bar{p} : E \rightarrow E_{<0}$ in the category $\mathrm{Ch}(k)$ of complexes of k -vector spaces. To see that different splittings p and q give the same map in the derived category $\mathbf{D}(k)$, we consider the canonical exact sequences of complexes

$$0 \rightarrow E_{<0} \rightarrow E \rightarrow E_{\geq 0} \rightarrow 0$$

and apply $\mathrm{Ext}^0(-, E_{<0})$, to get an exact sequence

$$\mathrm{Ext}^0(E_{\geq 0}, E_{<0}) \xrightarrow{a} \mathrm{Ext}^0(E, E_{<0}) \xrightarrow{b} \mathrm{Ext}^0(E_{<0}, E_{<0}).$$

Now, the class of the difference $(\bar{p} - \bar{q})$ in $\mathrm{Hom}_{\mathbf{D}(k)}(E, E_{<0}) = \mathrm{Ext}^0(E, E_{<0})$ is in the kernel of b , so it is enough to show that $\mathrm{Ext}^0(E_{\geq 0}, E_{<0}) = 0$. But $E_{\geq 0}$ is a bounded above complex of projectives, therefore (e.g. [39, Corollary 10.4.7]) $\mathrm{Ext}^0(E_{\geq 0}, E_{<0}) = 0$ is a quotient of $\mathrm{Hom}_{\mathrm{Ch}(k)}(E_{\geq 0}, E_{<0})$ which obviously consists of the zero morphism alone. \square

Proposition 4.5. *Let G be a derived group stack locally of finite presentation over a field k , $e : \mathrm{Spec} k \rightarrow G$ be its identity section, and $\mathfrak{g} := \mathbb{T}_e G$. Then there is a canonical map in $\mathrm{Ho}(\mathbf{dSt}_k)$*

$$\gamma(G) : \mathfrak{t}_0(G) \times \mathbb{R}\mathbf{Spec}(A) \rightarrow G$$

where $A := k \oplus (\mathfrak{g}^\vee)_{<0}$ is the commutative differential non-positively graded k -algebra which is the trivial square zero extension of k by the complex of k -vector spaces $(\mathfrak{g}^\vee)_{<0}$.

Proof. First observe that $\mathbb{R}\mathbf{Spec}(A)$ has a canonical k -point $x_0 : \mathrm{Spec} k \rightarrow \mathbb{R}\mathbf{Spec}(A)$, corresponding to the canonical projection $A \rightarrow k$. By definition of the derived cotangent com-

plex of a derived stack ([36, 1.4.1]), giving a map α such that the diagram

$$\begin{array}{ccc} \mathbb{R}\mathrm{Spec}(A) & \xrightarrow{\alpha} & G \\ & \swarrow x_0 \quad \searrow e & \\ & \mathrm{Spec} k & \end{array}$$

commutes in $\mathrm{Ho}(\mathbf{dSt}_{\mathbb{C}})$, is equivalent to giving a morphism in the derived category of complex of k -vector spaces

$$\alpha' : \mathbb{L}_{G,e} \simeq \mathfrak{g}^{\vee} \rightarrow (\mathfrak{g}^{\vee})_{<0}.$$

Since k is a field, we may take as α' the canonical map provided by Lemma 4.4, and define $\gamma(G)$ as the composition

$$t_0(G) \times \mathbb{R}\mathrm{Spec}(A) \xrightarrow{j \times \mathrm{id}} G \times \mathbb{R}\mathrm{Spec}(A) \xrightarrow{\mathrm{id} \times \alpha'} G \times G \xrightarrow{\mu} G$$

where μ is the product in G . □

Proposition 4.6. *Let S be a K3-surface over $k = \mathbb{C}$, and $G := \mathbb{R}\mathbf{Pic}(S)$ be its derived Picard group stack. Then the map $\gamma(G)$ defined in (the proof of) Proposition 4.5 is an isomorphism*

$$\gamma_S := \gamma(\mathbb{R}\mathbf{Pic}(S)) : \mathbf{Pic}(S) \times \mathbb{R}\mathrm{Spec}(\mathrm{Sym}(H^0(S, K_S)[1])) \xrightarrow{\sim} \mathbb{R}\mathbf{Pic}(S)$$

in $\mathrm{Ho}(\mathbf{dSt}_{\mathbb{C}})$, where K_S denotes the canonical bundle on S .

Proof. Since $G := \mathbb{R}\mathbf{Pic}(S)$ is a derived group stack, $\gamma(G)$ is an isomorphism if and only if it induces an isomorphism on truncations, and it is étale at e ([36, Theorem 2.2.2.6 and Lemma 2.2.1.1]), i.e. the induced map

$$\mathbb{T}_{(t_0(e), x_0)}(\gamma(G)) : \mathbb{T}_{(t_0(e), x_0)}(t_0(G) \times \mathbb{R}\mathrm{Spec}(A)) \rightarrow \mathbb{T}_e(G)$$

is an isomorphism in the derived category $D(k)$, where x_0 is the canonical k -point

$$\mathrm{Spec} \mathbb{C} \rightarrow \mathbb{R}\mathrm{Spec}(A),$$

corresponding to the canonical projection $A \rightarrow \mathbb{C}$. Since $\pi_0(A) \simeq \mathbb{C}$, $t_0(\gamma(G))$ is an isomorphism of stacks. So we are left to showing that $\gamma(G)$ induces an isomorphism between tangent spaces. Now,

$$\mathfrak{g} \equiv \mathbb{T}_e(G) = \mathbb{T}_e(\mathbb{R}\mathbf{Pic}(S)) \simeq \mathbb{R}\Gamma(S, \mathcal{O}_S)[1],$$

and, S being a K3-surface, we have

$$\mathfrak{g} \simeq \mathbb{R}\Gamma(S, \mathcal{O}_S)[1] \simeq H^0(S, \mathcal{O}_S)[1] \oplus H^2(S, \mathcal{O}_S)[-1]$$

so that

$$(\mathfrak{g}^{\vee})_{<0} \simeq H^2(S, \mathcal{O}_S)^{\vee}[1] \simeq H^0(S, K_S)[1]$$

(where we have used Serre duality in the last isomorphism). But $H^0(S, K_S)$ is free of dimension 1, so we have a canonical isomorphism

$$\mathbb{C} \oplus (\mathfrak{g}^{\vee})_{<0} \simeq \mathbb{C} \oplus H^0(S, K_S)[1] \simeq \mathrm{Sym}(H^0(S, K_S)[1])$$

in the homotopy category of commutative simplicial \mathbb{C} -algebras. Therefore

$$\mathbb{T}_{(t_0(e), x_0)}(\mathbf{Pic}(S) \times \mathbb{R}\mathrm{Spec}(A)) \simeq \mathfrak{g}_{\leq 0} \oplus \mathfrak{g}_{> 0} \simeq H^0(S, \mathcal{O}_S)[1] \oplus H^2(S, \mathcal{O}_S)[-1]$$

and $\mathbb{T}_{(t_0(e), x_0)}(\gamma(G))$ is obviously an isomorphism (given, in the notations of the proof of Proposition 4.5, by the sum of the dual of α' and the canonical map $\mathfrak{g}_{\leq 0} \rightarrow \mathfrak{g}$). \square

Using Proposition 4.6, we are now able to define the projection $\mathrm{pr}_{\mathrm{der}}$ of $\mathbb{R}\mathbf{Pic}(S)$ onto its full derived factor as the composite

$$\mathbb{R}\mathbf{Pic}(S) \xrightarrow{\gamma(S)^{-1}} \mathbf{Pic}(S) \times \mathbb{R}\mathrm{Spec}(\mathrm{Sym}(H^0(S, K_S)[1])) \xrightarrow{\mathrm{pr}_2} \mathbb{R}\mathrm{Spec}(\mathrm{Sym}(H^0(S, K_S)[1])).$$

Note that $\mathrm{pr}_{\mathrm{der}}$ yields on tangent spaces the canonical projection⁴⁾

$$\mathbb{T}_e(\mathbb{R}\mathbf{Pic}(S; \beta)) = \mathfrak{g} \rightarrow \mathfrak{g}_{> 0} = \mathbb{T}_{x_0}(\mathbb{R}\mathrm{Spec}(\mathrm{Sym}(H^0(S, K_S)[1]))) \simeq H^2(S, \mathcal{O}_S)[-1],$$

where x_0 is the canonical k -point $\mathrm{Spec} \mathbb{C} \rightarrow \mathrm{Spec}(\mathrm{Sym}(H^0(S, K_S)[1]))$, and

$$\mathfrak{g} \simeq H^0(S, \mathcal{O}_S)[1] \oplus H^2(S, \mathcal{O}_S)[-1].$$

4.3. The reduced derived stack of stable maps $\mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta)$. In this subsection we define the *reduced* version of the derived stack of stable maps of type (g, β) to S and describe the obstruction theory it induces on its truncation $\overline{\mathbf{M}}_g(S; \beta)$.

Let us define $\delta_1^{\mathrm{der}}(S, \beta)$, respectively, $\delta_1^{(n), \mathrm{der}}(S, \beta)$, as the composition (see Definition 3.7 and Definition 3.9)

$$\mathbb{R}\overline{\mathbf{M}}_g(S; \beta) \hookrightarrow \mathbb{R}\overline{\mathbf{M}}_g(S) \xrightarrow{\delta_1(S)} \mathbb{R}\mathbf{Pic}(S) \xrightarrow{\mathrm{pr}_{\mathrm{der}}} \mathbb{R}\mathrm{Spec}(\mathrm{Sym}(H^0(S, K_S)[1])),$$

resp., as the composition

$$\mathbb{R}\overline{\mathbf{M}}_{g,n}(S; \beta) \hookrightarrow \mathbb{R}\overline{\mathbf{M}}_{g,n}(S) \xrightarrow{\delta_1^{(n)}(S)} \mathbb{R}\mathbf{Pic}(S) \xrightarrow{\mathrm{pr}_{\mathrm{der}}} \mathbb{R}\mathrm{Spec}(\mathrm{Sym}(H^0(S, K_S)[1])).$$

Definition 4.7. The *reduced* derived stack of stable maps of genus g and class β to S , $\mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta)$, is defined by the following homotopy-cartesian square in $\mathbf{dSt}_{\mathbb{C}}$:

$$\begin{array}{ccc} \mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta) & \longrightarrow & \mathbb{R}\overline{\mathbf{M}}_g(S; \beta) \\ \downarrow & & \downarrow \delta_1^{\mathrm{der}}(S, \beta) \\ \mathrm{Spec} \mathbb{C} & \longrightarrow & \mathbb{R}\mathrm{Spec}(\mathrm{Sym}(H^0(S, K_S)[1])). \end{array}$$

The *reduced* derived stack of n -pointed stable maps of genus g and class β to S , $\mathbb{R}\overline{\mathbf{M}}_{g,n}^{\mathrm{red}}(S; \beta)$, is defined by the following homotopy-cartesian square in $\mathbf{dSt}_{\mathbb{C}}$:

$$\begin{array}{ccc} \mathbb{R}\overline{\mathbf{M}}_{g,n}^{\mathrm{red}}(S; \beta) & \longrightarrow & \mathbb{R}\overline{\mathbf{M}}_{g,n}(S; \beta) \\ \downarrow & & \downarrow \delta_1^{(n), \mathrm{der}}(S, \beta) \\ \mathrm{Spec} \mathbb{C} & \longrightarrow & \mathbb{R}\mathrm{Spec}(\mathrm{Sym}(H^0(S, K_S)[1])). \end{array}$$

⁴⁾ Recall that, if M is a \mathbb{C} -vector space, $\mathbb{T}_{x_0}(\mathbb{R}\mathrm{Spec}(\mathrm{Sym}(M[1]))) \simeq M^{\vee}[-1]$.

Since the truncation functor t_0 commutes with homotopy fiber products and

$$t_0(\mathbb{R}\mathrm{Spec}(\mathrm{Sym}(H^0(S, K_S)[1]))) \simeq \mathrm{Spec} \mathbb{C},$$

we get

$$t_0(\mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta)) \simeq \overline{\mathbf{M}}_g(S; \beta),$$

i.e. $\mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta)$ is a *derived extension* (Definition 1.1) of the usual stack of stable maps of type (g, β) to S , different from $\mathbb{R}\overline{\mathbf{M}}_g(S; \beta)$. Similarly in the pointed case.

We are now able to compute the obstruction theory induced, according to Section 1, by the closed immersion $j_{\mathrm{red}} : \overline{\mathbf{M}}_g(S; \beta) \hookrightarrow \mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta)$. We leave to the reader the straightforward modifications for the pointed case.

By applying Proposition 1.2 to the derived extension $\mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta)$ of $\overline{\mathbf{M}}_g(S; \beta)$, we get an obstruction theory

$$j_{\mathrm{red}}^* \mathbb{L}_{\mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta)} \rightarrow \mathbb{L}_{\overline{\mathbf{M}}_g(S; \beta)}$$

that we are now going to describe.

Let

$$\rho : \mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta) \rightarrow \mathbb{R}\overline{\mathbf{M}}_g(S; \beta)$$

be the canonical map. Since $\mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta)$ is defined by the homotopy pullback diagram in Definition 4.7 [above](#), we get an isomorphism in the derived category of $\mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta)$

$$\rho^* (\mathbb{L}_{\mathbb{R}\overline{\mathbf{M}}_g(S; \beta) / \mathbb{R}\mathrm{Spec}(\mathrm{Sym}(H^0(S, K_S)[1]))}) \simeq \mathbb{L}_{\mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta)}.$$

We will show below that $\mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta)$ is *quasi-smooth* so that, by Corollary 1.4,

$$j_{\mathrm{red}}^* \mathbb{L}_{\mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta)} \rightarrow \mathbb{L}_{\overline{\mathbf{M}}_g(S; \beta)}$$

is indeed a *perfect* obstruction theory on $\overline{\mathbf{M}}_g(S; \beta)$. Now, for any \mathbb{C} -point

$$\mathrm{Spec} \mathbb{C} \rightarrow \mathbb{R}\overline{\mathbf{M}}_g(S; \beta),$$

corresponding to a stable map $(f : C \rightarrow S)$ of type (g, β) , we get a distinguished triangle

$$\mathbb{L}_{\mathbb{R}\mathrm{Spec}(\mathrm{Sym}(H^0(S, K_S)[1])), x_0} \rightarrow \mathbb{L}_{\mathbb{R}\overline{\mathbf{M}}_g(S; \beta), (f : C \rightarrow S)} \rightarrow \mathbb{L}_{\mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta), (f : C \rightarrow S)}$$

(where we have denoted by $(f : C \rightarrow S)$ also the induced \mathbb{C} -point of $\mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta)$, and used that a derived stack and its truncation have the same classical points, i.e. points with values in usual commutative \mathbb{C} -algebras) in the derived category of complexes of \mathbb{C} -vector spaces. By dualizing, we get that the tangent complex

$$\mathbb{T}_{(f : C \rightarrow S)}^{\mathrm{red}} := \mathbb{T}_{(f : C \rightarrow S)}(\mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta))$$

of $\mathbb{R}\overline{\mathbf{M}}_g^{\mathrm{red}}(S; \beta)$ at the \mathbb{C} -point $(f : C \rightarrow S)$ of type (g, β) , sits into a distinguished triangle

$$\mathbb{T}_{(f : C \rightarrow S)}^{\mathrm{red}} \longrightarrow \mathbb{R}\Gamma(C, \mathrm{Cone}(\mathbb{T}_C \rightarrow f^* T_S)) \xrightarrow{\Theta_f} \mathbb{R}\Gamma(S, \mathcal{O}_S)[1] \xrightarrow{\mathrm{pr}} H^2(S, \mathcal{O}_S)[-1],$$

where Θ_f is the composite

$$\Theta_f : \mathbb{R}\Gamma(C, \mathrm{Cone}(\mathbb{T}_C \rightarrow f^* T_S)) \xrightarrow{\mathbb{T}_x \mathrm{Ax}} \mathbb{R}\mathrm{Hom}_S(\mathbb{R}f_* \mathcal{O}_C, \mathbb{R}f_* \mathcal{O}_C)[1] \xrightarrow{\mathrm{tr}_S} \mathbb{R}\Gamma(S, \mathcal{O}_S)[1],$$

and pr denotes the tangent map of $\mathrm{pr}_{\mathrm{der}}$ taken at the point $\delta_1(S)(f : C \rightarrow S)$. Note that the map pr obviously induces an isomorphism on H^1 .

4.4. Quasi-smoothness of $\mathbb{R}\overline{\mathbf{M}}_g^{\text{red}}(S; \beta)$ and comparison with Okounkov–Maulik–Pandharipande–Thomas reduced obstruction theory. In the case $\beta \neq 0$ is a curve class in $H^2(S, \mathbb{Z})$, we will prove *quasi-smoothness* of the derived stack $\mathbb{R}\overline{\mathbf{M}}_g^{\text{red}}(S; \beta)$, and compare the induced obstruction theory with that of Okounkov–Maulik–Pandharipande–Thomas (see Section 4.1.1 or [20, Section 2.2] and [25]).

Theorem 4.8. *Let $\beta \neq 0$ be a curve class in $H^2(S, \mathbb{Z}) \simeq H_2(S, \mathbb{Z})$, $f : C \rightarrow S$ be a stable map of type (g, β) , and*

$$\mathbb{T}_{(f:C \rightarrow S)}^{\text{red}} := \mathbb{T}_{(f:C \rightarrow S)}(\mathbb{R}\overline{\mathbf{M}}_g^{\text{red}}(S; \beta)) \rightarrow \mathbb{R}\Gamma(C, \text{Cone}(\mathbb{T}_C \rightarrow f^*T_S)) \rightarrow H^2(S, \mathcal{O}_S)[-1]$$

be the corresponding distinguished triangle. Then:

- (1) *the rightmost arrow in the triangle above induces on H^1 a map*

$$H^1(\Theta_f) : H^1(C, \text{Cone}(\mathbb{T}_C \rightarrow f^*T_S)) \rightarrow H^2(S, \mathcal{O}_S)$$

which is nonzero (hence surjective, since $H^2(S, \mathcal{O}_S)$ has dimension 1 over \mathbb{C}). Therefore the derived stack $\mathbb{R}\overline{\mathbf{M}}_g^{\text{red}}(S; \beta)$ is everywhere quasi-smooth,

- (2) *$H^0(\mathbb{T}_{(f:C \rightarrow S)}^{\text{red}})$ (resp., $H^1(\mathbb{T}_{(f:C \rightarrow S)}^{\text{red}})$) coincides with the reduced deformation space (resp., the reduced obstruction space) of O-M-P-T.*

Proof. Proof of quasi-smoothness. Let us prove quasi-smoothness first. It is clearly enough to prove that the composite

$$\begin{aligned} H^1(C, f^*T_S) &\longrightarrow H^1(C, \text{Cone}(\mathbb{T}_C \rightarrow f^*T_S)) \\ &\xrightarrow{H^1(\mathbb{T}_X A_X)} \text{Ext}_S^2(\mathbb{R}f_*\mathcal{O}_C, \mathbb{R}f_*\mathcal{O}_C) \\ &\xrightarrow{H^2(\text{tr}_S)} H^2(S, \mathcal{O}_S) \end{aligned}$$

is non-zero (hence surjective). Recall that $p : C \rightarrow \text{Spec } \mathbb{C}$ and $q : S \rightarrow \text{Spec } \mathbb{C}$ denote the structural morphisms, so that $p = q \circ f$. Now, the map

$$\mathbb{R}q_*T_S \rightarrow \mathbb{R}q_*\mathbb{R}f_*f^*T_S$$

induces a map $H^1(S, T_S) \rightarrow H^1(C, f^*T_S)$, and by Proposition 3.6 and Remark 3.8, the following diagram commutes:

$$\begin{array}{ccc} H^1(S, T_S) & \xrightarrow{\langle -, \text{at}_{\mathbb{R}f_*\mathcal{O}_C} \rangle} & \text{Ext}_S^2(\mathbb{R}f_*\mathcal{O}_C, \mathbb{R}f_*\mathcal{O}_C) \\ \downarrow & & \uparrow H^1(\mathbb{T}_X A_X) \\ H^1(C, f^*T_S) & \longrightarrow & H^1(C, \text{Cone}(\mathbb{T}_C \rightarrow f^*T_S)). \end{array}$$

So, we are reduced to proving that the composition

$$a : H^1(S, T_S) \xrightarrow{\langle -, \text{at}_{\mathbb{R}f_*\mathcal{O}_C} \rangle} \text{Ext}_S^2(\mathbb{R}f_*\mathcal{O}_C, \mathbb{R}f_*\mathcal{O}_C) \xrightarrow{H^2(\text{tr})} H^2(S, \mathcal{O}_S)$$

does not vanish. But, since the first Chern class is the trace of the Atiyah class (as in [12, (5.4.1) and Section 5.9]), this composition acts as follows (on maps in the derived category of S):

$$(\xi : \mathcal{O}_S \rightarrow T_S[1]) \longrightarrow (a(\xi) : \mathcal{O}_S \xrightarrow{c_1 \otimes \xi} \Omega_S^1 \otimes T_S[2] \xrightarrow{\langle -, - \rangle} \mathcal{O}_S[2])$$

where

$$c_1 := c_1(\mathbb{R}f_*\mathcal{O}_C) : \mathcal{O}_S \rightarrow \Omega_S^1[1]$$

is the first Chern class of the perfect complex $\mathbb{R}f_*\mathcal{O}_C$. What we have said so far, is true for any smooth complex projective scheme X in place of S . We now use the fact that S is a $K3$ -surface. Choose a non-zero section $\sigma : \mathcal{O}_S \rightarrow \Omega_S^2$ of the canonical bundle, and denote by

$$\varphi_\sigma : \Omega_S^1 \xrightarrow{\sim} T_S$$

the induced isomorphism. A straightforward linear algebra computation shows then that the composition

$$\mathcal{O}_S \xrightarrow{((\varphi_\sigma \circ c_1) \wedge \xi) \otimes \sigma} (T_S \wedge T_S \otimes \Omega_S^2)[2] \xrightarrow{\langle -, - \rangle[2]} \mathcal{O}_S[2]$$

coincides with $a(\xi)$. But, since $\beta \neq 0$, we have that $c_1 \neq 0$. The section σ is non-degenerate, so this composition cannot vanish for all ξ , and we conclude.

Alternatively, we could proceed as follows. By Serre duality, passing to dual vector spaces and maps, we are left to proving that the composite

$$H^0(S, \Omega_S^2) \xrightarrow{\text{tr}^\vee} \text{Ext}_S^0(\mathbb{R}\underline{\text{Hom}}(\mathbb{R}f_*\mathcal{O}_C, \mathbb{R}f_*\mathcal{O}_C), \Omega_S^2) \xrightarrow{\tau^\vee} \text{Ext}^0(\mathbb{R}f_*f^*T_S[-1], \Omega_S^2)$$

is non-zero. So it is enough to prove that the map obtained by further composing to the left with the adjunction map

$$\text{Ext}^0(\mathbb{R}f_*f^*T_S[-1], \Omega_S^2) \rightarrow \text{Ext}^0(T_S[-1], \Omega_S^2)$$

is nonzero. But this new composition acts as follows:

$$\begin{aligned} H^0(S, \Omega_S^2) \ni (\sigma : \mathcal{O}_S \rightarrow \Omega_S^2) &\mapsto (\sigma \circ \text{tr}) \\ &\mapsto (\sigma \circ \text{tr} \circ \text{at}) = (\sigma \circ c_1(\mathbb{R}f_*\mathcal{O}_C)) \in \text{Ext}^0(T_S[-1], \Omega_S^2) \end{aligned}$$

where $\text{at} : T_S[-1] \rightarrow \mathbb{R}\underline{\text{Hom}}(\mathbb{R}f_*\mathcal{O}_C, \mathbb{R}f_*\mathcal{O}_C)$ is the Atiyah class of $\mathbb{R}f_*\mathcal{O}_C$ (see Proposition 3.6 and Remark 3.8). Since $\beta \neq 0$, we have $c_1(\mathbb{R}f_*\mathcal{O}_C) \neq 0$, and we conclude.

Proof of the comparison. Let us move now to the second point of Theorem 4.8, i.e. the comparison statement about deformations and obstructions spaces. First of all it is clear that, for any β ,

$$H^0(\mathbb{T}_{\text{red}, (f:C \rightarrow S)}) \simeq H^0(\mathbb{T}_{\mathbb{R}\overline{\mathbf{M}}_g(S;\beta), (f:C \rightarrow S)}) \simeq H^0(C, \text{Cone}(\mathbb{T}_C \rightarrow f^*T_S))$$

therefore our deformation space is the same as O-M-P-T's one. Let us then concentrate on obstruction spaces.

We begin by noticing the following fact.

Lemma 4.9. *There is a canonical morphism in $D(\mathbb{C})$*

$$v : \mathbb{R}p_*\omega_C \otimes^{\mathbb{L}} H^2(S, \mathcal{O}_S) \rightarrow \mathbb{R}q_*\mathcal{O}_S[1]$$

inducing an isomorphism on H^1 .

Proof. To ease notation we will simply write \otimes for $\otimes^{\mathbb{L}}$. Recall that $p : C \rightarrow \text{Spec } \mathbb{C}$ and $q : S \rightarrow \text{Spec } \mathbb{C}$ denote the structural morphisms, so that $p = q \circ f$. Since S is a $K3$ -surface, the canonical map

$$\mathcal{O}_S \otimes H^0(S, \Omega_S^2) \rightarrow \Omega_S^2$$

is an isomorphism. Since $f^!$ preserves dualizing complexes, $\omega_S \simeq \Omega_S^2[2]$ and $\omega_C \simeq \omega_C[1]$, we have

$$\omega_C \simeq f^!\Omega_S^2[1] \simeq f^!(\mathcal{O}_S \otimes H^0(S, \Omega_S^2))[1].$$

By applying $\mathbb{R}p_*$ and using the adjunction map $\mathbb{R}f_*f^! \rightarrow \text{Id}$, we get a map

$$\begin{aligned} \mathbb{R}p_*\omega_C &\simeq \mathbb{R}q_*\mathbb{R}f_*\omega_C \simeq \mathbb{R}q_*\mathbb{R}f_*f^!(\mathcal{O}_S[1] \otimes H^0(S, \Omega_S^2)) \\ &\rightarrow \mathbb{R}q_*(\mathcal{O}_S[1] \otimes H^0(S, \Omega_S^2)) \simeq \mathbb{R}q_*\mathcal{O}_S[1] \otimes H^0(S, \Omega_S^2) \end{aligned}$$

(the last isomorphism being given by projection formula). Tensoring this map by **the isomorphism** $H^0(S, \Omega_S^2)^\vee \simeq H^2(S, \mathcal{O}_S)$ (a canonical isomorphism by Serre duality), and using the canonical evaluation map $V \otimes V^\vee \rightarrow \mathbb{C}$ for a \mathbb{C} -vector space V , we get the desired canonical map

$$v : \mathbb{R}p_*\omega_C \otimes H^2(S, \mathcal{O}_S) \rightarrow \mathbb{R}q_*\mathcal{O}_S[1].$$

The isomorphism on H^1 is obvious since the trace map $\mathbb{R}^1p_*\omega_C \rightarrow \mathbb{C}$ is an isomorphism (C is geometrically connected). \square

If $\sigma : \mathcal{O}_S \xrightarrow{\sim} \Omega_S^2$ is a nonzero element in $H^0(S, \Omega_S^2)$, and $\varphi_\sigma : T_S \simeq \Omega_S^1$ the induced isomorphism, the previous lemma gives us an induced map

$$v(\sigma) : \mathbb{R}p_*\omega_C \rightarrow \mathbb{R}q_*\mathcal{O}_S[1],$$

and an induced isomorphism

$$H^1(v(\sigma)) =: v_\sigma : H^1(C, \omega_C) \xrightarrow{\sim} H^2(S, \mathcal{O}_S).$$

Using the same notations as in Section 4.1.1, to prove that our reduced obstruction space

$$\ker(H^1(\Theta_f) : H^1(C, \text{Cone}(\mathbb{T}_C \rightarrow f^*T_S)) \rightarrow H^2(S, \mathcal{O}_S))$$

coincides with O-M-P-T's one, it will be enough to show that the following diagram is commutative:

$$\begin{array}{ccc} H^1(C, f^*T_S) & \xrightarrow{\text{can}} & H^1(C, \text{Cone}(\mathbb{T}_C \rightarrow f^*T_S)) \\ \text{can} \downarrow & & \downarrow H^1(\Theta_f) \\ H^1(C, \text{Cone}(\mathbb{T}_C \rightarrow f^*T_S)) & & \\ H^1(v) \downarrow & & \\ H^1(C, \omega_C) & \xrightarrow[\sim]{v_\sigma} & H^2(S, \mathcal{O}_S). \end{array}$$

But this follows from the commutativity of

$$\begin{array}{ccccccc}
 \mathbb{R}p_*f^*T_S[-1] & \xrightarrow{\mathbb{R}p_*(\varphi_\sigma)} & \mathbb{R}p_*f^*\Omega_S^1[-1] & \xrightarrow{\mathbb{R}p_*(s)} & \mathbb{R}p_*\Omega_C^1[-1] & \xrightarrow{\mathbb{R}p_*(t)} & \mathbb{R}p_*\omega_C[-1] \\
 \downarrow \text{id} & & & & & & \downarrow \nu(\sigma)[-1] \\
 \mathbb{R}p_*f^*T_S[-1] & \xrightarrow{\mathbb{T}_{xAX}} & \mathbb{R}q_*\mathbb{R}\underline{\text{Hom}}_S(\mathbb{R}f_*\mathcal{O}_C, \mathbb{R}f_*\mathcal{O}_C) & \xrightarrow{\text{tr}} & \mathbb{R}q_*\mathcal{O}_S & &
 \end{array}$$

that follows directly from the definitions of the maps involved. \square

Remark 4.10. Note that by Lemma 4.2, the second assertion of Theorem 4.8 implies the first one. Nonetheless, we have preferred to give an independent proof of the quasi-smoothness of $\mathbb{R}\overline{\mathbf{M}}_g^{\text{red}}(S; \beta)$ because we find it conceptually more relevant than the comparison with O-M-P-T, meaning that quasi-smoothness alone would in any case imply the existence of *some* perfect reduced obstruction theory on $\overline{\mathbf{M}}_g(S; \beta)$, regardless of its comparison with the one introduced and studied by O-M-P-T. A complete comparison with O-M-P-T would require not only Theorem 4.8 (2), but also a proof that all obstruction maps are the same. We think this is true, but we leave the task of verifying the details to the interested reader.

Moreover, we could only find in the literature a definition of O-M-P-T *global* reduced obstruction theory (relative to $\mathbf{M}_g^{\text{pre}}$) with values in the $\tau_{\geq -1}$ -truncation of the cotangent complex of the stack of stable maps⁵⁾, that uses a result on the semiregularity map whose proof is not completely convincing ([20, Section 2.2, formula (14)]); on the other hand there is a clean and complete description of the corresponding pointwise tangent and obstruction spaces. Therefore, our comparison is necessarily limited to these spaces. Our construction might also be seen as establishing such a reduced *global* obstruction theory – in the usual sense, i.e. with values in the full cotangent complex, and completely independent from any result on semiregularity maps.

Theorem 4.8 shows that the distinguished triangle

$$\mathbb{T}_{(f:C \rightarrow S)}^{\text{red}} := \mathbb{T}_{(f:C \rightarrow S)}(\mathbb{R}\overline{\mathbf{M}}_g^{\text{red}}(S; \beta)) \rightarrow \mathbb{R}\Gamma(C, \text{Cone}(\mathbb{T}_C \rightarrow f^*T_S)) \rightarrow H^2(S, \mathcal{O}_S)[-1]$$

induces isomorphisms

$$H^i(\mathbb{T}_{(f:C \rightarrow S)}^{\text{red}}) \simeq H^i(C, \text{Cone}(\mathbb{T}_C \rightarrow f^*T_S)),$$

for any $i \neq 1$, while in degree 1, it yields a short exact sequence

$$0 \rightarrow H^1(\mathbb{T}_{(f:C \rightarrow S)}^{\text{red}}) \rightarrow H^1(C, \text{Cone}(\mathbb{T}_C \rightarrow f^*T_S)) \rightarrow H^2(S, \mathcal{O}_S) \rightarrow 0.$$

So, the tangent complexes of $\mathbb{R}\overline{\mathbf{M}}_g^{\text{red}}(S; \beta)$ and $\mathbb{R}\overline{\mathbf{M}}_g(S; \beta)$ (hence our induced *reduced* and the *standard* obstruction theories) *only differ at the level of* H^1 where the former is the kernel of a 1-dimensional quotient of the latter: this is indeed the distinguished feature of a (codimension 1) reduced obstruction theory.

⁵⁾ The reason being that the authors use factorization through the cone, and therefore the resulting obstruction theory is only well-defined, without further arguments, if one considers it as having values in such a truncation.

The pointed case. In the pointed case, a completely analogous proof as that of Theorem 4.8 (1), yields

Theorem 4.11. *Let $\beta \neq 0$ be a curve class in $H^2(S, \mathbb{Z}) \simeq H_2(S, \mathbb{Z})$. The derived stack $\mathbb{R}\overline{\mathbf{M}}_{g,n}^{\text{red}}(S; \beta)$ of n -pointed stable maps of type (g, β) is everywhere quasi-smooth, and therefore the canonical map*

$$j^*(\mathbb{L}_{\mathbb{R}\overline{\mathbf{M}}_{g,n}(X; \beta)}) \rightarrow \mathbb{L}_{\overline{\mathbf{M}}_{g,n}(X; \beta)}$$

is a $[-1, 0]$ perfect obstruction theory on $\overline{\mathbf{M}}_{g,n}(X; \beta)$.

5. Moduli of perfect complexes

In this section we will define and study derived versions of various stacks of perfect complexes on a smooth projective variety X . If X is a K3-surface, by using the determinant map and the structure of $\mathbb{R}\mathbf{Pic}(X)$, we deduce that the derived stack of simple perfect complexes on X is smooth. This result was proved with different methods by Inaba in [13].

When X is a Calabi–Yau 3-fold, we prove that the derived stack of simple perfect complexes (with fixed determinant) is quasi-smooth, and then use an elaboration of the map

$$A_X^{(n)} : \mathbb{R}\overline{\mathbf{M}}_{g,n}(X) \rightarrow \mathbb{R}\mathbf{Perf}(X)$$

to compare the obstruction theories induced on the truncation stacks. This might be seen as a derived geometry approach to a naive, open version of the Gromov–Witten/Donaldson–Thomas comparison.

Definition 5.1. Let X be a smooth complex projective variety, \mathcal{L} be a line bundle on X , and $x_{\mathcal{L}} : \text{Spec } \mathbb{C} \rightarrow \mathbb{R}\mathbf{Pic}(X)$ be the corresponding point.

- The derived stack $\mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}$ of perfect complexes on X with fixed determinant \mathcal{L} is defined by the following homotopy cartesian diagram in $\mathbf{dSt}_{\mathbb{C}}$:

$$\begin{array}{ccc} \mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}} & \longrightarrow & \mathbb{R}\mathbf{Perf}(X) \\ \downarrow & & \downarrow \det \\ \text{Spec } \mathbb{C} & \xrightarrow{x_{\mathcal{L}}} & \mathbb{R}\mathbf{Pic}(X). \end{array}$$

We will write $\mathbb{R}\mathbf{Perf}(X)_0$ for $\mathbb{R}\mathbf{Perf}(X)_{\mathcal{O}_X}$, the derived stack of perfect complexes on X with trivial determinant.

- If $\mathbf{Perf}(X)^{\geq 0}$ denotes the open substack of $\mathbf{Perf}(X)$ consisting of perfect complexes F on X such that $\text{Ext}^i(F, F) = 0$ for $i < 0$, we define $\mathbb{R}\mathbf{Perf}(X)^{\geq 0} := \phi_{\mathbb{R}\mathbf{Perf}(X)}(\mathbf{Perf}(X)^{\geq 0})$ (as a derived open substack of $\mathbb{R}\mathbf{Perf}(X)$, see Proposition 2.1).
- If $\mathbf{Perf}(X)^{\text{si}, > 0}$ is the open substack of $\mathbf{Perf}(X)$ consisting of perfect complexes F on X for which $\text{Ext}^i(F, F) = 0$ for $i < 0$, and the trace map $\text{Ext}^0(F, F) \rightarrow H^0(X, \mathcal{O}_X) \simeq \mathbb{C}$ is an isomorphism, we define $\mathbb{R}\mathbf{Perf}(X)^{\text{si}, > 0} := \phi_{\mathbb{R}\mathbf{Perf}(X)}(\mathbf{Perf}(X)^{\text{si}, > 0})$ (as a derived open substack of $\mathbb{R}\mathbf{Perf}(X)$, see Proposition 2.1).

- The derived stack $\mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}^{\geq 0}$ is defined by the following homotopy cartesian diagram in $\mathbf{dSt}_{\mathbb{C}}$:

$$\begin{array}{ccc} \mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}^{\geq 0} & \longrightarrow & \mathbb{R}\mathbf{Perf}(X)^{\geq 0} \\ \downarrow & & \downarrow \det \\ \mathrm{Spec} \, \mathbb{C} & \xrightarrow{x_{\mathcal{L}}} & \mathbb{R}\mathbf{Pic}(X). \end{array}$$

As above, we will write $\mathbb{R}\mathbf{Perf}(X)_0^{\geq 0}$ for $\mathbb{R}\mathbf{Perf}(X)_{\mathcal{O}_X}^{\geq 0}$.

- The derived stack $\mathcal{M}_X \equiv \mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}^{\mathrm{si}, > 0}$ is defined by the following homotopy cartesian diagram in $\mathbf{dSt}_{\mathbb{C}}$:

$$\begin{array}{ccc} \mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}^{\mathrm{si}, > 0} & \longrightarrow & \mathbb{R}\mathbf{Perf}(X)^{\mathrm{si}, > 0} \\ \downarrow & & \downarrow \det \\ \mathrm{Spec} \, \mathbb{C} & \xrightarrow{x_{\mathcal{L}}} & \mathbb{R}\mathbf{Pic}(X). \end{array}$$

We will write $\mathbb{R}\mathbf{Perf}(X)_0^{\mathrm{si}, > 0}$ for $\mathbb{R}\mathbf{Perf}(X)_{\mathcal{O}_X}^{\mathrm{si}, > 0}$.

Proposition 5.2. *Let E be a perfect complex on X with determinant \mathcal{L} , and*

$$x_E : \mathrm{Spec} \, \mathbb{C} \rightarrow \mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}$$

be the corresponding point. The tangent complex of $\mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}$ at x_E is

$$\mathrm{Cone}(\mathrm{tr} : \mathbb{R}\mathrm{End}(E) \rightarrow \mathbb{R}\Gamma(X, \mathcal{O}_X)).$$

Proof. Let \mathbb{T} denote the tangent complex of $\mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}$ at the point x_E . By definition of $\mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}$, we have an exact triangle in the derived category $D(\mathbb{C})$ of \mathbb{C} -vector spaces

$$\mathbb{T} \rightarrow \mathbb{R}\mathrm{End}(E)[1] \xrightarrow{\mathrm{tr}} \mathbb{R}\Gamma(X, \mathcal{O}_X)[1]. \quad \square$$

Note that if $\chi(E) \neq 0$, we have that

$$\mathbb{T} \simeq \mathbb{R}\mathrm{End}(E)_0[1],$$

the shifted traceless derived endomorphisms complex of E ([11, Definition 10.1.4]), so that

$$H^i(\mathbb{R}\mathrm{End}(E)_0) = \ker(\mathrm{tr} : \mathrm{Ext}^i(E, E) \rightarrow H^i(X, \mathcal{O}_X)),$$

for any i . In fact the exact triangle in the proof above is split by $\chi(E)^{-1}\mathrm{id}$.

Remark 5.3. Since $\mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}^{\geq 0}$ and $\mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}^{\mathrm{si}, > 0}$ are derived open substacks of $\mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}$, Proposition 5.2 holds for their tangent complexes too.

5.1. On $K3$ -surfaces. By using the derived determinant map and the derived stack of perfect complexes, we are able to give another proof of a result by Inaba ([13, Theorem 3.2]) that generalizes an earlier work by Mukai ([22]). For simplicity, we prove this result for $K3$ -surfaces, the result for a general Calabi–Yau surface being similar.

Let S be a smooth projective $K3$ -surface, and let $\mathbb{R}\mathbf{Perf}(S)^{\text{si},>0}$ (Definition 5.1) be the open derived substack of $\mathbb{R}\mathbf{Perf}(S)$ consisting of perfect complexes F on S for which $\text{Ext}_S^i(F, F) = 0$ for $i < 0$, and the trace map

$$\text{Ext}_S^0(F, F) \rightarrow H^0(S, \mathcal{O}_S) \simeq \mathbb{C}$$

is an isomorphism. The truncation $\mathbf{Perf}(S)^{\text{si},>0}$ of $\mathbb{R}\mathbf{Perf}(S)^{\text{si},>0}$ is a stack whose coarse moduli space $\text{Perf}(S)^{\text{si},>0}$ is exactly the moduli space Inaba calls $\text{Splcp}_S^{\text{ét}}/\mathbb{C}$ in [13, Section 3].

As in Section 4.2, we consider the projection pr_{der} of $\mathbb{R}\mathbf{Pic}(S)$ onto its full derived factor

$$\mathbb{R}\mathbf{Pic}(S) \xrightarrow{\text{pr}_{\text{der}}} \mathbb{R}\text{Spec}(\text{Sym}(H^0(S, K_S)[1])).$$

Definition 5.4. The *reduced* derived stack $\mathbb{R}\mathbf{Perf}(S)^{\text{si},\text{red}}$ of simple perfect complexes on S is defined by the following homotopy pullback diagram:

$$\begin{array}{ccc} \mathbb{R}\mathbf{Perf}(S)^{\text{si},\text{red}} & \longrightarrow & \mathbb{R}\mathbf{Perf}(S)^{\text{si},>0} \\ \downarrow & & \downarrow \det_S \\ & & \mathbb{R}\mathbf{Pic}(S) \\ & & \downarrow \text{pr}_{\text{der}} \\ \text{Spec } \mathbb{C} & \xrightarrow{x_0} & \mathbb{R}\text{Spec}(\text{Sym}(H^0(S, K_S)[1])). \end{array}$$

Since the truncation functor commutes with homotopy pullbacks, the truncation of **the stack** $\mathbb{R}\mathbf{Perf}(S)^{\text{si},\text{red}}$ is the same as the truncation of $\mathbb{R}\mathbf{Perf}(S)^{\text{si},>0}$, i.e. $\mathbf{Perf}(S)^{\text{si},>0}$, therefore its coarse moduli space is again Inaba's $\text{Splcp}_S^{\text{ét}}/\mathbb{C}$ ([13, Section 3]).

Theorem 5.5. *The composite map*

$$\mathbb{R}\mathbf{Perf}(S)^{\text{si},>0} \xrightarrow{\det_S} \mathbb{R}\mathbf{Pic}(S) \xrightarrow{\text{pr}_{\text{der}}} \mathbb{R}\text{Spec}(\text{Sym}(H^0(S, K_S)[1]))$$

is smooth. Therefore the derived stack $\mathbb{R}\mathbf{Perf}(S)^{\text{si},\text{red}}$ is actually a smooth, usual (i.e. underived) stack, and

$$\mathbb{R}\mathbf{Perf}(S)^{\text{si},\text{red}} \simeq \text{t}_0(\mathbb{R}\mathbf{Perf}(S)^{\text{si},\text{red}}) \simeq \mathbf{Perf}(S)^{\text{si},>0}.$$

Under these identifications, the canonical map $\mathbb{R}\mathbf{Perf}(S)^{\text{si},\text{red}} \rightarrow \mathbb{R}\mathbf{Perf}(S)^{\text{si},>0}$ becomes isomorphic to the inclusion of the truncation $\mathbf{Perf}(S)^{\text{si},>0} \rightarrow \mathbb{R}\mathbf{Perf}(S)^{\text{si},>0}$.

Proof. Let E be a perfect complex on S such that $\text{Ext}_S^i(E, E) = 0$ for $i < 0$, and the trace map $\text{Ext}_S^0(E, E) \rightarrow H^0(S, \mathcal{O}_S) \simeq \mathbb{C}$ is an isomorphism. The homotopy fiber product defining $\mathbb{R}\mathbf{Perf}(S)^{\text{si},\text{red}}$ yields a distinguished triangle of tangent complexes

$$\mathbb{T}_E \mathbb{R}\mathbf{Perf}(S)^{\text{si},\text{red}} \rightarrow \mathbb{T}_E \mathbb{R}\mathbf{Perf}(S)^{\text{si},>0} \rightarrow H^0(S, K_S)^\vee[-1].$$

Since

$$\mathbb{T}_E \mathbb{R}\mathbf{Perf}(S)^{\text{si},>0} \simeq \mathbb{R}\text{End}_S(E)[1],$$

this complex is cohomologically concentrated in degrees $[-1, 1]$. Therefore, to prove the theo-

rem, it is enough to show that the map (induced by the above triangle on H^1)

$$\alpha : \mathrm{Ext}_S^2(E, E) \simeq H^1(\mathbb{T}_E \mathbb{R}\mathrm{Perf}(S)^{\mathrm{si}, > 0}) \rightarrow H^0(S, K_S)^\vee$$

is an isomorphism. If we denote by

$$\alpha' : \mathrm{Ext}_S^2(E, E) \xrightarrow{\alpha} H^0(S, K_S)^\vee \xrightarrow[\sim]{s} H^2(S, \mathcal{O}_S)$$

(the isomorphism s given by Serre duality), the following diagram:

$$\begin{array}{ccc} \mathrm{Ext}_S^2(E, E) & \xrightarrow{s} & \mathrm{Ext}_S^0(E, E)^\vee \\ \alpha' \downarrow & & \downarrow \mathrm{tr}_E^\vee \\ H^2(S, \mathcal{O}_S) & \xrightarrow{s} & H^0(S, \mathcal{O}_S)^\vee \end{array}$$

(where again, the s isomorphisms are given by Serre duality on S) is commutative. But, by hypothesis, tr_E is an isomorphism and we conclude. \square

The following corollary was first proved by Inaba [13, Theorem 3.2].

Corollary 5.6. *The coarse moduli space $\mathrm{Perf}(S)^{\mathrm{si}, > 0}$ of simple perfect complexes on a smooth projective K3-surface S is a smooth algebraic space.*

Proof. The stack $\mathbb{R}\mathrm{Perf}(S)^{\mathrm{si}, \mathrm{red}} \simeq \mathrm{Perf}(S)^{\mathrm{si}, > 0}$ is a \mathbb{G}_m -gerbe ([35, Corollary 3.22]), hence its smoothness is equivalent to the smoothness of its coarse moduli space (because the map to the coarse moduli space is smooth, being locally for the étale topology, given by the projection of $\mathrm{Perf}(S)^{\mathrm{si}, > 0} \times B\mathbb{G}_m$ onto the first factor). \square

Remark 5.7. Inaba shows in [13, Theorem 3.3] (again generalizing earlier results by Mukai in [22]) that the coarse moduli space $\mathrm{Perf}(S)^{\mathrm{si}, > 0}$ also carries a canonical *symplectic structure*. In [27] it is shown that in fact the whole derived stack $\mathbb{R}\mathrm{Perf}(S)$ carries a natural *derived symplectic structure* of degree 0, and that this induces on $\mathrm{Perf}(S)^{\mathrm{si}, > 0}$ the symplectic structure defined by Inaba.

5.2. On Calabi–Yau 3-folds. In this section, for X an arbitrary complex smooth projective variety, we first elaborate on the map

$$A_X^{(n)} : \mathbb{R}\overline{\mathbf{M}}_{g,n}(X) \rightarrow \mathbb{R}\mathrm{Perf}(X)$$

from Definition 3.9. This elaboration will give us a map $C_{X, \mathcal{L}}^{(n)}$ from a derived substack of $\mathbb{R}\overline{\mathbf{M}}_{g,n}(X)$ to the derived stack $\mathbb{R}\mathrm{Perf}(X)_{\mathcal{L}}^{\mathrm{si}, > 0}$, \mathcal{L} being a line bundle on X (see Definition 5.1). When we specialize to the case where X is a projective smooth *Calabi–Yau 3-fold* Y , we prove that $\mathbb{R}\mathrm{Perf}(Y)_{\mathcal{L}}^{\mathrm{si}, > 0}$ is *quasi-smooth* (Proposition 5.10), and that the map $C_{Y, \mathcal{L}}^{(n)}$ allows us to compare the induced obstruction theories on the truncations of its source and target.

To begin with, let X be a smooth complex projective variety. First of all, observe that taking tensor products of complexes induces an action of the derived group stack $\mathbb{R}\mathrm{Pic}(X)$ on $\mathbb{R}\mathrm{Perf}(X)$

$$\mu : \mathbb{R}\mathrm{Pic}(X) \times \mathbb{R}\mathrm{Perf}(X) \rightarrow \mathbb{R}\mathrm{Perf}(X).$$

Let $x_{\mathcal{L}} : \mathrm{Spec} \mathbb{C} \rightarrow \mathbb{R}\mathrm{Pic}(X)$ be the point corresponding to a line bundle \mathcal{L} on X .

Definition 5.8. Let $\sigma_{\mathcal{L}} : \mathbb{R}\mathbf{Pic}(X) \rightarrow \mathbb{R}\mathbf{Pic}(X)$ be the composite

$$\mathbb{R}\mathbf{Pic}(X) \xrightarrow{(\text{inv}, \times_{\mathcal{L}})} \mathbb{R}\mathbf{Pic}(X) \times \mathbb{R}\mathbf{Pic}(X) \xrightarrow{\times} \mathbb{R}\mathbf{Pic}(X)$$

where \times (resp., inv) denotes the product (resp., the inverse) map in $\mathbb{R}\mathbf{Pic}(X)$ (in other words, $\sigma_{\mathcal{L}}(\mathcal{L}_1) = \mathcal{L} \otimes \mathcal{L}_1^{-1}$).

- Define $A_{X, \mathcal{L}}^{(n)} : \mathbb{R}\overline{\mathbf{M}}_{g,n}(X) \rightarrow \mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}$ via the composite

$$\begin{aligned} \mathbb{R}\overline{\mathbf{M}}_{g,n}(X) &\xrightarrow{(\det \circ A_X^{(n)}, A_X^{(n)})} \mathbb{R}\mathbf{Pic}(X) \times \mathbb{R}\mathbf{Perf}(X) \\ &\xrightarrow{\sigma_{\mathcal{L}} \times \text{id}} \mathbb{R}\mathbf{Pic}(X) \times \mathbb{R}\mathbf{Perf}(X) \xrightarrow{\mu} \mathbb{R}\mathbf{Perf}(X) \end{aligned}$$

(in short, $A_{X, \mathcal{L}}^{(n)}(E) = E \otimes (\det E)^{-1} \otimes \mathcal{L}$).

- Define the derived open substack

$$\mathbb{R}\overline{\mathbf{M}}_{g,n}(X)^{\text{emb}} \hookrightarrow \mathbb{R}\overline{\mathbf{M}}_{g,n}(X)$$

as $\phi_{\mathbb{R}\overline{\mathbf{M}}_{g,n}(X)}(\overline{\mathbf{M}}_{g,n}(X)^{\text{emb}})$ (see Proposition 2.1) where $\overline{\mathbf{M}}_{g,n}(X)^{\text{emb}}$ is the open substack of the stack $\overline{\mathbf{M}}_{g,n}(X)$ consisting of pointed stable maps which are closed immersions.

- Define $C_{X, \mathcal{L}}^{(n)} : \mathbb{R}\overline{\mathbf{M}}_{g,n}(X)^{\text{emb}} \rightarrow \mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}^{\text{si}, > 0}$ via the composite

$$\mathbb{R}\overline{\mathbf{M}}_{g,n}(X)^{\text{emb}} \hookrightarrow \mathbb{R}\overline{\mathbf{M}}_{g,n}(X) \xrightarrow{A_{X, \mathcal{L}}^{(n)}} \mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}$$

(note that this composite indeed factors through $\mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}^{\text{si}, > 0}$, since

$$\text{tr} : \text{Ext}^0(\mathbb{R}f_*\mathcal{O}_C, \mathbb{R}f_*\mathcal{O}_C) \simeq \mathbb{C}$$

if the pointed stable map f is a closed immersion).

Remark 5.9. The map $C_{X, \mathcal{L}}^{(n)}$ is also defined on the a priori larger open derived substack consisting (in the sense of Proposition 2.1) of pointed stable maps f such that the trace map $\text{tr} : \text{Ext}^0(\mathbb{R}f_*\mathcal{O}_C, \mathbb{R}f_*\mathcal{O}_C) \rightarrow H^0(X, \mathcal{O}_X) \simeq \mathbb{C}$ is an isomorphism.

We would like to use the map $C_{X, \mathcal{L}}^{(n)}$ to induce a comparison map between the induced obstruction theories on the truncations of $\mathbb{R}\overline{\mathbf{M}}_{g,n}(X)^{\text{emb}}$ and of $\mathbb{R}\mathbf{Perf}(X)_{\mathcal{L}}^{\text{si}, > 0}$.

This is possible when we take X to be a Calabi–Yau 3-fold Y . In fact:

Theorem 5.10. *If Y is a smooth complex projective Calabi–Yau 3-fold, then the derived stack $\mathbb{R}\mathbf{Perf}(Y)_{\mathcal{L}}^{\text{si}, > 0}$ is quasi-smooth. Therefore, the closed immersion*

$$j : \mathbf{Perf}(Y)_{\mathcal{L}}^{\text{si}, > 0} \hookrightarrow \mathbb{R}\mathbf{Perf}(Y)_{\mathcal{L}}^{\text{si}, > 0}$$

induces a $[-1, 0]$ -perfect obstruction theory

$$j^* \mathbb{T}_{\mathbb{R}\mathbf{Perf}(Y)_{\mathcal{L}}^{\text{si}, > 0}} \rightarrow \mathbb{T}_{\mathbf{Perf}(Y)_{\mathcal{L}}^{\text{si}, > 0}}.$$

Proof. This is a corollary of Proposition 5.2. Let \mathbb{T}_E denote the tangent complex of $\mathbb{R}\mathbf{Perf}(Y)_{\mathcal{L}}^{\mathrm{si}, > 0}$ at a point corresponding to the perfect complex E . Now Y is Calabi–Yau of dimension 3, so

$$\Omega_Y^3 \equiv K_Y \simeq \mathcal{O}_Y;$$

but E is simple (i.e. the trace map $\mathrm{Ext}^0(F, F) \rightarrow H^0(X, \mathcal{O}_X) \simeq \mathbb{C}$ is an isomorphism), so Serre duality implies $\mathrm{Ext}^i(E, E)_0 = 0$ for $i \geq 3$ (and all $i \leq 0$). Therefore the perfect complex \mathbb{T}_E is concentrated in degrees $[0, 1]$, and $\mathbb{R}\mathbf{Perf}(Y)_{\mathcal{L}}^{\mathrm{si}, > 0}$ is quasi-smooth. The second assertion follows immediately from Proposition 1.2. \square

Remark 5.11. Note that the stack $\mathbf{Perf}(Y)_{\mathcal{L}}^{\mathrm{si}, > 0}$ is *not* proper over $\mathrm{Spec} \mathbb{C}$. However it receives maps from both Thomas moduli space $I_n(Y; \beta)$ of ideal sheaves (whose sub-schemes have Euler characteristic n and fundamental class $\beta \in H_2(Y, \mathbb{Z})$) – see [31] – and from Pandharipande–Thomas moduli space $P_n(Y; \beta)$ of stable pairs – see [26]. For example, the map from $P_n(Y; \beta)$ sends a pair to the pair itself, considered as a complex on Y . Moreover, at the points in the image of such maps, the tangent and obstruction spaces of these spaces, as considered in [26], are the same as those induced from the cotangent complex of our $\mathbb{R}\mathbf{Perf}(Y)_{\mathcal{L}}^{\mathrm{si}, > 0}$ ([26, Section 2.1]).

As showed in Section 1.2, the map

$$C_{Y, \mathcal{L}}^{(n)} : \mathbb{R}\overline{\mathbf{M}}_{g,n}(Y)^{\mathrm{emb}} \rightarrow \mathbb{R}\mathbf{Perf}(Y)_{\mathcal{L}}^{\mathrm{si}, > 0}$$

induces a comparison map between the two obstruction theories. More precisely, the commutative diagram in $\mathbf{dSt}_{\mathbb{C}}$

$$\begin{array}{ccc} \overline{\mathbf{M}}_{g,n}(Y)^{\mathrm{emb}} & \xrightarrow{t_0 C_{Y, \mathcal{L}}^{(n)}} & \mathbf{Perf}(Y)_{\mathcal{L}}^{\mathrm{si}, > 0} \\ j_{GW} \downarrow & & \downarrow j_{DT} \\ \mathbb{R}\overline{\mathbf{M}}_{g,n}(Y)^{\mathrm{emb}} & \xrightarrow{C_{Y, \mathcal{L}}^{(n)}} & \mathbb{R}\mathbf{Perf}(Y)_{\mathcal{L}}^{\mathrm{si}, > 0} \end{array}$$

(where each j is the closed immersion of the truncation of a derived stack into the full derived stack), induces a morphism of triangles

$$\begin{array}{ccccc} (t_0 C_{Y, \mathcal{L}}^{(n)})^* j_{DT}^* \mathbb{L}_{\mathbb{R}\mathbf{Perf}(Y)_{\mathcal{L}}^{\mathrm{si}, > 0}} & \rightarrow & (t_0 C_{Y, \mathcal{L}}^{(n)})^* \mathbb{L}_{\mathbf{Perf}(Y)_{\mathcal{L}}^{\mathrm{si}, > 0}} & \rightarrow & (t_0 C_{Y, \mathcal{L}}^{(n)})^* \mathbb{L}_{\mathbb{R}\mathbf{Perf}(Y)_{\mathcal{L}}^{\mathrm{si}, > 0} / \mathbf{Perf}(Y)_{\mathcal{L}}^{\mathrm{si}, > 0}} \\ \downarrow & & \downarrow & & \downarrow \\ j_{GW}^* \mathbb{L}_{\mathbb{R}\overline{\mathbf{M}}_{g,n}(Y)^{\mathrm{emb}}} & \longrightarrow & \mathbb{L}_{\overline{\mathbf{M}}_{g,n}(Y)^{\mathrm{emb}}} & \longrightarrow & \mathbb{L}_{\mathbb{R}\overline{\mathbf{M}}_{g,n}(Y)^{\mathrm{emb}} / \overline{\mathbf{M}}_{g,n}(Y)^{\mathrm{emb}}} \end{array}$$

– in the derived category of perfect complexes on $\overline{\mathbf{M}}_{g,n}(Y)^{\mathrm{emb}}$ – i.e. a morphism relating the two obstruction theories induced on the truncations stacks $\overline{\mathbf{M}}_{g,n}(Y)^{\mathrm{emb}}$ and $\mathbf{Perf}(Y)_{\mathcal{L}}^{\mathrm{si}, > 0}$. Note that, for the object in the upper left corner of the above diagram, we have a natural isomorphism

$$(t_0 C_{Y, \mathcal{L}}^{(n)})^* j_{DT}^* \mathbb{L}_{\mathbb{R}\mathbf{Perf}(Y)_{\mathcal{L}}^{\mathrm{si}, > 0}} \simeq j_{GW}^* (C_{Y, \mathcal{L}}^{(n)})^* \mathbb{L}_{\mathbb{R}\mathbf{Perf}(Y)_{\mathcal{L}}^{\mathrm{si}, > 0}}.$$

A. Derived stack of perfect complexes and Atiyah classes

We explain here the relationship between the tangent maps associated to morphisms to the derived stack of perfect complexes and Atiyah classes (of perfect complexes) used in the main text (see Section 3.2). As in the main text, we work over \mathbb{C} , even if most of what we say below holds true over any field of characteristic zero. As usual, all tensor products and fiber products will be implicitly derived, and we will simply write g^* for the derived pullback $\mathbb{L}g^*$, and g_* for the derived push-forward $\mathbb{R}g_*$, for any map g below.

If \mathcal{Y} is a derived geometric stack having a perfect cotangent complex ([36, Section 1.4]), and E is a perfect complex on \mathcal{Y} , then we will implicitly identify the Atiyah class map of E

$$\mathrm{at}_E : E \rightarrow \mathbb{L}\mathcal{Y} \otimes E[1]$$

with the corresponding map

$$\mathbb{T}_{\mathcal{Y}} \rightarrow E^\vee \otimes E[1]$$

via the bijection

$$[\mathbb{T}_{\mathcal{Y}}, E^\vee \otimes E[1]] \simeq [\mathbb{T}_{\mathcal{Y}} \otimes E, E[1]] \simeq [E, \mathbb{L}\mathcal{Y} \otimes E[1]]$$

given by the adjunction $(\otimes, \mathbb{R}\underline{\mathrm{Hom}})$, and perfectness of E and $\mathbb{L}\mathcal{Y}$ (where $[-, -]$ denotes the **Hom** set in the derived category of perfect complexes on \mathcal{Y}).

We start with a quite general situation. Let \mathcal{Y} be a derived geometric stack having a perfect cotangent complex, and $\mathbb{R}\mathbf{Perf}$ the stack of perfect complexes (see Section 3). Then, giving a map of derived stacks $\phi_E : \mathcal{Y} \rightarrow \mathbb{R}\mathbf{Perf}$ is the same thing as giving a perfect complex E on \mathcal{Y} , and

- $\phi_E^* \mathbb{T}_{\mathbb{R}\mathbf{Perf}} \simeq \mathbb{R}\underline{\mathrm{End}}_{\mathcal{Y}}(E)[1]$,
- the tangent map to ϕ_E

$$\mathbb{T}\phi_E : \mathbb{T}_{\mathcal{Y}} \rightarrow \phi_E^* \mathbb{T}_{\mathbb{R}\mathbf{Perf}} \simeq \mathbb{R}\underline{\mathrm{End}}_{\mathcal{Y}}(E)[1] \simeq E^\vee \otimes E[1]$$

is the Atiyah class map at_E of E .

Remark A.1. The second point above might be considered as a definition when \mathcal{Y} is a derived stack, and it coincides with Illusie's definition ([12, Chapter 4, Section 2.3.7]) when $\mathcal{Y} = Y$ is a quasi-projective scheme. In fact, in this case, the map ϕ_E factors through the stack of strict perfect complexes; thus the proof reduces immediately to the case where E is a vector bundle on Y , which is straightforward.

The above description applies in particular to a map of derived stacks of the form

$$\Phi_E : \mathcal{Y} := \mathcal{S} \times X \rightarrow \mathbb{R}\mathbf{Perf}$$

where X is a smooth projective scheme, \mathcal{S} is a derived geometric stack having a perfect cotangent complex, and E is a perfect complex on $\mathcal{S} \times X$: in the main text we are interested in $\mathcal{S} = \mathbb{R}\overline{\mathbf{M}}_g(X)$. Such a map corresponds, by adjunction, to a map

$$\Psi_E : \mathcal{S} \rightarrow \mathbb{R}\mathrm{HOM}(X, \mathbb{R}\mathbf{Perf}) = \mathbb{R}\mathbf{Perf}(X).$$

The tangent map of Ψ_E fits into the following commutative diagram:

$$\begin{array}{ccccc}
 \mathbb{T}_{\mathcal{S}} & \xrightarrow{\mathbb{T}\Psi_E} & \Psi_E^* \mathbb{T}_{\mathbb{R}\mathrm{Perf}(X)} & \xrightarrow{\sim} & \mathrm{pr}_{\mathcal{S},*}(E^\vee \otimes E)[1] \\
 \downarrow \mathrm{can} & & & & \uparrow \mathrm{pr}_{\mathcal{S},*}(\mathbb{T}\Phi_E) \\
 \mathrm{pr}_{\mathcal{S},*} \mathrm{pr}_{\mathcal{S}}^* \mathbb{T}_{\mathcal{S}} & \xrightarrow{\mathrm{can}} & \mathrm{pr}_{\mathcal{S},*}(\mathrm{pr}_{\mathcal{S}}^* \mathbb{T}_{\mathcal{S}} \oplus \mathrm{pr}_X^* \mathbb{T}_X) & \xrightarrow{\sim} & \mathrm{pr}_{\mathcal{S},*} \mathbb{T}_{\mathcal{S} \times X}
 \end{array}$$

where can denote obvious canonical maps, and we can identify $\mathrm{pr}_{\mathcal{S},*}(\mathbb{T}\Phi_E)$ with $\mathrm{pr}_{\mathcal{S},*}(\mathrm{at}_E)$, in the sense explained above. In other words, $\mathbb{T}\Psi_E$ is described in terms of the relative Atiyah class map

$$\mathrm{at}_{E/X} : \mathrm{pr}_{\mathcal{S}}^* \mathbb{T}_{\mathcal{S}} \simeq \mathbb{T}_{\mathcal{S} \times X/X} \rightarrow E^\vee \otimes^{\mathbb{L}} E[1]$$

of E relative to X , as the composition

$$\mathbb{T}\Psi_E : \mathbb{T}_{\mathcal{S}} \xrightarrow{\mathrm{can}} \mathrm{pr}_{\mathcal{S},*} \mathrm{pr}_{\mathcal{S}}^* \mathbb{T}_{\mathcal{S}} \xrightarrow{\sim} \mathrm{pr}_{\mathcal{S},*} \mathbb{T}_{\mathcal{S} \times X/X} \xrightarrow{\mathrm{pr}_{\mathcal{S},*}(\mathrm{at}_{E/X})} \mathrm{pr}_{\mathcal{S},*}(E^\vee \otimes E)[1].$$

Remark A.2. The map $\mathbb{T}\Psi_E$ might be viewed as a generalization of what is sometimes called the Kodaira–Spencer map associated to the \mathcal{S} -family E of perfect complexes over X (e.g. [15, formula (14)]).

In the main text, we are interested in the case $\mathcal{S} = \mathbb{R}\overline{\mathbf{M}}_g(X)$, $\mathrm{pr} := \mathrm{pr}_{\mathcal{S}}$, and E perfect of the form $\pi_* \mathcal{E}$, where

$$\pi : \mathbb{R}\mathcal{C}_{g;X} \rightarrow \mathbb{R}\overline{\mathbf{M}}_g(X) \times X$$

is the universal map and \mathcal{E} is a complex on $\mathbb{R}\mathcal{C}_{g;X}$, namely $\mathcal{E} = \mathcal{O}_{\mathbb{R}\mathcal{C}_{g;X}}$. In such cases, if we call $(f : C \rightarrow X)$ the stable map corresponding to the complex point x , we have a ladder of homotopy cartesian diagrams

$$\begin{array}{ccccc}
 C & \xrightarrow{\iota_f} & \mathbb{R}\mathcal{C}_{g;X} & & \\
 f \downarrow & & \downarrow \pi & & \\
 X & \xrightarrow{\underline{x}} & \mathbb{R}\overline{\mathbf{M}}_g(X) \times X & \xrightarrow{\mathrm{pr}_X} & X \\
 q \downarrow & & \downarrow \mathrm{pr} & & \downarrow q \\
 \mathrm{Spec} \mathbb{C} & \xrightarrow{x} & \mathbb{R}\overline{\mathbf{M}}_g(X) & \longrightarrow & \mathrm{Spec} \mathbb{C}
 \end{array}$$

and the base-change isomorphism (true in derived algebraic geometry with no need of flatness) gives us

$$\underline{x}^* E = \underline{x}^* \pi_* \mathcal{E} \simeq f_* \iota_f^* \mathcal{E}.$$

For $\mathcal{E} = \mathcal{O}_{\mathbb{R}\mathcal{C}_{g;X}}$, we then get

$$\underline{x}^* E = \underline{x}^* \pi_* \mathcal{O}_{\mathbb{R}\mathcal{C}_{g;X}} \simeq f_* \mathcal{O}_C.$$

Again by base-change formula, we get

$$x^* \mathrm{pr}_* \simeq q_* \underline{x}^*,$$

and therefore the tangent map to $A_X := \Psi_{\pi_* \mathcal{O}_{\mathbb{R}\mathcal{C}_{g;X}}}$ at the point $x = (f : C \rightarrow X)$ is the

composition

$$\begin{aligned} \mathbb{T}_x A_X : \mathbb{T}_x \mathbb{R}\overline{\mathbf{M}}_g(X) &\simeq \mathbb{R}\Gamma(C, \text{Cone}(\mathbb{T}_C \rightarrow f^* T_X)) \\ &\longrightarrow \mathbb{R}\Gamma(X, \underline{x}^* \mathbb{T}_{\mathbb{R}\overline{\mathbf{M}}_g(X) \times X}) \xrightarrow{\mathbb{R}\Gamma(X, \underline{x}^* \text{at}_E)} \mathbb{R}\text{End}_X(\mathbb{R}f_* \mathcal{O}_C)[1] \simeq \mathbb{T}_{f_* \mathcal{O}_C} \mathbb{R}\mathbf{Perf}(X). \end{aligned}$$

The following is the third **assertion** in Proposition 3.6, Section 3.2.

Proposition A.3. *The composition*

$$\begin{aligned} \mathbb{R}\Gamma(X, T_X) &\xrightarrow{\text{can}} \mathbb{R}\Gamma(X, f_* f^* T_X) \\ &\xrightarrow{\text{can}} \mathbb{R}\Gamma(X, \text{Cone}(f_* \mathbb{T}_C \rightarrow f_* f^* T_X)) \simeq \mathbb{T}_x \mathbb{R}\overline{\mathbf{M}}_g(X) \\ &\xrightarrow{\mathbb{T}_x A_X} x^* A_X^* \mathbb{T} \mathbb{R}\mathbf{Perf}(X) \simeq \mathbb{T}_{f_* \mathcal{O}_C} \mathbb{R}\mathbf{Perf}(X) \simeq \mathbb{R}\text{End}_X(f_* \mathcal{O}_C)[1] \end{aligned}$$

coincides with $\mathbb{R}\Gamma(X, \text{at}_{f_* \mathcal{O}_C})$.

Proof. We first observe that if \mathcal{F} is perfect complex on X , and $\mathbb{R}\mathbf{Aut}(X)$ is the derived stack of automorphisms of X , there are obvious maps of derived stacks

$$\rho_x : \mathbb{R}\mathbf{Aut}(X) \rightarrow \mathbb{R}\mathbf{HOM}_{\mathbf{dSt}_C}(C, X)$$

and

$$\sigma_{\mathcal{F}} : \mathbb{R}\mathbf{Aut}(X) \rightarrow \mathbb{R}\mathbf{Perf}(X)$$

induced by the natural action of $\mathbb{R}\mathbf{Aut}(X)$ by composition on maps and by pullbacks on perfect complexes, respectively. Moreover, the tangent map to $\sigma_{\mathcal{F}}$ at the identity $\text{Spec } \mathbb{C}$ -point of $\mathbb{R}\mathbf{Aut}(X)$

$$\mathbb{T}_{\text{id}_X} \sigma_{\mathcal{F}} : \mathbb{R}\Gamma(X, T_X) \simeq \mathbb{T}_{\text{id}_X} \mathbb{R}\mathbf{Aut}(X) \rightarrow \mathbb{T}_{\mathcal{F}} \mathbb{R}\mathbf{Perf}(X) \simeq \mathbb{R}\text{End}_X(\mathcal{F})[1]$$

is $\mathbb{R}\Gamma(X, \text{at}_{\mathcal{F}})$, where $\text{at}_{\mathcal{F}}$ is the Atiyah class map of \mathcal{F} . Then we observe that, by taking

$$\mathcal{F} := \underline{x}^* \pi_* \mathcal{O}_{\mathbb{R}\mathcal{C}_{g; X}}$$

– which is, by base-change formula, isomorphic to $f_* \mathcal{O}_C$ – we get that the composition

$$k_x : \mathbb{R}\mathbf{Aut}(X) \xrightarrow{\rho_x} \mathbb{R}\mathbf{HOM}_{\mathbf{dSt}_C}(C, X) \xrightarrow{\text{can}} \mathbb{R}\overline{\mathbf{M}}_g(X) \xrightarrow{A_X} \mathbb{R}\mathbf{Perf}(X)$$

coincides with $\sigma_{\mathcal{F}}$. But the map in the statement of the proposition is just $\mathbb{T}_{\text{id}_X} k_x$, and we conclude. \square

References

- [1] *K. Behrend*, Gromov–Witten invariants in algebraic geometry, *Invent. Math.* **127** (1997), 601–617.
- [2] *K. Behrend* and *J. Bryan*, Super-rigid Donaldson–Thomas invariants, *Math. Res. Lett.* **14** (2007), 559–571.
- [3] *K. Behrend* and *B. Fantechi*, The intrinsic normal cone, *Invent. Math.* **128** (1997), 45–88.
- [4] *K. Behrend* and *Y. Manin*, Stacks of stable maps and Gromov–Witten invariants, *Duke Math. J.* **85** (1996), no. 1, 1–60.

Note 10:
Please propose
where to add
references to [2, 14,
16, 18, 19, 28, 30]
(unicited). Or shall
I remove these
entries?

- [5] *P. Berthelot, A. Grothendieck and L. Illusie*, Séminaire de géométrie algébrique du Bois Marie 1966/67 (SGA 6). Théorie des intersections et théorème de Riemann-Roch, Lecture Notes in Math. **225**, Springer-Verlag, Berlin 1971.
- [6] *S. Bloch*, Semi-regularity and De Rham cohomology, *Invent. Math.* **17** (1972), 51–66.
- [7] *R.-O. Buchweitz and H. Flenner*, A semiregularity map for modules and applications to deformations, *Compos. Math.* **137** (2003), 135–210.
- [8] *I. Ciocan-Fontanine and M. Kapranov*, The derived Hilbert scheme, *J. Amer. Math. Soc.* **15** (2002), 787–815.
- [9] *B. Conrad*, Grothendieck duality and base change, *Lecture Notes in Math.* **1750**, Springer-Verlag, Berlin 2001.
- [10] *R. Hartshorne*, Residues and duality, *Lecture Notes in Math.* **20**, Springer-Verlag, Berlin 1966.
- [11] *D. Huybrechts and M. Lehn*, The geometry of moduli spaces of sheaves, 2nd ed., Cambridge University Press, Cambridge 2010.
- [12] *L. Illusie*, Complexe cotangent et déformations I, *Lecture Notes in Math.* **239**, Springer-Verlag, Berlin 1971.
- [13] *M. Inaba*, Smoothness of the moduli space of complexes of coherent sheaves on an abelian or projective $K3$ surface, *Adv. Math.* **227** (2011), no. 4, 1399–1412.
- [14] *S. Kleiman*, The Picard scheme, in: *Fundamental algebraic geometry. Grothendieck’s FGA explained*, *Math. Surveys Monogr.* **123**, American Mathematical Society, Providence 2005.
- [15] *A. Kuznetsov and D. Markushevich*, Symplectic structures on moduli spaces of sheaves via the Atiyah class, *J. Geom. Phys.* **59** (2009), 843–860.
- [16] *M. A. Mandell*, An inverse K -theory functor, *Doc. Math.* **15** (2010), 765–791.
- [17] *C. Manolache*, Virtual pullbacks, *J. Algebraic Geom.* **21** (2011), 201–245.
- [18] *D. Maulik, N. Nekrasov, A. Okounkov and R. Pandharipande*, Gromov–Witten theory and Donaldson–Thomas theory I, *Compos. Math.* **142** (2006), 1263–1285.
- [19] *D. Maulik, A. Oblomkov, A. Okounkov and R. Pandharipande*, Gromov–Witten/Donaldson–Thomas correspondence for toric 3-folds, *Invent. Math.* **186** (2011), 435–479.
- [20] *D. Maulik and R. Pandharipande*, Gromov–Witten theory and Noether–Lefschetz theory, preprint 2012, <http://arxiv.org/abs/0705.1653>.
- [21] *D. Maulik, R. Pandharipande and R. P. Thomas*, Curves on $K3$ -surfaces and modular forms, *J. Topol.* **3** (2010), 937–996.
- [22] *S. Mukai*, Symplectic structure of the moduli space of sheaves on an abelian or $K3$ surface, *Invent. Math.* **77** (1984), 101–116.
- [23] *A. Okounkov and R. Pandharipande*, Gromov–Witten theory, Hurwitz numbers, and Matrix models, in: *Algebraic geometry II* (Seattle 2005), part 2, *Proc. Sympos. Pure Math.* **80**, American Mathematical Society, Providence 2009.
- [24] *A. Okounkov and R. Pandharipande*, Quantum cohomology of the Hilbert scheme of points of the plane, *Invent. Math.* **179** (2010), 523–557.
- [25] *R. Pandharipande*, Maps, sheaves and $K3$ -surfaces, *Clay Research Lectures*, preprint 2008, <http://arxiv.org/abs/0808.0253>.
- [26] *R. Pandharipande and R. P. Thomas*, Curve counting via stable pairs in the derived category, *Invent. Math.* **178** (2009), 407–447.
- [27] *T. Pantev, B. Toën, G. Vezzosi and M. Vaquié*, Shifted symplectic structures, *Publ. Math. Inst. Hautes Études Sci.* **117** (2013), no. 1, 271–328.
- [28] *Z. Ran*, Semiregularity, obstructions and deformations of Hodge classes, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (4)* **28** (1999), 809–821.
- [29] *T. Schürg*, Deriving Deligne–Mumford stacks with perfect obstruction theories, Ph.D. thesis, Johannes Gutenberg Universität Mainz, 2011.
- [30] *C. Simpson*, Moduli of representations of the fundamental group of a smooth projective variety I, *Publ. Math. Inst. Hautes Études Sci.* **79** (1994), 47–129.
- [31] *R. P. Thomas*, A holomorphic Casson invariant for Calabi–Yau 3-folds, and bundles on $K3$ fibrations, *J. Differential Geom.* **54** (2000), no. 2, 367–438.
- [32] *B. Toën*, Champs affines, *Selecta Math. (N.S.)* **12** (2006), no. 1, 39–135.
- [33] *B. Toën*, Higher and derived stacks: A global overview, in: *Algebraic geometry I* (Seattle 2005), part 1, *Proc. Sympos. Pure Math.* **80**, American Mathematical Society, Providence (2009), 435–487.
- [34] *B. Toën*, Proper local complete intersection morphisms preserve perfect complexes, preprint 2012, <http://arxiv.org/abs/1210.2827>.
- [35] *B. Toën and M. Vaquié*, Moduli of objects in dg-categories, *Ann. Sci. Éc. Norm. Supér. (4)* **40** (2007), no. 3, 387–444.
- [36] *B. Toën and G. Vezzosi*, Homotopical algebraic geometry II: Geometric stacks and applications, *Mem. Amer. Math. Soc.* **193** (2008), no. 902.

- [37] *B. Toën and G. Vezzosi*, Algèbres simpliciales S^1 -équivariantes, théorie de de Rham et théorèmes HKR multiplicatifs, *Compos. Math.* **147** (2011), 1979–2000.
- [38] *F. Waldhausen*, Algebraic K -theory of spaces, in: Algebraic and geometric topology (New Brunswick 1983), *Lecture Notes in Math.* **1126**, Springer-Verlag, Berlin (1985), 318–419.
- [39] *C. Weibel*, An introduction to homological algebra, *Cambridge Stud. Adv. Math.* **38**, Cambridge University Press, Cambridge 1995.

Timo Schürg, **Max-Planck-Institut für Mathematik, P.O. Box 7280, 53072 Bonn, Germany**

Bertrand Toën, Institut de Mathématiques et de Modélisations de Montpellier, I3M UMR 5149,
Université de Montpellier 2, Case 051 Batiment 9, Place Eugène Bataillon, Montpellier Cedex 5, France

Gabriele Vezzosi, Institut de Mathématiques de Jussieu - UMR7586 Batiment Sophie Germain,
Université Paris Diderot, Case 7012, 75205 Paris Cedex 13, France

Eingegangen 4. November 2011, in revidierter Fassung 13. April 2013