



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Systematic investigations concerning *Saxifraga granulata* s.l. in the Tuscan Archipelago (northern Tyrrhenian sea): evidences for

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Systematic investigations concerning *Saxifraga granulata* s.l. in the Tuscan Archipelago (northern Tyrrhenian sea): evidences for describing new taxa in Capraia and Montecristo islands / G. Ferretti; M. Mannocci; V. Mazzoncini; G. Fiorini; B. Foggi; D. Viciani. - ELETTRONICO. - (2014), pp. 155-155. (109° Congresso della Società Botanica Italiana, International Plant Science Conference (IPSC), From Nature to

Availability:

The webpage <https://hdl.handle.net/2158/901164> of the repository was last updated on

Publisher:

Società Botanica Italiana

Terms of use:

Open Access


La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

La data sopra indicata si riferisce all'ultimo aggiornamento della scheda del Repository FloRe - The above-mentioned date refers to the last update of the record in the Institutional Repository FloRe

(Article begins on next page)

With the High Patronage of the President of the Italian Republic



109° Congresso della Società Botanica Italiana

**INTERNATIONAL PLANT SCIENCE CONFERENCE
(IPSC)**

FROM NATURE TO TECHNOLOGICAL EXPLOITATIONS
Florence, 2 - 5 September 2014

Programme

Index

Relations

Communications

Posters

Author Index

Authors


Scientific Committee

Local Committee

Sponsor

Supporter

Maps of Congressual sites



With the High Patronage of the President of the Italian Republic

Società Botanica Italiana
109° Congresso
International Plant Science Conference (IPSC)

Florence, 2 - 5 September 2014

Scientific Committee

Prof. Donato Chiatante (President)

Prof. Francesco Maria Raimondo (SBI President)

Prof. Pier Virgilio Arrigoni

Prof. Andrea Bennici

Prof.ssa Maria Beatrice Bitonti

Prof.ssa Paola Bonfante

Prof. Carlo Blasi

Prof. Alessandro Bruni

Prof. Ignazio Camarda

Prof.ssa Nicoletta Cannone

Prof. Guido Chelazzi

Prof.ssa Lucia Colombo

Prof. Giovanni Cristofolini

Prof. Giuseppe Dalessandro

Prof. Fabio Garbari

Prof. Paolo Grossoni

Prof.ssa Laura Maleci

Prof.ssa Michela Marignani

Prof.ssa Marta Mariotti Lippi

Prof. Guido Moggi

Prof. Enio Nardi

Prof. Ettore Pacini

Prof. Mauro Raffaelli

Prof. Bruno Romano

Prof.ssa Consolata Siniscalco

Home page

4.4 = SYSTEMATIC INVESTIGATIONS CONCERNING *SAXIFRAGA GRANULATA* S.L. IN THE TUSCAN ARCHIPELAGO (NORTHERN TYRRHENIAN SEA): EVIDENCES FOR DESCRIBING NEW TAXA IN CAPRAIA AND MONTECRISTO ISLANDS

GIULIO FERRETTI¹, MAIRO MANNOCCI², VIVIANO MAZZONCINI², GRAZIANA FIORINI¹, BRUNO FOGGI¹, DANIELE VICIANI¹

¹Department of Biology, University of Florence, Via G. La Pira 4, 50121 Firenze, Italy; ²Museo di Storia Naturale del Mediterraneo, Via Roma 234, 57127 Livorno, Italy

The genus *Saxifraga* L. (*Saxifragaceae*) has a holarctic distribution and counts more than 440 species, including about 120 living in Europe (1). Seven species with four additional subspecies are currently reported within the Series *Saxifraga* (2). Multi-year floristic and systematic research both in the wild in the Tuscan Archipelago (Central Italy) and in herbaria allowed to investigate some interesting populations of *Saxifraga* living in Capraia, Elba and Montecristo islands and known since the end of the nineteenth century. The territory covered by the three populations is located on the border between the distribution areas of *S. granulata* L. and *S. corsica* (Ser.) Gren. & Godr. Over the years the taxonomic treatment concerning these populations has undergone various interpretations, sometimes considering them as belonging to *S. granulata* group and sometimes as belonging to *S. corsica* group. The Capraia population was formally taken into account by Sommier (3), who described a new infra-specific taxon with the name *S. granulata* var. *brevicaulis* Sommier.

The three populations were analyzed both from the morphological and caryological points of view. Morphological investigation was carried out both on living cultivated plants and on dried specimens (exsiccata) stored in several herbaria (FI, FIAF, GDOR, GE, PIAGR, PI). Morphologically the three populations can be assigned to two distinct groups: the Elba population presents the typical features of *Saxifraga granulata* s.s., whereas the populations of Capraia and Montecristo certainly belong to the group of *S. granulata* s.l. (including *S. corsica*); however they show numerous peculiar characteristics that make it difficult to identify them within *S. granulata* s.s. or *S. corsica* s.s. (plant height less than 20 cm, corymbiform inflorescence, style-stigma complex shorter than the calyx, stigma flattened and elongated rather than rounded).

Cytological studies in the genus *Saxifraga* present many practical difficulties (4), further complicated by an extremely marked intraspecific aneuploidy, leading to different chromosome numbers within populations taxonomically considered as individual species (5). The ongoing caryological investigation showed low chromosome counts for the Elba and Montecristo populations and higher counts for the Capraia population. These results and other morphometric evidences as well highlight the differences between the populations of Capraia and Montecristo.

The affinity of the Elba population with *S. granulata* s.s., widely distributed in the Italian peninsula, can be well explained by the geographic contacts between the Elba Island and the mainland till the last glacial period. Similarly, the isolation of Capraia and Montecristo, dating back to the Upper Messinian, may have allowed the differentiation and stabilization of individuals distinguishable from the original pool.

In conclusion, based on the results of the present study, and on the geographical and ecological isolation, we propose to assign the populations of Capraia and Montecristo to two new taxa, that will be described in shortly, separated either from *S. granulata* s.s. or *S. corsica* s.s.

1) D.A. Webb (1993) *Saxifraga* L.. In: Tutin T.G., Burges N.A., Chater. A.O., Edmondson J.R., Heywood V.H., Moore D.M., Valentine D.H., Walters S.M., Webb D.A. (eds), 1993 - Flora Europaea. 2nd edition. Volume 1. Cambridge University Press, Cambridge, United Kingdom. pp. 437-458.

2) D.A. Webb, R.J. Gornall (1989) *Saxifrages of Europe*. Christopher Helm, London

3) S. Sommier (1898) Aggiunte alla florula di Capraia. *Nuovo Giorn Bot Ital* n.s. 5, 106–139

4) C. Favarger (1965) Notes de caryologie alpine, IV. *Bull. Soc. Neucheloise Sci. Nat.* 88, 15-23

5) S. Kumar, S. Kumari, R.C. Gupta, V.K. Sharma (2013) Additions to the cytology of *Saxifraga* (*Saxifragaceae*) from the Western Himalayas, India. *Botanica Serbica* 37(2), 147-154

