

As a library, NLM provides access to scientific literature. Inclusion in an NLM database does not imply endorsement, or agreement with, the contents by NLM or the National Institutes of Health. Learn more about our disclaimer.

Wiley Public Health Emergency Collection Public Health Emergency COVID-19 Initiative

J Dermatol. 2022 Aug 17 : 10.1111/1346-8138.16554. doi: 10.1111/1346-8138.16554 [Epub ahead of print]

PMCID: PMC9538601 PMID: 35975548

Five cases of new-onset pemphigus following vaccinations against coronavirus disease 2019

Alberto Corrá,<sup>1</sup> Francesca Barei,<sup>2-3</sup> Giovanni Genovese,<sup>2-3</sup> Martina Zussino,<sup>2</sup> Cristina B. Spigariolo,<sup>2,3</sup> Elena B. Mariotti,<sup>1</sup> Lavinia Quintarelli,<sup>4</sup> Alice Verdelli,<sup>4</sup> Marzia Caproni,<sup>4</sup> and Angelo V. Marzano<sup>2,3</sup>

Author information | Article notes | Copyright and License information | Disclaimer

Abstract

Go to:

Pemphigus is a group of blistering disorders characterized by the formation of intraepithelial blisters in skin and mucous membranes induced by the binding of circulating autoantibodies to intercellular adhesion molecules. The pathogenesis is complex and not fully understood; however, genetic predisposition and various triggers are widely accepted as key factors in pemphigus development. A few cases of new-onset pemphigus following coronavirus disease 2019 (COVID-19) vaccination have already been published. The present paper reports a total of two cases of pemphigus foliaceus and three cases of pemphigus vulgaris that occurred following vaccinations against COVID-19, with anamnestic, clinical, and diagnostic data collection suggesting assumptions over a possible causal correlation.

Keywords: COVID-19, mRNA-1273, mRNA BNT162b2, pemphigus foliaceus, pemphigus vulgaris

1. INTRODUCTION

Go to:

Pemphigus is a group of blistering disorders characterized by acantholysis, resulting in the formation of intraepithelial blisters in mucous membranes and skin. The process of acantholysis is induced by the binding of circulating autoantibodies to intercellular adhesion molecules (mainly desmoglein 1 and desmoglein 3). A few cases of new-onset pemphigus following coronavirus disease 2019 (COVID-19) vaccination have already been published. Herein, we present five cases, summarized in Tables 1 and 2, of new-onset pemphigus following vaccinations against COVID-19. Specifically, we describe two cases of pemphigus foliaceus (PF) and three cases of pemphigus vulgaris (PV).

TABLE 1 Demographics and clinical features of reported cases

Table with 9 columns: Case, Age/sex, Type of pemphigus, Latency, COVID-19 vaccine before clinical onset, Previous COVID-19 vaccination, COVID-19 vaccine after clinical onset, Anamnesis, Concomitant medication. Contains 5 rows of patient data.

Open in a separate window

Abbreviations: COVID-19, coronavirus disease 2019; mRNA, messenger RNA; PF, pemphigus foliaceus; PV, pemphigus vulgaris.

TABLE 2 Immunopathological features of reported cases

Table with 6 columns: Case, Histopathology, DIF, IIF, ELISA, Diagnosis. Contains 5 rows of immunopathological data.

Open in a separate window

Abbreviations: DIF, direct immunofluorescence; DSG1, desmoglein 1; DSG3, desmoglein 3; ELISA, enzyme-linked immunosorbent assay; IIF, indirect immunofluorescence; n.v., normal value; PF, pemphigus foliaceus; PV, pemphigus vulgaris.

2. CASE REPORT

Go to:

2.1. Case 1

A 61-year-old White woman presented to our institution for the appearance of a cutaneous rash 3 days after the COVID-19 vaccination booster dose with the messenger RNA (mRNA) BNT162b2 (Comirnaty, Pfizer-BioNTech) vaccine. She had received two previous doses of mRNA-1273 (Moderna) vaccine 8 and 6 months earlier, with both being well tolerated. The patient had arterial hypertension and undifferentiated connective tissue disease for which she was under chronic therapy with prednisone 2.5 mg/d, acetylsalicylic acid 100 mg/d, amlodipine 5 mg/d, and methotrexate 15 mg/wk (temporarily discontinued for vaccine injections). The patient denied any history of skin disease prior to vaccination. Dermatological examination revealed a few scattered superficial blisters and erythematous patches with scaly crusts involving the face and lower trunk (Figure 1a-c). No oral lesions were seen. Histopathological examination showed subepithelial intraepithelial cleavage with acantholysis. Direct immunofluorescence (DIF) demonstrated intercellular deposition of IgG and C3. Indirect immunofluorescence (IIF) showed intercellular IgG deposits. Enzyme-linked immunosorbent assay (ELISA) was positive for anti-desmoglein 1 (anti-DSG1) (106 IU/mL) and anti-desmoglein 3 (anti-DSG3) antibodies (15.5 IU/mL). A diagnosis of PV was made.




FIGURE 1 Clinical pictures of our cases. Case 1: scattered superficial blisters and erythematous patches with scaly crusts involving the face and lower trunk (a–c). Case 2: diffuse erythematous-squamous patches with scaly and crusted erosions involving the face and the trunk, following a seborrheic distribution (d, e). Case 3: a few erosive lesions, diffuse scales, and crusted erosions involving the trunk (f, g). Case 4: painful erosions on the gums and soft palate (h). Case 5: erosive lesions on the oral cavity, nose, right cheek, and abdomen (i, j).

2.2. Case 2

An 80-year-old White man developed an acute cutaneous eruption 17 days after his booster dose with the BNT162b2 vaccine. The patient had his first two doses 8 months (21 days apart) before the booster, which were both well tolerated. The patient had arterial hypertension and gastroesophageal reflux, for which he was under chronic therapy with amloride, hydrochlorothiazide, and esomeprazole. Physical examination revealed diffuse erythematous-squamous patches with scaly and crusted erosions involving the face and the trunk, following a seborrheic distribution (Figure 1d,e). Histopathological examination revealed subcorneal acantholysis with neutrophilic infiltration within the blister. DIF was negative, while IIF showed intercellular IgG deposits. ELISA was positive for anti-DSG1 antibodies (149 IU/mL), allowing for the diagnosis of PF.

2.3. Case 3

A 66-year-old White woman complained of a cutaneous eruption that appeared 4 weeks after her second dose of BNT162b2 vaccine. She had arterial hypertension, chronic coronary artery disease, gastroesophageal reflux, and hypercholesterolemia, for which she was taking treatment with rabeprazole, ticlopidine, atorvastatin, amlodipine, and hydrochlorothiazide. Physical examination revealed few erosive lesions, diffuse scales, and crusted erosions involving the trunk (Figure 1f,g). Histopathological examination revealed subcorneal acantholysis with neutrophils within the blister. DIF demonstrated intercellular deposition of IgG. IIF and ELISA were both negative. A diagnosis of PF was made.

2.4. Case 4

A 73-year-old White woman came to our attention for severe oral pain that occurred about 1 month after the booster dose of the BNT162b2 vaccine. The patient had received her first two doses of BNT162b2 vaccine 6 and 5 months earlier, which were both well tolerated. The patient had osteoporosis and senile dementia, for which he was under chronic therapy with alendronate, vitamin D supplementation, and haloperidol. Physical examination revealed painful erosions on the gums and soft palate (Figure 1h). No cutaneous lesions were observed. Histopathological examination showed erosive mucositis with diffuse acantholytic cells. DIF showed intercellular deposition of IgG and C3. IIF showed intercellular IgG deposits. ELISA was positive for anti-DSG3 antibodies (70 IU/mL). A diagnosis of PV was made.

2.5. Case 5

A 63-year-old White woman presented with painful oral erosions involving mainly the oral cavity and the skin of the face and trunk. Patients' medical history was unremarkable, except for arterial hypertension controlled with systemic therapy. She complained of three small erosions that appeared about 4 weeks after the first dose of Vaxzevria (ChAdOx1) nCoV-19 vaccine (Oxford-AstraZeneca), localized respectively on the floor of the mouth, soft palate, and right side of the buccal mucosa. Successively, she received the second vaccine dose 12 weeks after the first, experiencing a worsening of the disease. In fact, 4 days after vaccine injection, new erosions appeared on the floor of the mouth, tongue, and gums, while cutaneous erosions affected the nose, right cheek, and abdomen (Figure 1i,j). The patient therefore underwent a skin biopsy, which revealed acantholysis on histology and intercellular deposition of IgG and C3 on DIF. Serological investigation supported the diagnosis of PV showing intercellular deposition of IgG and positive staining for anti-DSG1 and anti-DSG3. ELISA was positive for anti-DSG1 and anti-DSG3 antibodies, with titers of 62.2 IU/mL and 157.3 IU/mL, respectively.

3. DISCUSSION

Go to:

Several cases of new-onset pemphigus following the anti-COVID-19 vaccine have already been published. A review by Calabria et al reported cases of severe autoimmune blistering diseases (AIBDs) associated with COVID-19 vaccination: they found six cases of PV and one case of PF. Coherently with our report, the median age of the cases of the pemphigus group was 60 years (interquartile range, 50–76 years) with no sex predilection (four women, three men). Among all of the cases of AIBDs, the majority (22 [62.9%]) developed after BNT162b2 vaccine administration, six (17.1%) after mRNA-1273 administration, three (8.6%) after ChAdOx1 administration, and three (8.6%) after CoronaVac administration.<sup>1</sup> According to Calabria et al's study and other studies in the literature, the Comirnaty vaccine seems to present the higher rate of association with onset of AIBDs, but this may probably be explained by the wider use of the BNT162b2 vaccine compared with the others in Western populations.<sup>2</sup> In a recent paper describing autoimmune skin disease flares following COVID-19 vaccine, full immunization with mRNA-1273 showed a higher likelihood of association with worsening of preexistent disease than BNT162b2, but the difference was not significant.<sup>3</sup>

The pathogenesis of autoimmune disorders following antiviral vaccinations is still under debate. One of the suggested mechanisms is immune cross-reactivity caused by molecular mimicry.<sup>4,5</sup> In fact, this phenomenon was advocated to explain several cases of pemphigus developed following exposure to bacterial and viral antigens, as well as pharmaceutical agents.<sup>6</sup> However, Kasperkiewicz et al<sup>7</sup> failed to find cross-reactivity between circulating anti-severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) antibodies and pemphigus or pemphigoid autoantigens in a 24-patient series. A relationship between autoimmune bullous diseases and COVID-19 vaccination has been hypothesized by other authors,<sup>8</sup> suggesting a transient immune activation in predisposed individuals with subclinical autoimmunity. The hypothesis of a nonspecific, nonselective mechanism of action may better explain the wide variety of immune-mediated diseases described following COVID-19 immunization, ranging from autoinflammatory to mixed or autoimmune manifestations, which may both be multisystemic or organ-specific such as pemphigus. Moreover, the onset of immune-mediated diseases seems to occur more frequently in individuals with a preexisting rheumatic or autoimmune background, thus supporting the importance of individual predisposition.<sup>9</sup> The role of personal susceptibility is even more supported by the fact that autoimmune bullous diseases with higher incidence in their idiopathic form, such as bullous pemphigoid, also seem to be more frequently triggered or boosted by COVID-19 vaccines.<sup>10</sup> In addition, Case 1 suggests that chronic immunosuppressive therapies prescribed for other concomitant diseases might maintain the autoimmune bullous disorder in a latent or subclinical stage until the balance is broken by strong precipitating factors such as vaccinations.

An innate inflammatory process is triggered by both BNT162b2 and ChAdOx1 vaccines caused by adjuvant function of mRNA itself and adenoviral proteins, respectively, which binds to different toll-like receptors, enhancing the production of type I interferon and several proinflammatory cytokines.<sup>11</sup> Type I interferon is well-known for its role in stimulating antigen presentation, B-cell differentiation, and IgG secretion.<sup>12,13,14</sup> However, a clear link between the vaccine's mechanism of action and pemphigus development is still far from being elucidated. In our case series, the timing of pemphigus onset is the only element that may suggest a causal relationship. The presence of a correlation between these events should be supported by larger population-based studies investigating an incidence variation of pemphigus and other autoimmune bullous diseases before and after large-scale vaccine administration. Until those kinds of studies are available in the literature, physicians should be aware of the possibility of autoimmune blistering diseases following SARS-CoV-2 vaccine, with the possibility of a causal relationship.

CONFLICT OF INTEREST

Go to:

The authors declare no conflicts of interest.

ACKNOWLEDGMENT

Go to:

The patients in this manuscript have given written informed consent to the publication of their case details. Open Access Funding provided by Università degli Studi di Firenze within the CRUI-CARE Agreement. Open Access Funding provided by Università degli Studi di Firenze within the CRUI-CARE Agreement.

Notes

Go to:

Corrá A, Barei F, Genovese G, Zussino M, Spigariolo CB & Mariotti EB et al. Five cases of new-onset pemphigus following vaccinations against coronavirus disease 2019. J Dermatol. 2022;00:1–5. 10.1111/1346-8138.16554 [PMC free article] [PubMed] [CrossRef] [Google Scholar]

Alberto Corrá and Francesca Barei contributed equally to this work.

REFERENCES

Go to:

1. Calabria E, Canfora F, Mascolo M, Varricchio S, Mignogna MD, Adamo D. Autoimmune mucocutaneous blistering diseases after SARS-Cov-2 vaccination: a case report of pemphigus vulgaris and a literature review. Pathol Res Pract. 2022;232:153834. [PMC free article] [PubMed] [Google Scholar]
2. Avallone G, Quaglino P, Cavallo F, Rocuzzo G, Riberio S, Zalaudek I, et al. SARS-CoV-2 vaccine-related cutaneous manifestations: a systematic review. Int J Dermatol. 2022. 10.1111/ijd.16063 [PMC free article] [PubMed] [CrossRef] [Google Scholar]
3. Sprow G, Afarideh M, Dan J, Feng R, Keyes E, Grinnell M, et al. Autoimmune skin disease exacerbations following COVID-19 vaccination. Front Immunol. 2022;13:899526. [PMC free article] [PubMed] [Google Scholar]
4. Segal Y, Shoenfeld Y. Vaccine-induced autoimmunity: the role of molecular mimicry and immune crossreaction. Cell Mol Immunol. 2018;15:586–94. [PMC free article] [PubMed] [Google Scholar]
5. Talotta R. Do COVID-19 RNA-based vaccines put at risk of immune-mediated diseases? In reply to "potential antigenic cross-reactivity between SARS-CoV-2 and human tissue with a possible link to an increase in autoimmune diseases". Clin Immunol. 2021;224:108665. [PMC free article] [PubMed] [Google Scholar]
6. Tchernev G, Orfanos CE. Antigen mimicry, epitope spreading and the pathogenesis of pemphigus. Tissue Antigens. 2006;68:280–6. [PubMed] [Google Scholar]
7. Kasperkiewicz M, Bednarek M, Tukaj S. Case report: circulating anti-SARS-CoV-2 antibodies do not cross-react with pemphigus or pemphigoid autoantigens. Front Med. 2021;8:807711. [PMC free article] [PubMed] [Google Scholar]
8. Tomayko MM, Damsky W, Fathy R, McMahon DE, Turner N, Valentini MN, et al. Subepidermal blistering eruptions, including bullous pemphigoid, following COVID-19 vaccination. J Allergy Clin Immunol. 2021;148:750–1. [PMC free article] [PubMed] [Google Scholar]
9. Watad A, De Marco G, Mahajna H, Druyan A, Elitty M, Hijazi N, et al. Immune-mediated disease flares or new-onset disease in 27 subjects following mRNA/DNA SARS-CoV-2 vaccination. Vaccines. 2021;9:435. [PMC free article] [PubMed] [Google Scholar]
10. Maronese CA, Caproni M, Moltrasio C, Genovese G, Vezzoli P, Sena P, et al. Bullous pemphigoid associated with COVID-19 vaccines: an Italian multicentre study. Front Med. 2022;9:841506. [PMC free article] [PubMed] [Google Scholar]
11. Teijaro JR, Farber DL. COVID-19 vaccines: modes of immune activation and future challenges. Nat Rev Immunol. 2021;21:195–7. [PMC free article] [PubMed] [Google Scholar]
12. Elkon K, Casali P. Nature and functions of autoantibodies. Nat Clin Pract Rheumatol. 2008;4:491–8. [PMC free article] [PubMed] [Google Scholar]
13. Anderson EI, Rouphael NG, Widge AT, Jackson LA, Roberts PC, Makhene M, et al. Safety and immunogenicity of SARS-CoV-2 mRNA-1273 vaccine in older adults. N Engl J Med. 2020;383:2427–38. [PMC free article] [PubMed] [Google Scholar]
14. Sahin U, Muik A, Derhovanessian E, Vogler I, Kranz LM, Vormehr M, et al. COVID-19 vaccine BNT162b1 elicits human antibody and TH1 T cell responses. Nature. 2020;586:594–9. Erratum in: Nature. 2021;590:E17. [PubMed] [Google Scholar]