



Carbonic anhydrase IX as a novel candidate in liquid biopsy

Ozen Ozensoy Guler, Claudiu. T. Supuran & Clemente Capasso

To cite this article: Ozen Ozensoy Guler, Claudiu. T. Supuran & Clemente Capasso (2020) Carbonic anhydrase IX as a novel candidate in liquid biopsy, Journal of Enzyme Inhibition and Medicinal Chemistry, 35:1, 255-260, DOI: [10.1080/14756366.2019.1697251](https://doi.org/10.1080/14756366.2019.1697251)

To link to this article: <https://doi.org/10.1080/14756366.2019.1697251>



© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 02 Dec 2019.



Submit your article to this journal [↗](#)



Article views: 2778



View related articles [↗](#)



View Crossmark data [↗](#)





Citing articles: 12 View citing articles [↗](#)

REVIEW ARTICLE



Carbonic anhydrase IX as a novel candidate in liquid biopsy

Ozen Ozensoy Guler^a, Claudiu. T. Supuran^b  and Clemente Capasso^c 

^aDepartment of Medical Biology, Faculty of Medicine, Yildirim Beyazit University, Ankara, Turkey; ^bDepartment of NEUROFARBA, Section of Pharmaceutical and Nutraceutical Sciences, University of Florence, Polo Scientifico, Firenze, Italy; ^cDepartment of Biology, Agriculture and Food Sciences, Institute of Biosciences and Bioresources, CNR, Napoli, Italy

ABSTRACT

Among the diagnostic techniques for the identification of tumour biomarkers, the liquid biopsy is considered one that offers future research on precision diagnosis and treatment of tumours in a non-invasive manner. The approach consists of isolating tumor-derived components, such as circulating tumour cells (CTC), tumour cell-free DNA (ctDNA), and extracellular vesicles (EVs), from the patient peripheral blood fluids. These elements constitute a source of genomic and proteomic information for cancer treatment. Within the tumour-derived components of the body fluids, the enzyme indicated with the acronym CA IX and belonging to the superfamily of carbonic anhydrases (CA, EC 4.2.1.1) is a promising aspirant for checking tumours. CA IX is a transmembrane-CA isoform that is strongly overexpressed in many cancers being not much diffused in healthy tissues except the gastrointestinal tract. Here, it is summarised the role of CA IX as tumour-associated protein and its putative relationship in liquid biopsy for diagnosing and monitoring cancer progression.

ARTICLE HISTORY

Received 5 November 2019
Revised 13 November 2019
Accepted 16 November 2019

KEYWORDS

Liquid biopsy; cancer; carbonic anhydrase IX; tumour markers; acidification

1. Introduction



1.1. Carbonic anhydrases (CAs)

A crucial physiological reaction for the survival of all living organisms is the pivotal CO₂ hydration/dehydration of the central metabolism. This reaction is connected with numerous metabolic pathways, such as photosynthesis and carboxylation reactions, and biochemical pathways including pH homeostasis, secretion of electrolytes, transport of CO₂ and bicarbonate, and so on^{1,2}. Moreover, the interconversion of CO₂ and HCO₃⁻ is spontaneously and precisely balanced form the living organisms to maintain the equilibrium between dissolved inorganic carbon dioxide (CO₂), carbonic acid (H₂CO₃), bicarbonate (HCO₃⁻) and carbonate (CO₃²⁻)³⁻⁶. The CO₂ hydration/dehydration is catalysed by a superfamily of metalloenzymes, known as carbonic anhydrases (CAs, EC 4.2.1.1)⁷⁻¹¹, which are categorised into eight genetically distinct families (or classes), named with the Greek letters: α , β , γ , δ , ζ , η , θ , and ι . The last three classes were recently discovered¹²⁻¹⁴. Moreover, members of each class possess multiple transcript variants and protein isoforms, which are characterised by different biochemical properties and have specific tissue/organ and sub-cellular localizations^{11,15-20}. CAs present in animals belong to α -class^{21,22}, plants and algae have α -, β -, γ -, δ - θ - and ι -classes; fungi encode for α - and β -CAs; protozoa for α -, β - and/or η -CAs; bacteria for α -, β -, γ - and ι -CA classes^{11,20,23-27}. Besides, a matrix protein called nacrein has been identified in the oyster *Pinctada fucata*. It participates in the formation of the nacreous layer and is characterised by a CA domain present at the N-terminus part of the polypeptide sequence²⁸. In mammals, 16 α -CA isoforms have been identified: five of them are cytosolic (CA I, CA II, CA III, CA VII, and CA XIII), five are membrane-bound (CA IV, CA IX, CA XII,

CA XIV, and CA XV), two are mitochondrial (CA VA and CA VB), and only one is secreted (CA VI), the last three (CA VIII, CA X, and CA XI) being devoid of catalytic activity and referred to as CA Related Proteins (CARPs)^{8,29,30}. CA active site includes a zinc ion (Zn²⁺), which plays a critical role in the catalytic enzyme function. In addition to the zinc, ζ - and γ -CAs reflect exceptions to this principle since they can use cadmium (ζ), iron (γ), or cobalt (γ)^{31,32}.

1.2. Tumour-associated CA IX

The glycolytic metabolism of cancer was evaluated for so many years to describe the fundamental role of tumour microenvironment and glycolysis in cancer growth and progression³³. The transcription factors of the glycolytic pathway affect cell proliferation, which is an essential feature of carcinogenesis³⁴. Different types of enzymes are produced by tumours or by the body in response to malignancy and used as cancer biomarkers³⁵. It has been shown that the expression levels of certain enzymes can vary in various types of cancer³⁵. CA IX is a transmembrane CA isoform expressed in healthy tissues. Figure 1 shows the catalytically active CA IX on the cellular membrane surface³⁶⁻³⁸. CA IX is highly overexpressed in many types of cancer^{39,40}. For example, its expression is increased considerably in solid tumours of uterus, kidney, lung, colon, breast, brain, and ovary⁴¹⁻⁴⁴. Tumour cells decrease their extracellular pH by lactic acid production and CO₂ hydration, which is catalysed by CAs (Figure 2). Since the tumour-associated CA IX is an efficient catalyst for the conversion of CO₂ in bicarbonate and protons, they contribute to the acidification of the tumour environment. Moreover, its activity leads to the acquisition of

CONTACT Clemente Capasso  clemente.capasso@ibbr.cnr.it  Department of Biology, Agriculture and Food Sciences, Institute of Biosciences and Bioresources, CNR, Via Pietro Castellino, 111 – 80131, Napoli, Italy

© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Structure of CA IX isoenzyme. (PG: Proteoglycan domain; TM: Transmembrane domain; IC: Intracellular domain).




Figure 2. pH regulation of carbonic anhydrase enzymes in tumour cells.

metastatic phenotypes and chemoresistance to weakly basic anti-cancer drugs²¹.

Interestingly, CA IX has a metastatic activity related to the extracellular acidity since it has been shown to promote migration and invasion in tumour cells^{45,46}. Furthermore, oxygen has a crucial function in the regulation of redox balance and energy production in tumour tissues. An inadequate amount of oxygen in the tissues causes hypoxia, which is a characteristic marker of the tumour microenvironment. Hypoxia regulates the expression of many genes inducing a phenotypic alteration of stromal cells in the tumour microenvironment and promoting the survival of cancer cells. Low oxygen activates the HIF-1, the hypoxia-inducible factor 1 (HIF-1)^{47,48}, which starts the transcription of several hypoxia-inducible genes, such as (Vascular Endothelial Growth Factor (VEGF), Glucose Transporter 1 (GLUT1), CA IX and CA XII)⁴⁹. CA IX is one of the most potent hypoxia-induced proteins, and the hypoxia-inducible proteins are important anti-cancer targets⁵⁰. As a result, it is readily apparent that CA IX is associated with many tumours and are involved in the cancer process³⁴. The specific inhibition of CA IX activity with selective inhibitors, such as sulphonamide derivatives, represents a good strategy for establishing the driving role of this isoform, as well as other CAs, for example, CA XII, in tumorigenesis⁵¹. Many CA inhibitors exist, which could be classified as inhibitors binding the metal ion, inhibitors anchoring to the water molecule/hydroxide ion coordinated to the metal, inhibitors occluding the active site entrance and inhibitors linking out of the active site⁵². Many of these inhibitors can be used to reduce the proliferation and invasion capacity of cancer cells⁵³⁻⁵⁵.

Since pH-related cancer growth and metastasis might be dependent on the enzyme activity of the CA IX isoform, the identification of CA IX as tumour biomarker in the liquid biopsy is crucial for a new approach concerning the treatment of malignancy. The role of cancer markers in carcinogenesis is pivotal for early diagnosis, promotion of suitable procedures, and leading right management strategies⁵⁶.

2. Tumour-derived components in liquid biopsy

Recently, liquid biopsy has begun an exciting approach in terms of early detection of tumor^{57,58}. The technique, starting from the blood or body fluids, isolates tumour-derived components, such as circulating tumour cells (CTCs), circulating tumour DNA (ctDNA), and extracellular vesicles (EVs), as a source of genomic and proteomic information in patients with cancer⁵⁷. Liquid biopsy is an appropriate technique for correct treatments and figuring out the genetic changes in the tumour. It can be applied to many types of cancer, analysing the blood or body fluids of the patient taken in a non-invasive manner compared to the biopsy and without even detecting the symptoms⁵⁹.

2.1. Circulating tumour cells (CTCs)

Thomas Ashworth investigated CTCs in the 1860s, considering that tumour cells could have crossed the vessel wall and enter the bloodstream⁶⁰. These cells are released from primary or metastatic cells. A few steps are required for the metastatic process. First, CTCs are separated from the tumour and incorporated into the bloodstream. The circulating CTCs are then protected from the immune cells to perform extravasation. After that, these cells adapt to the microenvironment of the new tissue and to form metastatic lesions. CTCs are found in circulation as single CTCs or CTC clusters. In the literature, it has been reported that tumour progression and patient survival is correlated with the numbers of CTCs⁶¹. The cont of CTCs is also useful in the treatment response process. The technics known as immunophenotypic identification of cytokeratin, enzymatic methods, and RT-qPCR (reverse transcriptase quantitative polymerase chain reaction) are used to detect CTCs⁶²⁻⁶⁶. The cell surface glycoprotein, EPCAM (Epithelial Cell Adhesion Molecule), which is highly expressed in epithelial cancer cells, serves as the primary antigen for CTCs detection. Moreover, with this method, prognostic information can be obtained in metastatic breast, colon, and prostate cancers. CTCs are subjected to genomic mapping⁶²⁻⁶⁶, allowing a detailed analysis of the genes responsible for the uncontrolled growth of cells. Of course, the major drawbacks of this technique are (i) to distinguish cancer cells from millions of healthy cells; (ii) the detection of the kind of cancer.

2.2. Circulating tumour DNA (ctDNA)

The presence of ctDNA in the blood is dated back in 1948⁶⁷. Although circulating cell-tumour DNA was first identified in 1948, it has only recently been investigated in "liquid biopsy" as cancer biomarkers⁶⁸. Tumours release fragments of DNA into the circulatory system, which are detectable and specific to cancer. There are several advantages to assessing ctDNA. Sampling is non-invasive and inexpensive compared to the tissue biopsy. Besides, ctDNA testing can be easily and frequently repeated to monitor changes that occur during treatment, serving as an early indicator of recurrence, resistance, or metastasis⁶⁹. Liquid biopsy is believed

Table 1. Comparison of CTCs and ctDNA.

Comparison	CTCs	ctDNA
Origin	Intact cells	Released from apoptotic and necrotic cells
Description	Originates from primary tumours or metastasis	DNA fragments in blood circulation
Detection techniques	Density/size based immunomagnetic and microfluidic techniques	PCR or sequencing based
Advantages	<ul style="list-style-type: none"> • Allows for DNA, RNA, protein research • Research at single-cell level • Captured cells can be used for <i>in vitro</i> or <i>in vivo</i> studies • Clinically validated technology available (CellSearch System) 	<ul style="list-style-type: none"> • Easily isolated with kits • Can be stored for a long time • Gives more precise results • Clinically validated for EGFR mutations in non-small cell lung cancer
Disadvantages	<ul style="list-style-type: none"> • A small number of cells are obtained in non-metastatic conditions • Can not be stored for a long time • Detection steps are expensive 	<ul style="list-style-type: none"> • Prognostics and predictability are unclear • Only DNA sequence analysis can be performed • Known target mutations are needed

to exhibit tumour heterogeneity, as cells circulate from different regions of the tumour, and to obtain information in a shorter time⁶⁹. CTCs and ctDNAs derived from primary or metastatic cells are abundant in blood. Both CTC and ctDNA provide prognostic information based on the number and level of events detected^{70–72} (Table 1).

2.3. Extracellular vesicles (EVs)

Extracellular Vesicles (EVs) are lipid structures released from cells. EVs contain proteins and nucleic acids and play a role in cellular communication, immune regulation, and microenvironmental modulation^{73,74}. Nanosized exosomes (70–150 nm) are the most prominent members of these so-called extracellular vesicles (EVs) and are released from body fluids such as urine, ascites, and plasma^{73,74}. Moreover, they are liberated over all kinds of body cells (epithelium cells, haematopoietic cells, adipocytes, healthy and malignant cells)⁷⁵, released in almost all cell types under physiological and pathophysiological conditions and mediate intercellular contacts⁷⁶. A theranostic solution could be represented by the nanosized EVs, which may transmit biomarkers of diseases and/or vectors of therapeutic molecules, offering a unique opportunity to use a combination of different markers specifically expressed for tumour-derived EVs^{74,76,77}. For example, Prostate-Specific Antigen (PSA) does not differentiate between benign prostatic hyperplasia (BPH) and a Prostate Cancer (PC), resulting in large numbers of unnecessary biopsies and missed diagnosis of cancer. Since exosomes are directly detectable in patient plasma, the plasmatic exosomes expressing PSA have the potential in distinguishing healthy individuals, BPH, and PC⁷⁷. Recently, it has been demonstrated that neurodegenerative disorders, including Alzheimer's disease (AD), Parkinson, and amyotrophic lateral sclerosis, are correlated with extracellular vesicles. EVs have also been investigated in relation to infection caused by viruses, bacteria, fungi, protozoa, and helminths⁷⁶. Such pathogens secrete EVs, and prions were even present in EVs. Finally, EVs seem to play key roles in autoimmune diseases⁷⁶. EVs circulating in body fluids are valuable liquid biopsy biomarkers. Additionally, their protein concentration is higher in patients with advanced tumours⁷³.

3. Tumour-associated CA IX as biomarker in liquid biopsy

The tumour-associated CA IX may be used as a cancer biomarker in the liquid biopsy technique. Carbonic anhydrase IX is a transmembrane enzyme⁷⁸, and it is involved in the growth and development of tumour cell adhesion^{79,80}. There is an association between elevated serum levels of CA IX and CTCs⁴⁷. This suggests a relationship between hypoxia and tumour cell circulation in the

bloodstream. In the peripheral venous blood, it is also possible to find the soluble form of CA IX, which is released by proteolytic cleavage^{47,50,81}. For example, a high level of soluble CA IX was found in the serum of patients with renal cancer⁵⁰. Müller et al. investigated the relationship between serum levels of CA IX and CTCs in metastatic breast cancer⁴⁷. Their findings suggested that the CA IX activity level was higher in cancer types with a high number of CTCs. In this condition, it is expected a decrease in the patient's overall survival. Besides, CA IX disrupts cell-cell and cell-matrix interactions by triggering tumour acidification. As a consequence, CTCs separate from the primary tumour, and invasion occurs. Probably, CA IX inhibition could slow down the invasion process of CTCs.

During the tumour progression, exosomes and the metalloenzyme CA IX affect the growth and proliferation of the tumour. The relationship between exosomes and CA IX has been investigated using an *in vitro* cellular model of human prostate carcinoma cell line cultured in different pH conditions. The results showed that the acidic microenvironment increased both the expression and activity of CA IX in cancer cells⁸². Besides, the number of exosomes released by the cancer cells was raised together with the upregulation of the CA IX⁸². These data strongly support that exosomes and CA IX are tumour-associated components and the enzyme CA IX is a cancer biomarker that could be used as valuable target of the liquid biopsy⁸². Horie et al. demonstrated that CA IX exosomes were released from renal carcinoma cells⁴⁹. The quantity of exosomal CA IX is increased in hypoxia response, promoting upregulation of MMP-2, migration and tube formation, and may induce angiogenesis in the tumour microenvironment⁴⁹. Dorai and co-workers analysed the effect of increasing expression levels of CA IX in renal cancer cells. They showed that the level of released gangliocytes was positively correlated to that of exosomal CA IX expression⁸³. Gangliosides play a critical role in cell adhesion, migration, and cell signalling⁸⁴. Since CA IX induces the release of gangliocyte-containing exosomes, the exosomal CA IX may be a valuable biomarker of the carcinogenesis process.

CA IX expression is regulated exclusively by HIF-1 α , rapidly increases in response to hypoxia, and is very important for maintaining the acidic pH of the tumor⁸⁵. Brown-Glaberman proposed that the circulating CA IX could be considered as a biomarker for detecting the level of hypoxia and the upregulation of HIF-1 α ^{86,87}. Circulating CA IX can be easily isolated from body fluids and considered a biomarker for different stages of cancers and the differentiation of local/advanced tumours. Malentacchi and coworkers to validate circulating CA IX as a tumour biomarker measuring the CA IX mRNA in the urine sediments of patients affected by kidney, prostate, and bladder cancers⁸⁸. As a result, they associated the mRNA CA IX expression in the tumour of urogenital origin. Liu et al. have found that the combination of the CA IX/CD147 antibodies achieved higher efficiency in the NanoVelcro platform

compared to EPCAM-based methods for capturing circulating cells coming from the renal carcinoma⁸⁹.

4. Conclusion

Liquid biopsy technology allows the detection of solid tumours, such as those involving lung, breast, and pancreatic, using the blood or other body fluids. Liquid biopsy can detect cancer-specific markers even in lesions that are too small to be recognised by other available methods, indicating that this method can be used early in cancer diagnosis. Among the known biomarkers of the liquid biopsy, the CA IX isoenzyme could be a promising candidate for tumour detection. In fact, the CA IX activity level is higher in cancer types with a high number of CTCs; the quantity of exosomal CA IX is increased in hypoxia response; the release of the exosomal ganglioside is correlated to the exosomal CA IX expression increase; the circulating CA IX is associated to tumours of urogenital origin; CA IX disrupts cell-cell and cell-matrix interactions by triggering tumour acidification. In this context, the tumour-associated CA IX could be considered a valid biomarker of the non-invasive liquid biopsy, which is viewed as a technique that offers future research on precision diagnosis and treatment of tumours in a non-invasive manner. Moreover, CA IX could be a valid molecular target in the treatment of cancer.

Disclosure statement

No potential conflict of interest was reported by the authors.

ORCID

Claudiu. T. Supuran  <http://orcid.org/0000-0003-4262-0323>
Clemente Capasso  <http://orcid.org/0000-0003-3314-2411>

References

- Johnson X, Alric J. Interaction between starch breakdown, acetate assimilation, and photosynthetic cyclic electron flow in *Chlamydomonas reinhardtii*. *J Biol Chem* 2012;287:26445–52.
- Tcherkez G, Boex-Fontvieille E, Mahe A. Respiratory carbon fluxes in leaves. *Curr Opin Plant Biol* 2012;15:308–14.
- Smith KS, Ferry JG. Prokaryotic carbonic anhydrases. *FEMS Microbiol Rev* 2000;24:335–66.
- Maeda S, Price GD, Badger MR, et al. Bicarbonate binding activity of the CmpA protein of the cyanobacterium *Synechococcus* sp. strain PCC 7942 involved in active transport of bicarbonate. *J Biol Chem* 2000;275:20551–5.
- Joseph P, Ouahrani-Bettache S, Montero JL, et al. A new beta-carbonic anhydrase from *Brucella suis*, its cloning, characterization, and inhibition with sulfonamides and sulfamates, leading to impaired pathogen growth. *Bioorg Med Chem* 2011;19:1172–8.
- Joseph P, Turtaut F, Ouahrani-Bettache S, et al. Cloning, characterization, and inhibition studies of a beta-carbonic anhydrase from *Brucella suis*. *J Med Chem* 2010;53:2277–85.
- Annunziato G, Angeli A, D'Alba F, et al. Discovery of new potential anti-infective compounds based on carbonic anhydrase inhibitors by rational target-focused repurposing approaches. *ChemMedChem* 2016;11:1904–14.
- Ozensoy Guler O, Capasso C, Supuran CT. A magnificent enzyme superfamily: carbonic anhydrases, their purification and characterization. *J Enzyme Inhib Med Chem* 2016;31:689–94.
- Del Prete S, Vullo D, De Luca V, et al. Sulfonamide inhibition studies of the beta-carbonic anhydrase from the pathogenic bacterium *Vibrio cholerae*. *Bioorg Med Chem* 2016;24:1115–20.
- Del Prete S, De Luca V, De Simone G, et al. Cloning, expression and purification of the complete domain of the eta-carbonic anhydrase from *Plasmodium falciparum*. *J Enzyme Inhib Med Chem* 2016;31:54–9.
- Capasso C, Supuran CT. An overview of the carbonic anhydrases from two pathogens of the oral cavity: *Streptococcus mutans* and *Porphyromonas gingivalis*. *Curr Top Med Chem* 2016;16:2359–68.
- Jensen EL, Clement R, Kosta A, et al. A new widespread subclass of carbonic anhydrase in marine phytoplankton. *Isme J* 2019;13:2094–106.
- Kikutani S, Nakajima K, Nagasato C, et al. Thylakoid luminal theta-carbonic anhydrase critical for growth and photosynthesis in the marine diatom *Phaeodactylum tricorutum*. *Proc Natl Acad Sci USA* 2016;113:9828–33.
- Del Prete S, Vullo D, Fisher GM, et al. Discovery of a new family of carbonic anhydrases in the malaria pathogen *Plasmodium falciparum*—the eta-carbonic anhydrases. *Bioorg Med Chem Lett* 2014;24:4389–96.
- Supuran CT, Capasso C. Biomedical applications of prokaryotic carbonic anhydrases. *Expert Opin Ther Pat* 2018;28:745–54.
- Supuran CT. Carbonic anhydrase activators. *Future Med Chem* 2018;10:561–73.
- Supuran CT, Capasso C. An overview of the bacterial carbonic anhydrases. *Metabolites* 2017;7:56–74.
- Supuran CT, Capasso C. Carbonic anhydrase from *Porphyromonas gingivalis* as a drug target. *Pathogens* 2017;6:30–42.
- Supuran CT. Advances in structure-based drug discovery of carbonic anhydrase inhibitors. *Expert Opin Drug Discov* 2017;12:61–88.
- Capasso C, Supuran CT. Bacterial, fungal and protozoan carbonic anhydrases as drug targets. *Expert Opin Ther Targets* 2015;19:1689–704.
- Aspatwar A, Tolvanen ME, Ortutay C, et al. Carbonic anhydrase related proteins: molecular biology and evolution. *Subcell Biochem* 2014;75:135–56.
- Supuran CT. Carbonic anhydrases as drug targets—an overview. *Curr Top Med Chem* 2007;7:825–33.
- Supuran CT, Capasso C. The eta-class carbonic anhydrases as drug targets for antimalarial agents. *Expert Opin Ther Targets* 2015;19:551–63.
- Capasso C, Supuran CT. An overview of the selectivity and efficiency of the bacterial carbonic anhydrase inhibitors. *Curr Med Chem* 2015;22:2130–9.
- Capasso C, Supuran CT. An overview of the alpha-, beta- and gamma-carbonic anhydrases from bacteria: can bacterial carbonic anhydrases shed new light on evolution of bacteria? *J Enzyme Inhib Med Chem* 2015;30:325–32.
- Capasso C, Supuran CT. Sulfa and trimethoprim-like drugs – antimetabolites acting as carbonic anhydrase, dihydropterolate synthase and dihydrofolate reductase inhibitors. *J Enzyme Inhib Med Chem* 2014;29:379–87.
- Capasso C, Supuran CT. Anti-infective carbonic anhydrase inhibitors: a patent and literature review. *Expert Opin Ther Pat* 2013;23:693–704.

28. Miyamoto H, Miyashita T, Okushima M, et al. A carbonic anhydrase from the nacreous layer in oyster pearls. *Proc Natl Acad Sci USA* 1996;93:9657–60.
29. Supuran CT. Carbonic anhydrases—an overview. *Curr Pharm Des* 2008;14:603–14.
30. Supuran CT. Carbonic anhydrase inhibition and the management of hypoxic tumors. *Metabolites* 2017;7:48–60.
31. Lane TW, Saito MA, George GN, et al. Biochemistry: a cadmium enzyme from a marine diatom. *Nature* 2005;435:42.
32. Ferry JG. The gamma class of carbonic anhydrases. *Biochim Biophys Acta* 2010;1804:374–81.
33. Jang M, Kim SS, Lee J. Cancer cell metabolism: implications for therapeutic targets. *Exp Mol Med* 2013;45:e45.
34. Petrou A, Geronikaki A, Terzi E, et al. Inhibition of carbonic anhydrase isoforms I, II, IX and XII with secondary sulfonamides incorporating benzothiazole scaffolds. *J Enzyme Inhib Med Chem* 2016;31:1306–11.
35. Neri D, Supuran CT. Interfering with pH regulation in tumours as a therapeutic strategy. *Nat Rev Drug Discov* 2011;10:767–77.
36. Supuran CT, Di Fiore A, Alterio V, et al. Recent advances in structural studies of the carbonic anhydrase family: the crystal structure of human CA IX and CA XIII. *Curr Pharm Des* 2010;16:3246–54.
37. Hilvo M, Baranauskienė L, Salzano AM, et al. Biochemical characterization of CA IX, one of the most active carbonic anhydrase isozymes. *J Biol Chem* 2008;283:27799–809.
38. Di Fiore A, De Simone G, Menchise V, et al. Carbonic anhydrase inhibitors: X-ray crystal structure of a benzenesulfonamide strong CA II and CA IX inhibitor bearing a pentafluorophenylaminothioureido tail in complex with isozyme II. *Bioorg Med Chem Lett* 2005;15:1937–42.
39. Winum JY, Rami M, Scozzafava A, et al. Carbonic anhydrase IX: a new druggable target for the design of antitumor agents. *Med Res Rev* 2008;28:445–63.
40. Winum JY, Pastorekova S, Jakubickova L, et al. Carbonic anhydrase inhibitors: synthesis and inhibition of cytosolic/tumor-associated carbonic anhydrase isozymes I, II, and IX with bis-sulfamates. *Bioorg Med Chem Lett* 2005;15:579–84.
41. Eldehna WM, Abo-Ashour MF, Berrino E, et al. SLC-0111 enaminone analogs, 3/4-(3-aryl-3-oxopropenyl) aminobenzenesulfonamides, as novel selective subnanomolar inhibitors of the tumor-associated carbonic anhydrase isoform IX. *Bioorg Chem* 2019;83:549–58.
42. Mboge MY, Chen Z, Wolff A, et al. Selective inhibition of carbonic anhydrase IX over carbonic anhydrase XII in breast cancer cells using benzene sulfonamides: Disconnect between activity and growth inhibition. *PLoS One* 2018;13:e0207417.
43. Supuran CT. Carbonic anhydrase inhibitors as emerging agents for the treatment and imaging of hypoxic tumors. *Expert Opin Investig Drugs* 2018;27:963–70.
44. Eldehna WM, Nocentini A, Al-Rashood ST, et al. Tumor-associated carbonic anhydrase isoform IX and XII inhibitory properties of certain isatin-bearing sulfonamides endowed with *in vitro* antitumor activity towards colon cancer. *Bioorg Chem* 2018;81:425–32.
45. Liao SY, Lerman MI, Stanbridge EJ. Expression of transmembrane carbonic anhydrases, CAIX and CAXII, in human development. *BMC Dev Biol* 2009;9:22.
46. Supuran CT. Carbonic anhydrases: novel therapeutic applications for inhibitors and activators. *Nat Rev Drug Discov* 2008;7:168–81.
47. Muller V, Riethdorf S, Rack B, et al. Prospective evaluation of serum tissue inhibitor of metalloproteinase 1 and carbonic anhydrase IX in correlation to circulating tumor cells in patients with metastatic breast cancer. *Breast Cancer Res* 2011;13:R71.
48. Aspatwar A, Tolvanen ME, Ortutay C, et al. Carbonic anhydrase related protein VIII and its role in neurodegeneration and cancer. *Curr Pharm Des* 2010;16:3264–76.
49. Horie K, Kawakami K, Fujita Y, et al. Exosomes expressing carbonic anhydrase 9 promote angiogenesis. *Biochem Biophys Res Commun* 2017;492:356–61.
50. Zavada J, Zavadova Z, Zat'ovicova M, et al. Soluble form of carbonic anhydrase IX (CA IX) in the serum and urine of renal carcinoma patients. *Br J Cancer* 2003;89:1067–71.
51. Pan PW, Waheed A, Sly WS, et al. Carbonic anhydrases in the mouse harderian gland. *J Mol Histol* 2010;41:411–7.
52. Supuran CT. How many carbonic anhydrase inhibition mechanisms exist? *J Enzyme Inhib Med Chem* 2016;31:345–60.
53. Parkkila S, Rajaniemi H, Parkkila AK, et al. Carbonic anhydrase inhibitor suppresses invasion of renal cancer cells *in vitro*. *Proc Natl Acad Sci USA* 2000;97:2220–4.
54. Guler O, Simone G, Supuran C. Drug design studies of the novel antitumor targets carbonic anhydrase IX and XII. *Curr Med Chem* 2010;17:1516–26.
55. Monti SM, Supuran CT, De Simone G. Anticancer carbonic anhydrase inhibitors: a patent review (2008–2013). *Expert Opin Ther Pat* 2013;23:737–49.
56. Nishiumi S, Yoshida M. [Possibility of metabolite biomarkers for early detection of cancer]. *Gan to Kagaku Ryoho* 2018;45:894–8.
57. Han X, Wang J, Sun Y. Circulating tumor DNA as biomarkers for cancer detection. *Genomics Proteomics Bioinformatics* 2017;15:59–72.
58. Zhang W, Xia W, Lv Z, et al. Liquid biopsy for cancer: circulating tumor cells, circulating free DNA or exosomes? *Cell Physiol Biochem* 2017;41:755–68.
59. Offin M, Chabon JJ, Razavi P, et al. Capturing genomic evolution of lung cancers through liquid biopsy for circulating tumor DNA. *J Oncol* 2017;2017:1.
60. Ashworth TR. A case of cancer in which cells similar to those in the tumours were seen in the blood after death. *Austr Med J* 1869;14:146–7.
61. Sheng W, Ogunwobi OO, Chen T, et al. Capture, release and culture of circulating tumor cells from pancreatic cancer patients using an enhanced mixing chip. *Lab Chip* 2014;14:89–98.
62. Suo Y, Gu Z, Wei X. Advances of *in vivo* flow cytometry on cancer studies. *Cytometry A* 2019. [Epub ahead of print]. doi:10.1002/cyto.a.23851
63. Khetani S, Mohammadi M, Nezhad AS. Filter-based isolation, enrichment, and characterization of circulating tumor cells. *Biotechnol Bioeng* 2018;115:2504–29.
64. Werbin JL, Nordberg JJ, Tzucker J, et al. RareCyte® CTC analysis step 2: detection of circulating tumor cells by CyteFinder® automated scanning and semiautomated image analysis. *Methods Mol Biol* 2017;1634:173–80.
65. Hassan EM, Willmore WG, DeRosa MC. Aptamers: promising tools for the detection of circulating tumor cells. *Nucleic Acid Ther* 2016;26:335–47.
66. Ghossein RA, Bhattacharya S. Molecular detection and characterization of circulating tumor cells and micrometastases in prostatic, urothelial, and renal cell carcinomas. *Semin Surg Oncol* 2001;20:304–11.

67. Mandel P, Metais P. Les acides nucléiques du plasma sanguin chez l'homme. *C R Seances Soc Biol Fil* 1948;142:241–3.
68. Kanwar N, Hu P, Bedard P, et al. Identification of genomic signatures in circulating tumor cells from breast cancer. *Int J Cancer* 2015;137:332–44.
69. Li H, Jing C, Wu J, et al. Circulating tumor DNA detection: a potential tool for colorectal cancer management. *Oncol Lett* 2019;17:1409–16.
70. Heitzer E, Auer M, Ulz P, et al. Circulating tumor cells and DNA as liquid biopsies. *Genome Med* 2013;5:73.
71. Lim M, Kim CJ, Sunkara V, et al. Liquid biopsy in lung cancer: clinical applications of circulating biomarkers (CTCs and ctDNA). *Micromachines (Basel)* 2018;9. doi:10.3390/mi9030100
72. Neumann MHD, Bender S, Krahn T, et al. ctDNA and CTCs in liquid biopsy - current status and where we need to progress. *Comput Struct Biotechnol J* 2018;16:190–5.
73. Shang M, Ji JS, Song C, et al. Extracellular vesicles: a brief overview and its role in precision medicine. *Methods Mol Biol* 2017;1660:1–14.
74. Campanella C, Caruso Bavisotto C, Logozzi M, et al. On the choice of the extracellular vesicles for therapeutic purposes. *Int J Mol Sci* 2019;20. doi:10.3390/ijms20020236
75. Colombo M, Giannandrea D, Lesma E, et al. Extracellular vesicles enhance multiple myeloma metastatic dissemination. *Int J Mol Sci* 2019;20. doi: 10.3390/ijms20133236
76. Fais S, O'Driscoll L, Borrás FE, et al. Evidence-based clinical use of nanoscale extracellular vesicles in nanomedicine. *ACS Nano* 2016;10:3886–99.
77. Logozzi M, Angelini DF, Giuliani A, et al. Increased plasmatic levels of PSA-expressing exosomes distinguish prostate cancer patients from benign prostatic hyperplasia: a prospective study. *Cancers (Basel)* 2019;11. doi:10.3390/cancers11101449
78. De Simone G, Supuran CT. Carbonic anhydrase IX: Biochemical and crystallographic characterization of a novel antitumor target. *Biochim Biophys Acta* 2010;1804:404–9.
79. Svastova E, Witarski W, Csaderova L, et al. Carbonic anhydrase IX interacts with bicarbonate transporters in lamellipodia and increases cell migration via its catalytic domain. *J Biol Chem* 2012;287:3392–402.
80. Supuran CT, Alterio V, Di Fiore A, et al. Inhibition of carbonic anhydrase IX targets primary tumors, metastases, and cancer stem cells: Three for the price of one. *Med Res Rev* 2018;38:1799–836.
81. Ilie M, Mazure NM, Hofman V, et al. High levels of carbonic anhydrase IX in tumour tissue and plasma are biomarkers of poor prognostic in patients with non-small cell lung cancer. *Br J Cancer* 2010;102:1627–35.
82. Logozzi M, Capasso C, Di Raimo R, et al. Prostate cancer cells and exosomes in acidic condition show increased carbonic anhydrase IX expression and activity. *J Enzyme Inhib Med Chem* 2019;34:272–8.
83. Dorai T, Sawczuk IS, Pastorek J, et al. The role of carbonic anhydrase IX overexpression in kidney cancer. *Eur J Cancer* 2005;41:2935–47.
84. Murai T. The role of lipid rafts in cancer cell adhesion and migration. *Int J Cell Biol* 2012;2012:1.
85. Kaluz S, Kaluzova M, Liao SY, et al. Transcriptional control of the tumor- and hypoxia-marker carbonic anhydrase 9: a one transcription factor (HIF-1) show? *Biochim Biophys Acta* 2009;1795:162–72.
86. Brown-Glaberman U, Marron M, Chalasani P, et al. Circulating carbonic anhydrase IX and antiangiogenic therapy in breast cancer. *Dis Markers* 2016;2016:1.
87. Finkelmeier F, Canli O, Peiffer KH, et al. Circulating hypoxia marker carbonic anhydrase IX (CA9) in patients with hepatocellular carcinoma and patients with cirrhosis. *PLoS One* 2018;13:e0200855.
88. Malentacchi F, Vinci S, Melina AD, et al. Urinary carbonic anhydrase IX splicing messenger RNA variants in urogenital cancers. *Urol Oncol* 2016;34:292 e9–292 e16.
89. Liu S, Tian Z, Zhang L, et al. Combined cell surface carbonic anhydrase 9 and CD147 antigens enable high-efficiency capture of circulating tumor cells in clear cell renal cell carcinoma patients. *Oncotarget* 2016;7:59877–91.