



The prescribed mean curvature equation in weakly regular domains

Gian Paolo Leonardi  and Giorgio Saracco 

Abstract. We show that the characterization of existence and uniqueness up to vertical translations of solutions to the prescribed mean curvature equation, originally proved by Giusti in the smooth case, holds true for domains satisfying very mild regularity assumptions. Our results apply in particular to the non-parametric solutions of the capillary problem for perfectly wetting fluids in zero gravity. Among the essential tools used in the proofs, we mention a *generalized Gauss–Green theorem* based on the construction of the weak normal trace of a vector field with bounded divergence, in the spirit of classical results due to Anzellotti, and a *weak Young’s law* for (Λ, r_0) -minimizers of the perimeter.

Mathematics Subject Classification. Primary 49K20, 35J93; Secondary 49Q20.

Keywords. Prescribed mean curvature, Capillarity, Weak normal trace, Perimeter.

1. Introduction

Let Ω be an open bounded set in \mathbb{R}^n and let $H : \Omega \rightarrow \mathbb{R}$ be a Lipschitz continuous function. A classical solution to the Prescribed Mean Curvature equation is a function $u : \Omega \rightarrow \mathbb{R}$ of class C^2 satisfying

$$\operatorname{div} \left(\frac{\nabla u(x)}{\sqrt{1 + |\nabla u(x)|^2}} \right) = H(x) \quad \forall x \in \Omega. \quad (\text{PMC})$$

The left-hand side of (PMC) corresponds to the mean curvature of the graph of u at the point $(x, u(x))$. The existence and the properties of solutions to

G. P. Leonardi and G. Saracco have been supported by GNAMPA projects: *Problemi isoperimetrici e teoria della misura in spazi metrici* (2015) and *Variational problems and geometric measure theory in metric spaces* (2016).

G. Saracco has also been supported by the DFG Grant n. GZ:PR 1687/1–1.

(PMC), possibly satisfying some given boundary conditions, have been the object of extensive studies in the past, also due to the close connection between (PMC) and capillarity. After the pioneering works by Young [57], Laplace [36], and Gauss [28], it is nowadays a well-known fact that the mean curvature of a capillary surface in a cylindrical container with cross-section Ω is determined by the surface tension, by the wetting properties of the fluid with respect to the container, and by the presence of external forces such as gravity. The modern theory of capillarity has its roots in a series of fundamental papers by Finn [25], Concus–Finn [13–15], Emmer [20, 21], Gerhardt [29–31], Giaquinta [32], Giusti [33, 34], and many others (see [26] and the references therein). Other contributions to the theory have been obtained in various directions, see for instance Tam [54, 55], Finn [27], Concus–Finn [16], Caffarelli and Friedman [5], as well as more recent works by De Philippis and Maggi [19], Caffarelli and Mellet [6] and Lancaster [35]. However the above list is far from being complete.

A necessary condition on the pair (Ω, H) for the existence of a solution to (PMC) can be easily found by integrating (PMC) on any relatively compact set $A \subset \Omega$ with smooth boundary. Indeed, by applying the divergence theorem we get

$$\left| \int_A H \, dx \right| \leq \int_{\partial A} |\langle Tu, \nu \rangle| \, d\mathcal{H}^{n-1},$$

where ν is the exterior normal to ∂A and \mathcal{H}^{n-1} is the Hausdorff $(n-1)$ -dimensional measure in \mathbb{R}^n . Then using the fact that the vector field

$$Tu(x) := \frac{\nabla u(x)}{\sqrt{1 + |\nabla u(x)|^2}}$$

has modulus less than 1 on Ω , we obtain for every such A the strict inequality

$$\left| \int_A H \, dx \right| < P(A), \tag{1}$$

where $P(A)$ denotes the perimeter of A (when ∂A is smooth, $P(A) = \mathcal{H}^{n-1}(\partial A)$; more generally, $P(A)$ has to be understood in the sense of Definition 2.1).

Notice that whenever H is a non-negative constant on Ω one obtains the necessary condition $H < \frac{P(A)}{|A|}$ for all relatively compact subsets $A \subset \Omega$ with positive volume. Hence, the existence of solutions to (PMC) is closely related to the so-called *Cheeger problem*, which consists in minimizing the quotient $\frac{P(A)}{|A|}$ among all $A \subset \subset \Omega$ (see for instance the review papers [37] and [46], and references therein).

In the fundamental paper [34], Giusti proved that the necessary condition (1) is also sufficient for the existence of solutions to (PMC) in any bounded connected open set Ω with Lipschitz boundary. More specifically, he showed that if (1) holds together with the strict inequality

$$\left| \int_{\Omega} H \, dx \right| < P(\Omega) \tag{2}$$

then one can find many variational solutions (see [32]) attaining any given Dirichlet $L^1(\partial\Omega)$ boundary datum in a weak sense. On the other hand, a much more subtle situation occurs when the equality

$$\left| \int_{\Omega} H \, dx \right| = P(\Omega) \quad (3)$$

holds, as it corresponds to the so-called *extremal case*. Whenever the pair (Ω, H) is such that both (1) and (3) are satisfied, we will call the pair *extremal*.

Concerning the existence of solutions to (PMC) in the extremal case, one can essentially consider a suitably translated sequence of variational (non-extremal) solutions u_i of (PMC), defined on subsets Ω_i that converge to Ω both in volume and in perimeter, as $i \rightarrow \infty$. Then, one obtains a so-called *generalized solution* u defined on Ω as the limit of u_i (in the sense of the L^1 -convergence of the subgraphs, see [45]). The extremal case is particularly relevant because it corresponds to capillarity for a perfectly wetting fluid under zero-gravity conditions. By definition of perfect wetting, the fluid-gas interface meets the (smooth) boundary of the cylindrical container with a zero contact angle; in other words one expects that any solution u in the extremal case automatically satisfies the boundary condition of Neumann type

$$\langle Tu, \nu \rangle = 1 \quad \text{on } \partial\Omega. \quad (4)$$

At the same time, one also experimentally observes that the solution u is unique up to additive constants. This is what Giusti showed to be a consequence of a more general equivalence result (see Theorem 2.1 in [34]) that he proved for the extremal case under the strong regularity assumption $\partial\Omega \in C^2$. Later, Finn observed that the regularity requirements on $\partial\Omega$ can be reduced to piece-wise Lipschitz (see [26, Chapter 6]) if one is interested in the existence of solutions to (PMC) in the 2-dimensional case, and to “ C^1 up to a \mathcal{H}^{n-1} -negligible set” if uniqueness up to vertical translations has to be shown in the extremal case. However the question about the validity of Giusti’s result under weaker assumptions on $\partial\Omega$ is still not completely answered.

In this paper we prove Giusti’s characterization of existence and uniqueness of solutions to (PMC) under very mild regularity hypotheses on Ω , see Theorems 4.3, 4.7, and 5.1. In particular, our results are valid for domains with inner cusps or with some porosity (see Example 5.5), which of course fall outside of the Lipschitz class.

Specifically, we assume that $\Omega \subset \mathbb{R}^n$ is an open, bounded set with finite perimeter, satisfying the following properties. First, we require that Ω coincides with its measure-theoretic interior (roughly speaking, we do not allow Ω to have “measure-zero holes”). Then we assume the existence of $k = k(\Omega) > 0$ such that

$$\min\{P(E; \Omega^c), P(\Omega \setminus E; \Omega^c)\} \leq k P(E; \Omega) \quad (5)$$

for all $E \subset \Omega$. Finally, we require that

$$P(\Omega) = \mathcal{H}^{n-1}(\partial\Omega). \quad (6)$$

Whenever an open set Ω satisfies (5) and (6), we say that Ω is *weakly regular*. We stress that weak regularity can be regarded as a minimal assumption in the following sense. On one hand, if one assumes (6), then (5) is equivalent to the existence of a continuous and surjective trace operator from $BV(\Omega)$ to $L^1(\partial\Omega)$, by well-known results about traces of functions in $BV(\Omega)$ (see Theorem 2.6). On the other hand, by Federer's Structure Theorem (see Theorem 2.5), (6) amounts to requiring that the set of points of $\partial\Omega$ that are of density 0 or 1 for Ω is \mathcal{H}^{n-1} -negligible, which can be considered as a very mild regularity assumption on $\partial\Omega$. Moreover, in the extremal case one can show that (5) is automatically satisfied by Ω , thus only (6) needs to be assumed (see [49]).

The proofs of the above-mentioned theorems require some facts and preliminary results of independent interest.

One of the key tools that we shall systematically use in our proofs is Theorem 2.8 about the interior approximation of an open set Ω with finite perimeter satisfying (6), by means of sequences of smooth sets that converge to Ω in measure and in perimeter. This result has been proved by Schmidt [51], here we only add to the statement the useful observation that, being Ω connected, one can find a sequence of connected smooth sets with the above-mentioned property. Another, more technical tool is the recent characterization of $W_0^{1,1}(\Omega)$ as the space of functions in $W^{1,1}(\Omega)$ having zero trace at $\partial\Omega$, due to Swanson [53] (see Theorem 2.7).

In Sect. 3 we introduce some notions and prove some results, that will be needed in the following sections. Under the assumptions (5) and (6), we prove Theorem 3.2 which states a generalized Gauss–Green formula valid for bounded continuous vector fields with bounded divergence and for BV functions on Ω . We recall that very general forms of the Gauss–Green Theorem have been already obtained by several authors, see for instance [18, 22, 23], [4, 56], [1, 58], and [47, 48]. We recall in particular the extensions of the divergence theorem for bounded, divergence-measure vector fields on sets with finite perimeter [7–10]. These last results rely on a notion of *weak normal trace* of a bounded, divergence-measure vector field ξ on the reduced boundary of E , where $E \subset\subset \Omega$ is a set of finite perimeter and Ω is the domain of the vector field, see [11, 17, 42]. This notion of trace already appears in [1], in the special case of E being an open bounded set with Lipschitz boundary. A crucial tool used in [10] (see also [12]) is the approximation of E by smooth sets which are “mostly” contained in the measure-theoretic interior of E with respect to the measure $\mu = \operatorname{div} \xi$. Actually, this is the main reason why E needs to be compactly contained in the domain of the vector field ξ . On the other hand, if such a domain Ω has finite perimeter and $P(\Omega) = \mathcal{H}^{n-1}(\partial\Omega)$ then one can consider the vector field $\hat{\xi}$ defined as $\hat{\xi} = \xi$ on Ω and $\hat{\xi} = 0$ on $\mathbb{R}^n \setminus \Omega$, so that by relying on Theorem 2.8 it is possible to show that $\operatorname{div} \hat{\xi}$ is a finite measure on \mathbb{R}^n . Then by applying [10, Theorem 25.1] one might show the validity of the divergence theorem for the field ξ on $E = \Omega$, which in turn leads to the generalized Gauss–Green formula

$$\int_{\Omega} \varphi \operatorname{div} \xi + \int_{\Omega} \nabla \varphi \cdot \xi = \int_{\partial\Omega} \varphi [\xi \cdot \nu] d\mathcal{H}^{n-1}, \quad (7)$$

where ν is the exterior weak normal to $\partial^*\Omega$, $[\xi \cdot \nu]$ denotes the weak normal trace of ξ on $\partial^*\Omega$, and $\varphi \in C_c^\infty(\mathbb{R}^n)$. However, also in view of the results of Sect. 5, in Sect. 3.1 we give a very direct proof of (7) when $\Omega \subset \mathbb{R}^n$ is an open bounded set satisfying (5) and (6). This will be accomplished by adapting the construction proposed by Anzellotti in [1] (see also [3, 50]). More precisely, we will show that (7) holds for every bounded continuous vector field ξ with divergence in $L^\infty(\Omega)$ and for any $\varphi \in BV(\Omega)$. We remark that the extra assumptions on ξ that we are requiring reflect the properties of the vector field Tu when u is a solution of (PMC) on Ω . We also stress that all bounded and connected Lipschitz domains, as well as some domains with inner cusps or with some controlled porosity (see for instance Example 5.5), are weakly regular and therefore (7) holds on them.

Finally, in the proof of Theorem 5.1 we shall use a so-called *weak Young's law* for (Λ, r_0) -minimizers of the perimeter, Theorem 3.5, that was originally shown in [38, Proposition 2.5] in the special case of Cheeger sets.

Some final observations about the stability of the solution to (PMC) in the extremal case are made. On one hand it is well-known in capillarity theory that even small and smooth deformations of Ω typically produce discontinuous changes in the solution of the capillary problem in Ω , and even the existence of such a solution in the non-parametric setting may instantaneously drop (see [26]). On the other hand, in Proposition 5.4 we give an answer to the question whether or not it is possible to obtain some stability result for the solution $u = u_\Omega$ of (PMC) when the pair (Ω, H) is extremal. Then, by coupling Proposition 5.4 with the construction described in Example 5.5, a sequence of non-smooth perturbations of a 2-dimensional disk can be constructed, in such a way that the corresponding sequence of solutions to the capillary problem for perfectly wetting fluids in zero gravity converge (up to suitable translations, and in the sense of L^1_{loc} -convergence of the epigraphs) to the solution of the problem in the disk.

2. Preliminaries

We first introduce some basic notations. We fix $n \geq 2$ and denote by \mathbb{R}^n the Euclidean n -space. Let $E \subset \mathbb{R}^n$, then we denote by χ_E the characteristic function of E . For any $x \in \mathbb{R}^n$ and $r > 0$ we denote by $B_r(x)$ the Euclidean open ball of center x and radius r . Given two sets E, F , we denote by $E\Delta F = (E \setminus F) \cup (F \setminus E)$ their symmetric difference. In order to define rescalings of sets, we conveniently introduce the notation $E_{x,r} = r^{-1}(E - x)$, where $E \subset \mathbb{R}^n$, $x \in \mathbb{R}^n$, and $r > 0$. Let $E \subset \Omega \subset \mathbb{R}^n$ with Ω open; we write $E \subset\subset \Omega$ whenever the topological closure of E , \overline{E} , is a compact subset of Ω . Given a Borel set E we denote by $|E|$ its n -dimensional Lebesgue measure. Whenever a measurable function, or vector field, f is defined on \mathbb{R}^n , we set $\|f\|_\infty$ for the L^∞ -norm of f on \mathbb{R}^n .

Definition 2.1. (Perimeter) Let E be a Borel set in \mathbb{R}^n . We define the perimeter of E in an open set $\Omega \subset \mathbb{R}^n$ as

$$P(E; \Omega) := \sup \left\{ \int_{\Omega} \chi_E(x) \operatorname{div} g(x) \, dx : g \in C_c^1(\Omega; \mathbb{R}^n), \|g\|_{\infty} \leq 1 \right\}.$$

We set $P(E) = P(E; \mathbb{R}^n)$. If $P(E; \Omega) < \infty$ we say that E is a set of finite perimeter in Ω . In this case (see [41]) one has that the perimeter of E coincides with the total variation $|D\chi_E|$ of the vector-valued Radon measure $D\chi_E$ (the distributional gradient of χ_E), which is defined for all Borel subsets of Ω thanks to Riesz Theorem.

Definition 2.2. (Points of density α) Let E be a Borel set in \mathbb{R}^n , $x \in \mathbb{R}^n$. If the limit

$$\theta(E)(x) := \lim_{r \rightarrow 0^+} \frac{|E \cap B_r(x)|}{\omega_n r^n}$$

exists, it is called the density of E at x . We define the set of points of density $\alpha \in [0, 1]$ of E as

$$E^{(\alpha)} := \{x \in \mathbb{R}^n : \theta(E)(x) = \alpha\}.$$

We also define the essential boundary $\partial^e E := \mathbb{R}^n \setminus (E^{(0)} \cup E^{(1)})$.

Definition 2.3. (Approximate limit) Let f be a measurable function or vector field defined on Ω . Given $z \in \bar{\Omega}$ we write

$$\operatorname{ap}\text{-}\lim_{x \rightarrow z} f(x) = w$$

if for every $\alpha > 0$ the set $\{x \in \Omega : |f(x) - w| \geq \alpha\}$ has density 0 at z .

Theorem 2.4. (De Giorgi Structure Theorem) *Let E be a set of finite perimeter and let $\partial^* E$ be the reduced boundary of E defined as*

$$\partial^* E := \left\{ x \in \partial^e E : \lim_{r \rightarrow 0^+} \frac{D\chi_E(B_r(x))}{|D\chi_E|(B_r(x))} = -\nu_E(x) \in \mathbb{S}^{n-1} \right\}.$$

Then,

- (i) $\partial^* E$ is countably \mathcal{H}^{n-1} -rectifiable in the sense of Federer [24];
- (ii) for all $x \in \partial^* E$, $\chi_{E_{x,r}} \rightarrow \chi_{H_{\nu_E(x)}}$ in $L^1_{loc}(\mathbb{R}^n)$ as $r \rightarrow 0^+$, where $H_{\nu_E(x)}$ denotes the half-space through 0 whose exterior normal is $\nu_E(x)$;
- (iii) for any Borel set A , $P(E; A) = \mathcal{H}^{n-1}(A \cap \partial^* E)$, thus in particular $P(E) = \mathcal{H}^{n-1}(\partial^* E)$;
- (iv) $\int_E \operatorname{div} g = \int_{\partial^* E} g \cdot \nu_E \, d\mathcal{H}^{n-1}$ for any $g \in C_c^1(\mathbb{R}^n; \mathbb{R}^n)$.

Theorem 2.5. (Federer’s Structure Theorem) *Let E be a set of finite perimeter. Then, $\partial^* E \subset E^{(1/2)} \subset \partial^e E$ and one has*

$$\mathcal{H}^{n-1}(\partial^e E \setminus \partial^* E) = 0.$$

In what follows, Ω will always denote a *domain* of \mathbb{R}^n , i.e., an open connected set coinciding with its measure-theoretic interior, that is, we assume that any point $x \in \mathbb{R}^n$, such that there exists a radius $r > 0$ with the property $|B_r(x) \setminus \Omega| = 0$, is necessarily contained in Ω .

The next result combines [43, Theorem 9.6.4] and [2, Theorem 10 (a)].

Theorem 2.6. *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with $P(\Omega) = \mathcal{H}^{n-1}(\partial\Omega) < +\infty$. Then the following are equivalent:*

(i) there exists $k = k(\Omega)$ such that for all $E \subset \Omega$

$$\min\{P(E; \Omega^c), P(\Omega \setminus E; \Omega^c)\} \leq kP(E; \Omega);$$

(ii) there exists a continuous trace operator from $BV(\Omega)$ to $L^1(\partial\Omega)$ with the following property: for any $\varphi \in L^1(\partial\Omega)$ there exists $\Psi \in W^{1,1}(\mathbb{R}^n)$ such that φ is the trace of Ψ on $\partial\Omega$.

Another useful result is the characterization of $W_0^{1,1}(\Omega)$ as the space of functions in $W^{1,1}(\Omega)$ having zero trace at $\partial\Omega$.

Theorem 2.7. ([53, Theorem 5.2]) *Let $\Omega \subset \mathbb{R}^n$ be an open set and let $u \in W^{1,1}(\Omega)$. Then, $u \in W_0^{1,1}(\Omega)$ if and only if*

$$\lim_{r \rightarrow 0} \frac{1}{r^n} \int_{B_r(x) \cap \Omega} |u(y)| dy = 0$$

for \mathcal{H}^{n-1} -almost all $x \in \partial\Omega$.

The following approximation theorem is essentially due to Schmidt [51] and will play a crucial role in the paper.

Theorem 2.8. (Interior smooth approximation) *Suppose that Ω is a bounded open set in \mathbb{R}^n such that $P(\Omega) = \mathcal{H}^{n-1}(\partial\Omega) < +\infty$. Then, for every $\delta > 0$ there exist an open set Ω_δ with smooth boundary in \mathbb{R}^n such that*

$$\begin{aligned} \Omega_\delta \subset\subset \Omega, \quad \Omega \setminus \Omega_\delta \subset (\mathcal{N}_\delta(\partial\Omega) \cap \mathcal{N}_\delta(\partial\Omega_\delta)), \quad |\Omega \setminus \Omega_\delta| < \delta, \\ P(\Omega_\delta) \leq P(\Omega) + \delta, \end{aligned} \tag{8}$$

where $\mathcal{N}_\delta(A)$ denotes the δ -tubular neighborhood of $A \subset \mathbb{R}^n$. Moreover, Ω_δ can be chosen connected as soon as Ω is connected.

Proof. The existence of Ω_δ satisfying (8) is proved in [51]. In order to show the last part of the statement, we fix a compact set $K \subset \Omega$ such that $|\Omega \setminus K| < \delta$, then setting $d = \min\{\text{dist}(x, \partial\Omega) : x \in K\}$ we take a finite covering of K by balls of radius $d/2$ and let x_1, \dots, x_N denote their centers. By connectedness, for any $h, k \in \{1, \dots, N\}$ there exists a path $\Gamma_{hk} \subset \Omega$ connecting x_h to x_k , so that the set

$$\tilde{K} = \bigcup_{h=1}^N B_{d/2}(x_h) \cup \bigcup_{h,k=1}^N \Gamma_{hk}$$

is contained in Ω , connected, compact, and such that $|\Omega \setminus \tilde{K}| < \delta$. Let now $\tilde{\delta} = \min(\min\{\text{dist}(x, \partial\Omega) : x \in \tilde{K}\}, \delta) > 0$, then by (8) with $\tilde{\delta}$ replacing δ we get an open set $\Omega_{\tilde{\delta}}$ which necessarily has a connected component A containing \tilde{K} , so that (8) and the last part of the statement are satisfied by setting $\Omega_\delta = A$. \square

3. Some technical tools

We collect in this section some key notions and results that will be later needed. Our first aim is to prove the Gauss–Green Theorem 3.2, on which the main

results of Sect. 5 are based. For this we shall introduce the *weak normal trace* of a vector field ξ on $\partial\Omega$, denoted as $[\xi \cdot \nu]$, as a suitable extension of the usual scalar product between the trace of ξ and the normal to $\partial\Omega$, whenever the former exists. It is indeed quite easy to prove that whenever the approximate limit of the vector field $\xi(x)$ exists as $x \rightarrow z \in \partial^*\Omega$, then $[\xi \cdot \nu](z)$ equals the scalar product between that limit and the outer normal to $\partial^*\Omega$ at z , see Proposition 3.3.

Our second tool is represented by a *weak Young's law* for perimeter quasi-minimizers, Theorem 3.5, that will be needed in Sect. 5 for the proof of the implication $(U) \Rightarrow (E)$ in Theorem 5.1. A slightly less general form of this lemma has been proved in [38], in the context of Cheeger sets. Roughly speaking, it says that the inner boundary of any (Λ, r_0) -minimizer of the perimeter in a domain $\bar{\Omega}$ must meet the reduced boundary of Ω in a tangential way.

3.1. The weak normal trace

Let $\Omega \subset \mathbb{R}^n$ be open, bounded, and weakly regular, i.e., satisfying (5) and (6). We denote by $X(\Omega)$ the collection of vector fields $\xi \in L^\infty(\Omega; \mathbb{R}^n) \cap C^0(\Omega; \mathbb{R}^n)$ such that $\operatorname{div} \xi \in L^\infty(\Omega)$. Following Anzellotti [1], for every $u \in BV(\Omega)$ we define the pairing

$$\langle \xi, u \rangle_{\partial\Omega} = \int_{\Omega} u \operatorname{div} \xi + \int_{\Omega} \xi \cdot Du. \quad (9)$$

The map $\langle \cdot, \cdot \rangle_{\partial\Omega} : X(\Omega) \times BV(\Omega) \rightarrow \mathbb{R}$ is bilinear. If $u, v \in W^{1,1}(\Omega)$ have the same trace on $\partial\Omega$ then by Theorem 2.7 there exists a sequence $\{g_j\}$ of functions in $C_c^\infty(\Omega)$ such that $g_j \rightarrow u - v$ weakly in $BV(\Omega)$, so that we have

$$\begin{aligned} \langle \xi, u - v \rangle_{\partial\Omega} &= \int_{\Omega} (u - v) \operatorname{div} \xi + \int_{\Omega} \xi \cdot D(u - v) \\ &= \lim_j \int_{\Omega} g_j \operatorname{div} \xi + \int_{\Omega} \xi \cdot \nabla g_j = 0. \end{aligned}$$

This shows that the pairing defined in (9) only depends on the trace of u on $\partial\Omega$. Then by Anzellotti–Giaquinta's approximation in $BV(\Omega)$ and by Theorem 2.6 we infer that $\langle \xi, u \rangle_{\partial\Omega} = \langle \xi, v \rangle_{\partial\Omega}$ whenever $u, v \in BV(\Omega)$ have the same trace on $\partial\Omega$.

At this point we can show the continuity of the pairing (9) in the topology of $L^\infty(\Omega; \mathbb{R}^n) \times L^1(\partial\Omega)$. The following, key lemma extends [1, Lemma 5.5].

Lemma 3.1. *Let Ω be weakly regular. Then for every $u \in L^1(\partial\Omega)$ and $\varepsilon > 0$ there exists $w_\varepsilon \in BV(\Omega) \cap C^\infty(\Omega)$ such that*

- (i) *the trace of w_ε on $\partial\Omega$ equals u \mathcal{H}^{n-1} -almost everywhere on $\partial\Omega$,*
- (ii) $\int_{\Omega} |\nabla w_\varepsilon| \leq \int_{\partial\Omega} |u| + \varepsilon$,
- (iii) $w_\varepsilon(x) = 0$ whenever $\operatorname{dist}(x, \partial\Omega) > \varepsilon$,
- (iv) $\int_{\Omega} |w_\varepsilon| \leq \varepsilon$,
- (v) $\|w_\varepsilon\|_{L^\infty(\Omega)} \leq \|u\|_{L^\infty(\partial\Omega)}$.

Proof. Let us fix $\varepsilon > 0$. By Theorem 2.6 (ii) there exists $\Psi \in W^{1,1}(\mathbb{R}^n)$ such that its trace on $\partial\Omega$ coincides with u . Up to an application of Meyer–Serrin's

approximation theorem, we can additionally assume that $\Psi \in C^\infty(\Omega)$. Moreover we fix a sequence $\{\Psi_j\}_j$ of smooth functions such that $\|\Psi - \Psi_j\|_{W^{1,1}(\mathbb{R}^n)} \rightarrow 0$ as $j \rightarrow \infty$. Again by Theorem 2.6 (ii) we have that the trace operator from $BV(\Omega)$ to $L^1(\partial\Omega)$ is continuous, hence

$$\int_{\partial\Omega} |\Psi_j| d\mathcal{H}^{n-1} \rightarrow \int_{\partial\Omega} |\Psi| d\mathcal{H}^{n-1} = \int_{\partial\Omega} |u| d\mathcal{H}^{n-1} \quad \text{as } j \rightarrow \infty.$$

Given $\delta, \eta > 0$ we define $\chi_{\delta,\eta}(x) = \chi_{\Omega_\delta} * \rho_\eta(x)$, where ρ_η is a standard symmetric mollifier with support in $B_\eta(0)$, while $\Omega_\delta \subset\subset \Omega$ is obtained in virtue of Theorem 2.8, so that the Hausdorff distance between $\partial\Omega_\delta$ and $\partial\Omega$ is smaller than δ and $|P(\Omega_\delta) - P(\Omega)| \leq \delta$. We note that up to choosing δ and η small enough we get $\text{spt}(\chi_{\delta,\eta}) \subset\subset \Omega$, $\chi_{\delta,\eta} = 1$ on the set $\{x \in \Omega : \text{dist}(x, \partial\Omega) > \varepsilon\}$, and $\left| \int_\Omega |\nabla \chi_{\delta,\eta}| - P(\Omega_\delta) \right| \leq \delta$. Then we define $w_{\delta,\eta}(x) = \Psi(x)(1 - \chi_{\delta,\eta}(x))$ and, for any fixed vector field $g \in C^1(\mathbb{R}^n; \mathbb{R}^n)$ with $\|g\|_\infty \leq 1$ and compact support in Ω , up to choosing δ and η small enough as well as j sufficiently large we obtain

$$\begin{aligned} & \int_\Omega \nabla w_{\delta,\eta} \cdot g \, dx \\ &= \int_\Omega (1 - \chi_{\delta,\eta}) \nabla \Psi \cdot g \, dx - \int_\Omega \Psi \nabla \chi_{\delta,\eta} \cdot g \, dx \\ &\leq \int_\Omega (1 - \chi_{\delta,\eta}) |\nabla \Psi| - \int_\Omega \Psi_j \nabla \chi_{\delta,\eta} \cdot g \, dx - \int_\Omega (\Psi - \Psi_j) \nabla \chi_{\delta,\eta} \cdot g \, dx \\ &\leq \frac{\varepsilon}{4} + \int_\Omega |\Psi_j| |\nabla \chi_{\delta,\eta}| \, dx + \int_\Omega \chi_{\delta,\eta} \left(\nabla(\Psi - \Psi_j) \cdot g + (\Psi - \Psi_j) \text{div } g \right) \, dx \\ &\leq \frac{\varepsilon}{4} + \int_\Omega |\Psi_j| |\nabla \chi_{\delta,\eta}| \, dx + (1 + \|\text{div } g\|_\infty) \int_\Omega \left(|D(\Psi - \Psi_j)| + |\Psi - \Psi_j| \right) \, dx \\ &\leq \int_\Omega |\Psi_j| |\nabla \chi_{\delta,\eta}| \, dx + \frac{\varepsilon}{2} \leq \int |\Psi_j| d|D\chi_\Omega| + \frac{3}{4}\varepsilon \leq \int_{\partial\Omega} |u| d\mathcal{H}^{n-1} + \varepsilon. \end{aligned}$$

We finally set $w_\varepsilon = w_{\delta,\eta}$ and, by taking the supremum over g , we find

$$\int_\Omega |\nabla w_\varepsilon| \, dx \leq \int_{\partial\Omega} |u| d\mathcal{H}^{n-1} + \varepsilon,$$

which proves (ii). Finally, (i), (iii) and (v) are immediate from the construction, while (iv) is easily shown to hold up to possibly taking smaller δ and η . \square

Now, given $\varepsilon > 0$ and $\varphi \in BV(\Omega) \cap L^\infty(\Omega)$, taking w_ε as in Lemma 3.1 (with $u = \varphi$ on $\partial\Omega$), and setting $\Omega_\varepsilon = \{x \in \Omega : \text{dist}(x, \partial\Omega) \geq \varepsilon\}$ we obtain

$$\begin{aligned} |\langle \xi, \varphi \rangle_{\partial\Omega}| &= |\langle \xi, w_\varepsilon \rangle_{\partial\Omega}| \\ &\leq \|\varphi\|_{L^\infty(\Omega)} \int_{\Omega \setminus \Omega_\varepsilon} |\text{div } \xi| + \|\xi\|_{L^\infty(\Omega)} \int_\Omega |\nabla w_\varepsilon| \\ &\leq \|\varphi\|_{L^\infty(\Omega)} \int_{\Omega \setminus \Omega_\varepsilon} |\text{div } \xi| + \|\xi\|_{L^\infty(\Omega)} \left(\int_{\partial\Omega} |\varphi| + \varepsilon \right), \end{aligned}$$

which by the arbitrary choice of ε leads to

$$|\langle \xi, \varphi \rangle_{\partial\Omega}| \leq \|\xi\|_{L^\infty(\Omega)} \int_{\partial\Omega} |\varphi|. \tag{10}$$

One can check by a truncation argument that (10) holds for each $\varphi \in BV(\Omega)$. An immediate consequence of (10) is the fact that the linear functional $N_\xi : L^1(\partial\Omega) \rightarrow \mathbb{R}$ defined as $N_\xi(u) = \langle \xi, u \rangle_{\partial\Omega}$ is continuous on $L^1(\partial\Omega)$, thus it can be represented by a function in $L^\infty(\partial\Omega)$, hereafter denoted by $[\xi \cdot \nu]$. This function is the so-called *weak normal trace* of the vector field $\xi \in X(\Omega)$ on $\partial\Omega$. Another immediate consequence of (10) is the following L^∞ -estimate of the weak normal trace:

$$\|[\xi \cdot \nu]\|_{L^\infty(\partial\Omega)} \leq \|\xi\|_{L^\infty(\Omega)}. \tag{11}$$

Summing up, we have proved that (9) can be rewritten in the form of the generalized Gauss–Green formula stated in the next theorem.

Theorem 3.2. *Let $\Omega \subset \mathbb{R}^n$ be open, bounded and weakly regular. Let $\xi \in X(\Omega)$ and $\varphi \in BV(\Omega)$, then*

$$\int_\Omega \varphi \operatorname{div} \xi + \int_\Omega \xi \cdot D\varphi = \int_{\partial\Omega} \varphi [\xi \cdot \nu] d\mathcal{H}^{n-1}. \tag{12}$$

The next proposition shows that the weak normal trace is a proper extension of the normal component of the usual trace of ξ on $\partial\Omega$, whenever such a trace exists in measure-theoretic sense.

Proposition 3.3. *Let $\Omega \subset \mathbb{R}^n$ be open, bounded and weakly regular. Let $\xi \in X(\Omega)$ and let $z \in \partial^*\Omega$ be a Lebesgue point for the weak normal trace $[\xi \cdot \nu]$. Assume*

$$\operatorname{ap}\text{-}\lim_{x \rightarrow z} \xi(x) = w, \tag{13}$$

then

$$[\xi \cdot \nu](z) = w \cdot \nu(z). \tag{14}$$

Proof. We can assume that $z = 0$ up to a translation. We fix a sequence $r_i \downarrow 0$ as $i \rightarrow \infty$. Given any function (or vector field) f defined in Ω , we set

$$\Omega_i = r_i^{-1}\Omega, \quad f_i(y) = f(r_i y).$$

We note that $Df_i(y) = r_i Df(r_i y)$ in the sense of distributions. By (13) we infer that for all $\alpha > 0$ the set

$$N_i(\alpha) = r_i^{-1}N(\alpha) = \{y \in \Omega_i : |\xi_i(y) - w| \geq \alpha\}$$

satisfies

$$\lim_{i \rightarrow \infty} |N_i(\alpha) \cap B_1| = 0. \tag{15}$$

On the other hand, the fact that $z = 0$ is by assumption a Lebesgue point for $[\xi \cdot \nu]$ implies that

$$[\xi \cdot \nu](0) = \lim_{i \rightarrow \infty} \mu_i^{-1} \int_{\partial\Omega_i \cap B_1} [\xi \cdot \nu]_i(y) d\mathcal{H}^{n-1}(y), \tag{16}$$

where $\mu_i = \mathcal{H}^{n-1}(\partial\Omega_i \cap B_1)$. Now we take $\delta \in (0, 1)$ and set $\alpha = \delta^2$ and

$$\varphi(y) = \max(0, \min(1, (1 - |y|)/\delta)).$$

By Theorem 2.4(ii), setting $H = H_{\nu(0)}$ for brevity, we obtain

$$\begin{aligned} \left| \int_{\Omega_i \cap B_1} D\varphi(x) dx - \int_{H \cap B_1} D\varphi(x) dx \right| &\leq \delta^{-1} |(\Omega_i \Delta H) \cap B_1| \\ &= m_i(\delta) \rightarrow 0 \quad \text{as } i \rightarrow \infty. \end{aligned} \quad (17)$$

Moreover by Theorem 3.2 we get for a suitable constant $C > 0$

$$\begin{aligned} \left| \int_{H \cap B_1} D\varphi(x) dx - \omega_{n-1} \nu(0) \right| &= \left| \int_{\partial H \cap B_1} \varphi(x) d\mathcal{H}^{n-1}(x) - \omega_{n-1} \right| \\ &= \omega_{n-1} \int_0^1 [1 - (1 - \delta t)^{n-1}] dt \\ &\leq C\delta. \end{aligned} \quad (18)$$

Then by (16), (17), (18), and Theorem 2.4 (ii), we find

$$\begin{aligned} &\omega_{n-1} \left| [\xi \cdot \nu](0) - w \cdot \nu(0) \right| \\ &\leq \left| \lim_{i \rightarrow \infty} \int_{\partial\Omega_i \cap B_1} [\xi \cdot \nu]_i d\mathcal{H}^{n-1} - w \cdot \int_{H \cap B_1} D\varphi(x) dx \right| + C\delta \\ &\leq \limsup_{i \rightarrow \infty} \left| \int_{\Omega_i \cap B_1} \varphi \operatorname{div} \xi_i \right| + \left| \int_{\Omega_i \cap B_1} (\xi_i - w) \cdot D\varphi \right| \\ &\quad + m_i(\delta) + C\delta \\ &= \limsup_{i \rightarrow \infty} (A_i + B_i + m_i(\delta)) + C\delta. \end{aligned} \quad (19)$$

Then we notice that $A_i + m_i(\delta) \rightarrow 0$ as $i \rightarrow \infty$, while

$$\begin{aligned} B_i &= \left| \int_{(\Omega_i \cap B_1) \setminus N_i(\alpha)} (\xi_i - w) \cdot D\varphi + \int_{N_i(\alpha) \cap B_1} (\xi_i - w) \cdot D\varphi \right| \\ &\leq \frac{\omega_n \alpha}{\delta} + \frac{2\|\xi\|_\infty}{\delta} |N_i(\alpha) \cap B_1| \\ &\leq \omega_n \delta + \frac{2\|\xi\|_\infty}{\delta} |N_i(\alpha) \cap B_1|. \end{aligned}$$

Therefore by passing to the limit as $i \rightarrow \infty$ in (19) and using (15) we finally get

$$\omega_{n-1} \left| [\xi \cdot \nu](0) - w \cdot \nu(0) \right| \leq (\omega_n + C)\delta,$$

which implies (14) at once by the arbitrary choice of $\delta \in (0, 1)$. \square

In general, the weak normal trace $[\xi \cdot \nu]$ of a vector field $\xi \in X(\Omega)$ at $x \in \partial\Omega$ does not coincide to any pointwise, almost-everywhere, or measure-theoretic limit of the scalar product $\xi(y) \cdot \nu(x)$, as $y \rightarrow x$. However, one should expect some weak-type convergence of the normal component of ξ to the value of $[\xi \cdot \nu]$ at any Lebesgue point $x_0 \in \partial^*\Omega$. More precisely, let Ω_h be a sequence of relatively compact, open subsets of Ω with smooth boundary, that converge

to Ω both in perimeter and volume (see Theorem 2.8). We can consider the corresponding sequence of Radon measures $\mu_h = \langle \xi, \nu_h \rangle \mathcal{H}^{n-1} \llcorner \partial\Omega_h$. By Theorem 3.2 one easily checks that μ_h weakly- $*$ converges to $\mu = [\xi \cdot \nu] \mathcal{H}^{n-1} \llcorner \partial^*\Omega$ as $h \rightarrow \infty$.

By a similar application of Theorem 3.2 (simply take $\varphi = \chi_{B_r(x_0)}$) one can more explicitly characterize the weak normal trace at \mathcal{H}^{n-1} -almost every point $x_0 \in \partial^*\Omega$ as the following limit of spherical averages, as pointed out for instance in [10]:

$$[\xi \cdot \nu](x_0) = \lim_{r \rightarrow 0} \frac{1}{\omega_{n-1} r^{n-1}} \int_{\partial B_r(x_0) \cap \Omega} \xi(x) \cdot \frac{x - x_0}{|x - x_0|} d\mathcal{H}^{n-1}(x).$$

Nevertheless, such a characterization of the weak normal trace is not fully satisfactory, as one would expect to obtain coincidence with the classical trace in some special cases (see in particular the characterization of extremality discussed in Section 5). A more specific study of weak normal traces will appear in [39].

3.2. The weak Young’s law for (Λ, r_0) -minimizers

Let us start recalling the definition of (Λ, r_0) -minimizer of the perimeter.

Definition 3.4. Let $\Omega \subset \mathbb{R}^n$ be an open set of locally finite perimeter, and let E be a measurable subset of Ω . We say that E is a (Λ, r_0) -perimeter minimizer in $\overline{\Omega}$ if there exist two constants $\Lambda \in [0, +\infty)$ and $r_0 > 0$ such that for every $x \in \mathbb{R}^n$, every Borel set F such that $F \Delta E$ is compactly contained in $B_r(x) \cap \overline{\Omega}$, and every $r < r_0$, one has

$$P(E; B_r(x)) \leq P(F; B_r(x)) + \Lambda |F \Delta E|.$$

Theorem 3.5. (Weak Young’s Law) *Let Ω be an open set with locally finite perimeter and let E be a (Λ, r_0) -minimizer in $\overline{\Omega}$. Then $\partial E \cap \Omega$ meets $\partial^*\Omega$ in a tangential way, i.e., for any $x \in \partial^*\Omega \cap \overline{(\partial E \cap \Omega)}$ one has that $x \in \partial^*E$ and $\nu_E(x) = \nu_\Omega(x)$.*

Proof. Let us fix a point $x \in \partial^*\Omega \cap \partial E$ and let $x+H$ be the half space obtained by blowing up Ω around x . We divide the proof in three steps. In the first one we prove that E and Ω have the same tangential space at x , while in the third one we prove that x is in ∂^*E and that the outward normal is equal to the one outward Ω . Step 2 provides a tool to prove Step 3.

Step 1. Let us prove that E has the same tangent space $x+H$ at x . In order to do so, we need to prove perimeter and volume density estimates for $E \subset \Omega$ at x . Fix $m(r) := |E \cap B_r(x)|$ so that one has $P(E; \partial B_r(x)) = 0$, $m'(r) = P(E \cap B_r(x), \partial B_r(x))$ and $m(r) > 0$ for almost every $r > 0$. Being E a (Λ, r_0) -minimizer, for any $r < r_0$ and any competitor F , such that $F \Delta E \subset\subset B_r(x) \cap \overline{\Omega}$, one obtains

$$P(E; B_r(x)) \leq P(F; B_r(x)) + \Lambda |F \Delta E|.$$

Fix two radii, $r_2 < r_1 < r_0$ and consider as competitor in $B_{r_1}(x) \cap \bar{\Omega}$ the set $F := E \setminus B_{r_2}$. Therefore, exploiting the Λ -minimality one has

$$P(E; B_{r_1}(x)) \leq P(F; B_{r_1}(x)) + \Lambda|E \Delta F| \leq P(E; B_{r_1}(x) \setminus B_{r_2}(x)) + m'(r_2) + \Lambda m(r_2).$$

Thus

$$P(E; B_{r_2}(x)) = P(E; B_{r_1}(x)) - P(E; B_{r_1}(x) \setminus B_{r_2}(x)) \leq \Lambda m(r_2) + m'(r_2). \tag{20}$$

Due to the latter and to the isoperimetric inequality, it follows

$$\begin{aligned} c_1 m(r_2)^{\frac{n-1}{n}} &= c_1 |E \cap B_{r_2}(x)|^{\frac{n-1}{n}} \leq P(E \cap B_{r_2}(x)) \\ &= P(E; B_{r_2}(x)) + P(E \cap B_{r_2}(x); \partial B_{r_2}(x)) \leq \Lambda m(r_2) + 2m'(r_2). \end{aligned} \tag{21}$$

Hence for r_2 small enough and for some uniform constant c_2 we have

$$\frac{m'(r_2)}{m(r_2)^{\frac{n-1}{n}}} \geq c_2.$$

By integrating this inequality on $(\rho/2, \rho)$ we obtain for ρ small enough the volume density estimate

$$m(\rho) \geq c_3 \rho^n,$$

where c_3 is a uniform constant.

Regarding the perimeter, directly from (20) one can infer that $P(E; B_{r_2}) \leq \Lambda \omega_n r_2^n + m'(r_2)$, which, for r_2 small enough implies

$$P(E; B_{r_2}) \leq c_4 r_2^{n-1},$$

which then yields the perimeter density estimate.

Now blowing up E at x we find a limit set E_∞ contained in the half-space $x + H$ with $x \in \partial E_\infty$. It can be shown that E_∞ is not empty and minimizes the perimeter without volume constraint with respect to any compact variation contained in $x + H$. By convexity of H and by a maximum principle argument [52] one infers that E admits $x + H$ as unique blow up at the point x .

Step 2. Let us prove that

$$\lim_{r \rightarrow 0} \frac{P(E; B_r(x))}{r^{n-1}} = \omega_{n-1} \tag{22}$$

holds. Let E_r be $r^{-1}(E - x)$. Since the blow up of E at x is the half space $x + H$ one has the L^1_{loc} -convergence $\chi_{E_r} \rightarrow \chi_H$ as r goes to 0. By the lower semi-continuity of the perimeter we have

$$\liminf_{r \rightarrow 0} \frac{P(E; B_r(x))}{r^{n-1}} = \liminf_{r \rightarrow 0} P(E_r; B_1(0)) \geq P(H; B_1(0)) \geq \omega_{n-1},$$

therefore to prove (22) it is enough to show that

$$\limsup_{r \rightarrow 0} P(E_r; B_1(0)) \leq \omega_{n-1}. \tag{23}$$

Argue by contradiction and suppose there exists a sequence of radii r_i going to 0 such that

$$P(E_{r_i}; B_1(0)) \geq \omega_{n-1} + \varepsilon. \quad (24)$$

Recall that $x \in \partial^* \Omega$, therefore for r_i small enough one has

$$P(\Omega_{r_i}; B_s(0)) \leq s^{n-1} \omega_{n-1} + \varepsilon/3, \quad \text{for all } 1 < s < 2, \quad (25)$$

where Ω_{r_i} is defined in the same manner of E_{r_i} . Due to the L^1 -convergence in $B_2(0)$ of $\chi_{E_{r_i}}$ to χ_H and by coarea formula one can find a suitable

$$t \in \left(1, \left(\frac{\omega_{n-1} + \varepsilon/2}{\omega_{n-1} + \varepsilon/3} \right)^{\frac{1}{n-1}} \right)$$

such that

$$P(\Omega_i; \partial B_t(0)) = P(E_i; \partial B_t(0)) = 0 \quad (26)$$

$$\mathcal{H}^{n-1}(E_i \Delta \Omega_i \cap \partial B_t(0)) < \frac{\varepsilon}{4} \quad (27)$$

hold. Consider now the sets $F_i := (E \cup B_{tr_i}(x)) \cap \Omega$, for which, due to the previous, one has

$$\begin{aligned} P(F_i, B_{r_0}(x)) &= P(E; (\Omega \cap B_{r_0}(x)) \setminus B_{tr_i}(x)) + P(\Omega; B_{tr_i}(x)) \\ &\quad + r_i^{n-1} \mathcal{H}^{n-1}(E_i \Delta \Omega_i \cap \partial B_t(0)). \end{aligned}$$

For r_i small enough that $tr_i < r_0$, the set F_i is a competitor to E in B_{r_0} , therefore

$$\begin{aligned} r^{n-1}(\omega_{n-1} + \varepsilon) &\leq P(E; B_{r_i}(x)) \leq P(E; B_{r_0}(x)) - P(E; (\Omega \cap B_{r_0}(x)) \setminus B_{tr_i}(x)) \\ &\leq P(F; B_{r_0}(x)) - P(E; (\Omega \cap B_{r_0}(x)) \setminus B_{tr_i}(x)) + \Lambda |F \Delta E| \\ &\leq P(F; B_{r_0}(x)) - P(E; (\Omega \cap B_{r_0}(x)) \setminus B_{tr_i}(x)) + \Lambda |E \cap B_{tr_i}(x)| \\ &\leq P(\Omega; B_{tr_i}(x)) + r_i^{n-1} \frac{\varepsilon}{4} + \Lambda \omega_n (tr_i)^n \\ &\leq (tr_i)^{n-1} (\omega_{n-1} + \varepsilon/3) + r_i^{n-1} \frac{\varepsilon}{4} + \Lambda \omega_n (tr_i)^n \\ &< r_i^{n-1} (\omega_{n-1} + \varepsilon/2) + r_i^{n-1} \frac{\varepsilon}{2} \leq r^{n-1} (\omega_{n-1} + \varepsilon), \end{aligned} \quad (28)$$

which leads to a contradiction.

Step 3. Owing to (22), in order to show that $x \in \partial^* E$ and that $\nu_E(x) = \nu_\Omega(x)$ it is enough to prove that

$$\lim_{r \rightarrow 0} \frac{D\chi_E(B_r(x)) \cdot v}{\omega_{n-1} r^{n-1}} = 1, \quad (29)$$

where we have set $v = -\nu_\Omega(x)$. In virtue of Theorem 2.4 (iv), for almost every $r > 0$ one has

$$\begin{aligned} D\chi_E(B_r(x)) \cdot v &= \int_{E \cap \partial B_r(x)} v \cdot N \, d\mathcal{H}^{n-1} = \int_{H \cap \partial B_r(0)} v \cdot N \, d\mathcal{H}^{n-1} + A(x, r) \\ &= \omega_{n-1} r^{n-1} + A(x, r), \end{aligned} \quad (30)$$

where N is the outward normal to $\partial B_r(x)$ and

$$\begin{aligned} |A(x, r)| &= \left| v \cdot \int_{\partial B_r(x)} (\chi_E(y) - \chi_{x+H}(y)) N(y) d\mathcal{H}^{n-1}(y) \right| \\ &\leq \int_{\partial B_r(x)} |\chi_E(y) - \chi_{x+H}(y)| d\mathcal{H}^{n-1}(y). \end{aligned}$$

Now for any fixed $\delta > 0$, define the set $\Sigma(x, \delta) \subseteq (0, +\infty)$ of radii $r > 0$ such that $A(x, r) > \delta r^{n-1}$. Hence, by the L^1_{loc} -convergence of $r^{-1}(E - x)$ to the half-space H we infer that

$$\lim_{\rho \rightarrow 0^+} \frac{\mathcal{H}^1(\Sigma(x, \delta) \cap (0, \rho))}{\rho} = 0.$$

Therefore, for any decreasing infinitesimal sequence of radii $\{r_i\}_i$ we can find another sequence $\{\rho_i\}_i$ such that $\rho_i \notin \Sigma(x, \delta)$ for all i and $\rho_i = r_i + o(r_i)$ as $i \rightarrow \infty$. Suppose by contradiction that (29) does not hold. Then, there exist $\alpha > 0$ and a decreasing infinitesimal sequence $\{r_i\}_i$ such that

$$\left| \frac{D\chi_E(B_{r_i}(x)) \cdot v}{\omega_{n-1} r_i^{n-1}} \right| \geq \alpha, \tag{31}$$

for all $i \in \mathbb{N}$. By suitably choosing δ as $\omega_{n-1}\alpha/2$ and considering the sequence ρ_i defined above, one gets in (30) with the substitution $r = \rho_i$

$$|D_{\chi_E}(B_{\rho_i}(x)) \cdot v - \omega_{n-1} \rho_i^{n-1}| = |A(x, \rho_i)| \leq \frac{\alpha}{2} \omega_{n-1} \rho_i^{n-1}.$$

On the other hand, by (22), we also have

$$\begin{aligned} |D_{\chi_E}(B_{\rho_i}(x)) - D_{\chi_E}(B_{r_i}(x))| &\leq P(E; B_{\rho_i}(x) \Delta B_{r_i}(x)) \\ &\leq \omega_{n-1} |\rho_i^{n-1} - r_i^{n-1}| + o(r_i^{n-1}) \end{aligned}$$

as $i \rightarrow \infty$. Combining these two latter inequalities yields to

$$\begin{aligned} |D_{\chi_E}(B_{r_i}(x)) \cdot v - \omega_{n-1} r_i^{n-1}| &\leq \frac{\alpha}{2} \omega_{n-1} \rho_i^{n-1} + o(r_i^{n-1}) \\ &= \frac{\alpha}{2} \omega_{n-1} r_i^{n-1} + o(r_i^{n-1}), \end{aligned}$$

which contradicts (31) for i large enough. □

4. Existence theorems

This section is devoted to the proof of existence of solutions to the prescribed mean curvature equation (PMC), that we recall here:

$$\operatorname{div} Tu(x) = H(x), \quad x \in \Omega.$$

In what follows we show that the weak regularity assumption, i.e. the validity of (5) and (6), coupled with the necessary condition (1) is enough to ensure existence of solutions to (PMC).

We will follow the argument of [32, 34], which is based on the minimization of the functional

$$\mathcal{J}[u] = \int_{\Omega} \sqrt{1 + |\nabla u|^2} \, dx + \int_{\Omega} Hu \, dx + \int_{\partial\Omega} |u - \varphi| \, d\mathcal{H}^{n-1}, \quad (32)$$

defined on $BV(\Omega)$, for a given $\varphi \in L^1(\partial\Omega)$. Note that the Euler–Lagrange equation of \mathcal{J} , obtained by perturbations with compact support in Ω , is precisely equation (PMC). By Theorem 2.6, the last term in (32) is well-defined.

In the existence proof we will have first to discuss the easier *non-extremal case*, in which the necessary condition (1) holds for the domain Ω as well, and then the more involved *extremal case*, that is when (3) is satisfied.

First we need some preliminary results. The first one shows how to extend the necessary condition (1) to all measurable $A \subset \Omega$ such that $0 < |A| < |\Omega|$.

Proposition 4.1. *Let $\Omega \subset \mathbb{R}^n$ be a domain satisfying condition (6). Assume that the necessary condition (1) holds for every $A \subset\subset \Omega$, then it also holds for every $A \subset \Omega$ such that $0 < |A| < |\Omega|$.*

Proof. Let us fix a measurable set $A \subset \Omega$ with $0 < |A| < |\Omega|$ and finite perimeter. By Theorem 2.8 there exists a sequence $\{\Omega_j\}_{j \in \mathbb{N}}$ of relatively compact, smooth open subsets of Ω , such that $|\Omega \setminus \Omega_j| \rightarrow 0$ and $P(\Omega_j) \rightarrow P(\Omega)$ as $j \rightarrow \infty$. Now take $A_j = A \cap \Omega_j$ and notice that $A_j \subset\subset \Omega$, $P(A_j) < +\infty$, and $A_j \rightarrow A$ in L^1 as $j \rightarrow \infty$. Since

$$P(A_j) + P(A \cup \Omega_j) \leq P(A) + P(\Omega_j),$$

and owing to the fact that $A \cup \Omega_j \rightarrow \Omega$ in L^1 as $j \rightarrow \infty$, we deduce that

$$\begin{aligned} P(A) &\leq \liminf_j P(A_j) \leq \limsup_j P(A_j) \leq \limsup_j \left(P(A) + P(\Omega_j) - P(A \cup \Omega_j) \right) \\ &= P(A) + P(\Omega) - \liminf_j P(A \cup \Omega_j) \leq P(A) + P(\Omega) - P(\Omega) = P(A), \end{aligned}$$

which proves that

$$\lim_j P(A_j) = P(A). \quad (33)$$

Now we observe that $P(A; \Omega) > 0$, which follows from the connectedness of Ω coupled with the fact that $0 < |A| < |\Omega|$. Therefore owing to (8) we can assume that $P(A_j; \Omega_{j_0}) \geq c > 0$ for a suitably large j_0 and for all $j \geq j_0$, which means that

$$\begin{aligned} \left| \int_{A_j} H \, dx \right| &= \left| \int_{\partial^* A_j} \langle Tu, \nu \rangle \, d\mathcal{H}^{n-1} \right| \leq P(A_j; \mathbb{R}^n \setminus \Omega_{j_0}) \\ &\quad + \int_{\partial^* A_j \cap \Omega_{j_0}} |\langle Tu, \nu \rangle| \, d\mathcal{H}^{n-1} \\ &\leq P(A_j; \mathbb{R}^n \setminus \Omega_{j_0}) + \alpha P(A_j; \Omega_{j_0}) = P(A_j) - (1 - \alpha)c, \end{aligned}$$

where $\alpha < 1$ is the supremum of $|\langle Tu, \nu \rangle|$ on Ω_{j_0} . Since $|A_j| \rightarrow |A|$ as $j \rightarrow \infty$, by the necessary condition written for A_j , and passing to the limit as $j \rightarrow \infty$, we get by (33)

$$\left| \int_A H \, dx \right| \leq P(A) - (1 - \alpha)c < P(A), \tag{34}$$

whence the conclusion follows. □

The next lemma corresponds to [34, Lemma 1.1], thus we omit its proof.

Lemma 4.2. *Let Ω be a domain such that $|\int_A H \, dx| < P(A)$ holds for all $A \subset \Omega$ with the property that $|A| > 0$. Then there exists $\varepsilon_0 > 0$ such that the stronger inequality*

$$\left| \int_A H \, dx \right| \leq (1 - \varepsilon_0)P(A)$$

holds for all such A .

Theorem 4.3. (Existence, non-extremal case) *Let Ω be a weakly regular domain. If the necessary condition (1) holds also for Ω , that is, we have the non-extremal condition (2), that is*

$$\left| \int_\Omega H \, dx \right| < P(\Omega),$$

then the functional \mathcal{J} defined in (32) is minimized in $BV(\Omega)$.

Proof. Fix a ball B containing Ω and extend the function H to 0 in $B \setminus \Omega$. Fix a function $\Phi \in W_0^{1,1}(B)$ such that $\Phi = \varphi$ on $\partial\Omega$ (this can be done according to Theorem 2.6). Then minimizing \mathcal{J} on $BV(\Omega)$ is equivalent to minimizing $\tilde{\mathcal{J}}$ defined as

$$\tilde{\mathcal{J}} : u \mapsto \int_B \sqrt{1 + |\nabla u|^2} \, dx + \int_B H u \, dx,$$

in $K = \{u \in BV(B) \mid u = \Phi \text{ in } B \setminus \Omega\}$, which is a closed subset of $BV(B)$. Owing to Proposition 4.1 and by the assumption on Ω we can apply Lemma 4.2 and get the lower bound

$$\int_\Omega H u \, dx \geq -(1 - \varepsilon_0) \int_B |Du| - c \int_{\partial\Omega} |\varphi| \, d\mathcal{H}^{n-1}$$

for some $\varepsilon_0 > 0$, whence

$$\tilde{\mathcal{J}}[u] \geq \varepsilon_0 \int_B |Du| \, dx - c \int_{\partial\Omega} |\varphi| \, d\mathcal{H}^{n-1}. \tag{35}$$

Exploiting Poincaré’s inequality on the ball B one finally shows the coercivity of $\tilde{\mathcal{J}}$ in $L^1(\Omega)$. Since it is also lower semi-continuous within respect to the L^1 -norm we infer the existence of a minimizer of $\tilde{\mathcal{J}}$ in K , hence of a minimizer of \mathcal{J} in $BV(\Omega)$. □

In order to prove the existence of minimizers in the extremal case (3), following [45] we introduce the notion of *generalized solution* of (PMC). For technical reasons, we consider the epigraph of u instead of its subgraph, therefore the definition is slightly offset from the one in [45] (but of course equivalent up to changing the minus sign in (36)).

Definition 4.4. A function $u : \Omega \rightarrow [-\infty, +\infty]$ is said to be a generalized solution to (PMC) if the epigraph of u

$$U = \{(x, y) \in \Omega \times \mathbb{R} : y > u(x)\},$$

minimizes the functional

$$P(U) - \int_U H \, dx \, dy, \tag{36}$$

locally in $\Omega \times \mathbb{R}$.

It is clear that any classical solution to (PMC) is also a generalized solution. Moreover, any generalized solution of (PMC) can be shown to satisfy some key properties, that we collect in the following proposition (see [34] and [44, 45] for the proof).

Proposition 4.5. *Let u be a generalized solution of (PMC) and define $N_{\pm} = \{x \in \Omega : u(x) = \pm\infty\}$. Then the following properties hold.*

- (i) *If $x \in N_{\pm}$ then $|N_{\pm} \cap B_r(x)| > 0$ for all $r > 0$.*
- (ii) *The set N_{\pm} minimizes the functional*

$$E \mapsto P(E) \pm \int_E H \, dx$$

locally in Ω .

- (iii) *The function u is smooth on $\Omega \setminus (N_+ \cup N_-)$.*
- (iv) *Given a sequence $\{u_k\}$ of generalized solutions of (PMC), then up to subsequences the epigraphs U_k of u_k converge to an epigraph U of a function u locally in $L^1(\Omega \times \mathbb{R})$, moreover u is a generalized solution of (PMC).*
- (v) *If u is locally bounded, then u is a classical solution of (PMC).*

The next lemma is a straightforward adaptation of [34, Lemma 1.2]. The proof is the same up to choosing a sequence $\{\Omega_j\}_j$ as provided by Theorem 2.8 with $\varepsilon = 1/j$.

Lemma 4.6. *Let Ω and $H(x)$ be such that (1), (6) and (3) hold. Let $E \subset \Omega$ be a set of finite perimeter minimizing the functional*

$$P(E) - \int_E H \, dx$$

locally in Ω . Then either $E = \emptyset$ or $E = \Omega$, up to null sets.

We now come to the existence of solutions of (PMC) in the extremal case.

Theorem 4.7. (Existence, extremal case) *Let Ω be a weakly regular domain. Assume that (1) is satisfied and that the extremal condition (3) holds. Then there exists a solution u of (PMC).*

Proof. By Theorem 2.8 we find a sequence of smooth, connected sets $\Omega_j \subset\subset \Omega$, such that $|\Omega \setminus \Omega_j| \rightarrow 0$ and $P(\Omega_j) \rightarrow P(\Omega)$ as $j \rightarrow +\infty$. Since (1) holds for any $A \subset \Omega_j$ (and in particular for $A = \Omega_j$), in virtue of Theorem 4.3 (existence in

the non-extremal case) we find a minimizer $u_j \in BV(\Omega_j)$ of \mathcal{J} restricted to $BV(\Omega_j)$, as every Ω_j satisfies (5). Setting

$$t_j = \inf \left\{ t : |\{x \in \Omega_j : u_j(x) \geq t\}| \leq |\Omega_j|/2 \right\}$$

we obtain

$$\min(|\{x \in \Omega_j : u_j(x) \geq t_j\}|, |\{x \in \Omega_j : u_j(x) \leq t_j\}|) \geq |\Omega_j|/2 \geq |\Omega|/4$$

for all j large enough. Therefore, we can consider the sequence of vertically translated functions $\{u_j(x) - t_j\}_j$ defined for $x \in \Omega_j$, and relabel it as $\{u_j\}_j$, so that

$$\min(|\{x \in \Omega_j : u_j(x) \geq 0\}|, |\{x \in \Omega_j : u_j(x) \leq 0\}|) \geq |\Omega|/4 \quad (37)$$

for all j large enough. Then, by applying Proposition 4.5 (iv) on Ω_{j_0} for any fixed $j_0 \in \mathbb{N}$, and by a diagonal argument, we infer that u_j locally converges up to subsequences to a generalized solution u as $j \rightarrow \infty$, in the sense that the epigraph U_j locally converges to the epigraph of u in $L^1_{loc}(\Omega \times \mathbb{R})$ as $j \rightarrow \infty$. Let us set $N_{\pm} = \{x \in \Omega : u(x) = \pm\infty\}$ as in Proposition 4.5. We claim that N_{\pm} are both empty, which in turn implies by Proposition 4.5 (v) that u is a classical solution of (PMC). Indeed by Proposition 4.5 (ii) the set N_- minimizes the functional $P(E) - \int_E H dx$ defined for $E \subset \Omega$, thus by Lemma 4.6 we have either $N_- = \emptyset$ or $N_- = \Omega$. Similarly, the set $\Omega \setminus N_+$ minimizes $P(E) - \int_E H dx$ (this follows from the fact that N_+ minimizes $P(E) + \int_E H dx$), hence either $N_+ = \Omega$ or $N_+ = \emptyset$. By (37) we conclude that $N_{\pm} = \emptyset$, which proves our claim. \square

5. Characterization of extremality

We have seen in the previous section that, given a domain Ω and a prescribed mean curvature function H , the condition (1) is necessary and sufficient for the existence of solutions to (PMC), however the proof of this fact is different depending on the validity or not of the *extremality condition* (3) (compare Theorems 4.3 and 4.7). While in the non-extremal case the existence of solutions is genuinely variational, in the extremal case one recovers a solution as a limit of variational solutions defined on subdomains. Since extremality arises in physical models of capillarity for perfectly wetting fluids, the uniqueness and the stability of solutions with respect to suitable perturbations of the domain are of special interest in this case.

In [33] Giusti showed that, assuming C^2 regularity of $\partial\Omega$ and (1), the extremality condition (3) is equivalent to a series of facts, and in particular to the uniqueness of the solution of (PMC) up to vertical translations.

Here we obtain essentially the same result only assuming that Ω is weakly regular. Before stating our main result, we present a list of properties using the same labels as those appearing in [34].

(E) (*Extremality*) The pair (Ω, H) satisfies (3), i.e., $|\int_{\Omega} H dx| = P(\Omega)$.

(U) (*Uniqueness*) The solution of (PMC) is unique up to vertical translations.

- (M) (*Maximality*) Ω is maximal, i.e. no solution of (PMC) can exist in any domain strictly containing Ω .
- (V) (*weak Verticality*) There exists a solution u of (PMC) which is weakly vertical at $\partial\Omega$, i.e.

$$[Tu \cdot \nu] = 1 \quad \mathcal{H}^{n-1}\text{-a.e. on } \partial\Omega,$$

where $[Tu \cdot \nu]$ is the weak normal trace of Tu on $\partial\Omega$.

- (V') (*integral Verticality*) There exists a solution u of (PMC) and a sequence $\{\Omega_i\}_i$ of smooth subdomains, such that $\Omega_i \subset\subset \Omega$, $|\Omega \setminus \Omega_i| \rightarrow 0$, $P(\Omega_i) \rightarrow P(\Omega)$, and

$$\lim_{i \rightarrow \infty} \int_{\partial\Omega_i} Tu(x) \cdot \nu \, d\mathcal{H}^{n-1} = P(\Omega),$$

as $i \rightarrow \infty$.

Then we come to the main result of this section.

Theorem 5.1. *Let Ω and H be given, such that Ω is weakly regular and (1) holds. Then the properties (E), (U), (M), (V) and (V') are equivalent.*

Before proving Theorem 5.1 some further comments about properties (V) and (V') above are in order. In [34] the property (V) is stated in the stronger, pointwise form $Tu(x) = \nu(x)$ for all $x \in \partial\Omega$ (moreover $\partial\Omega$ is assumed of class C^2 , hence Tu can be continuously extended on $\partial\Omega$ owing to well-known regularity results, see [21]) while (V') is stated by using the one-parameter family of inner parallel sets (which is again well-defined owing to the C^2 -smoothness of $\partial\Omega$).

The Maximum Principle Lemma that we state hereafter has been originally proved in [25] and then in [34]. We remark that it remains valid under the weaker assumptions guaranteeing the interior smooth approximation property, in the sense of Theorem 2.8.

Lemma 5.2. (Maximum Principle) *Let $\Omega \subset \mathbb{R}^n$ be open, bounded, connected and weakly regular. Let u and v be two functions of class $C^2(\Omega)$, such that $\operatorname{div}(Tu) \leq \operatorname{div}(Tv)$ in Ω . Assume that $\partial\Omega = \Gamma_1 \cup \Gamma_2$ with Γ_1 relatively open in $\partial\Omega$, and $u, v \in C^0(\Omega \cup \Gamma_1)$ with $u \geq v$ on Γ_1 . Assume further that*

$$\lim_{i \rightarrow \infty} \int_{\partial\Omega_i \setminus A} (1 - Tu \cdot \nu) \, d\mathcal{H}^{n-1} = 0$$

for every open set $A \supset \Gamma_1$, where $\{\Omega_i\}_{i \in \mathbb{N}}$ is a sequence of smooth and relatively compact open subsets of Ω , such that $|\Omega \setminus \Omega_i| \rightarrow 0$ and $P(\Omega_i) \rightarrow P(\Omega)$ as $i \rightarrow \infty$. Then

- (a) if $\Gamma_1 \neq \emptyset$ then $u \geq v$ in Ω ;
- (b) if $\Gamma_1 = \emptyset$ then $u = v + c$.

Proof. In order to prove case (a) we first assume that $u > v$ on Γ_1 . By the Gauss–Green formula on Ω_i , for any positive function $\varphi \in W^{1,\infty}(\Omega_i)$ one

obtains

$$\begin{aligned} \int_{\Omega_i} (Tu - Tv) \cdot \nabla \varphi &= - \int_{\Omega_i} \varphi (\operatorname{div} Tu - \operatorname{div} Tv) + \int_{\partial\Omega_i} \varphi (Tu - Tv) \cdot \nu \\ &\geq \int_{\partial\Omega_i} \varphi (Tu - Tv) \cdot \nu \geq \int_{\partial\Omega_i} \varphi (Tu \cdot \nu - 1). \end{aligned}$$

Fix a positive constant $M > 0$ and define the function $\varphi_M(x) = \max\left(0, \min(v - u, M)\right)$. Of course $\varphi_M \in W^{1,\infty}(\Omega_i)$ for all i and $0 \leq \varphi_M \leq M$. Moreover, we can find an open set A containing Γ_1 and such that $\varphi_M = 0$ on $A \cap \Omega$. We also notice that

$$(Tu - Tv) \cdot \nabla \varphi_M = \begin{cases} (Tu - Tv) \cdot (\nabla v - \nabla u) & \text{if } 0 < v - u < M, \\ 0 & \text{elsewhere,} \end{cases}$$

hence by a straightforward computation

$$(Tu - Tv) \cdot \nabla \varphi_M \leq (|\nabla v| - |\nabla u|) \left(\frac{|\nabla u|}{\sqrt{1 + |\nabla u|^2}} - \frac{|\nabla v|}{\sqrt{1 + |\nabla v|^2}} \right) \leq 0.$$

Consequently, we obtain

$$\int_{\partial\Omega_i \setminus A} \varphi_M (Tu \cdot \nu - 1) \leq \int_{\Omega_i \setminus A} (Tu - Tv) \cdot \nabla \varphi_M \leq 0,$$

thus by taking the limit as $i \rightarrow \infty$ we find

$$\int_{\Omega} (Tu - Tv) \cdot \nabla \varphi_M = 0$$

for all $M > 0$. Therefore, setting $\varphi = \max(v - u, 0)$ we find $\nabla \varphi = 0$ on Ω , which means that φ is constant on Ω . However, since $\varphi = 0$ on $A \cap \Omega$ we deduce that $\varphi = 0$, hence that $u \geq v$, on the whole Ω . The full proof of case (a) is then completed by considering $v_\varepsilon = v + \varepsilon$ in place of v and then letting $\varepsilon \rightarrow 0^+$.

Finally, for the proof of case (b) we fix $x_0 \in \Omega$ and assume $v(x_0) = u(x_0) + 1$ up to a vertical translation. Arguing exactly as in the proof of case (a), we end up with φ constant on Ω , where as before we set $\varphi = \max(v - u, 0)$. Since $\varphi = \varphi(x_0) = 1$ we conclude that $v = u + 1$ on Ω , as wanted. \square

We finally come to the proof of Theorem 5.1.

Proof of Theorem 5.1. We shall split the proof in five steps.

Step one: $(E) \Rightarrow (V')$. Owing to (E) we have

$$\begin{aligned} P(\Omega) &\stackrel{(E)}{=} \int_{\Omega} H \, dx = \lim_{i \rightarrow \infty} \int_{\Omega_i} H \, dx = \lim_{i \rightarrow \infty} \int_{\Omega_i} \operatorname{div}(Tu) \, dx \\ &= \lim_{i \rightarrow \infty} \int_{\partial\Omega_i} Tu(x) \cdot \nu \, d\mathcal{H}^{n-1}, \end{aligned}$$

which implies (V') .

Step two: (E) ⇔ (M). Let us start by showing (E)⇒(M). We argue by contradiction and suppose there exists a solution u of (PMC) defined on $\tilde{\Omega} \supsetneq \Omega$. Then Proposition 4.1 gives

$$\left| \int_{\Omega} H \, dx \right| < P(\Omega),$$

which immediately contradicts (E). Let us now show the implication (M)⇒(E). Again by contradiction we assume that

$$\left| \int_A H \, dx \right| < P(A)$$

for all $A \subset \Omega$. By Lemma 4.2 there exists $\varepsilon_0 > 0$ such that

$$\left| \int_A H \, dx \right| < (1 - \varepsilon_0)P(A) \tag{38}$$

for all $A \subset \Omega$. Now we claim that (compare with Lemma 2.1 in [34]) given a ball B such that $\Omega \subset\subset B$, for all $0 < \varepsilon < \varepsilon_0$ one can find an open set $\Omega_\varepsilon \subset B$ with smooth boundary, such that $\Omega \subset\subset \Omega_\varepsilon$ and

$$\left| \int_A H \, dx \right| < (1 - \varepsilon)P(A), \quad \forall A \subset \Omega_\varepsilon. \tag{39}$$

Of course, the validity of (39) would allow us to apply Theorem 4.3 on Ω_ε , which in turn would contradict our assumption (M). In order to show (39) we argue again by contradiction, i.e., we assume that there exists $\varepsilon \in (0, \varepsilon_0)$ such that, for every U with smooth boundary satisfying $\Omega \subset\subset U$, one can find $A \subset U$ for which (39) fails. In particular, for every $k \in \mathbb{N}$ we may choose a suitable U_k as specified below, such that $\Omega \subset\subset U_k$, $|U_k \setminus \Omega| < 1/k$, ∂U_k is smooth and there exists $A_k \subset U_k$ for which

$$\left| \int_{A_k} H \, dx \right| \geq (1 - \varepsilon)P(A_k) \tag{40}$$

holds. By (40) we have that

$$P(A_k) \leq \frac{|B| \sup_B |H|}{1 - \varepsilon} \quad \forall k \in \mathbb{N},$$

hence we can extract a not relabeled subsequence A_k converging to some $A \subset B$ in L^1 . On the other hand, since $|A_k \setminus \Omega| \leq |U_k \setminus \Omega| \rightarrow 0$ as $k \rightarrow \infty$, we infer that $A \subset \Omega$ up to null sets. By (40), by the lower semi-continuity of the perimeter and by the continuity of the term $\int_{A_k} H \, dx$ with respect to L^1 -convergence, we conclude that

$$\left| \int_A H \, dx \right| \geq (1 - \varepsilon)P(A)$$

which is in contrast with (38). We are left to prove that such a sequence U_k exists. To this aim we consider the open set $V = B \setminus \bar{\Omega}$ and notice that $P(V) = P(B) + P(\Omega) = \mathcal{H}^{n-1}(\partial B) + \mathcal{H}^{n-1}(\partial \Omega) = \mathcal{H}^{n-1}(\partial V)$ owing to the assumption on Ω . We can now apply Theorem 2.8 to V with $\delta_k = \min(\text{dist}(\partial B, \partial \Omega)/3, 1/k)$ and set $U_k = B \setminus (V_{\delta_k} \cup \mathcal{N}_{2\delta_k}(\partial B))$. Thanks to (8) we find that ∂U_k is smooth, $\Omega \subset\subset U_k$ and $|U_k \setminus \Omega| < \delta_k \leq 1/k$, as wanted.

Step three: $(V') \Rightarrow (U)$. We consider two solutions u, v of (PMC), then if we take $\Gamma_1 = \emptyset$ and thanks to the property $P(\Omega_i) \rightarrow P(\Omega)$ as $i \rightarrow \infty$, we infer that the assumptions of Lemma 5.2(b) are satisfied. Consequently there exists a constant $c \in \mathbb{R}$ such that $u = v + c$.

Step four: $(U) \Rightarrow (E)$. Let u be the unique solution of $\operatorname{div}(Tu) = H$ on Ω , up to vertical translations. By contradiction we suppose that

$$\int_{\Omega} H \, dx < P(\Omega).$$

Arguing as in Step two we find a bounded and smooth domain $\tilde{\Omega} \supsetneq \Omega$ for which (1) holds. By Theorems 4.3 and 4.7 there exists a solution \tilde{u} of $\operatorname{div}(T\tilde{u}) = H$ on $\tilde{\Omega}$. Then (U) implies the existence of $t \in \mathbb{R}$ such that $u = \tilde{u} + t$ on Ω . By internal regularity of \tilde{u} , we infer that $u \in C^1(\bar{\Omega})$. Fix now a function $\varphi \in C^2(\mathbb{R}^n)$ such that

$$\mathcal{H}^{n-1}(\{x \in \partial\Omega : \varphi(x) - u(x) \neq s\}) > 0 \quad \forall s \in \mathbb{R}. \tag{41}$$

The choice of φ satisfying (41) can be easily made as follows: if u is constant on $\partial\Omega$, then one can choose any smooth function φ taking different values on two distinct points of $\partial\Omega$; conversely, if u is not constant on $\partial\Omega$ then one can take $\varphi = 0$. Now we consider a minimizer w of the functional

$$\int_{\Omega} \sqrt{1 + |Dw|^2} + \int_{\Omega} Hw + \int_{\partial\Omega} |w - \varphi| \, d\mathcal{H}^{n-1},$$

then w necessarily satisfies (PMC). By the assumed uniqueness up to translations one has that $w = u + s$ for some $s \in \mathbb{R}$. Then it follows that

$$|Tu(x_0)| = |Tw(x_0)| < 1. \tag{42}$$

Moreover by (41) we have that $w \neq \varphi$ on some set $K \subset \partial^*\Omega$ with $\mathcal{H}^{n-1}(K) > 0$. Fix now a point $x_0 \in K$ and assume without loss of generality that $\varphi(x_0) > w(x_0)$. Set now $\mathcal{C} = \Omega \times \mathbb{R}$, $p_0 = (x_0, w(x_0)) \in \partial\mathcal{C}$, and notice that by the continuity of w and φ on $\partial\Omega$ there exists $R > 0$ such that the subgraph of φ contains the ball $B_R(p_0) \subset \mathbb{R}^{n+1}$. Owing to the choice of $B_R(p_0)$, the epigraph

$$W := \{p = (x, y) \in \mathcal{C} : y > w(x)\}$$

necessarily minimizes the functional

$$P(W; B_R(p_0)) - \int_{W \cap B_R(p_0)} H$$

with obstacle $\mathbb{R}^{n+1} \setminus \mathcal{C}$ inside $B_R(p_0)$. In other words, for any set U that coincides with W outside the set $A := B_R(p_0) \cap \bar{\mathcal{C}}$, one has

$$P(W; B_R(p_0)) - \int_{W \cap B_R(p_0)} H \leq P(U; B_R(p_0)) - \int_{U \cap B_R(p_0)} H. \tag{43}$$

It is then easy to show that W is a (Λ, R) -perimeter minimizer in $\bar{\mathcal{C}}$ (see Definition 3.4), where R is the radius of the ball defined above and $\Lambda =$

$\sup_{\Omega} |H|$. Indeed for any ball $B_r \subset B_R(p_0)$ and any set U such that $U\Delta W \subset \subset B_r \cap \bar{C}$, by (43) one has that

$$\begin{aligned} P(W; B_r) &= P(W; B_R(p_0)) - P(W; B_R(p_0) \setminus \bar{B}_r) \\ &\leq P(U; B_R(p_0)) - P(U; B_R(p_0) \setminus \bar{B}_r) - \int_{B_R(p_0)} H(\chi_U - \chi_W) \\ &\leq P(U; B_r) + \sup_{\Omega} |H| |U\Delta W|, \end{aligned}$$

which proves the (Λ, R) -minimality of W in \bar{C} . Then by Theorem 3.5 we infer that $\nu_W(p_0) = \nu_C(p_0)$, which contradicts (42).

Step five: (V) and (V') are equivalent. We can consider the sequence Ω_j of Theorem 2.8 and apply Theorem 3.2 to get

$$\begin{aligned} \int_{\Omega \setminus \Omega_j} H(x) dx &= \int_{\Omega \setminus \Omega_j} \operatorname{div} Tu(x) dx \\ &= \int_{\partial\Omega} [Tu \cdot \nu] d\mathcal{H}^{n-1} - \int_{\partial\Omega_j} Tu \cdot \nu_j d\mathcal{H}^{n-1}. \end{aligned} \quad (44)$$

Now, observing that the left-hand side of (44) is infinitesimal as $j \rightarrow \infty$ the equivalence between (V) and (V') is immediate.

The proof is finally completed by combining the previous five steps. \square

We now show a well-known consequence of Lemma 5.2, which can be obtained by arguing as in Step two of the proof of Theorem 5.1.

Proposition 5.3. *Assume that u is a solution of (PMC) on Ω and that either (V) or (V') holds. Then u is bounded from below.*

Proof. Let B denote a ball compactly contained in Ω and consider the open set $S = \Omega \setminus \bar{B}$. By Lemma 4.2 and arguing as in Step two of the proof of Theorem 5.1 we find a solution w of (PMC) which is of class $C^1(\bar{S})$. Since in particular $u \in C^2(\bar{B})$ we can assume that $w \leq u$ on ∂B up to a vertical translation, hence by Lemma 5.2(a) we deduce that $w \leq u$ on S , which gives the conclusion at once. \square

We conclude the section with some remarks about the stability of solutions of (PMC) in the extremal case. One might ask whether or not there exists some perturbation $(\Omega_\varepsilon, H_\varepsilon)$ of an extremal pair (Ω, H) , such that $(\Omega_\varepsilon, H_\varepsilon)$ satisfies the necessary condition (1) and the solution u_ε of (PMC) on Ω_ε is in a suitable sense a small perturbation of u up to translations, as soon as ε is small. The following proposition contains a result in this direction.

Proposition 5.4. *(Stability) Let $\{\Omega_j\}_j$ be a sequence of bounded domains and $\{H_j\}_j$ a sequence of Lipschitz functions, such that Ω_j is weakly regular and the pair (Ω_j, H_j) is extremal. Assume moreover that $\Omega_j \rightarrow \Omega_\infty$ in L^1 and $P(\Omega_j) \rightarrow P(\Omega_\infty)$, as $j \rightarrow \infty$, with Ω_∞ weakly regular, and that H_j uniformly converges to H_∞ such that the pair $(\Omega_\infty, H_\infty)$ is extremal as well. Then the sequence of unique (up to translations) solutions $\{u_j\}_j$ to the (PMC) problem for the pair (Ω_j, H_j) converges to a solution u_∞ of (PMC) for the pair $(\Omega_\infty, H_\infty)$, in the sense of the L^1_{loc} -convergence of the epigraphs.*




FIGURE 1. The “Swiss cheese” set $O_{a,\delta,\epsilon}$ constructed in Example 5.5

Proof. Due to our hypotheses, the existence of a solution u_j to (PMC) for the pair (Ω_j, H_j) (also for $j = \infty$) is guaranteed by Theorem 4.7. Arguing as in Theorem 4.7, for any j large enough we can find a suitable t_j such that the translated solution $u_j + t_j$ which we just rename u_j satisfies

$$\min \left(|\{x \in \Omega_j : u_j(x) \geq 0\}|, |\{x \in \Omega_j : u_j(x) \leq 0\}| \right) \geq |\Omega|/4.$$

Then we find that the epigraphs U_j of u_j converge in $L^1_{loc}(\mathbb{R}^{n+1})$ to a set U_∞ which is the epigraph of a classical solution u_∞ defined on Ω_∞ . By Theorem 5.1 we have that $u_\infty = u_\infty$ up to a translation, thus the thesis follows. \square

In the recent paper [40], an explicit example of an extremal pair (Ω, H) and of a sequence of extremal pairs (Ω_j, H_j) satisfying the hypotheses of Proposition 5.4 is constructed, for the special case $H_j = P(\Omega_j)/|\Omega_j|$, by removing a sequence of smaller and smaller disks from the unit disk in \mathbb{R}^2 , in such a way that it looks like a sort of Swiss cheese with holes accumulating towards a portion of its boundary (see Fig. 1 and Example 5.5 below; for a more complete discussion we refer to [40]).

This shows the following, remarkable fact: while a generic small and smooth perturbation of the unit disk may produce a dramatic change in the capillary solution (and even end up with non-existence of a solution), there exist some non-smooth perturbations that, instead, preserve both existence and stability.

Example 5.5. Let $0 < \delta < \epsilon < 1$ and $a > 1$ be fixed. For $i \geq 1$ and $j = 1, \dots, i$ we set

$$\rho_{ij} = 1 - \frac{\epsilon}{a^{i^2+j}}, \quad r_{ij} = \frac{\delta}{a^{2i^2+2j}}, \quad \theta_{ij} = \frac{\pi}{2} \frac{j}{i+1},$$

Then we define

$$O_{a,\delta,\epsilon} = B_1 \setminus \bigcup_{i,j} \overline{B_{ij}},$$

where $B_1 \subset \mathbb{R}^2$ is the unit disk centered at the origin, and

$$B_{ij} := B_{r_{ij}}((\rho_{ij} \cos(\theta_{ij}), \rho_{ij} \sin(\theta_{ij}))$$

(see Fig. 1). We prove in [40] that for a suitable choice of parameters a, δ, ϵ the open set $O_{a,\delta,\epsilon}$ fulfils the hypotheses of Theorem 5.1.

One can then build a sequence of non-smooth perturbations of the unit disk by simply filling one hole of the Swiss cheese at a time: indeed this operation creates a sequence of subdomains of the unit disk that satisfy the hypotheses of Proposition 5.4 with the choice $H_j = P(\Omega_j)/|\Omega_j|$ and with $(B_1, 2)$ as the limit extremal pair.

References

- [1] Anzellotti, G.: Pairings between measures and bounded functions and compensated compactness. *Ann. Mat. Pura Appl.* **135**(1), 293–318 (1983)
- [2] Anzellotti, G., Giaquinta, M.: Funzioni bv e tracce. *Rend. Sem. Mat. Univ. Padova* **60**, 1–21 (1978)
- [3] Beck, L., Schmidt, T.: Convex duality and uniqueness for bv -minimizers. *J. Funct. Anal.* **268**(10), 3061–3107 (2015)
- [4] Burago, Y., Maz'ya, V.: *Potential Theory and Function Theory for Irregular Regions*. Translated from Russian. *Seminars in Mathematics, V. A.* Translated from Russian. *Seminars in Mathematics, V.A.* Steklov Mathematical Institute, Leningrad, vol. 3. Consultants Bureau, New York (1969)
- [5] Caffarelli, L.A., Friedman, A.: Regularity of the boundary of a capillary drop on an inhomogeneous plane and related variational problem. *Rev. Mat. Iberoam.* **1**(1), 61–84 (1985)
- [6] Caffarelli, L.A., Mellet, A.: Capillary drops: contact angle hysteresis and sticking drops. *Calc. Var. PDE* **29**(2), 141–160 (2007)
- [7] Chen, G.-Q., Frid, H.: Divergence-measure fields and hyperbolic conservation laws. *Arch. Rational Mech. Anal.* **147**(2), 89–118 (1999)
- [8] Chen, G.-Q., Frid, H.: Extended divergence-measure fields and the euler equations for gas dynamics. *Commun. Math. Phys.* **236**(2), 251–280 (2003)
- [9] Chen, G.-Q., Torres, M.: Divergence-measure fields, sets of finite perimeter, and conservation laws. *Arch. Rational Mech. Anal.* **175**(2), 245–267 (2005)
- [10] Chen, G.-Q., Torres, M., Ziemer, W.P.: Gauss–Green theorem for weakly differentiable vector fields, sets of finite perimeter, and balance laws. *Commun. Pure Appl. Math.* **62**(2), 242–304 (2009)
- [11] Comi, G.E., Payne, K.R.: On locally essentially bounded divergence measure fields and sets of locally finite perimeter. *Adv. Calc. Var.* (2017). <https://doi.org/10.1515/acv-2017-0001>
- [12] Comi, G.E., Torres, M.: One-sided approximation of sets of finite perimeter. *Rend. Lincei Mat. Appl.* **28**(1), 181–190 (2017)
- [13] Concus, P., Finn, R.: On the behavior of a capillary surface in a wedge. *Proc. Natl. Acad. Sci.* **63**(2), 292–299 (1969)

- [14] Concus, P., Finn, R.: On capillary free surfaces in a gravitational field. *Acta Math.* **132**, 207–223 (1974)
- [15] Concus, P., Finn, R.: On capillary free surfaces in the absence of gravity. *Acta Math.* **132**, 177–198 (1974)
- [16] Concus, P., Finn, R.: Exotic containers for capillary surfaces. *J. Fluid Mech.* **224**, 383–394 (1991)
- [17] Crasta, G., De Cicco, V.: Anzellotti’s pairing theory and the Gauss–Green theorem. Preprint [arXiv:1708.00792v1](https://arxiv.org/abs/1708.00792v1) (2017)
- [18] De Giorgi, E.: *Frontiere orientate di misura minima*. Seminario di Matematica della Scuola Normale Superiore di Pisa, 1960–61. Editrice Tecnico Scientifica, Pisa (1961)
- [19] De Philippis, G., Maggi, F.: Regularity of free boundaries in anisotropic capillarity problems and the validity of Young’s law. *Arch. Rational Mech. Anal.* **216**(2), 473–568 (2015)
- [20] Emmer, M.: Esistenza, unicità e regolarità nelle superfici di equilibrio nei capillari. *Ann. Univ. Ferrara* **18**(1), 79–94 (1973)
- [21] Emmer, M.: Superfici di curvatura media assegnata con ostacolo. *Ann. di Mat. Pura ed Appl. Ser. Quarta* **109**, 371–389 (1976)
- [22] Federer, H.: The Gauss–Green theorem. *Trans. Amer. Math. Soc.* **58**, 44–76 (1945)
- [23] Federer, H.: A note on the Gauss–Green theorem. *Proc. Am. Math. Soc.* **9**, 447–451 (1958)
- [24] Federer, H.: *Geometric Measure Theory*. Die Grundlehren der mathematischen Wissenschaften, Band 153. Springer, New York (1969)
- [25] Finn, R.: Remarks relevant to minimal surfaces, and to surfaces of prescribed mean curvature. *J. d’Anal. Math.* **14**, 139–160 (1965)
- [26] Finn, R.: *Equilibrium Capillary Surfaces*, 1st edn. Springer, New York (1986)
- [27] Finn, R.: Nonuniqueness and uniqueness of capillary surfaces. *Manuscr. Math.* **61**(3), 347–372 (1988)
- [28] Gauss, C.F.: *Principia generalia theoriae figurae fluidorum in statu aequilibrii*. *Comment. Soc. Regiae Scient. Gottingensis Rec.* **7**, 1–53 (1830)
- [29] Gerhardt, C.: Existence, regularity, and boundary behaviour of generalized surfaces of prescribed mean curvature. *Math. Zeit.* **139**(2), 173–198 (1974)
- [30] Gerhardt, C.: On the capillarity problem with constant volume. *Ann. Sc. Norm. Sup. Pisa* **2**(2), 303–320 (1975)
- [31] Gerhardt, C.: Global regularity of the solutions to the capillarity problem. *Ann. Sc. Norm. Sup. Pisa* **3**(1), 157–175 (1976)

- [32] Giaquinta, M.: On the Dirichlet problem for surfaces of prescribed mean curvature. *Manuscr. Math.* **12**, 73–86 (1974)
- [33] Giusti, E.: Boundary value problems for non-parametric surfaces of prescribed mean curvature. *Ann. Sc. Norm. Sup. Pisa Cl.* **3**, 501–548 (1976)
- [34] Giusti, E.: On the equation of surfaces of prescribed mean curvature. *Invent. Math.* **46**, 111–137 (1978)
- [35] Lancaster, K.E.: A proof of the Concus–Finn conjecture. *Pac. J. Math.* **247**(1), 75–108 (2010)
- [36] Laplace, P.S.: *Traité de mécanique céleste: suppléments au Livre X*. Gauthier-Villars, Paris (1806)
- [37] Leonardi, G.P.: An overview on the Cheeger problem. In: *New Trends in Shape Optimization*, volume 166 of *International Series of Numerical Mathematics*, pp. 117–139. Springer Int. Publ. (2015)
- [38] Leonardi, G.P., Pratelli, A.: On the Cheeger sets in strips and non-convex domains. *Calc. Var. PDE* **55**(1), 1–28 (2016)
- [39] Leonardi, G.P., Saracco, G.: Rigidity and trace properties of divergence-measure vector fields. Preprint <https://arxiv.org/abs/1708.01393> (2017)
- [40] Leonardi, G.P., Saracco, G.: Two examples of minimal Cheeger sets in the plane. *Ann. Mat. Pura Appl.* accepted paper preprint [arXiv:1709.00851](https://arxiv.org/abs/1709.00851) (2018)
- [41] Maggi, F.: *Sets of Finite Perimeter and Geometric Variational Problems*. Cambridge Studies in Advanced Mathematics (2012)
- [42] Maniglia, S., Ambrosio, L., Crippa, G.: Traces and fine properties of a BD class of vector fields and applications. *Ann. Fac. Sci. Toulouse* **14**(4), 527–561 (2005)
- [43] Maz’ya, V.: *Sobolev Spaces with Applications to Elliptic Partial Differential Equations*. Number 342 in *Grundlehren der mathematischen Wissenschaften.*, 2nd edn. Springer, Berlin (2011)
- [44] Miranda, M.: Superfici cartesiane generalizzate. *Ann. Sc. Norm. Sup. Pisa* **18**(4), 515–542 (1964)
- [45] Miranda, M.: Superfici minime illimitate. *Ann. Sc. Norm. Sup. Pisa* **4**(4), 313–322 (1977)
- [46] Parini, E.: An introduction to the Cheeger problem. *Surv. Math. its Appl.* **6**, 9–21 (2011)
- [47] Pfeffer, W.F.: The Gauss–Green theorem in the context of Lebesgue integration. *Bull. Lond. Math. Soc.* **37**(1), 81–94 (2005)
- [48] Pfeffer, W.F., De Pauw, T.: The Gauss–Green theorem and removable sets for PDEs in divergence form. *Adv. Math.* **183**(1), 155–182 (2004)
- [49] Saracco, G.: Weighted Cheeger sets are domains of isoperimetry. *Manuscripta Math.* pp. 1–11 (2017). <https://doi.org/10.1007/s00229-017-0974-z>

- [50] Scheven, C., Schmidt, T.: BV supersolutions to equations of 1-Laplace and minimal surface type. *J. Differ. Equ.* **261**(3), 1904–1932 (2016)
- [51] Schmidt, T.: Strict interior approximation of sets of finite perimeter and functions of bounded variation. *Proc. Am. Math. Soc.* **143**(5), 2069–2084 (2014)
- [52] Simon, L.: A strict maximum principle for area minimizing hypersurfaces. *J. Differ. Geom.* **26**(2), 327–335 (1987)
- [53] Swanson, D.: Area, coarea, and approximation in $W^{1,1}$. *Ark. Mat.* **45**, 381–399 (2007)
- [54] Tam, L.-F.: On existence criteria for capillary free surfaces without gravity. *Pac. J. Math.* **125**(2), 469–485 (1986)
- [55] Tam, L.-F.: Regularity of capillary surfaces over domains with corners: borderline case. *Pac. J. Math.* **124**(2), 469–482 (1986)
- [56] Vol’pert, A.-I.: The spaces BV and quasilinear equations. *Mat. Sb.* **115**(2), 255–302 (1967)
- [57] Young, T.: An essay on the cohesion of fluids. *Philos. Trans. Roy. Soc. Lond.* **95**, 65–87 (1805)
- [58] Ziemer, W.P.: Cauchy flux and sets of finite perimeter. *Arch. Rational Mech. Anal.* **84**(3), 189–201 (1983)

Gian Paolo Leonardi
Dipartimento di Scienze Fisiche,
Informatiche e Matematiche
Università degli Studi di Modena
e Reggio Emilia
Via Campi 213/b
41125 Modena
Italy
e-mail: gianpaolo.leonardi@unimore.it

Giorgio Saracco
Department Mathematik
Universität Erlangen-Nürnberg
Cauerst. 11
91058 Erlangen
Germany
e-mail: saracco@math.fau.de

Received: 12 July 2017.

Accepted: 8 February 2018.