

INHOMOGENEOUS RANDOM GRAPHS WITH INFINITE-MEAN FITNESS VARIABLES

LUCA AVENA,^{***} *Leiden University and Università degli Studi di Firenze*

DIEGO GARLASCHELLI,^{****} *Lorentz Institute for Theoretical Physics, IMT School for Advanced Studies, INdAM-GNAMPa Istituto Nazionale di Alta Matematica*

RAJAT SUBHRA HAZRA,^{*****} *Leiden University*

MARGHERITA LALLI,^{*****} *IMT School of Advanced Studies*

Abstract

We consider an inhomogeneous Erdős–Rényi random graph ensemble with exponentially decaying random disconnection probabilities determined by an independent and identically distributed field of variables with heavy tails and infinite mean associated with the vertices of the graph. This model was recently investigated in the physics literature (Garuccio, Lalli, and Garlaschelli 2023) as a scale-invariant random graph within the context of network renormalization. From a mathematical perspective, the model fits in the class of scale-free inhomogeneous random graphs whose asymptotic geometrical features have recently attracted interest. While for this type of graph several results are known when the underlying vertex variables have finite mean and variance, here instead we consider the case of one-sided stable variables with necessarily infinite mean. To simplify our analysis, we assume that the variables are sampled from a Pareto distribution with parameter $\alpha \in (0, 1)$. We start by characterizing the asymptotic distributions of the typical degrees and some related observables. In particular, we show that the degree of a vertex converges in distribution, after proper scaling, to a mixed Poisson law. We then show that correlations among degrees of different vertices are asymptotically non-vanishing, but at the same time a form of asymptotic tail independence is found when looking at the behavior of the joint Laplace transform around zero. Moreover, we present some findings concerning the asymptotic density of wedges and triangles, and show a cross-over for the existence of dust (i.e. disconnected vertices).

Keywords: Generalized random graphs; inhomogeneous Erdős–Rényi random graph; infinite mean

2010 Mathematics Subject Classification: Primary 60G15

Secondary 82B20; 82B41; 60G70

Received 18 July 2025; accepted 18 July 2025.

* Postal address: Leiden University, Mathematical Institute, Niels Bohrweg 1 2333 CA, Leiden, The Netherlands.

** Postal address: Dipartimento di Matematica e Informatica "Ulisse Dini" – Università degli Studi di Firenze, Italy.

Email: luca.avena@unifi.it

*** Postal address: Lorentz Institute for Theoretical Physics, Leiden University, P.O. Box 9504, 2300 RA Leiden, The Netherlands. INdAM-GNAMPa Istituto Nazionale di Alta Matematica, Italy. Email: garlaschelli@lorentz.leidenuniv.nl

**** Postal address: IMT School for Advanced Studies, Piazza S. Francesco 19, 55100 Lucca, Italy.

***** Email: r.s.hazra@math.leidenuniv.nl

***** Email: margherita.lalli@alumni.imtlucca.it

© The Author(s), 2025. Published by Cambridge University Press on behalf of Applied Probability Trust. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

We consider a class of inhomogeneous Erdős–Rényi random graphs on n vertices. Our vertex set V is denoted by $[n] = \{1, 2, \dots, n\}$ and on each vertex we assign independent weights (or ‘fitness’ variables) $(W_i)_{i \in [n]}$ distributed according to a common distribution $F_W(\cdot)$ with $1 - F_W(x) \sim x^{-\alpha}$ for some $\alpha \in (0, 1)$. Therefore, the weights have infinite mean. Conditioned on the weights, an edge between two distinct vertices i and j is drawn independently with probability

$$p_{ij} = 1 - \exp(-\varepsilon W_i W_j), \quad (1.1)$$

where ε is a parameter tuning the overall density of edges in the graph and playing a crucial role in the analysis of the model. This inhomogeneous random graph model with infinite-mean weights has recently been proposed in [19] in the statistical physics literature, where it was studied as a scale-invariant random graph under hierarchical coarse-graining of vertices. In particular, the model follows from the fundamental requirement that both the connection probability p_{ij} and the fitness distribution $F_W(x)$ retain the same mathematical form when applied to the ‘renormalized’ graph obtained by agglomerating vertices into equally sized ‘supervertices’, where the fitness of each supervertices is defined as the sum of the fitnesses of the constituent vertices [19]. This agglomeration process provides a renormalization scheme for the graph [16] that, by admitting any homogeneous partition of vertices, does not require the notion of vertex coordinates in some underlying metric space, unlike other models based on the idea of geometric renormalization where ‘closer’ vertices are merged [7, 17]. We summarize the main ideas behind the original model later in this paper.

More generally, (1.1) represents a special example of connection probability between vertices i and j defined as $\kappa_n(W_i, W_j)$, where $\kappa_n: [0, \infty)^2 \rightarrow [0, 1]$ is a well-behaved function and the weights are drawn independently from a certain distribution. In the physics literature, these are called ‘fitness’ or ‘hidden variable’ network models [8, 10, 18]. In the mathematical literature, a well-known example is the generalized random graph model [11, 31]. In most cases considered in the literature so far, due to the integrability conditions on κ_n and moment properties of F_W , these models have a locally tree-like structure. We refer to [32, Chapter 6] for the properties of the degree distribution and to [9, 33] for further geometric structures. Models with exactly the same connection probability as in (1.1), but with finite-mean weights, have previously been considered [3, 4, 24, 27, 31]. In this article we are instead interested in the non-standard case of infinite-mean weights, corresponding to the choice $\alpha \in (0, 1)$ as mentioned above. The combination of the specific form of the connection probability (1.1) and these heavy-tailed weights makes the model interesting. We believe that certain mathematical features of an *ultra-small world* network, where the degrees exhibit infinite variance, can be captured by this model. In this case, due to the absence of a finite mean, the typical distances may be much slower than the doubly logarithmic behavior (in relation to the graph’s size) observed in ultra-small networks (refer to [36] for further discussion on this).

Another model where a connection probability similar to (1.1) occurs, but with an additional notion of embedding geometry, is the scale-free percolation model on $\mathbb{Z}^d$. The vertex set in this graph is no longer a finite set of points, and the connection probabilities also depend on the spatial positions of the vertices. Here we also start with independent weights $(W_x)_{x \in \mathbb{Z}^d}$ distributed according to $F_W(\cdot)$, where F_W has a power law index of $\beta \in (0, \infty)$; conditioned on

the weights, vertices x and y are connected independently with probability

$$\tilde{p}_{xy} = 1 - \exp\left(-\frac{\lambda W_x W_y}{\|x - y\|^s}\right),$$

where s and λ are some positive parameters. The model was introduced in [13], where it was shown that the degree of distribution has a power-law exponent of parameter $-\tau = -s\beta/d$. The asymptotics of the maximum degree was derived recently in [5] and further properties of the chemical distances were studied in [14, 21, 35]. In some cases the degree can be infinite too. The mixing properties of the scale-free percolation on a torus of side length n was studied in [12]. In our model, the distance term $\|x - y\|^{-s}$ is not present and hence on one hand the form becomes easier, but on the other hand many useful integrability properties are lost due to the fact that interactions do not decay with distance.

In this paper we show some important properties of our model. In particular, we show that the average degree grows like $\log n$ if we choose the specific scaling $\varepsilon = n^{-1/\alpha}$. In this case, the cumulative degree distribution behaves roughly like a power law with exponent -1 . In the literature for random graphs with degree sequences having infinite mean, this falls in the critical case of exponent $\tau = 1$. The configuration model with given degree sequence $(D_i)_{i \in [n]}$, independently and identically distributed with law D having a power-law exponent $\tau \in (0, 1)$, was studied in [30]. It was shown that the typical distance between two randomly chosen points is either 2 or 3. It was also shown that for $\tau = 1$ similar ultra-small-world behavior is true. Instead of the configuration model, we study the properties of the degree distribution for the model, which also naturally gives rise to degree distributions with power-law exponent -1 . Additionally, we investigate certain dependencies between the degrees of different vertices, the asymptotic density of wedges and triangles, and some first observations on the subtle connectivity properties of the random graph.

The rest of the paper is organized as follows: in Section 2 we state our main results, in Section 3 we discuss the connection to the original model [19], and finally in Sections 4, 5, and 6 we prove our results.

2. Model and main results

The formal definition of the model considered here reads as follows. Let the vertex set be given by $[n] = \{1, 2, \dots, n\}$ and let $\varepsilon = \varepsilon_n > 0$ be a parameter that depends on n . For notational simplicity, we will drop the subscript n from ε_n throughout the paper, except where the context needs it. The random graph with law $\mathbb{P}$ is constructed in the following way:

- (i) Sample n independent weights (W_i) , under $\mathbb{P}$, according to a Pareto distribution with parameter $\alpha \in (0, 1)$, i.e.

$$1 - F_W(w) = \mathbb{P}(W_i > w) = \begin{cases} w^{-\alpha}, & w > 1, \\ 1, & 0 < w \leq 1. \end{cases} \quad (2.1)$$

- (ii) For all $n \geq 1$, given the weights $(W_i)_{i \in [n]}$, construct the random graph G_n by joining edges independently with probability given by (1.1). That is,

$$p_{ij} := \mathbb{P}(i \leftrightarrow j \mid W_i, W_j) = 1 - \exp(-\varepsilon W_i W_j), \quad (2.2)$$

where the event $\{i \leftrightarrow j\}$ means that vertices i and j are connected by an edge in the graph.

We will denote the above random graph by $\mathbf{G}_n(\alpha, \varepsilon)$ as it depends on the parameters α and ε . Self-loops and multi-edges are not allowed, and hence the final graph is given by a simple graph on n vertices. Note that in choosing the distribution of the weights in (2.1) we could alternatively have started with a regularly varying random variable with power-law exponent $-\alpha$, i.e. $\mathbb{P}(W_i > w) = w^{-\alpha} L(w)$ where $L(\cdot)$ is a slowly varying function, i.e., for any $t > 0$,

$$\lim_{w \rightarrow \infty} \frac{L(wt)}{L(w)} = 1.$$

It is our belief that most of the results stated in this article will go through in the presence of a slowly varying function, even if the analysis is more involved. In particular, as we explain in Section 3, in the approach of [19] the weights are drawn from a one-sided α -stable distribution with scale parameter γ , and not from a Pareto (the α -stable following from the requirement of invariance of the fitness distribution under graph renormalization). We expect that the computations will go through if we assume $\mathbb{P}(W_i > w) \sim w^{-\alpha}$ as $w \rightarrow \infty$, which is the case for α -stable distributions. In this work, however, we refrain from going into this technical side.

In terms of notation, convergence in distribution and convergence in probability will be denoted respectively by $\xrightarrow{d}$ and $\xrightarrow{P}$. $\mathbb{E}[\cdot]$ is the expectation with respect to $\mathbb{P}$, and the conditional expectation with respect to the weight W of a typical vertex is denoted by $\mathbb{E}_W[\cdot] = \mathbb{E}[\cdot | W]$. We write $X | W$ to denote the distribution of the random variable X conditioned on the variable W . Let $(a_{ij})_{1 \leq i, j \leq n}$ be the indicator variables $(\mathbf{1}_{i \leftrightarrow j})_{1 \leq i, j \leq n}$. As standard, as $n \rightarrow \infty$ we will write $f(n) \sim g(n)$ if $f(n)/g(n) \rightarrow 1$, $f(n) = o(g(n))$ if $f(n)/g(n) \rightarrow 0$, and $f(n) = O(g(n))$ if $f(n)/g(n) \leq C$ for some $C > 0$, for n large enough. Lastly, $f(n) \asymp g(n)$ denotes that there exist positive constants c_1 and C_2 such that

$$c_1 \leq \liminf_{n \rightarrow \infty} \frac{f(n)}{g(n)} \leq \limsup_{n \rightarrow \infty} \frac{f(n)}{g(n)} \leq C_2.$$

Finally, we use $A \vee B$ to represent $\max\{A, B\}$, the maximum of two numbers A and B .

2.1. Degrees

Our first theorem characterizes the behavior of a typical degree and of the joint distribution of the degrees. Consider the degree of vertex $i \in [n]$, defined as $D_n(i) = \sum_{j \neq i} a_{ij}$, where a_{ij} denotes the entries of the adjacency matrix of the graph, i.e.

$$a_{ij} = \begin{cases} 1 & \text{if } i \leftrightarrow j, \\ 0 & \text{otherwise.} \end{cases}$$

Theorem 2.1 (Scaling and asymptotics of the degrees.) *Consider the graph $\mathbf{G}_n(\alpha, \varepsilon)$ and let $D_n(i)$ be the degree of the vertex $i \in [n]$.*

(i) *Expected degree: The expected degree of a vertex i scales as*

$$\mathbb{E}[D_n(i)] \sim -(n-1)\Gamma(1-\alpha)\varepsilon^\alpha \log \varepsilon^\alpha$$

as $\varepsilon \downarrow 0$. In particular, if $\varepsilon = n^{-1/\alpha}$ then we have

$$\mathbb{E}[D_n(i)] \sim \Gamma(1-\alpha) \log n \quad \text{as } n \rightarrow \infty. \quad (2.3)$$

- (ii) *Asymptotic degree distribution:* Let $\varepsilon = n^{-1/\alpha}$. Then, for all $i \in [n]$, $D_n(i) \xrightarrow{d} D_\infty$ as $n \rightarrow \infty$, where D_∞ is a mixed Poisson random variable with parameter $\Lambda = \Gamma(1 - \alpha)W^\alpha$, where W has distribution (2.1). Additionally, we have

$$\mathbb{P}(D_\infty > x) \sim \Gamma(1 - \alpha)x^{-1} \quad \text{as } x \rightarrow \infty. \quad (2.4)$$

- (iii) *Asymptotic joint degree behavior:* Let $D_\infty(i)$ and $D_\infty(j)$ be the asymptotic degree distributions of two arbitrary distinct vertices $i, j \in \mathbb{N}$. Then

$$\mathbb{E}[t^{D_\infty(i)}s^{D_\infty(j)}] \neq \mathbb{E}[t^{D_\infty(i)}]\mathbb{E}[s^{D_\infty(j)}] \quad \text{for fixed } t, s \in (0, 1), \quad (2.5)$$

and for s, t sufficiently close to 1 we have

$$\begin{aligned} & |\mathbb{E}[t^{D_\infty(i)}s^{D_\infty(j)}] - \mathbb{E}[t^{D_\infty(i)}]\mathbb{E}[s^{D_\infty(j)}]| \\ & \leq O\left((1-s)(1-t) \log\left(\left(1 + \frac{1}{\Gamma(1-\alpha)(1-s)}\right)\left(1 + \frac{1}{\Gamma(1-\alpha)(1-t)}\right)\right)\right) \\ & \quad + C((1-t) + (1-s)) \end{aligned} \quad (2.6)$$

for some constant $C \in (0, \infty)$.

We prove Theorem 2.1 in Section 4. The first part of the result shows that, in the chosen regime, the average degree of the graph diverges logarithmically. This indeed rules out any kind of local weak limit of the graph. We also see in the second part that, asymptotically, degrees have cumulative power-law exponent -1 , for any $\alpha \in (0, 1)$. This rigorously proves a result that was observed with different analytical and numerical arguments in the original paper [19], as we further discuss in Section 3. It is expected that when $\varepsilon = n^{-1/\alpha}$, we should have $\mathbb{P}(D_n(i) > x) \asymp x^{-1}$ as $x \rightarrow \infty$.

The third part of the result deserves further comment. Indeed, (2.5) shows that $D_\infty(i)$ and $D_\infty(j)$ are *not independent*. In the generalized random graph model, this is a surprising phenomenon. If we consider a generalized random graph with weights as described in (2.1) and

$$\tilde{p}_{ij} = \frac{W_i W_j}{n^{1/\alpha} + W_i W_j},$$

then it follows from [32, Theorem 6.14] that the asymptotic degree distribution has the same behavior as our model and the asymptotic degree distributions are independent. Although there is no independence as (2.5) shows, we still believe that

$$|\mathbb{P}(D_\infty(i) > x, D_\infty(j) > x) - \mathbb{P}(D_\infty(i) > x)\mathbb{P}(D_\infty(j) > x)| = o(\mathbb{P}(D_\infty(i) > x)\mathbb{P}(D_\infty(j) > x)) \quad (2.7)$$

holds, and hence the limiting vector will be *asymptotically tail independent*. Although not provided with a rigorous proof yet, this conjecture is supported by numerical simulations (see Fig. 1). Such a property of limiting degree was observed and proved using a multivariate version of Karamata's Tauberian theorem for preferential attachment models, see [26]. In our case, (2.7) would be valid, given an explicit characterization of the complete joint distribution of the asymptotic degrees. Currently, we have not been able to verify the conditions outlined in [26] for the application of their general multivariate Tauberian theorem. We hope to address this question in the future.




FIGURE 1. Asymptotic tail independence between degrees. Scatterplot of the degrees of the vertices with labels 1 and 2 (assigned randomly but fixed for every realization in the ensemble). Each point in the plot corresponds to one of 2000 realizations of a network of $N = 2000$ vertices, each generated as described at the beginning of Section 2 (see (2.1) and (2.2)).

2.2. Wedges, triangles, and clustering

Our second result concerns the number of wedges and triangles associated with a typical vertex $i \in [n]$, defined respectively as

$$\mathbb{W}_n(i) := \frac{1}{2} \sum_{j \neq i} \sum_{k \neq i, j} a_{ij} a_{ik}, \quad \Delta_n(i) = \frac{1}{6} \sum_{j \neq i} \sum_{k \neq i, j} a_{ij} a_{ik} a_{jk}.$$

Theorem 2.2. (Triangles and wedges of typical vertices.) *Consider the graph $\mathbf{G}_n(\alpha, \varepsilon)$ and let $\mathbb{W}_n(i)$ and $\Delta_n(i)$ be the numbers of wedges and triangles at vertex $i \in [n]$. Then:*

- (i) *Average number of wedges: $\mathbb{E}[\mathbb{W}_n(i)] \asymp \varepsilon^\alpha n^2$ as $\varepsilon \downarrow 0$. In particular, when $\varepsilon = n^{-1/\alpha}$, $\mathbb{E}[\mathbb{W}_n(i)] \asymp n$.*
- (ii) *Asymptotic distribution of wedges: Let $\varepsilon = n^{-1/\alpha}$. Then $\mathbb{W}_n(i) \xrightarrow{d} \mathbb{W}_\infty(i)$, where $\mathbb{W}_\infty(i) = D_\infty(i)(D_\infty(i) - 1)$ with $D_\infty(i)$ as in Theorem 2.1. Also,*

$$\mathbb{P}(\mathbb{W}_\infty(i) > x) \sim \Gamma(1 - \alpha)x^{-1/2} \quad \text{as } x \rightarrow \infty.$$
- (iii) *Average number of triangles: Let $i \in [n]$. The average number of triangles grows as $\mathbb{E}[\Delta_n(i)] \asymp \varepsilon^{3\alpha/2} n^2$ as $\varepsilon \downarrow 0$. In particular, when $\varepsilon = n^{-1/\alpha}$ we have $\mathbb{E}[\Delta_n(i)] \asymp \sqrt{n}$ as $n \rightarrow \infty$.*
- (iv) *Concentration for the number of triangles: Let $\varepsilon = n^{-1/\alpha}$ and $\Delta_n = \sum_{i \in [n]} \Delta_n(i)$ be the total number of triangles. Then*

$$\frac{\Delta_n}{\mathbb{E}[\Delta_n]} \xrightarrow{\mathbb{P}} 1, \quad \frac{\Delta_n(i)}{\mathbb{E}[\Delta_n(i)]} \xrightarrow{\mathbb{P}} 1.$$

Remark 2.1. (*Global and local clustering.*) Let $\mathbb{W}_n = \sum_{i \in [n]} \mathbb{W}_n(i)$ be the total number of wedges. We see from Theorem 2.2 that $\mathbb{E}[\Delta_n]/\mathbb{E}[\mathbb{W}_n] \asymp \varepsilon^{\alpha/2}$ as $\varepsilon \rightarrow 0$. This shows in a quantitative form that the graph is not highly clustered from the point of view of the *global* count of triangles. In particular, in the scale of $\varepsilon = n^{-1/\alpha}$, the above ratio goes to zero like $n^{-1/2}$. However, this does not mean that the graph is not highly clustered from the point of view of the *local* count of triangles around individual vertices. Indeed, simulations in [19] of the average local clustering coefficient suggest that the graph is locally clustered (see also Section 3). A dissimilarity in the behavior of local and global clustering coefficients has also been observed in different inhomogeneous random graph models; see, for example, [23, 34, 37]. We do not consider the local clustering here.

2.3. Connectedness: Some first observations

Connectivity properties of inhomogeneous random graphs were studied in the sparse setting in [9]. The connectivity properties when the connection probabilities are of the form $\min\{1, \kappa(W_i, W_j) \log n/n\}$ with κ being a square-integrable kernel were studied in [15]. Note that due to the dependency of ε in our p_{ij} , this does not fall in this setting; as such, connectivity properties of this ensemble would deserve a new detailed analysis, which will be addressed elsewhere. We close this first rigorous work on this ensemble by pointing out that connectivity properties heavily depend on the ε regime considered, as can be already appreciated by looking at the presence of *dust* (i.e. isolated points in the graph). Indeed, the following and previous statements show a cross-over for the absence of dust at the ε scale $(\log n/n)^{1/\alpha}$.

Proposition 2.1. (No-dust regime.) *Consider the graph $\mathbf{G}_n(\alpha, \varepsilon)$. Let N_0 be the number of isolated vertices, i.e.*

$$N_0 = \sum_{i=1}^n \mathbf{1}_{\{i \text{ is isolated}\}}. \quad (2.8)$$

Then $\mathbb{E}[N_0] \sim n\mathbb{E}[e^{-(n-1)\Gamma(1-\alpha)\varepsilon^\alpha W_1^\alpha}]$. In particular, if $\varepsilon \downarrow 0$ and $\varepsilon^\alpha n/\log n \rightarrow \infty$, then

$$\mathbb{P}(N_0 = 0) \rightarrow 1. \quad (2.9)$$

If $\varepsilon = n^{-1/\alpha}$, then a positive fraction of points are isolated, i.e. $\mathbb{E}[N_0]/n \rightarrow \mathbb{E}[e^{-\Gamma(1-\alpha)W_1^\alpha}]$.

3. The multi-scale model (MSM)

In this section we discuss the connection between our results and the model introduced in [19].

3.1. Motivation for the MSM

The motivation for the MSM arises from statistical physics, where the concept of *renormalization* [22, 39] plays a central role. In the context of networks, renormalization involves selecting a *coarse-graining* approach for a graph, which essentially means projecting a larger ‘microscopic’ graph onto a ‘reduced’ graph with fewer vertices [16]. This reduction is determined by a non-overlapping partition of the vertices of the original microscopic graph into ‘clusters’ or ‘supervertices’, which then become the vertices of the reduced graph. The edges of the reduced graph are defined according to specific rules, usually aimed at preserving certain structural features of the original network. This renormalization process can be iterated, potentially infinitely, in the case of an infinite graph.

For example, when the network is a regular lattice (or geometric graph) embedded in a specific metric space, a straightforward renormalization scheme exists. However, for generic (non-geometric) graphs, the absence of an explicit metric embedding makes the choice of renormalization significantly more challenging. Proposed approaches include borrowing box-covering techniques from fractal analysis [25, 29], employing spectral coarse-graining methods [20, 38], and utilizing graph embedding techniques to infer optimal vertex coordinates in an imposed (usually hyperbolic) latent space [7, 17]. For a recent review on network renormalization, see [16].

Regardless of the method used to find the optimal sequence of coarse-grainings for a given graph, the MSM has been introduced as a random graph model that remains consistently applicable to both the original graph and any reduced versions of it [19]. This implies that, assuming the probability of generating a graph at the microscopic level follows a specific function of the model parameters, the probability of generating any reduced version of the graph should have the same functional form, potentially with renormalized parameters.

The MSM can be explicitly obtained as the model that fulfills the requirement that the random graph ensemble is scale-invariant under a renormalization process that accepts any partition of vertices. Importantly, this renormalization scheme is non-geometric by design, as it doesn't rely on the notion of vertex coordinates in an underlying metric space, unlike the previously mentioned models based on the concept of geometric renormalization, where 'closer' vertices are merged.

3.2. Construction of the MSM

The renormalization framework allows for the same random graph ensemble to be observed at different hierarchical levels $\ell = 0, 1, 2, \dots$. Let us start with the 'microscopic' level $\ell = 0$ and consider a random graph on n_0 vertices where, adopting the notation used in [19], each vertex (labeled as $i_0 = 1, \dots, n_0$) has a weight X_{i_0} . To move to level $\ell = 1$, we specify a partition of the original n_0 vertices into $n_1 < n_0$ blocks (which here, for simplicity, we assume to be all equal in size and composed of b vertices, so that $n_1 = n_0/b$). The blocks of the partition, labeled $i_1 = 1, \dots, n_1$, become the vertices of the graph at the new level and each pair of blocks is connected if there existed at least an edge between any two original vertices placed across the two blocks. At this new level, the weights of all vertices inside a block i_1 get summed to produce the (renormalized) weight for that block, denoted as $X_{i_1} \equiv \sum_{i_0 \in i_1} X_{i_0}$. The process can continue to higher levels $\ell > 1$ by progressively reducing the graph to one with $n_{\ell+1} = n_\ell/b = \dots = n_0/b^{\ell+1}$ vertices and renormalizing the weights as $X_{i_{\ell+1}} \equiv \sum_{i_\ell \in i_{\ell+1}} X_{i_\ell}$.

To define the MSM, we enforce the requirement that, under the coarse-graining process defined above, the probability distribution of the graph preserves the same functional form across all levels. This scale-invariant requirement becomes particularly simple if we consider the family of random graph models with independent edges, which are entirely specified by a function $p_{i_\ell j_\ell}$ of the parameters, representing the probability that the vertices i_ℓ and j_ℓ are connected. For this family, the connection probability $p_{i_{\ell+1} j_{\ell+1}}$ between two vertices $i_{\ell+1}$ and $j_{\ell+1}$ defined at the level $\ell + 1$ is related to the connection probabilities $\{p_{i_\ell j_\ell}\}_{i_\ell, j_\ell}$ between the vertices at the previous level ℓ via

$$p_{i_{\ell+1} j_{\ell+1}} = 1 - \prod_{i_\ell \in i_{\ell+1}} \prod_{j_\ell \in j_{\ell+1}} (1 - p_{i_\ell j_\ell}). \quad (3.1)$$

Assuming that the connection probability depends on a global parameter $\delta > 0$ as well as on the additive vertex weights $\{X_{i_\ell}\}_{i_\ell}$ introduced above, the simplest non-trivial expression consistent

with (3.1) is given by

$$p_{i\ell j\ell} = 1 - \exp(-\delta X_{i\ell} X_{j\ell}). \quad (3.2)$$

At this point, we may require that the weights are either deterministic parameters assigned to the vertices, so that the only source of randomness lies in the realization of the graph ('quenched' variant of the MSM), or that they are random variables themselves, thus adding a second layer of randomness ('annealed' variant of the MSM). In the latter case, it is natural to subject the weights to the same scale-invariant requirement as the random graph, i.e. to demand that the weights are drawn from the same probability density function (with possibly renormalized parameters) at all hierarchical levels. Since the weights are chosen to be additive upon renormalization, this requirement immediately implies that they must be drawn from an α -stable distribution. Moreover, the positivity of the weights and the concurrent requirement that the support of their pdf is the non-negative real axis, irrespective of the hierarchical level, imply that they should be *one-sided* α -stable random variables with (scale-invariant) parameter $\alpha \in (0, 1)$ and some (scale-dependent) scale parameter γ_ℓ . In this way, if the blocks of the partition are always of size b , as assumed above, then the vertex weights at level ℓ are one-sided α -stable random variables with rescaled parameter $\gamma_\ell = b^{1/\alpha} \gamma_{\ell-1} = \dots = b^{\ell/\alpha} \gamma_0 = (n_0/n_\ell)^{1/\alpha} \gamma_0$. This completes the definition of the annealed MSM, along with its renormalization rules for both the weights of vertices and all the other model parameters.

3.3. Connection between the MSM and the model studied in this paper

Despite the obvious relationship between the model studied in this paper and the original annealed version of the MSM recalled above (in particular, between (1.1) and (3.2)), there are apparently some differences that require further discussion. First, here we have considered weights drawn from a Pareto distribution with tail exponent α , rather than a one-sided α -stable distribution; second, here the other parameters of the weight distribution are fixed and the scale parameter ε is n -dependent, while in the original model the other parameters (γ) of the weight distribution are ℓ -dependent (hence also n -dependent) and the scale parameter δ is fixed; third, here we have not exploited the scale-invariant nature of the MSM under coarse-graining. We now clarify the close relationship between the two variants of the model, these apparent differences notwithstanding.

Let us start by recalling that, for large values of the argument x , a one-sided α -stable distribution $\mathbb{P}(X > x)$ with scale parameter γ is well approximated by a pure power-law (Pareto) distribution $\mathbb{P}(X > x) \sim C_{\alpha,\gamma} x^{-\alpha}$ with a prefactor $C_{\alpha,\gamma}$ that depends on the parameters of the stable law [28]:

$$C_{\alpha,\gamma} \equiv \gamma^\alpha c_\alpha, \quad \text{with } c_\alpha \equiv \frac{2\Gamma(\alpha)}{\pi} \sin \frac{\pi\alpha}{2}. \quad (3.3)$$

Then, we note that, besides n_0 and b , the three remaining parameters of the original annealed MSM are $\alpha \in (0, 1)$, $\gamma_0 \in (0, \infty)$, and $\delta \in (0, \infty)$. However, of these three parameters, only α and the combination $\delta\gamma_0^2$ are independent. Indeed, it is easy to realize that rescaling γ_0 to γ_0/λ (which is equivalent to rescaling $X_{i\ell}$ to $X_{i\ell}/\lambda$, for some $\lambda > 0$) while simultaneously rescaling δ to $\lambda^2\delta$ leaves the connection probability unchanged. In combination with the scale-invariant nature of the MSM, this property can be exploited to map the quantities $(\{X_{i\ell}\}_{i\ell=1}^{n_\ell}, \delta)$ introduced in the original model to the quantities $(\{W_i\}_{i=1}^n, \varepsilon)$ used here by choosing a level ℓ such that the number n_ℓ of vertices in the MSM equals the one desired here, i.e. $n = n_\ell$, and

defining the weight of vertex i as $W_i \equiv c_\alpha^{-1/\alpha} \gamma_\ell^{-1} X_{i_\ell}$ for $i = 1, \dots, n$, so that

$$\lim_{x \rightarrow \infty} \mathbb{P}(W_i > x) x^\alpha = 1, \quad i = 1, \dots, n, \quad (3.4)$$

irrespective of ℓ . This implies that, while the distribution of X_{i_ℓ} depends on ℓ through the parameter γ_ℓ (see (3.3)), the distribution of W_i is actually ℓ -independent in the tail, which is why we could drop the subscript ℓ in redefining W_i . Note that this procedure yields weights that are only asymptotically ℓ -independent, as expressed in (3.4). Nevertheless, in this way we can keep the connection probability unchanged (i.e. $1 - e^{-\delta X_{i_\ell} X_{j_\ell}} \equiv 1 - e^{-\varepsilon_{n_\ell} W_i W_j}$) while moving the scale-dependence from $\{X_{i_\ell}\}_{i_\ell=1}^{n_\ell}$ to ε_{n_ℓ} by redefining the latter in one of the following equivalent ways:

$$\varepsilon_{n_\ell} \equiv c_\alpha^{2/\alpha} \gamma_\ell^2 \delta = c_\alpha^{2/\alpha} \left(\frac{n_0}{n_\ell} \right)^{2/\alpha} \gamma_0^2 \delta = c_\alpha^{2/\alpha} b^{2\ell/\alpha} \gamma_0^2 \delta = b^{2\ell/\alpha} \varepsilon_{n_0}, \quad \text{where } \varepsilon_{n_0} \equiv c_\alpha^{2/\alpha} \gamma_0^2 \delta, \quad (3.5)$$

where we now make explicit the dependence of ε on n_ℓ to track variations across scales ℓ . In other words, our formulation here can be thought of as deriving from an equivalent MSM where, rather than having a scale-dependent fitness distribution and a scale-independent global parameter δ , we have a scale-independent fitness distribution (with asymptotically the same tail as the Pareto in (2.1)) and a scale-dependent global parameter $\varepsilon_n = \varepsilon_{n_\ell}$, for an implied hierarchical level ℓ . According to (3.5), since δ , α , and b are finite constants, achieving a certain scaling of ε_n with n in the model considered here corresponds to achieving a corresponding scaling of γ_ℓ with n_ℓ (or equivalently of γ_0 with n_0), or ultimately to finding an appropriate ℓ , in the original model. Results that we obtain for a specific range of values of ε can therefore be thought of as applying to a corresponding specific range of hierarchical levels in the original model. In particular, Theorems 2.1 and 2.2 reveal that at the specific scale $\varepsilon = \lambda n^{-1/\alpha}$ the model undergoes drastic structural changes (e.g. concerning typical degrees and wedges).

3.4. Implications of our results for the MSM

Some topological properties of the original MSM were investigated in [19] numerically, and either analytically (for $\alpha = \frac{1}{2}$, corresponding to the Lévy distribution, which is the only one-sided α -stable distribution that can be written in explicit form) or semi-analytically (for generic $\alpha \in (0, 1)$). Notably, it was found that networks sampled from the MSM feature an empirical degree distribution $P(k)$ exhibiting a scale-free region, characterized by a universal power-law decay $\propto k^{-2}$ (corresponding to a cumulative distribution with decay $\propto k^{-1}$) irrespective of the value of $\alpha \in (0, 1)$, followed by a density-dependent cut-off.

The results obtained here provide significant additional insights. With regard to the degrees, we have identified the specific scaling (or equivalently, as explained in Section 3.3, the specific hierarchical level) for which the density-dependent cut-off disappears and the tail of the cumulative degree distribution can be rigorously proven (through an independent proof) to become a pure power law with exponent -1 , for any $\alpha \in (0, 1)$. Secondly, we have provided a rigorous evaluation of the overall number of triangles and wedges, valid for any α , that supports the outcome of simulations shown in the original paper, which illustrated the vanishing of the global clustering coefficient in the sparse limit (as opposed to the local clustering coefficient, which remains bounded away from zero as recalled above).

4. Proof of Theorem 2.1: Typical degrees

Since Karamata's Tauberian theorem is used here as a key tool in the analysis of the degree distribution and later analysis, it is first worth recalling those results.

Theorem 4.1. (Karamata's Tauberian theorem [6, Theorem 8.1.6].) *Let X be a non-negative random variable with distribution F and Laplace transform $\widehat{F}(s) = \mathbb{E}[e^{-sX}]$, $s \geq 0$. Let L be a slowly varying function and $\alpha \in (0, 1)$; then the following are equivalent:*

$$\begin{aligned} 1 - \widehat{F}(s) &\sim \Gamma(1 - \alpha)s^\alpha L(1/s) \quad \text{as } s \downarrow 0, \\ 1 - F(x) &\sim x^{-\alpha} L(x) \quad \text{as } x \rightarrow \infty. \end{aligned}$$

Then, another property of the tails of products of regularly varying distributions will be needed. A general statement about the product of n independent and identically distributed (i.i.d.) random variables with Pareto tails can be found in [1, Lemma 4.1 (4)]. For completeness, a proof for two random variables is given here, which is useful in our analysis.

Lemma 4.1. *Let W_1 and W_2 be independent random variables satisfying the tail assumptions (2.1). Then $\mathbb{P}(W_1 W_2 \geq x) \sim \alpha x^{-\alpha} \log x$ as $x \rightarrow \infty$.*

Proof. Consider the random variable $\log(W_1)$, which follows an exponential distribution, or alternatively a Gamma distribution with shape parameter $k = 1$ and scale $\theta = 1/\alpha$. Then, the random variable $Z = \log(W_1) + \log(W_2)$ follows a Gamma distribution with shape parameter 2 and scale θ . This means that

$$\mathbb{P}(\log(W_1) + \log(W_2) > x) = \frac{\alpha^2}{\Gamma(2)} \int_x^\infty y e^{-\alpha y} dy.$$

Therefore

$$\mathbb{P}(W_1 W_2 > x) = \mathbb{P}(\log W_1 + \log W_2 > \log x) = \alpha^2 \int_{\log x}^\infty y e^{-\alpha y} dy = \alpha^2 \int_x^\infty \log(t) t^{-\alpha-1} dt. \quad (4.1)$$

Then, applying Karamata's theorem (see [1, Theorem 12]),

$$\mathbb{P}(W_1 W_2 > x) \sim \alpha^2 \frac{x^{-\alpha} \log x}{\alpha},$$

which proves the statement. $\square$

Remark 4.1. Theorem 4.1 remains true if W_1 and W_2 are not exactly Pareto, but asymptotically tail-equivalent to a Pareto distribution, i.e. under the assumption $\mathbb{P}(W_1 > x) \sim c x^{-\alpha}$ as $x \rightarrow \infty$. See [1, Lemma 4.1] for a proof.

Proof of Theorem 2.1. For (i), we begin by evaluating the asymptotics of the expected degree, which is an easy consequence of Lemma 4.1 and Theorem 4.1. Indeed, we can write

$$\mathbb{E}[D_n(i)] = \sum_{j \neq i} \mathbb{E}[1 - \exp(-\varepsilon W_i W_j)] = (n-1) \mathbb{E}[1 - \exp(-\varepsilon W_1 W_2)], \quad (4.2)$$

where the last equality is due to the i.i.d. nature of the weights. It follows from Lemma 4.1 that $\mathbb{P}(W_1 W_2 > x) \sim \alpha x^{-\alpha} \log x$. Therefore, using Theorem 4.1, we have

$$\mathbb{E}[1 - \exp(-\varepsilon W_1 W_2)] \sim \Gamma(1 - \alpha) \alpha \varepsilon^\alpha \log \frac{1}{\varepsilon} \quad \text{as } \varepsilon \downarrow 0,$$

which together with (4.2) gives the claim.

For (ii), by following the line of the proof of [32, Theorem 6.14], we can prove our statement by showing that the probability-generating function of $D_n(i)$ in the limit $n \rightarrow \infty$ corresponds to the probability-generating function of a mixed Poisson random variable. Let $t \in (0, 1)$; the probability-generating function of the degree $D_n(i)$ reads

$$\mathbb{E}[t^{D_n(i)}] = \mathbb{E}[t^{\sum_{j \neq i} a_{ij}}] = \mathbb{E}\left[\prod_{j \neq i} t^{a_{ij}}\right],$$

where a_{ij} are the entries of the adjacency matrix related to the graph $\mathbf{G}_n(\alpha, \varepsilon)$, i.e. Bernoulli random variables with parameter p_{ij} as in (2.2). Conditioned on the weights, these variables are independent. Recall that we denoted by $\mathbb{E}_{W_i}[\cdot]$ the conditional expectation given the weight W_i . Then

$$\begin{aligned} \mathbb{E}_{W_i}[t^{D_n(i)}] &= \mathbb{E}_{W_i}\left[\prod_{j \neq i} ((1-t)e^{-\varepsilon W_j W_i} + t)\right] = \prod_{j \neq i} \mathbb{E}_{W_i}[(1-t)e^{-\varepsilon W_j W_i} + t] \\ &= \prod_{j \neq i} \mathbb{E}_{W_i}[\varphi_{W_i}(\varepsilon W_j)], \end{aligned}$$

where we have used the independence of the weights and introduced the function

$$\varphi_{W_i}(x) := (1-t)e^{-W_i x} + t.$$

To simplify our expression we introduce the notation $\psi_n(W_i) := \mathbb{E}_{W_i}[\varphi_{W_i}(\varepsilon W_j)]$ for some $j \neq i$. Using exchangeability and the tower property of the conditional expectation, the moment-generating function of $D_n(i)$ can be written as

$$\mathbb{E}[t^{D_n(i)}] = \mathbb{E}\left[\prod_{j \neq i} \mathbb{E}_{W_i}[\varphi_{W_i}(\varepsilon W_j)]\right] = \mathbb{E}[\psi_n(W_i)^{n-1}]. \quad (4.3)$$

Consider now a differentiable function $h: [0, \infty) \rightarrow \mathbb{R}$ such that $h(0) = 0$. By integration by parts, we can show that

$$\mathbb{E}[h(W_j)] = \int_0^\infty h'(x) \mathbb{P}(W_j > x) dx. \quad (4.4)$$

By using (4.4) with $h(w) = \varphi_{W_i}(\varepsilon w) - 1$, we have

$$\begin{aligned} \psi_n(W_i) &= 1 + \mathbb{E}[\varphi_{W_i}(\varepsilon w) - 1] \\ &= 1 + \int_0^\infty \varepsilon \varphi'_{W_i}(\varepsilon w)(1 - F_W(w)) dw \\ &= 1 + \int_0^\infty \varphi'_{W_i}(y)(1 - F_W(\varepsilon^{-1}y)) dy \\ &= 1 + \int_0^\varepsilon \varphi'_{W_i}(y) dy + \int_\varepsilon^\infty \varphi'_{W_i}(y)(1 - F_W(\varepsilon^{-1}y)) dy \\ &= 1 + \varphi_{W_i}(\varepsilon) - \varphi_{W_i}(0) + \varepsilon^\alpha \int_\varepsilon^\infty (t-1)W_i e^{-yW_i} y^{-\alpha} dy. \end{aligned} \quad (4.5)$$

In particular, for $\varepsilon = n^{-1/\alpha}$, combining (4.3) and (4.5) gives

$$\begin{aligned}\mathbb{E}[t^{D_n(i)}] &= \mathbb{E}[\psi_n(W_i)^{n-1}] \\ &= \mathbb{E}\left[\left(1 + \varphi_{W_i}(n^{-1/\alpha}) - \varphi_{W_i}(0) + \frac{1}{n} \int_{n^{-1/\alpha}}^{\infty} (t-1)W_i e^{-yW_i} y^{-\alpha} dy\right)^{n-1}\right].\end{aligned}$$

Note that for a fixed realization of W_i , using the change of variable $z = W_i y$ we have, as $n \rightarrow \infty$,

$$(1-t) \int_{n^{-1/\alpha}}^{\infty} W_i e^{-yW_i} y^{-\alpha} dy \rightarrow (1-t)W_i^{\alpha} \Gamma(1-\alpha)$$

and $\varphi_{W_i}(n^{-1/\alpha}) \rightarrow \varphi_{W_i}(0) = 1$.

Observe that $\varphi_{W_i}(n^{-1/\alpha}) - \varphi(0) = -(1-t)(1 - e^{-W_i n^{1/\alpha}}) \leq 0$ and

$$0 \leq (1-t) \int_{n^{-1/\alpha}}^{\infty} W_i e^{-yW_i} y^{-\alpha} dy \leq (1-t)W_i^{\alpha} \Gamma(1-\alpha).$$

Hence, using $(1-x/n)^n \leq e^{-x}$ we have

$$\begin{aligned}&\left(1 + \varphi_{W_i}(n^{-1/\alpha}) - \varphi_{W_i}(0) - (1-t)\frac{1}{n} \int_{n^{-1/\alpha}}^{\infty} W_i e^{-yW_i} y^{-\alpha} dy\right)^{n-1} \\ &\leq \exp(-(1-t)W_i^{\alpha} \Gamma(1-\alpha)) \leq 1.\end{aligned}$$

Thus, we can apply the dominated convergence theorem to claim that

$$\lim_{n \rightarrow \infty} \mathbb{E}[t^{D_n(i)}] = \mathbb{E}[\exp(-(1-t)W_i^{\alpha} \Gamma(1-\alpha))],$$

so the generating function of the graph degree $D_n(i)$ asymptotically corresponds to the generating function of a mixed Poisson random variable with parameter $\Gamma(1-\alpha)W_i^{\alpha}$. Therefore,

the variable $D_n(i) \xrightarrow{d} D_{\infty}(i)$, where $D_{\infty}(i) \mid W_i \stackrel{d}{=} \text{Poisson}(\Gamma(1-\alpha)W_i^{\alpha})$.

In particular, we have the following tail of the distribution of the random variable $D_{\infty}(i)$:

$$\begin{aligned}\mathbb{P}(D_{\infty}(i) \geq k) &= \int_0^{\infty} \mathbb{P}(\text{Poisson}(\Gamma(1-\alpha)w^{\alpha}) \geq k \mid W_i = w) F_{W_i}(dw) \\ &= \int_0^{\infty} \sum_{m \geq k} \frac{e^{-\Gamma(1-\alpha)w^{\alpha}} \Gamma(1-\alpha)^m w^{\alpha m}}{m!} F_{W_i}(dw) \\ &= \sum_{m \geq k} \frac{1}{m!} \int_1^{\infty} e^{-\Gamma(1-\alpha)w^{\alpha}} \Gamma(1-\alpha)^m w^{\alpha m} \alpha w^{-\alpha-1} dw.\end{aligned}$$

Introducing the new variable $y = \Gamma(1-\alpha)w^{\alpha}$, we have

$$\begin{aligned}&\int_1^{\infty} e^{-\Gamma(1-\alpha)w^{\alpha}} \Gamma(1-\alpha)^m w^{\alpha m} \alpha w^{-\alpha-1} dw \\ &= \int_1^{\infty} e^{-\Gamma(1-\alpha)w^{\alpha}} \Gamma(1-\alpha)^{m-1} w^{\alpha m} w^{-2\alpha} \alpha \Gamma(1-\alpha) w^{\alpha-1} dw \\ &= \Gamma(1-\alpha) \int_{\Gamma(1-\alpha)}^{\infty} e^{-y} y^{m-2} dy \\ &= \Gamma(1-\alpha) \Gamma(m-1) - \Gamma(1-\alpha) \int_0^{\Gamma(1-\alpha)} e^{-y} y^{m-2} dy.\end{aligned}\tag{4.6}$$

The first integral is dominant with respect to the second one. To show this, we can use a trivial bound:

$$\int_0^{\Gamma(1-\alpha)} e^{-y} y^{m-2} dy \leq \Gamma(1-\alpha)^m.$$

Since $m! \geq (m/e)^m$, the following inequalities hold true:

$$\sum_{m \geq k} \frac{\Gamma(1-\alpha)^m}{m!} \leq \sum_{m \geq k} \frac{(e\Gamma(1-\alpha))^m}{m^m} \leq C(e\Gamma(1-\alpha))^k k^{-k}$$

for some positive constant C . Note that in last step we used the fact that k is large enough (it is at least greater than $e\Gamma(1-\alpha)$). Note that

$$k \sum_{m \geq k} \frac{\Gamma(1-\alpha)^m}{m!} \leq C e^{\log k - k \log k + k e \Gamma(1-\alpha)} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Using (4.6), we therefore obtain

$$\begin{aligned} \sum_{m \geq k} \frac{1}{m!} \int_1^\infty e^{-\Gamma(1-\alpha)w^\alpha} \Gamma(1-\alpha)^m w^{\alpha m} \alpha w^{-\alpha-1} dw &= \Gamma(1-\alpha) \sum_{m \geq k} \frac{\Gamma(m-1)}{m!} + o(k^{-1}) \\ &= \Gamma(1-\alpha) \sum_{m \geq k} \frac{(m-2)!}{m!} + o(k^{-1}) \\ &= \Gamma(1-\alpha) \sum_{m \geq k} \frac{1}{m(m-1)} + o(k^{-1}) \\ &\sim \frac{\Gamma(1-\alpha)}{k}. \end{aligned}$$

This shows that $\mathbb{P}(D_\infty(i) \geq k) \sim \Gamma(1-\alpha)k^{-1}$ as $k \rightarrow \infty$.

For (iii), fix $t, s \in (0, 1)$. Due to the exchangeability of vertices, without loss of generality we just consider vertices 1 and 2:

$$\begin{aligned} &\mathbb{E}[t^{D_n(1)} s^{D_n(2)}] \\ &= \mathbb{E}[t^{\sum_{j \neq 1} a_{1j}} s^{\sum_{j \neq 2} a_{2j}}] \\ &= \mathbb{E}\left[\prod_{j \neq 1, 2} t^{a_{1j}} s^{a_{2j}} (ts)^{a_{12}}\right] \\ &= \mathbb{E}\left[\left((1-ts)e^{-\varepsilon W_1 W_2} + ts\right) \prod_{j \neq 1, 2} \left((1-t)e^{-\varepsilon W_1 W_j} + t\right) \left((1-s)e^{-\varepsilon W_2 W_j} + s\right)\right], \quad (4.7) \end{aligned}$$

where we have used the independence of the connection probabilities *given* the weights. In order to simplify the notation, we can introduce the functions

$$\begin{aligned} \phi_a^b(x) &:= (1-b)e^{-\varepsilon ax} + b, \\ \psi_n(W_1, W_2) &:= \mathbb{E}_{W_1, W_2}[\phi_{W_1}^t(W_j) \phi_{W_2}^s(W_j)] \quad \text{for some } j \neq 1, 2, \end{aligned}$$

where $a, b > 0$ and, as customary throughout this paper, $\mathbb{E}_{W_1, W_2}[\cdot] := \mathbb{E}[\cdot | W_1, W_2]$.

Using the tower property of conditional expectation, (4.7) reads

$$\begin{aligned}\mathbb{E}[t^{D_n(1)}s^{D_n(2)}] &= \mathbb{E}\left[\phi_{W_1}^{ts}(W_2) \prod_{j \neq 1,2} \phi_{W_1}^t(W_j)\phi_{W_2}^s(W_j)\right] \\ &= \mathbb{E}\left[\phi_{W_1}^{ts}(w_2)\mathbb{E}_{W_1, W_2}\left[\prod_{j \neq 1,2} \phi_{W_1}^t(w_j)\phi_{W_2}^s(w_j)\right]\right] \\ &= \mathbb{E}\left[\phi_{W_1}^{ts}(W_2) \prod_{j \neq 1,2} \mathbb{E}_{W_1, W_2}[\phi_{W_1}^t(W_j)\phi_{W_2}^s(W_j)]\right] \\ &= \mathbb{E}[\phi_{W_1}^{ts}(W_2)\psi_n(W_1, W_2)^{n-2}],\end{aligned}$$

where we used conditional independence in the second-last step and exchangeability in the last step. The function ψ_n can be processed as follows. Just as in the one-dimensional case, using $\varepsilon = n^{-1/\alpha}$ we get, $\mathbb{P}$ -almost surely (a.s.),

$$\begin{aligned}\psi_n(W_1, W_2) - 1 &= \mathbb{E}_{W_1, W_2}[\phi_{W_1}^t(W_3)\phi_{W_2}^s(W_3) - 1] \\ &\rightarrow -\Gamma(1-\alpha)[(1-t)(1-s)(W_1 + W_2)^\alpha + (1-t)sW_1^\alpha + t(1-s)W_2^\alpha] \\ &= -\Gamma(1-\alpha)\{(1-t)(1-s)[(W_1 + W_2)^\alpha - W_1^\alpha - W_2^\alpha] + (1-t)W_1^\alpha + (1-s)W_2^\alpha\}, \quad (4.8)\end{aligned}$$

where in the second step we used (4.4) with $h(x) := \phi_{W_1}^t(x)\phi_{W_2}^s(x) - 1$. Observe that $\phi_{W_1}^{ts}(W_2) \leq 1$ and $1 - \psi_n(W_1, W_2) \geq 0$, and hence we can use the dominated convergence theorem as in the single-vertex case. Therefore, using $\phi_{W_1}^{ts}(W_2) \rightarrow 1$ and (4.8), we get

$$\begin{aligned}\mathbb{E}[t^{D_\infty(1)}s^{D_\infty(2)}] &= \lim_{n \rightarrow \infty} \mathbb{E}[t^{D_n(1)}s^{D_n(2)}] \\ &= \mathbb{E}\left[e^{-\Gamma(1-\alpha)\{(1-t)(1-s)[(W_1 + W_2)^\alpha - W_1^\alpha - W_2^\alpha] + (1-t)W_1^\alpha + (1-s)W_2^\alpha\}}\right] \\ &= \mathbb{E}\left[e^{-\Gamma(1-\alpha)(1-t)(1-s)[(W_1 + W_2)^\alpha - W_1^\alpha - W_2^\alpha]}e^{-\Gamma(1-\alpha)(1-t)W_1^\alpha}e^{-\Gamma(1-\alpha)(1-s)W_2^\alpha}\right].\end{aligned}$$

It is straightforward to note that in the limit $t \rightarrow 1$ and for fixed $s \in (0, 1)$ we recover the correct moment-generating function of $D_\infty(1)$, and the inverse holds true as well. Finally, since $(W_1 + W_2)^\alpha \neq W_1^\alpha + W_2^\alpha$ $\mathbb{P}$ -a.s., (2.5) follows. We next move to the proof of (2.6), for which we abbreviate $\eta = 1 - t$ and $\gamma = 1 - s$, and show that

$$\lim_{\substack{\eta \rightarrow 0 \\ \gamma \rightarrow 0}} \left| \mathbb{E}[(1-\eta)^{D_\infty(1)}(1-\gamma)^{D_\infty(2)}] - \mathbb{E}[(1-\eta)^{D_\infty(1)}]\mathbb{E}[(1-\gamma)^{D_\infty(2)}] \right| = 0:$$

$$\begin{aligned}
& \left| \mathbb{E}[(1-\eta)^{D_\infty(1)}(1-\gamma)^{D_\infty(2)}] - \mathbb{E}[(1-\eta)^{D_\infty(1)}]\mathbb{E}[(1-\gamma)^{D_\infty(1)}] \right| \\
&= \left| \mathbb{E}[e^{-\Gamma(1-\alpha)\eta\gamma[(w_1+w_2)^\alpha - w_1^\alpha - w_2^\alpha]} e^{-\Gamma(1-\alpha)\eta w_1^\alpha} e^{-\Gamma(1-\alpha)\gamma w_2^\alpha}] \right. \\
&\quad \left. - \mathbb{E}[e^{-\Gamma(1-\alpha)\eta w_1^\alpha}] \mathbb{E}[e^{-\Gamma(1-\alpha)\gamma w_2^\alpha}] \right| \\
&= \left| \mathbb{E}[e^{-\Gamma(1-\alpha)\eta\gamma[(w_1+w_2)^\alpha - w_1^\alpha - w_2^\alpha]} - 1] e^{-\Gamma(1-\alpha)\eta w_1^\alpha} e^{-\Gamma(1-\alpha)\gamma w_2^\alpha} \right| \\
&= \left| \int_1^\infty \left(\sum_{k \geq 1} \frac{(\Gamma(1-\alpha)\eta\gamma)^k}{k!} [-(x+y)^\alpha + x^\alpha + y^\alpha]^k \right) \right. \\
&\quad \left. \times e^{-\Gamma(1-\alpha)\eta x^\alpha} e^{-\Gamma(1-\alpha)\gamma y^\alpha} \alpha^2 (xy)^{-\alpha-1} dx dy \right| \\
&\leq \int_1^\infty \sum_{k=1}^\infty \frac{(\Gamma(1-\alpha)\eta\gamma)^k}{k!} |-(x+y)^\alpha + x^\alpha + y^\alpha|^k e^{-\Gamma(1-\alpha)\eta x^\alpha} e^{-\Gamma(1-\alpha)\gamma y^\alpha} \alpha^2 (xy)^{-\alpha-1} dx dy.
\end{aligned}$$

Now, since $(x+y)^\alpha \leq (x^\alpha + y^\alpha)$ for all $\alpha \in (0, 1)$, we get

$$|x^\alpha + y^\alpha - (x+y)^\alpha|^k \leq (x^\alpha + y^\alpha)^k \leq 2^{k-1} (x^{\alpha k} + y^{\alpha k}). \quad (4.9)$$

Therefore, using Fubini we can bring the summation out of the integral and, using the inequality (4.9), we are left with the following quantity (which we will show to be converging to zero):

$$\alpha^2 \sum_{k=1}^\infty \frac{(\Gamma(1-\alpha)\eta\gamma)^k}{k!} 2^{k-1} \int_1^\infty (x^{\alpha k} + y^{\alpha k}) e^{-\Gamma(1-\alpha)\eta x^\alpha} e^{-\Gamma(1-\alpha)\gamma y^\alpha} (xy)^{-\alpha-1} dx dy.$$

Let us consider the different terms of the sum separately. In the following we will consider the exponential integral $E_1(x) = \int_1^\infty e^{-tx}/t dt$ and the related inequality $E_1(x) < e^{-x} \ln(1 + (1/x))$ for any $x > 0$.

When $k = 1$,

$$\begin{aligned}
& \alpha^2 \Gamma(1-\alpha) \eta \gamma \int_1^\infty \int_1^\infty (x^\alpha + y^\alpha) e^{-\Gamma(1-\alpha)\eta x^\alpha} e^{-\Gamma(1-\alpha)\gamma y^\alpha} (xy)^{-\alpha-1} dx dy \\
&= \Gamma(1-\alpha) \eta \gamma \left[E_1(\Gamma(1-\alpha)\eta) \int_1^\infty \frac{e^{-\Gamma(1-\alpha)\gamma z}}{z^2} dz + E_1(\Gamma(1-\alpha)\gamma) \int_1^\infty \frac{e^{-\Gamma(1-\alpha)\eta z}}{z^2} dz \right] \\
&\leq \Gamma(1-\alpha) \eta \gamma [E_1(\Gamma(1-\alpha)\eta) + E_1(\Gamma(1-\alpha)\gamma)] \\
&< \Gamma(1-\alpha) \left[\eta \gamma e^{-\Gamma(1-\alpha)\eta} \log \left(1 + \frac{1}{\Gamma(1-\alpha)\eta} \right) + \eta \gamma e^{-\Gamma(1-\alpha)\gamma} \log \left(1 + \frac{1}{\Gamma(1-\alpha)\gamma} \right) \right].
\end{aligned}$$

When $k = 2$,

$$\begin{aligned}
& (\alpha \Gamma(1-\alpha) \eta \gamma)^2 \int_1^\infty \int_1^\infty (x^{2\alpha} + y^{2\alpha}) e^{-\Gamma(1-\alpha)\eta x^\alpha} e^{-\Gamma(1-\alpha)\gamma y^\alpha} (xy)^{-\alpha-1} dx dy \\
&= \Gamma(1-\alpha) (\eta \gamma)^2 \left[\frac{e^{-\Gamma(1-\alpha)\eta}}{\eta} \int_1^\infty \frac{e^{-\Gamma(1-\alpha)\gamma z}}{z^2} dz + \int_1^\infty \frac{e^{-\Gamma(1-\alpha)\eta z}}{z^2} dz \frac{e^{-\Gamma(1-\alpha)\gamma}}{\gamma} \right] \\
&\leq \Gamma(1-\alpha) (\eta \gamma)^2 \left[\frac{e^{-\Gamma(1-\alpha)\eta}}{\eta} + \frac{e^{-\Gamma(1-\alpha)\gamma}}{\gamma} \right].
\end{aligned}$$

When $k > 2$,

$$\begin{aligned}
 & \alpha^2 \frac{(\Gamma(1-\alpha)\eta\gamma)^k}{k!} 2^{k-1} \int_1^\infty (x^{\alpha k} + y^{\alpha k}) e^{-\Gamma(1-\alpha)\eta x^\alpha} e^{-\Gamma(1-\alpha)\gamma y^\alpha} (xy)^{-\alpha-1} dx dy \\
 &= \alpha^2 \frac{(\Gamma(1-\alpha)\eta\gamma)^k}{k!} 2^{k-1} \left[\int_1^\infty \frac{e^{-\Gamma(1-\alpha)\eta x^\alpha}}{x^{\alpha(1-k)+1}} dx \int_1^\infty \frac{e^{-\Gamma(1-\alpha)\gamma y^\alpha}}{y^{\alpha+1}} dy \right. \\
 &\quad \left. + \int_1^\infty \frac{e^{-\Gamma(1-\alpha)\eta x^\alpha}}{x^{\alpha+1}} dx \int_1^\infty \frac{e^{-\Gamma(1-\alpha)\gamma y^\alpha}}{y^{\alpha(1-k)+1}} dy \right] \\
 &= \alpha^2 \frac{(\Gamma(1-\alpha)\eta\gamma)^k}{k!} 2^{k-1} \\
 &\quad \times \left[\int_{\Gamma(1-\alpha)\eta}^\infty \left(\frac{z}{\Gamma(1-\alpha)\eta} \right)^{k-2} e^{-z} \frac{dz}{\alpha\Gamma(1-\alpha)\eta} \int_1^\infty \frac{e^{-\Gamma(1-\alpha)\gamma z}}{z^2} dz \frac{1}{\alpha} \right. \\
 &\quad \left. + \int_1^\infty \frac{e^{-\Gamma(1-\alpha)\eta z}}{z^2} dz \frac{1}{\alpha} \int_{\Gamma(1-\alpha)\gamma}^\infty \left(\frac{z}{\Gamma(1-\alpha)\gamma} \right)^{k-2} e^{-z} \frac{dz}{\alpha\Gamma(1-\alpha)\gamma} \right] \\
 &\leq \alpha^2 \frac{(\Gamma(1-\alpha)\eta\gamma)^k}{k!} 2^{k-1} \left[\left(\frac{1}{\Gamma(1-\alpha)\eta} \right)^{k-1} \frac{1}{\alpha^2} \int_{\Gamma(1-\alpha)\eta}^\infty z^{k-2} e^{-z} dz \right. \\
 &\quad \left. + \left(\frac{1}{\Gamma(1-\alpha)\gamma} \right)^{k-1} \frac{1}{\alpha^2} \int_{\Gamma(1-\alpha)\gamma}^\infty z^{k-2} e^{-z} dz \right] \\
 &\leq \frac{\Gamma(1-\alpha)}{k!} 2^{k-1} \Gamma(k-1) (\eta\gamma)^k \left[\left(\frac{1}{\eta} \right)^{k-1} + \left(\frac{1}{\gamma} \right)^{k-1} \right] \\
 &= \Gamma(1-\alpha) \frac{2^{k-1}}{k(k-1)} (\eta\gamma^k + \eta^k\gamma).
 \end{aligned}$$

Combining together all the bounds, we have

$$\begin{aligned}
 & \left| \mathbb{E}[(1-\eta)^{D_\infty(1)}(1-\gamma)^{D_\infty(2)}] - \mathbb{E}[(1-\eta)^{D_\infty(1)}] \mathbb{E}[(1-\gamma)^{D_\infty(2)}] \right| \\
 &< \Gamma(1-\alpha) \left[\eta\gamma \log \left(1 + \frac{1}{\Gamma(1-\alpha)\eta} \right) + \eta\gamma \log \left(1 + \frac{1}{\Gamma(1-\alpha)\gamma} \right) \right. \\
 &\quad \left. + \frac{1}{2} \sum_{k=2}^\infty \frac{2^k}{k(k-1)} (\eta\gamma^k + \eta^k\gamma) \right].
 \end{aligned}$$

Since $x \log(1 + (1/x)) \rightarrow 0$ as $x \rightarrow 0$, the above quantity goes to 0 as $\eta, \gamma \rightarrow 0$. This completes the proof of Theorem 2.1. $\square$

5. Proof of Theorem 2.2: Wedges and triangles

Proof of Theorem 2.2. For (i), start from the equality

$$\begin{aligned}
 2\mathbb{E}[\mathbb{W}_n(i)] &= \mathbb{E} \left[\sum_{j \neq i} \sum_{k \neq i, j} a_{ij} a_{ik} \right] \\
 &= (n-1)(n-2)\alpha^3 \int_1^\infty \int_1^\infty \int_1^\infty \frac{(1-e^{-\varepsilon xy})(1-e^{-\varepsilon xz})}{(xyz)^{\alpha+1}} dx dy dz.
 \end{aligned}$$

We split the terms into two integrals by restricting the x variable to take values between $(0, 1/\varepsilon)$ and $(1/\varepsilon, \infty)$, and write $2\mathbf{E}[\mathbb{W}_n(i)] = (n-1)(n-2)(A+B)$, where

$$\begin{aligned} A &= \alpha^3 \int_1^{1/\varepsilon} \int_1^\infty \int_1^\infty \frac{(1 - e^{-\varepsilon xy})(1 - e^{-\varepsilon xz})}{(xyz)^{\alpha+1}} dy dz dx, \\ B &= \alpha^3 \int_{1/\varepsilon}^\infty \int_1^\infty \int_1^\infty \frac{(1 - e^{-\varepsilon xy})(1 - e^{-\varepsilon xz})}{(xyz)^{\alpha+1}} dy dz dx. \end{aligned} \quad (5.1)$$

We first provide a bound for B . Note that $x \geq 1/\varepsilon$ and $y \geq 1$, so we have $1 - e^{-1} \leq 1 - e^{-\varepsilon xy} \leq 1$. Hence,

$$B = \alpha^3 \int_{1/\varepsilon}^\infty \frac{1}{x^{\alpha+1}} \left(\int_1^\infty \frac{(1 - e^{-\varepsilon xy})}{y^{\alpha+1}} dy \right)^2 \leq \alpha^3 \int_{1/\varepsilon}^\infty \frac{dx}{x^{\alpha+1}} \left(\int_1^\infty \frac{dy}{y^{\alpha+1}} \right)^2 = \varepsilon^\alpha.$$

A lower bound for B can be obtained similarly:

$$B \geq \alpha^3 (1 - e^{-1})^2 \int_{1/\varepsilon}^\infty \frac{1}{x^{\alpha+1}} \left(\int_1^\infty \frac{dy}{y^{\alpha+1}} \right)^2 dx = (1 - e^{-1})^2 \varepsilon^\alpha.$$

So we have proved that $B \asymp \varepsilon^\alpha$.

Now we bound the term A in (5.1):

$$A = \alpha^3 \int_1^{1/\varepsilon} \frac{1}{x^{\alpha+1}} \left(\int_1^\infty \frac{1 - e^{-\varepsilon xy}}{y^{\alpha+1}} dy \right)^2 dx = \alpha^3 \int_1^{1/\varepsilon} \frac{1}{x^{\alpha+1}} \left[\frac{1}{\alpha} - (\varepsilon x)^\alpha \Gamma(-\alpha; \varepsilon x) \right]^2 dx,$$

where $\Gamma(-s; \varepsilon)$ is the incomplete Gamma function. When ε is small, the following expansion can be used [2]:

$$\Gamma(s; \varepsilon) = \Gamma(s) - \sum_{k=0}^\infty (-1)^k \frac{\varepsilon^{s+k}}{k!(s+k)}, \quad s \neq 0, -1, -2, -3, \dots \quad (5.2)$$

Thus

$$\begin{aligned} A &= \alpha^3 \int_1^{1/\varepsilon} \frac{1}{x^{\alpha+1}} \left[-(\varepsilon x)^\alpha \Gamma(-\alpha) + \sum_{l=1}^\infty \frac{(-1)^l}{l!(l-\alpha)} (\varepsilon x)^l \right]^2 dx \\ &= \alpha^3 \int_1^{1/\varepsilon} \frac{1}{x^{\alpha+1}} \left[\Gamma^2(-\alpha) (\varepsilon x)^{2\alpha} - 2\Gamma(-\alpha) \sum_{l=1}^\infty \frac{(-1)^l}{l!(l-\alpha)} (\varepsilon x)^{l+\alpha} + \left(\sum_{l=1}^\infty \frac{(-1)^l}{l!(l-\alpha)} (\varepsilon x)^l \right)^2 \right] dx \\ &= \alpha^3 \Gamma^2(-\alpha) \varepsilon^{2\alpha} \left[\frac{x^\alpha}{\alpha} \right]_1^{1/\varepsilon} - 2\alpha^3 \Gamma(-\alpha) \sum_{l=1}^\infty \frac{(-1)^l}{l!(l-\alpha)} \varepsilon^{l+\alpha} \left[\frac{x^l}{l} \right]_1^{1/\varepsilon} + C(\varepsilon) \\ &= \alpha^2 \Gamma^2(-\alpha) (\varepsilon^\alpha - \varepsilon^{2\alpha}) - 2\alpha^3 \Gamma(-\alpha) \sum_{l=1}^\infty \frac{(-1)^l}{l \cdot l!(l-\alpha)} (\varepsilon^\alpha - \varepsilon^{l+\alpha}) + C(\varepsilon) \\ &= O(\varepsilon^\alpha) + C(\varepsilon) \end{aligned}$$

as $\varepsilon \rightarrow 0+$, where we have introduced

$$\begin{aligned} C(\varepsilon) &= \alpha^3 \int_1^{1/\varepsilon} \frac{1}{x^{\alpha+1}} \left(\sum_{l=1}^{\infty} \frac{(-1)^l}{l!(l-\alpha)} (\varepsilon x)^l \right)^2 dx \\ &= \sum_{l=1}^{\infty} d_{l,\alpha} \varepsilon^{2l} [x^{2l-\alpha}]_1^{1/\varepsilon} + \sum_{\substack{l,k=1 \\ k \neq l}}^{\infty} d_{l,k,\alpha} \varepsilon^{l+k} [x^{l+k-\alpha}]_1^{1/\varepsilon} \\ &= \sum_{l=1}^{\infty} d_{l,\alpha} (\varepsilon^\alpha - \varepsilon^{2l}) + \sum_{\substack{l,k=1 \\ k \neq l}}^{\infty} d_{l,k,\alpha} (\varepsilon^\alpha - \varepsilon^{l+k}) = O(\varepsilon^\alpha), \end{aligned}$$

with $d_{l,\alpha}$ and $d_{l,k,\alpha}$ suitable constants depending on l, k , and α . So, we can deduce that $A + B = O(\varepsilon^\alpha)$. Hence we have shown that $\mathbf{E}[\mathbb{W}_n(i)] \asymp n^2 \varepsilon^\alpha$, as desired.

For (ii), we now assume that $\varepsilon = n^{-1/\alpha}$. From Theorem 2.1 we know that $D_n(i) \xrightarrow{d} D_\infty(i)$. Using the continuous mapping $x \mapsto x(x-1)$ we have, by the continuous mapping theorem, convergence in distribution of the number of wedges $\mathbb{W}_n(i)$:

$$\mathbb{W}_n(i) = D_n(i)(D_n(i) - 1) \xrightarrow{d} D_\infty(i)(D_\infty(i) - 1) \equiv \mathbb{W}_\infty(i).$$

We then show that the tail satisfies $\mathbb{P}(\mathbb{W}_\infty(i) > x) \sim \Gamma(1-\alpha)x^{-1/2}$ as $x \rightarrow \infty$ by bounding the ratio

$$\frac{\mathbb{P}(D_\infty(i)^2 - D_\infty(i) > x)}{\mathbb{P}(D_\infty(i)^2 > x)},$$

where, by (2.4), $\mathbb{P}(D_\infty(i)^2 > x) \sim \Gamma(1-\alpha)x^{-1/2}$. Let $\delta > 0$. Then

$$\begin{aligned} \mathbb{P}(D_\infty(i)^2 - D_\infty(i) > x) &= \mathbb{P}(D_\infty(i)^2 > x + D_\infty(i), D_\infty(i) > x + \delta) \\ &\quad + \mathbb{P}(D_\infty(i)^2 > x + D_\infty(i), D_\infty(i) \leq x + \delta) \\ &\leq \mathbb{P}(D_\infty(i) > x + \delta) + \mathbb{P}(D_\infty(i)^2 > x). \end{aligned}$$

This implies that, for any $\delta > 0$,

$$\frac{\mathbb{P}(\mathbb{W}_\infty(i) > x)}{\mathbb{P}(D_\infty(i)^2 > x)} \leq \frac{\mathbb{P}(D_\infty(i) > x + \delta)}{\mathbb{P}(D_\infty(i)^2 > x)} + 1 \sim \frac{\Gamma(1-\alpha)(x+\delta)^{-1}}{\Gamma(1-\alpha)x^{-1/2}} + 1 \xrightarrow{x \rightarrow \infty} 1.$$

A similar break-up yields the lower bound:

$$\begin{aligned} \mathbb{P}(D_\infty(i)^2 - D_\infty(i) > x) &\geq \mathbb{P}(D_\infty(i)^2 > (1+\delta)x, D_\infty(i) \leq \delta x) \\ &\geq \mathbb{P}(D_\infty(i)^2 > (1+\delta)x) - \mathbb{P}(D_\infty(i) > \delta x), \end{aligned}$$

which yields

$$\frac{\mathbb{P}(D_\infty(i)^2 - D_\infty(i) > x)}{\mathbb{P}(D_\infty(i)^2 > x)} \geq \frac{((1+\delta)x)^{-1/2} - (\delta x)^{-1}}{x^{-1/2}} \xrightarrow{x \rightarrow \infty} \frac{1}{\sqrt{1+\delta}}.$$

Hence,

$$\limsup_{x \rightarrow \infty} \frac{\mathbb{P}(\mathbb{W}_\infty(i) > x)}{\Gamma(1-\alpha)x^{-1/2}} \leq 1; \quad \liminf_{x \rightarrow \infty} \frac{\mathbb{P}(\mathbb{W}_\infty(i) > x)}{\Gamma(1-\alpha)x^{-1/2}} \geq \frac{1}{\sqrt{(1+\delta)}}.$$

The result follows by taking $\delta \rightarrow 0$.

For (iii), we focus on the average number of triangles, whose evaluation will require integral asymptotics similar to those used for the wedges:

$$\begin{aligned} 6\mathbb{E}[\Delta_n(i)] &= \mathbb{E}\left[\sum_{j \neq i} \sum_{k \neq i,j} a_{ij}a_{ik}a_{jk}\right] \\ &= \sum_{j \neq i} \sum_{k \neq i,j} \mathbb{E}[p_{ij}p_{ik}p_{jk}] \\ &= (n-1)(n-2)\alpha^3 \int_1^\infty \int_1^\infty \int_1^\infty \frac{(1-e^{-\varepsilon xy})(1-e^{-\varepsilon xz})(1-e^{-\varepsilon yz})}{(xyz)^{\alpha+1}} dx dy dz. \end{aligned}$$

First, we provide an upper bound for the above integral. To do this, we use the variables $A = xy$, $B = xz$, and $C = yz$. Note that by setting $x = \sqrt{AC/B}$, $y = \sqrt{AB/C}$, and $z = \sqrt{BC/A}$, and considering that the Jacobian of this transformation is $\frac{1}{4}(ABC)^{-1/2}$, we have

$$\begin{aligned} &\alpha^3 \int_1^\infty \int_1^\infty \int_1^\infty \frac{(1-e^{-\varepsilon xy})(1-e^{-\varepsilon xz})(1-e^{-\varepsilon yz})}{(xyz)^{\alpha+1}} dx dy dz \\ &= \alpha^3 \int_1^\infty \int_1^\infty \int_{(A/B) \vee (B/A)}^{AB} \frac{(1-e^{-\varepsilon A})(1-e^{-\varepsilon B})(1-e^{-\varepsilon C})}{(\sqrt{ABC})^{\alpha+1}} \frac{1}{4\sqrt{ABC}} dC dB dA \\ &= \frac{\alpha^3}{4} \int_1^\infty \int_1^\infty \int_{(A/B) \vee (B/A)}^{AB} \frac{(1-e^{-\varepsilon A})(1-e^{-\varepsilon B})(1-e^{-\varepsilon C})}{(ABC)^{\alpha/2+1}} dC dB dA. \end{aligned}$$

Observing that for $A, B \geq 1$ the interval $((A/B) \vee (B/A), AB)$ is contained in $[1, \infty)$, the last integral can be upper bounded by

$$\begin{aligned} &\frac{\alpha^3}{4} \int_1^\infty \int_1^\infty \int_{(A/B) \vee (B/A)}^{AB} \frac{(1-e^{-\varepsilon A})(1-e^{-\varepsilon B})(1-e^{-\varepsilon C})}{(ABC)^{\alpha/2+1}} dC dB dA \\ &\leq \frac{\alpha^3}{4} \int_1^\infty \int_1^\infty \int_1^\infty \frac{(1-e^{-\varepsilon A})(1-e^{-\varepsilon B})(1-e^{-\varepsilon C})}{(ABC)^{\alpha/2+1}} dC dB dA \\ &= \frac{\alpha^3}{4} \left(\int_1^\infty \frac{1-e^{-\varepsilon A}}{A^{\alpha/2+1}} dA \right)^3 \\ &= \frac{\alpha^3}{4} \left[\frac{2}{\alpha} - \varepsilon^{\alpha/2} \Gamma\left(-\frac{\alpha}{2}; \varepsilon\right) \right]^3 \\ &= \frac{\alpha^3}{4} \left[-\varepsilon^{\alpha/2} \Gamma\left(-\frac{\alpha}{2}\right) + O(\varepsilon) \right]^3 = -\frac{n^2 \alpha^3}{2} \varepsilon^{3\alpha/2} \Gamma\left(-\frac{\alpha}{2}\right)^3 + O(n^2 \varepsilon^3), \end{aligned}$$

where in the last step we used the expansion approximating the incomplete Gamma function (5.2). By our assumption, since $\alpha < 2$ and $\varepsilon \rightarrow 0$, we have $\varepsilon^3 = o(\varepsilon^{3\alpha/2})$, and hence $\mathbb{E}[\Delta_n(i)] = O(n^2 \varepsilon^{3\alpha/2})$.

To complete the proof, we now show a lower bound that matches the correct order. Specifically, we provide a lower bound for the integral

$$I = \int_1^\infty \int_1^\infty \int_1^\infty \frac{(1-e^{-\varepsilon xy})(1-e^{-\varepsilon xz})(1-e^{-\varepsilon yz})}{(xyz)^{\alpha+1}} dx dy dz.$$

First, we perform a change of variables with $x = u/\sqrt{\varepsilon}$, $y = v/\sqrt{\varepsilon}$, and $z = w/\sqrt{\varepsilon}$. Then we obtain

$$I = \varepsilon^{3\alpha/2} \int_{\sqrt{\varepsilon}}^{\infty} \int_{\sqrt{\varepsilon}}^{\infty} \int_{\sqrt{\varepsilon}}^{\infty} \frac{(1 - e^{-uv})(1 - e^{-vw})(1 - e^{-uw})}{(uvw)^{\alpha+1}} du dv dw.$$

Since $\varepsilon < 1$, it is not difficult to see that

$$I \geq \varepsilon^{3\alpha/2} \int_1^{\infty} \int_1^{\infty} \int_1^{\infty} \frac{(1 - e^{-uv})(1 - e^{-vw})(1 - e^{-uw})}{(uvw)^{\alpha+1}} du dv dw = c_{\alpha} \varepsilon^{3\alpha/2},$$

where $c_{\alpha} \in (0, \infty)$ denotes the value of the above integral. This completes the proof of the lower bound.

For (iv), let $\varepsilon = n^{-1/\alpha}$; then $\mathbb{E}[\Delta_n(i)] \asymp n^{1/2}$. The above computations also show that $\Delta_n = \sum_{i=1}^n \Delta_n(i)$ behaves as

$$\mathbb{E}[\Delta_n] \asymp n^{3/2}. \quad (5.3)$$

To study the concentration of the latter quantity, we start by evaluating the second moment:

$$\mathbb{E}[\Delta_n^2] = \mathbb{E} \left[\sum'_{i,j,k} \sum'_{u,v,w} a_{ij} a_{ik} a_{jk} a_{uv} a_{uw} a_{vw} \right] = A + B + C + D,$$

where A represents the term in which there is no intersection between the triples of indices of the two summations $((u, v, w) \neq (i, j, k))$, i.e. $|\{u, v, w\} \cap \{i, j, k\}| = 0$; B is the term in which there is an intersection of one index, i.e. $|\{u, v, w\} \cap \{i, j, k\}| = 1$; C is an intersection of two indices, i.e. $|\{u, v, w\} \cap \{i, j, k\}| = 2$; and D is the term in which all the indices coincide, $|\{u, v, w\} \cap \{i, j, k\}| = 3$. $\sum'_{i,j,k}$ means sum over distinct indices.

For A , with no common indices,

$$\begin{aligned} A &= \mathbb{E} \left[\sum'_{i,j,k} \sum'_{(u,v,w) \neq (i,j,k)} a_{ij} a_{ik} a_{jk} a_{uv} a_{uw} a_{vw} \right] \\ &= \mathbb{E} \left[\sum'_{i,j,k} a_{ij} a_{ik} a_{jk} \right] \mathbb{E} \left[\sum'_{u,v,w} a_{uv} a_{uw} a_{vw} \right] = \mathbb{E}[\Delta_n]^2. \end{aligned}$$

For B , with one common index,

$$\begin{aligned} B &= \mathbb{E} \left[\sum'_{i,j,k} \sum'_{1 \text{ intersection}} a_{ij} a_{ik} a_{jk} a_{uv} a_{uw} a_{vw} \right] \\ &= \mathbb{E} \left[\sum_{i,j,k} a_{ij} a_{ik} a_{jk} 3 \sum_{v,w} a_{vw} (a_{iv} a_{iw} + a_{jv} a_{jw} + a_{kv} a_{kw}) \right] \\ &\leq \mathbb{E} \left[\sum_{i,j,k} a_{ij} a_{ik} a_{jk} 9 \sum_{v,w} a_{vw} \right] = 9n \mathbb{E}[\Delta_n] \mathbb{E}[D_n(i)]. \end{aligned}$$

For C , with exactly two common indices,

$$C = \mathbf{E} \left[\sum'_{i,j,k} \sum'_{2 \text{ intersections}} a_{ij} a_{ik} a_{jk} a_{uv} a_{uw} a_{vw} \right] \leq 6n \mathbf{E} \left[\sum'_{i,j,k} a_{ij} a_{ik} a_{jk} \right] = 6n \mathbf{E}[\Delta_n].$$

For D , with all indices matching,

$$D = \mathbf{E} \left[\sum'_{i,j,k} a_{ij} a_{ik} a_{jk} a_{ij} a_{ik} a_{jk} \right] = \mathbf{E} \left[\sum'_{i,j,k} a_{ij} a_{ik} a_{jk} \right] = \mathbf{E}[\Delta_n].$$

Therefore,

$$\begin{aligned} \frac{\text{Var}(\Delta_n)}{\mathbf{E}[\Delta_n]^2} &= \frac{\mathbf{E}[\Delta_n^2] - \mathbf{E}[\Delta_n]^2}{\mathbf{E}[\Delta_n]^2} = \frac{B + C + D}{\mathbf{E}[\Delta_n]^2} \\ &\leq \frac{9n \mathbf{E}[\Delta_n] \mathbf{E}[D_n(i)] + 6n \mathbf{E}[\Delta_n] + \mathbf{E}[\Delta_n]}{\mathbf{E}[\Delta_n]^2} \\ &\leq \frac{c_7 (n^{5/2} \log n + n^{5/2} + n^{3/2})}{c_6 n^3} = O\left(\frac{\log n}{n^{1/2}}\right), \end{aligned}$$

where c_6 and c_7 are positive constants, taken respectively from (5.3) and (2.3). Now, using Chebyshev's inequality it follows that for any $\delta > 0$,

$$\mathbb{P}\left(\left|\frac{\Delta_n}{\mathbf{E}[\Delta_n]} - 1\right| \geq \delta\right) \leq \frac{\text{Var}(\Delta_n)}{\delta^2 \mathbf{E}[\Delta_n]^2} = O\left(\frac{\log n}{n^{1/2}}\right).$$

This completes the proof of the first statement in part (iv). The second one follows from the very same computations. $\square$

6. Absence of dust: Proof of Proposition 2.1

We begin with some nice bounds for the incomplete Gamma function.

Lemma 6.1. *Let $s = (1 - \alpha) \in (0, 1)$ and let $\Gamma(s, x) = \int_x^\infty e^{-z} z^{s-1} dz$, $x > 0$. Then*

$$e^{-x}(1+x)^{s-1} \leq \Gamma(s, x) \leq e^{-x} \Gamma(s) \left(1 + \frac{x}{s}\right)^{s-1}.$$

Proof. The proof is a standard use of Jensen's inequality but we provide the details for completeness. First we show the lower bound. Let

$$\Gamma(s, x) = \int_x^\infty z^{s-1} e^{-z} dz = e^{-x} \int_0^\infty (z+x)^{s-1} e^{-z} dz = e^{-x} \mathbb{E}[(Z+x)^{s-1}],$$

where Z is an exponential random variable with parameter 1. Define the function $g(t) = (t+x)^{s-1}$. As $s \in (0, 1)$, $g(t)$ is a convex function. Hence, by Jensen's inequality we have

$$\Gamma(s, x) = e^{-x} \mathbb{E}[g(Z)] \geq e^{-x} g(\mathbb{E}[Z]) = e^{-x} g(1),$$

which gives the lower bound. For the upper bound, we again represent the incomplete Gamma function as

$$\Gamma(s, x) = e^{-x} \int_0^\infty \left(1 + \frac{x}{z}\right)^{s-1} z^{s-1} e^{-z} dz = e^{-x} \Gamma(s) \mathbb{E} \left[\left(1 + \frac{x}{G}\right)^{s-1} \right],$$

where G is a Gamma distribution with shape parameter s and rate 1. Using the function $h(t) = (1 + x/t)^{s-1}$, which is a concave function, and using the reverse Jensen's inequality we have

$$\Gamma(s, x) = e^{-x} \Gamma(s) \mathbb{E}[h(G)] \leq e^{-x} \Gamma(s) h(\mathbb{E}[G]) = e^{-x} \Gamma(s) h(s),$$

where we used $\mathbb{E}[G] = s$. This gives the upper bound. $\square$

The following is a non-standard bound that improves the well-known bound $1 - e^{-x} \leq x$.

Lemma 6.2. For $\alpha \in [0, 1]$ and $x > 0$, $1 - e^{-x} \leq x^\alpha$.

Proof. The proof follows by considering the function $h(\alpha, x) = 1 - e^{-x} - x^\alpha$ and studying the derivative with respect to α , $\partial h(\alpha, x)/\partial \alpha = -x^\alpha \ln x$. $\square$

The next lemma gives a bound on the conditional degree distribution.

Lemma 6.3. For $i \in [n]$, $j \neq i$, and $w > 1$,

$$\varepsilon^\alpha w^\alpha e^{-(1+\alpha)\varepsilon w} \leq \mathbb{E}[p_{ij} \mid W_i = w] \leq \varepsilon^\alpha w^\alpha (1 + \Gamma(1 - \alpha) e^{-\varepsilon w}).$$

In particular, for fixed $w > 1$,

$$\mathbb{E}[p_{ij} \mid W_i = w] \sim \Gamma(1 - \alpha) \varepsilon^\alpha w^\alpha \quad \text{as } \varepsilon \downarrow 0. \quad (6.1)$$

Proof.

$$\begin{aligned} \mathbb{E}[p_{ij} \mid W_i = w] &= \mathbb{E}[(1 - e^{-\varepsilon W_j w})] \\ &= \int_0^\infty (1 - e^{-\varepsilon w u}) F_W(du) = \int_0^\infty \int_0^u e^{-\varepsilon w z} (\varepsilon w) dz F_W(du) \\ &= \varepsilon w \int_0^\infty e^{-\varepsilon w z} \mathbb{P}(W_j > z) dz, \end{aligned}$$

where we used Fubini's theorem to swap the integrals. Using the fact that the W_j are supported on $(1, \infty)$ we get

$$\begin{aligned} \mathbb{E}[p_{ij} \mid W_i = w] &= \varepsilon w \int_0^1 e^{-\varepsilon w z} dz + \varepsilon w \int_1^\infty e^{-\varepsilon w z} z^{-\alpha} dz \\ &= (1 - e^{-\varepsilon w}) + \varepsilon^\alpha w^\alpha \int_{\varepsilon w}^\infty e^{-z} z^{-\alpha} dz = (1 - e^{-\varepsilon w}) + \varepsilon^\alpha w^\alpha \Gamma(1 - \alpha, \varepsilon w), \end{aligned} \quad (6.2)$$

where $\Gamma(1 - \alpha, \varepsilon w)$ is an incomplete Gamma function. Now we can use the bounds from Lemmas 6.1 and 6.2. For the upper bound we have

$$\mathbb{E}[p_{ij} \mid W_i = w] \leq \varepsilon^\alpha w^\alpha + \varepsilon^\alpha w^\alpha \Gamma(1 - \alpha) e^{-\varepsilon w} \left(1 + \frac{\varepsilon w}{1 - \alpha}\right)^{-\alpha}.$$

Now using the trivial bounds $1 + \varepsilon w/(1 - \alpha) \geq 1$ we have

$$\mathbb{E}[p_{ij} \mid W_i = w] \leq \varepsilon^\alpha w^\alpha (1 + \Gamma(1 - \alpha)e^{-\varepsilon w}).$$

For the lower bound, we ignore the first term in (6.2) and use the lower bound in Lemma 6.1. We have

$$\mathbb{E}[p_{ij} \mid W_i = w] \geq \varepsilon^\alpha w^\alpha \Gamma(1 - \alpha, \varepsilon w) \geq \varepsilon^\alpha w^\alpha e^{-\varepsilon w} (1 + \varepsilon w)^{-\alpha} \geq \varepsilon^\alpha w^\alpha e^{-\varepsilon w} e^{-\alpha \varepsilon w},$$

where we used $1 + x \leq e^x$ for $x > 0$, and hence $(1 + x)^{-\alpha} \geq e^{-\alpha x}$. This completes the proof of the lower bound. The asymptotic estimate in (6.1) follows from (6.2). $\square$

Proof of Proposition 2.1. From the definition of the number of isolated vertices in (2.8), we compute

$$\begin{aligned} \mathbb{E}[N_0] &= \sum_{i=1}^n \mathbf{P}(i \text{ is isolated}) = \sum_{i=1}^n \mathbb{E} \left[\prod_{k \neq i} (1 - p_{ik}) \right] \\ &= \sum_{i=1}^n \mathbb{E} \left[\mathbb{E}_{W_i} \left[\prod_{k \neq i} e^{-\varepsilon W_i W_k} \right] \right] \\ &= \sum_{i=1}^n \mathbb{E} \left(\prod_{k \neq i} \mathbb{E}_{W_i} [e^{-\varepsilon W_i W_k}] \right) \\ &= n \mathbb{E} [(\mathbb{E}_{W_1} [e^{-\varepsilon W_1 W_2}])^{n-1}], \end{aligned} \quad (6.3)$$

where we have used the property of the weights being independent and identically distributed.

We can then estimate asymptotically the last expression by means of (6.1) to obtain

$$\mathbb{E}[(\mathbb{E}_{W_1} [e^{-\varepsilon W_1 W_2}])^{n-1}] = \mathbb{E}[e^{(n-1) \log(1 - \mathbb{E}_{W_1}(p_{12}))}] \sim \mathbb{E}[e^{-(n-1)\Gamma(1-\alpha)\varepsilon^\alpha W_1^\alpha}].$$

Plugging this estimate into (6.3) we get

$$\mathbb{E}[N_0] \sim n \mathbb{E}[e^{-(n-1)\Gamma(1-\alpha)\varepsilon^\alpha W_1^\alpha}] = \mathbb{E} \left[\exp \left\{ -(n-1)\varepsilon^\alpha \left(W_1^\alpha - \frac{\log n}{(n-1)\varepsilon^\alpha} \right) \right\} \right].$$

By the Markov inequality $P(N_0 \geq 1) \leq \mathbb{E}[N_0]$, and noticing that the right-hand side goes to zero as n grows, as soon as $\varepsilon^\alpha \gg (\log n)/(n-1)$ the claim in (2.9) follows. $\square$

Acknowledgements

The authors thank the referees for their valuable comments. We also thank Elena Matteini for spotting a mistake in the previous version of the article.

Funding information

Part of this work started during M. Lalli's internship in Lorentz Institute of Theoretical Physics in Leiden, which was supported by the ERASMUS+ Grant Agreement no. 2020-1-IT02-KA103-077946/03. L. Avena and R. S. Hazra were supported by NWO Gravitation Grant 024.002.003-NETWORKS. D. Garlaschelli acknowledges support from the Dutch Econophysics Foundation (Stichting Econophysics, Leiden, the Netherlands). His work is

also supported by the European Union – NextGenerationEU – National Recovery and Resilience Plan (Piano Nazionale di Ripresa e Resilienza, PNRR), projects ‘SoBigData.it – Strengthening the Italian RI for Social Mining and Big Data Analytics’ – Grant IR0000013 (no. 3264, 28/12/2021) (<https://pnrr.sobigdata.it/>) and ‘Reconstruction, Resilience and Recovery of Socio-Economic Networks’ RECON-NET EP_FAIR_005 – PE0000013 ‘FAIR’ – PNRR M4C2 Investment 1.3.

Competing interests

There were no competing interests to declare which arose during the preparation or publication process of this article.

References

- [1] ANDERS HEDEGAARD JESSEN, T. M. (2006). Regularly varying functions. *Publ. Inst. Math.* **80**, 171–192.
- [2] BENDER, C. M., ORSZAG, S. AND ORSZAG, S. A. (1999). *Advanced Mathematical Methods for Scientists and Engineers I: Asymptotic Methods and Perturbation Theory*, vol. 1. Springer, New York.
- [3] BHAMIDI, S., VAN DER HOFSTAD, R. AND SEN, S. (2018). The multiplicative coalescent, inhomogeneous continuum random trees, and new universality classes for critical random graphs. *Prob. Theory Relat. Fields* **170**, 387–474.
- [4] BHAMIDI, S., VAN DER HOFSTAD, R. AND VAN LEEUWAARDEN, J. (2010). Scaling limits for critical inhomogeneous random graphs with finite third moments. *Electron. J. Prob.* **15**, 1682–1702.
- [5] BHATTACHARJEE, C. AND SCHULTE, M. (2022). Large degrees in scale-free inhomogeneous random graphs. *Ann. Appl. Prob.* **32**, 696–720.
- [6] BINGHAM, N. H., GOLDIE, C. M., TEUGELS, J. L. AND TEUGELS, J. (1989). *Regular Variation*. Cambridge University Press.
- [7] BOGUNA, M., BONAMASSA, I., DE DOMENICO, M., HAVLIN, S., KRIOUKOV, D. AND SERRANO, M. (2021). Network geometry. *Nature Rev. Phys.* **3**, 114–135.
- [8] BOGUÑA, M. AND PASTOR-SATORRAS, R. (2003). Class of correlated random networks with hidden variables. *Phys. Rev. E* **68**, 036112.
- [9] BOLLOBÁS, B., JANSON, S. AND RIORDAN, O. (2007). The phase transition in inhomogeneous random graphs. *Random Structures Algorithms* **31**, 3–122.
- [10] CALDARELLI, G., CAPOCCI, A., DE LOS RIOS, P. AND MUNOZ, M. A. (2002). Scale-free networks from varying vertex intrinsic fitness. *Phys. Rev. Lett.* **89**, 258702.
- [11] CHUNG, F. AND LU, L. (2002). The average distances in random graphs with given expected degrees. *Proc. Nat. Acad. Sci.* **99**, 15879–15882.
- [12] CIPRIANI, A. AND SALVI, M. (2021). Scale-free percolation mixing time. Preprint, [arXiv:2111.05201](https://arxiv.org/abs/2111.05201).
- [13] DEIJFEN, M., VAN DER HOFSTAD, R. AND HOOGHIEMSTRA, G. (2013). Scale-free percolation. *Ann. Inst. H. Poincaré Prob. Statist.* **49**, 817–838.
- [14] DEPREZ, P., HAZRA, R. S. AND WÜTHRICH, M. V. (2015). Inhomogeneous long-range percolation for real-life network modeling. *Risks* **3**, 1–23.
- [15] DEVROYE, L. AND FRAIMAN, N. (2014). Connectivity of inhomogeneous random graphs. *Random Structures Algorithms* **45**, 408–420.
- [16] GABRIELLI, A., GARLASCELLI, D., PATIL, S. P. AND SERRANO, M. Á. (2025). Network renormalization. *Nature Rev. Phys.* **7**, 203–219.
- [17] GARCA-PÉREZ, G., BOGUÑA, M. AND SERRANO, M. (2018). Multiscale unfolding of real networks by geometric renormalization. *Nature Phys.* **14**, 583–589.
- [18] GARLASCELLI, D., CAPOCCI, A. AND CALDARELLI, G. (2007). Self-organized network evolution coupled to extremal dynamics. *Nature Phys.* **3**, 813–817.
- [19] GARUCCIO, E., LALLI, M. AND GARLASCELLI, D. (2023). Multiscale network renormalization: Scale-invariance without geometry. *Phys. Rev. Res.* **5**, 043101.
- [20] GFELLER, D. AND DE LOS RIOS, P. (2007). Spectral coarse graining of complex networks. *Phys. Rev. Lett.* **99**, 038701.
- [21] HEYDENREICH, M., HULSHOF, T. AND JORRITSMA, J. (2017). Structures in supercritical scale-free percolation. *Ann. Appl. Prob.* **27**, 2569–2604.
- [22] KADANOFF, L. P. (2000). *Statistical Physics: Statics, Dynamics and Renormalization*. World Scientific, Singapore.

- [23] MICHIELAN, R., LITVAK, N. AND STEGEHUIS, C. (2022). Detecting hyperbolic geometry in networks: Why triangles are not enough. *Phys. Rev. E* **106**, 054303.
- [24] NORROS, I. AND REITTU, H. (2006). On a conditionally Poissonian graph process. *Adv. Appl. Prob.* **38**, 59–75.
- [25] RADICCHI, F., BARRAT, A., FORTUNATO, S. AND RAMASCO, J. J. (2009). Renormalization flows in complex networks. *Phys. Rev. E* **79**, 026104.
- [26] RESNICK, S. AND SAMORODNITSKY, G. (2015). Tauberian theory for multivariate regularly varying distributions with application to preferential attachment networks. *Extremes* **18**, 349–367.
- [27] RODGERS, G. J., AUSTIN, K., KAHNG, B. AND KIM, D. (2005). Eigenvalue spectra of complex networks. *J. Phys. A* **38**, 9431.
- [28] SAMORODNITSKY, G. AND TAQQU, M. (1994). *Stable Non-Gaussian Random Processes. Stochastic Models with Infinite Variance*. Chapman & Hall, New York.
- [29] SONG, C., HAVLIN, S. AND MAKSE, H. A. (2006). Origins of fractality in the growth of complex networks. *Nature Phys.* **2**, 275–281.
- [30] VAN DEN ESKEER, H., VAN DER HOFSTAD, R., HOOGHIEMSTRA, G. AND ZNAMENSKI, D. (2005). Distances in random graphs with infinite mean degrees. *Extremes* **8**, 111–141.
- [31] VAN DER HOFSTAD, R. (2013). Critical behavior in inhomogeneous random graphs. *Random Structures Algorithms* **42**, 480–508.
- [32] VAN DER HOFSTAD, R. (2016). *Random Graphs and Complex Networks*, vol. 1 (Camb. Ser. Statist. Prob. Math. **43**). Cambridge University Press.
- [33] VAN DER HOFSTAD, R. (2022). *Random Graphs and Complex Networks*, vol. 2. Available at: <https://www.win.tue.nl/rhofstad/NotesRGCNII.pdf>.
- [34] VAN DER HOFSTAD, R., JANSSEN, A. J. E. M., VAN LEEUWAARDEN, J. S. H. AND STEGEHUIS, C. (2017). Local clustering in scale-free networks with hidden variables. *Phys. Rev. E* **95**, 022307.
- [35] VAN DER HOFSTAD, R. AND KOMJATHY, J. (2017). Explosion and distances in scale-free percolation. Preprint, [arXiv:1706.02597](https://arxiv.org/abs/1706.02597).
- [36] VAN DER HOFSTAD, R. AND KOMJÁTHY, J. (2017). When is a scale-free graph ultra-small? *J. Statist. Phys.* **169**, 223–264.
- [37] VAN DER HOFSTAD, R., VAN DER HOORN, P., LITVAK, N. AND STEGEHUIS, C. (2020). Limit theorems for assortativity and clustering in null models for scale-free networks. *Adv. Appl. Prob.* **52**, 1035–1084.
- [38] VILLEGAS, P., GILI, T., CALDARELLI, G. AND GABRIELLI, A. (2023). Laplacian renormalization group for heterogeneous networks. *Nature Phys.* **19**, 445–450.
- [39] WILSON, K. G. (1983). The renormalization group and critical phenomena. *Rev. Mod. Phys.* **55**, 583.