



UNIVERSITÀ
DEGLI STUDI
FIRENZE

FLORE

Repository istituzionale dell'Università degli Studi di Firenze

Infinite families of homogeneous Bismut Ricci flat manifolds

Questa è la versione Preprint (Submitted version) della seguente pubblicazione:

Original Citation:

Infinite families of homogeneous Bismut Ricci flat manifolds / Fabio Podesta'; Alberto Rafferò. - In: COMMUNICATIONS IN CONTEMPORARY MATHEMATICS. - ISSN 1793-6683. - STAMPA. - ---:(2022), pp. 00-00. [10.1142/S0219199722500754]

Availability:

This version is available at: 2158/1290244 since: 2023-03-30T10:16:56Z

Published version:

DOI: 10.1142/S0219199722500754

Terms of use:

Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (<https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf>)

Publisher copyright claim:

(Article begins on next page)

INFINITE FAMILIES OF HOMOGENEOUS BISMUT RICCI FLAT MANIFOLDS

FABIO PODESTÀ AND ALBERTO RAFFERO

ABSTRACT. Starting from compact symmetric spaces of inner type, we provide infinite families of compact homogeneous spaces carrying invariant non-flat Bismut connections with vanishing Ricci tensor. These examples turn out to be generalized symmetric spaces of order 4 and (up to coverings) can be realized as minimal submanifolds of the Bismut flat model spaces, namely compact Lie groups. This construction generalizes the standard Cartan embedding of symmetric spaces.

1. INTRODUCTION

On a Riemannian n -manifold (M, g) , metric connections ∇ are completely characterized by their torsion tensor

$$T_X Y = \nabla_X Y - \nabla_Y X - [X, Y], \quad X, Y \in \Gamma(TM),$$

and they can be classified into eight families determined by the irreducible $O(n)$ -module decomposition of the space $\mathbb{R}^n \otimes \Lambda^2 \mathbb{R}^n$ [Car, TV]. The class of metric connections with totally skew-symmetric torsion is made up of those connections for which the 3-covariant tensor

$$H(X, Y, Z) := g(T_X Y, Z)$$

is totally skew-symmetric. Every such connection is related to the Levi Civita connection ∇^g of (M, g) as follows

$$(1.1) \quad g(\nabla_X Y, Z) = g(\nabla_X^g Y, Z) + \frac{1}{2}H(X, Y, Z),$$

and it has the same geodesics as ∇^g .

Consistently with [GFS], given a 3-form $H \in \Omega^3(M)$, we shall refer to the metric connection ∇ defined in (1.1) as the *Bismut connection* induced by the pair (g, H) .

In complex non-Kähler geometry, a typical example is given by the Bismut connection of a Hermitian manifold (M, g, J) , which is the unique metric connection with totally skew-symmetric torsion preserving the complex structure J [Bis, Gau]. In such a case, the torsion is given by $H = d^c \omega$, where $\omega = g(J \cdot, \cdot)$ is the fundamental 2-form. If the torsion is also closed, so that $dd^c \omega = 0$, one obtains the widely studied class of strong Kähler with torsion complex manifolds (SKT). On the other hand, Riemannian manifolds carrying a Bismut connection ∇ with ∇ -parallel torsion have gained a lot of attention in the literature, since they are naturally associated with various geometrically meaningful structures as

2020 *Mathematics Subject Classification.* 53C25, 53C07, 53C30, 53B05, 53E20.

Key words and phrases. Bismut connection, Ricci flat connection, homogeneous space.

naturally reductive spaces, nearly Kähler and Sasakian structures as well as nearly parallel G_2 -structures among others, see e.g. [Agr, AFF, CMS, FI].

Metric connections with totally skew symmetric torsion are also an important object of interest in theoretical and mathematical physics, e.g. in Type II string theory or in supergravity theories, see [Agr, FI, IP] for more details and references.

In this paper, we will focus on Bismut connections ∇ with closed torsion form H and zero Ricci tensor Ric^∇ . In generalized Riemannian geometry, Bismut connections with closed torsion play a prominent role, as they are naturally associated with generalized metrics on exact Courant algebroids. Moreover, the vanishing of Ric^∇ implies that the associated generalized metric is generalized Einstein, see [GF, GFS].

Since the torsion of ∇ is non-vanishing, the Ricci tensor Ric^∇ is not symmetric, and one has (see [GFS, Prop. 3.18])

$$\text{Ric}^\nabla = \text{Ric}_g - \frac{1}{4}H^2 - \frac{1}{2}\delta_g H,$$

where Ric_g denotes the Ricci tensor of ∇^g , δ_g is the formal adjoint of d , and the symmetric 2-tensor H^2 is defined as $H^2(X, Y) := g(\iota_X H, \iota_Y H)$, for every $X, Y \in \Gamma(TM)$.

Therefore, a Bismut connection ∇ with closed torsion form H has zero Ricci tensor if and only if H is g -harmonic and the Ricci tensor of g satisfies the equation $\text{Ric}_g = \frac{1}{4}H^2$. In this context, we shall say that (g, H) is a *Bismut Ricci flat pair* (*BRF pair* for short) if H is closed and the corresponding Bismut connection ∇ is Ricci flat.

A further motivation to consider BRF pairs relies on the fact that they provide fixed points of the *generalized Ricci flow*, a geometric flow evolving a family of Riemannian metrics g_t and 2-forms b_t as follows

$$\begin{cases} \frac{\partial}{\partial t} g_t = -2 \text{Ric}_{g_t} + \frac{1}{2} H_t^2, \\ \frac{\partial}{\partial t} b_t = -\delta_{g_t} H_t, \end{cases}$$

where $H_t = H + db_t$ for a given background closed 3-form H . This flow was introduced in [CFMP, OSW] in the context of renormalization group flows of two-dimensional nonlinear sigma models, and it can be considered as a generalization of Hamilton's Ricci flow to Bismut connections with closed torsion form [Str], and as a flow of generalized metrics on exact Courant algebroids [GF, GFS, Str2]. Moreover, it is also related to some geometric flows in Hermitian Geometry, like the pluriclosed flow and the generalized Kähler Ricci flow, see e.g. [GJS, Str3, ST1, ST2, ST3].

A key example of (homogeneous) manifold with a BRF pair is provided by a semisimple compact Lie group G endowed with the bi-invariant metric g_G induced by $-B$, where B denotes its Cartan-Killing form, together with the *standard* harmonic 3-form $H_G(X, Y, Z) = g_G([X, Y], Z)$, where X, Y, Z are left-invariant vector fields. Indeed, it is well-known that the Bismut connection on G induced by the pair (g_G, H_G) is flat. Viceversa, it has been proved that a compact simply connected Riemannian manifold carrying a flat Bismut connection with closed 3-form is isometric to the product of compact simple Lie groups, see [AF, CS].

In [GFS], the authors asked whether an invariant Bismut connection with zero Ricci tensor on a homogeneous manifold should be flat, generalizing the well-known Alekseevsky-Kimelfeld Theorem in the Riemannian case [AK]. In [PR], we answered this question negatively, proving the existence of invariant non-flat BRF pairs on a series of 5-dimensional homogeneous manifolds $M_{p,q}$ parametrized by a pair of positive integers with $\gcd(p, q) = 1$. These manifolds can be represented as quotient spaces $(\mathrm{SU}(2) \times \mathrm{SU}(2))/\mathrm{T}_{\mathrm{diag}}^1$, for some suitably embedded torus $\mathrm{T}^1 \subset \mathrm{SU}(2) \times \mathrm{SU}(2)$.

The aim of this paper is to generalize the previous result by providing a construction which allows to obtain infinite families of non-flat BRF pairs on compact homogeneous manifolds. These families include the manifold $M_{1,1}$, but not $M_{p,q}$ when $p, q \neq 1$. The new examples, which are constructed starting from compact symmetric spaces of inner type, admit a subcover which can be isometrically and minimally embedded into a compact semisimple Lie group in such a way that the harmonic 3-form coincides with the pull-back of the standard harmonic 3-form on the group.

Our main result can be stated as follows.

Main Theorem. *Let G be a compact, connected semisimple Lie group and let σ be an involutive inner automorphism of G . If $K \subset G$ is a compact subgroup with $(G^\sigma)^\circ \subseteq K \subseteq G^\sigma$, then the homogeneous space $M = (G \times G)/K_{\mathrm{diag}}$, where $K_{\mathrm{diag}} = \{(k, k) \in G \times G \mid k \in K\}$, is endowed with an invariant non-flat BRF pair (\bar{g}, \bar{H}) .*

Moreover, if $K = G^\sigma$, there exists a $G \times G$ -equivariant minimal embedding $\iota : M \hookrightarrow G \times G$ so that $(\bar{g}, \bar{H}) = \iota^(g_G \oplus g_G, H_G \oplus H_G)$.*

We remark that the pair (G, K) is known as a *symmetric pair*, and that the involution σ is inner if and only if G^σ has maximal rank in G (see e.g. [Hel, Ch. IX, Thm 5.6]). We also recall that every compact semisimple Lie group admits at least one involutive inner automorphism. The list of all symmetric pairs $(\mathfrak{g}, \mathfrak{k})$ of compact type with \mathfrak{g} simple and $\mathrm{rank} \mathfrak{g} = \mathrm{rank} \mathfrak{k}$ can be deduced from [Hel, Ch. X, Table V]. For the sake of completeness, we list them in Table 1.

As an immediate consequence, we have the following.

Corollary. *There exist infinitely many compact homogeneous spaces admitting an invariant non-flat BRF pair (g, H) .*

We note that the space $(G \times G)/K_{\mathrm{diag}}$ is diffeomorphic to $G \times (G/K)$ (see details in Section 3), but this diffeomorphism is not isometric when the latter is endowed with the product of g_G and the standard metric on the symmetric space G/K . We also remark that the homogeneous space $M = (G \times G)/G_{\mathrm{diag}}^\sigma$ is a 4-symmetric space defined by means of the order four automorphism of $G \times G$ given by $(g_1, g_2) \mapsto (g_2, \sigma(g_1))$, see e.g. [Jim]. It is a well-known fact that for any symmetric space G/G^σ the Cartan embedding $\phi : G/G^\sigma \rightarrow G$ given by $\phi(aG^\sigma) = \sigma(a)a^{-1}$ is totally geodesic. Moreover, the pull-back of the 3-form H_G vanishes on G/G^σ . Our embedding of the 4-symmetric space M into $G \times G$ is equivalent to the embedding of $G \times (G/G^\sigma)$ into $G \times G$ via $\mathrm{Id} \times \phi$, and it gives a generalization of the Cartan embedding for 2-symmetric spaces, with minimality in place of the total geodesic feature. Finally, we point out that \bar{H} is never zero, as otherwise (M, \bar{g}) would be Ricci flat and thus flat by [AK].

class	\mathfrak{g}	\mathfrak{k}	class	\mathfrak{g}	\mathfrak{k}
<i>A III</i>	$\mathfrak{su}(n)$	$\mathfrak{s}(\mathfrak{u}(p) + \mathfrak{u}(n-p))$	<i>E V</i>	\mathfrak{e}_7	$\mathfrak{su}(8)$
<i>BD I</i>	$\mathfrak{so}(2n), n \geq 4$	$\mathfrak{so}(2p) + \mathfrak{so}(2(n-p))$	<i>E VI</i>	\mathfrak{e}_7	$\mathfrak{so}(12) + \mathfrak{su}(2)$
<i>BD I</i>	$\mathfrak{so}(2n+1), n \geq 2$	$\mathfrak{so}(2p) + \mathfrak{so}(2(n-p) + 1)$	<i>E VII</i>	\mathfrak{e}_7	$\mathfrak{e}_6 + \mathbb{R}$
<i>D III</i>	$\mathfrak{so}(2n), n \geq 3$	$\mathfrak{u}(n)$	<i>E VIII</i>	\mathfrak{e}_8	$\mathfrak{so}(16)$
<i>C I</i>	$\mathfrak{sp}(n), n \geq 3$	$\mathfrak{u}(n)$	<i>E IX</i>	\mathfrak{e}_8	$\mathfrak{e}_7 + \mathfrak{su}(2)$
<i>C II</i>	$\mathfrak{sp}(p+q)$	$\mathfrak{sp}(p) + \mathfrak{sp}(q)$	<i>F I</i>	\mathfrak{f}_4	$\mathfrak{sp}(3) + \mathfrak{su}(2)$
<i>E II</i>	\mathfrak{e}_6	$\mathfrak{su}(6) + \mathfrak{su}(2)$	<i>F II</i>	\mathfrak{f}_4	$\mathfrak{so}(9)$
<i>E III</i>	\mathfrak{e}_6	$\mathfrak{so}(10) + \mathbb{R}$	<i>G</i>	\mathfrak{g}_2	$\mathfrak{su}(2) + \mathfrak{su}(2)$

TABLE 1. Symmetric pairs of compact type $(\mathfrak{g}, \mathfrak{k})$ with \mathfrak{g} simple and $\text{rank } \mathfrak{g} = \text{rank } \mathfrak{k}$

The Main Theorem is proved in Section 2. We first consider the case when $K = G^\sigma$, where $\sigma = \tau_z$ is the conjugation by some element $z \in G$. We construct an action of $G \times G$ on itself preserving the flat BRF pair $(g_G \oplus g_G, H_G \oplus H_G)$, and we identify a particular minimal orbit with M . We then show that the induced metric \bar{g} and 3-form \bar{H} provide an invariant BRF pair on M whose corresponding Bismut connection $\bar{\nabla}$ is not flat. To conclude the proof, it is then sufficient to observe that $(G \times G)/K_{\text{diag}}$ is a finite cover of $(G \times G)/G_{\text{diag}}^\sigma$. In Proposition 2.11, we also prove that the torsion \bar{H} is not parallel with respect to $\bar{\nabla}$.

Some remarks on the geometry of M are discussed in Section 3, where we prove the following.

Proposition 1.1. *The manifold M has finite fundamental group and $b_3(M) = \ell$, where ℓ is the number of simple factors of G . Moreover*

- 1) *if G is simple, then $b_3(M) = 1$ and M is not diffeomorphic to the product of two manifolds carrying a BRF pair with non-trivial torsion;*
- 2) *if G is semisimple not simple, then M is finitely covered by the product of ℓ factors of the form $(G' \times G')/K'_{\text{diag}}$, with G' simple and (G', K') an inner symmetric pair.*

Notation. Throughout the paper, Lie groups will be denoted by capital letters and their Lie algebras will be denoted by the respective gothic letters. When a Lie group G acts on a manifold M , the vector field associated to any $X \in \mathfrak{g}$ will be denoted by \hat{X} .

2. PROOF OF THE MAIN THEOREM

We keep the same notation as in Section 1, and we start considering the Lie group $N := G \times G$ endowed with the biinvariant product metric $g := g_G \oplus g_G$ together with the harmonic 3-form $H := H_G \oplus H_G$. We denote by ∇ the flat Bismut connection corresponding

to (g, H) . The Lie group $L := G \times G$ acts isometrically on (N, g) preserving H as follows

$$(g_1, g_2) \cdot (x_1, x_2) = (g_1 x_1 g_2^{-1}, g_1 x_2 g_2^{-1}).$$

The involution σ of G is inner and therefore there exists $z \in G$ so that σ coincides with the conjugation τ_z . The subgroup G^σ coincides with the centralizer $C := C_G(z) = \{a \in G \mid az = za\}$, and if we consider the L -orbit through the point $p = (e, z)$ in N , then the stabilizer L_p of p is given by

$$L_p = C_{\text{diag}} = \{(a, a) \mid a \in C\},$$

and the orbit is then

$$L \cdot p \cong L/L_p = (G \times G)/C_{\text{diag}} = M.$$

We denote by \bar{g} the metric induced on M as a submanifold of (N, g) and by \bar{H} the pull-back to M of the 3-form H .

In order to prove that the pair (\bar{g}, \bar{H}) on M induces a Bismut connection $\bar{\nabla}$ that is Ricci flat and non-flat, we split the proof into several steps which are dealt with in separate subsections.

2.1. Basic facts on the induced metric \bar{g} . Let $\mathfrak{c} = \{u \in \mathfrak{g} \mid \text{Ad}(z)u = u\}$ denote the Lie algebra of C and let $\mathfrak{q} = \{u \in \mathfrak{g} \mid \text{Ad}(z)u = -u\}$ denote its B -orthogonal complement in \mathfrak{g} . We put $k := \dim \mathfrak{c}$, $q := \dim \mathfrak{q}$, so that $m := \dim M = k + 2q$ and $n := \dim N = 2k + 2q$.

We can consider the following decomposition of the Lie algebra $\mathfrak{l} = \mathfrak{g} \oplus \mathfrak{g}$ of L

$$\mathfrak{l} = \mathfrak{c}_{\text{diag}} \oplus \mathfrak{p} \oplus \mathfrak{m}_1 \oplus \mathfrak{m}_2,$$

where

$$\mathfrak{c}_{\text{diag}} = \{(u, u) \mid u \in \mathfrak{c}\}, \quad \mathfrak{p} = \{(u, -u) \mid u \in \mathfrak{c}\}, \quad \mathfrak{m}_1 = \{(v, 0) \mid v \in \mathfrak{q}\}, \quad \mathfrak{m}_2 = \{(0, v) \mid v \in \mathfrak{q}\},$$

and there is a natural identification $\mathfrak{p} \oplus \mathfrak{m}_1 \oplus \mathfrak{m}_2 \cong T_p M$ by means of

$$V \mapsto \widehat{V}_p = \left. \frac{d}{dt} \right|_{t=0} \exp(tV) \cdot p.$$

For every $V = (v_1, v_2) \in \mathfrak{l}$ and $(x_1, x_2) \in N$, we have

$$\begin{aligned} \widehat{V}_{(x_1, x_2)} &= \left. \frac{d}{dt} \right|_{t=0} (\exp(tv_1) x_1 \exp(-tv_2), \exp(tv_1) x_2 \exp(-tv_2)) \\ (2.1) \quad &= (dR_{x_1}(v_1) - dL_{x_1}(v_2), dR_{x_2}(v_1) - dL_{x_2}(v_2)) \\ &= \left((v_1^R - v_2^L)_{x_1}, (v_1^R - v_2^L)_{x_2} \right) =: \left(\widehat{V}_{x_1}^{(1)}, \widehat{V}_{x_2}^{(2)} \right), \end{aligned}$$

where v^R and v^L denote, respectively, the right-invariant and left-invariant vector field induced by $v \in \mathfrak{g}$ on G , and $\widehat{V}^{(1)}$ and $\widehat{V}^{(2)}$ are the projections of \widehat{V} onto the first and second factor of $TG \times TG$. Moreover, for $V = (v_1, v_2)$, $W = (w_1, w_2)$, $U = (u_1, u_2) \in \mathfrak{l}$, we obtain

$$\begin{aligned} (2.2) \quad g \left(\widehat{V}_{(x_1, x_2)}, \widehat{W}_{(x_1, x_2)} \right) &= g_G \left(\widehat{V}_{x_1}^{(1)}, \widehat{W}_{x_1}^{(1)} \right) + g_G \left(\widehat{V}_{x_2}^{(2)}, \widehat{W}_{x_2}^{(2)} \right) \\ &= -B \left(dL_{x_1}^{-1} \widehat{V}_{x_1}^{(1)}, dL_{x_1}^{-1} \widehat{W}_{x_1}^{(1)} \right) - B \left(dL_{x_2}^{-1} \widehat{V}_{x_2}^{(2)}, dL_{x_2}^{-1} \widehat{W}_{x_2}^{(2)} \right), \end{aligned}$$

and

$$\begin{aligned}
(2.3) \quad H\left(\widehat{V}_{(x_1, x_2)}, \widehat{W}_{(x_1, x_2)}, \widehat{U}_{(x_1, x_2)}\right) &= H_G\left(\widehat{V}_{x_1}^{(1)}, \widehat{W}_{x_1}^{(1)}, \widehat{U}_{x_1}^{(1)}\right) + H_G\left(\widehat{V}_{x_2}^{(2)}, \widehat{W}_{x_2}^{(2)}, \widehat{U}_{x_2}^{(2)}\right) \\
&= -B\left([dL_{x_1}^{-1}\widehat{V}_{x_1}^{(1)}, dL_{x_1}^{-1}\widehat{W}_{x_1}^{(1)}], dL_{x_1}^{-1}\widehat{U}_{x_1}^{(1)}\right) \\
&\quad -B\left([dL_{x_2}^{-1}\widehat{V}_{x_2}^{(2)}, dL_{x_2}^{-1}\widehat{W}_{x_2}^{(2)}], dL_{x_2}^{-1}\widehat{U}_{x_2}^{(2)}\right).
\end{aligned}$$

For every subspace $\mathfrak{v} \subseteq \mathfrak{p} \oplus \mathfrak{m}_1 \oplus \mathfrak{m}_2$, we let $\widehat{\mathfrak{v}}|_p := \{\widehat{V}_p \in T_p M \mid V \in \mathfrak{v}\}$. The following lemma easily follows from (2.1).

Lemma 2.1. *At the point $p = (e, z)$, we have*

$$\widehat{\mathfrak{p}}|_p = \{(2u, dL_z(2u)) \mid u \in \mathfrak{c}\}, \quad \widehat{\mathfrak{m}}_i|_p = \{((-1)^{i+1}v, -dL_z(v)) \mid v \in \mathfrak{q}\}, \quad i = 1, 2.$$

Proof. If $V = (u, -u) \in \mathfrak{p}$, then using $\text{Ad}(z^{-1})|_{\mathfrak{c}} = \text{Id}_{\mathfrak{c}}$, we get

$$\widehat{V}_{(e, z)} = (2u, dR_z(u) - dL_z(-u)) = (2u, dL_z(\text{Ad}(z^{-1})u + u)) = (2u, dL_z(2u)).$$

If $V = (v, 0) \in \mathfrak{m}_1$, then

$$\widehat{V}_{(e, z)} = (v, dR_z(v)) = (v, dL_z \text{Ad}(z^{-1})v) = (v, -dL_z(v)),$$

since $\text{Ad}(z^{-1})|_{\mathfrak{q}} = -\text{Id}_{\mathfrak{q}}$. Finally, if $V = (0, v) \in \mathfrak{m}_2$, then

$$\widehat{V}_{(e, z)} = (-v, -dL_z(v)).$$

□

From (2.2), we see that the Riemannian metric g at the point $p = (e, z)$ is given by

$$g_{(e, z)}\left(\widehat{V}_{(e, z)}, \widehat{W}_{(e, z)}\right) = -B\left(\widehat{V}_e^{(1)}, \widehat{W}_e^{(1)}\right) - B\left(dL_z^{-1}\widehat{V}_z^{(2)}, dL_z^{-1}\widehat{W}_z^{(2)}\right),$$

and we obtain the following.

Lemma 2.2. *The decomposition $T_p M = \widehat{\mathfrak{p}}|_p \oplus \widehat{\mathfrak{m}}_1|_p \oplus \widehat{\mathfrak{m}}_2|_p$ is \bar{g} -orthogonal. Moreover,*

a) *if $V = (u, -u)$, $W = (y, -y) \in \mathfrak{p}$, then $\bar{g}(\widehat{V}, \widehat{W})_{(e, z)} = -8B(u, y)$;*

b) *if $V = (v, 0)$, $W = (w, 0) \in \mathfrak{m}_1$, then $\bar{g}(\widehat{V}, \widehat{W})_{(e, z)} = -2B(v, w)$;*

c) *if $V = (0, v)$, $W = (0, w) \in \mathfrak{m}_2$, then $\bar{g}(\widehat{V}, \widehat{W})_{(e, z)} = -2B(v, w)$.*

Finally, the g -orthogonal complement of $T_p M \subset T_p N$ is given by

$$T_p M^\perp = \{(u, -dL_z(u)) \mid u \in \mathfrak{c}\}.$$

Proof. We first note that $\alpha := \text{Ad}_L((z, z))$ acts isometrically on $\mathfrak{p} \oplus \mathfrak{m}_1 \oplus \mathfrak{m}_2$ with $\alpha|_{\mathfrak{p}} = \text{Id}$, $\alpha|_{\mathfrak{m}_i} = -\text{Id}$, so that $\bar{g}(\widehat{\mathfrak{p}}|_p, \widehat{\mathfrak{m}}_i|_p) = 0$, for $i = 1, 2$. As for a), we have

$$g_{(e, z)}\left(\widehat{V}_{(e, z)}, \widehat{W}_{(e, z)}\right) = -B(2u, 2y) - B(2u, 2y) = -8B(u, y).$$

b) is proved as follows

$$g_{(e, z)}\left(\widehat{V}_{(e, z)}, \widehat{W}_{(e, z)}\right) = -B(v, w) - B(-v, -w) = -2B(v, w),$$

and c) is proved similarly. Finally, if $V = (v, 0) \in \mathfrak{m}_1$ and $W = (0, w) \in \mathfrak{m}_2$, then

$$g_{(e,z)} \left(\widehat{V}_{(e,z)}, \widehat{W}_{(e,z)} \right) = -B(v, -w) - B(-v, -w) = 0.$$

The last claim follows from Lemma 2.1 and $\text{codim}_{\mathbb{N}}\mathbb{M} = \dim \mathfrak{c}$. \square

2.2. The second fundamental form of the immersion and the curvature of \mathbb{M} .

We begin stating some general facts on the induced Bismut connection on submanifolds, giving a formula for its curvature and the relative Ricci tensor.

Let (\mathbb{N}, g) be a Riemannian manifold of dimension $n = m + k$, consider a 3-form $H \in \Omega^3(\mathbb{N})$ and let ∇ be the Bismut connection associated with the pair (g, H) . The torsion tensor T of ∇ is related to H via the identity $H(X, Y, Z) = g(T_X Y, Z)$, for all $X, Y, Z \in \Gamma(T\mathbb{N})$, and thus

$$\nabla_X Y = \nabla_X^g Y + \frac{1}{2} T_X Y.$$

Let \mathbb{M} be an m -dimensional submanifold of \mathbb{N} , denote by $\iota : \mathbb{M} \rightarrow \mathbb{N}$ the corresponding injective immersion and by $\bar{g} = \iota^* g$ the Riemannian metric induced by g . If $\nu\mathbb{M}$ denotes the normal bundle over \mathbb{M} , then for any pair of vector fields $X, Y \in \Gamma(T\mathbb{M})$ arbitrarily extended to \mathbb{N} , we have

$$\nabla_X^g Y = \bar{\nabla}_X^{\bar{g}} Y + h(X, Y),$$

where $h \in \Gamma(S^2(T^*\mathbb{M}) \otimes \nu\mathbb{M})$ denotes the second fundamental form of \mathbb{M} and $\bar{\nabla}^{\bar{g}}$ is the Levi Civita connection of \bar{g} .

The submanifold \mathbb{M} has a natural Bismut connection $\bar{\nabla}$ induced by the pair $(\bar{g}, \bar{H} = \iota^* H)$. It is related to the Bismut connection ∇ on \mathbb{N} as follows

$$\nabla_X Y = \bar{\nabla}_X Y + h(X, Y) + \frac{1}{2} (T_X Y)^\perp,$$

where $(T_X Y)^\perp$ denotes the normal component of $T_X Y$.

We can now determine the relation between the curvature tensor R^∇ of ∇ and the curvature tensor $R^{\bar{\nabla}}$ of $\bar{\nabla}$. At each point p of \mathbb{M} , we consider a g -orthonormal basis (ξ_1, \dots, ξ_k) of the normal space $\nu_p \mathbb{M}$, and a standard computation shows that for $X, Y, Z, U \in T_p \mathbb{M}$

$$\begin{aligned} g(R_{X,Y}^\nabla Z, U) &= g(\nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X,Y]} Z, U) \\ &= \bar{g} \left(R_{X,Y}^{\bar{\nabla}} Z, U \right) - \left(h_i(Y, Z) + \frac{1}{2} g(T_Y Z, \xi_i) \right) \left(h_i(X, U) + \frac{1}{2} g(\xi_i, T_X U) \right) \\ &\quad + \left(h_i(X, Z) + \frac{1}{2} g(T_X Z, \xi_i) \right) \left(h_i(Y, U) + \frac{1}{2} g(\xi_i, T_Y U) \right). \end{aligned}$$

As for the relation between the Ricci tensor Ric^∇ of ∇ and the Ricci tensor $\text{Ric}^{\bar{\nabla}}$ of $\bar{\nabla}$, we consider a \bar{g} -orthonormal basis (e_1, \dots, e_m) of $T_p M$ and we compute

$$\begin{aligned} \text{Ric}^\nabla(Y, Z) &= g\left(R_{e_j, Y}^\nabla Z, e_j\right) \\ &= \text{Ric}^{\bar{\nabla}}(Y, Z) - \left(h_i(Y, Z) + \frac{1}{2}g(T_Y Z, \xi_i)\right) \left(h_i(e_j, e_j) + \frac{1}{2}g(\xi_i, T_{e_j} e_j)\right) \\ &\quad + \left(h_i(e_j, Z) + \frac{1}{2}g(T_{e_j} Z, \xi_i)\right) \left(h_i(Y, e_j) + \frac{1}{2}g(\xi_i, T_Y e_j)\right) \\ &= \text{Ric}^{\bar{\nabla}}(Y, Z) - g\left(h(Y, Z) + \frac{1}{2}T_Y Z, h(e_j, e_j)\right) + g(h(Y, e_j), h(Z, e_j)) \\ &\quad + \frac{1}{2}g(h(Z, e_j), T_Y e_j) + \frac{1}{2}g(h(Y, e_j), T_{e_j} Z) - \frac{1}{4}g(T_Y e_j, \xi_i)g(T_Z e_j, \xi_i). \end{aligned}$$

Using this last expression, and recalling that $H(X, Y, Z) = g(T_X Y, Z)$, we obtain the following.

Proposition 2.3. *If the Bismut connection ∇ on N is flat, then for any $Y, Z \in T_p M$ we have*

$$\begin{aligned} \text{Ric}^{\bar{\nabla}}(Y, Z) &= g(h(Y, Z), \mu) - g(h(Y, e_j), h(Z, e_j)) + \frac{1}{4}H(Y, e_j, \xi_i)H(Z, e_j, \xi_i) \\ &\quad + \frac{1}{2}H(Y, Z, \mu) + \frac{1}{2}h_i(Y, e_j)H(Z, e_j, \xi_i) - \frac{1}{2}h_i(Z, e_j)H(Y, e_j, \xi_i), \end{aligned}$$

where (e_1, \dots, e_m) is a \bar{g} -orthonormal basis of $T_p M$ and $\mu := h(e_j, e_j)$ is the mean curvature vector of M .

We now go back to the proof of the Main Theorem, considering $M = (G \times G)/C_{\text{diag}}$ and $N = G \times G$. In the following, we compute the second fundamental form h of the immersion $\iota : M \hookrightarrow N$.

We begin with some preliminary observations. Some well-known facts are summarized in the next lemma.

Lemma 2.4. *Let $v, w \in \mathfrak{g}$ and denote by v^R and v^L the right-invariant and left-invariant vector fields induced by v on G , respectively. Then*

- 1) $[v^R, w^L] = 0$;
- 2) $[v^R, w^R] = -[v, w]^R$;
- 3) $\nabla_{v^L}^{gG} w^L = \frac{1}{2}[v, w]^L$.

As for the Levi Civita connection ∇^g at $p = (e, z)$, for every pair of vectors $V = (v_1, v_2), W = (w_1, w_2) \in \mathfrak{l}$, we have

$$\begin{aligned} \nabla_V^g \widehat{W}_{(e,z)} &= \left(\nabla_{v_1^R - v_2^L}^{gG} (w_1^R - w_2^L)_e, \nabla_{v_1^R - v_2^L}^{gG} (w_1^R - w_2^L)_z \right) \\ &= \left(\nabla_{v_1^R}^{gG} w_1^R + \nabla_{v_2^L}^{gG} w_2^L - \nabla_{v_1^R}^{gG} w_2^L - \nabla_{v_2^L}^{gG} w_1^R \Big|_e, \nabla_{v_1^R}^{gG} w_1^R + \nabla_{v_2^L}^{gG} w_2^L - \nabla_{v_1^R}^{gG} w_2^L - \nabla_{v_2^L}^{gG} w_1^R \Big|_z \right). \end{aligned}$$

Some useful identities are collected in the next result.

Lemma 2.5. *Let $v, w, u \in \mathfrak{g}$ and let $x \in \mathbf{G}$, then*

- 1) $g_{\mathbf{G}}(\nabla_{v^R}^{g_{\mathbf{G}}} w^L, u^L)_x = -\frac{1}{2}B([w, u], \text{Ad}(x^{-1})v) = g_{\mathbf{G}}(\nabla_{w^L}^{g_{\mathbf{G}}} v^R, u^L)_x$;
- 2) $g_{\mathbf{G}}(\nabla_{v^R}^{g_{\mathbf{G}}} w^R, u^L)_x = \frac{1}{2}B([v, w], \text{Ad}(x)u)$.

Proof.

1) We have

$$\begin{aligned} 2g_{\mathbf{G}}(\nabla_{v^R}^{g_{\mathbf{G}}} w^L, u^L) &= v^R g_{\mathbf{G}}(w^L, u^L) + w^L g_{\mathbf{G}}(u^L, v^R) - u^L g_{\mathbf{G}}(v^R, w^L) \\ &\quad - g_{\mathbf{G}}(w^L, [v^R, u^L]) - g_{\mathbf{G}}(u^L, [w^L, v^R]) + g_{\mathbf{G}}(v^R, [u^L, w^L]) \\ &= w^L g_{\mathbf{G}}(u^L, v^R) - u^L g_{\mathbf{G}}(v^R, w^L) + g_{\mathbf{G}}(v^R, [u^L, w^L]), \end{aligned}$$

where we used that $g_{\mathbf{G}}(w^L, u^L)$ is constant and $[v^R, u^L] = 0 = [w^L, v^R]$. Now, both w^L and u^L are Killing fields for the biinvariant metric $g_{\mathbf{G}}$, so we have

$$\begin{aligned} w^L g_{\mathbf{G}}(u^L, v^R) &= g_{\mathbf{G}}([w^L, u^L], v^R) + g_{\mathbf{G}}(u^L, [w^L, v^R]) = g_{\mathbf{G}}([w^L, u^L], v^R), \\ u^L g_{\mathbf{G}}(v^R, w^L) &= g_{\mathbf{G}}([u^L, v^R], w^L) + g_{\mathbf{G}}(v^R, [u^L, w^L]) = -g_{\mathbf{G}}([w^L, u^L], v^R). \end{aligned}$$

Therefore

$$2g_{\mathbf{G}}(\nabla_{v^R}^{g_{\mathbf{G}}} w^L, u^L)_x = g_{\mathbf{G}}([w^L, u^L], v^R)_x = g_{\mathbf{G}}(dL_x[w, u], dR_x v) = g_{\mathbf{G}}([w, u], \text{Ad}(x^{-1})v)_e.$$

2) With similar computations as in the previous point, we have

$$\begin{aligned} 2g_{\mathbf{G}}(\nabla_{v^R}^{g_{\mathbf{G}}} w^R, u^L) &= v^R g_{\mathbf{G}}(w^R, u^L) + w^R g_{\mathbf{G}}(u^L, v^R) - g_{\mathbf{G}}(u^L, [w^R, v^R]) \\ &= g_{\mathbf{G}}([v^R, w^R], u^L) = -g_{\mathbf{G}}([v, w]^R, u^L), \end{aligned}$$

thus

$$2g_{\mathbf{G}}(\nabla_{v^R}^{g_{\mathbf{G}}} w^R, u^L)_x = -g_{\mathbf{G}}(dR_x[v, w], dL_x u) = -g_{\mathbf{G}}([v, w], \text{Ad}(x)u)_e.$$

□

Let $(\kappa_1, \dots, \kappa_k)$ be a B -orthogonal basis of \mathfrak{c} such that the vectors $\xi_i := (\kappa_i, -dL_z \kappa_i)$, $i = 1, \dots, k$, form a g -orthonormal basis of $\nu_p M$. Then, using 3) of Lemma 2.4, Lemma 2.5, and recalling that $g_{\mathbf{G}}$ is biinvariant and $\text{Ad}(z)\kappa_i = \kappa_i$, we obtain

$$\begin{aligned} h_i(\widehat{V}, \widehat{W})_{(e,z)} &= g\left(\nabla_{\widehat{V}}^g \widehat{W}, \xi_i\right)_{(e,z)} \\ &= g_{\mathbf{G}}\left(\nabla_{v_1^R}^{g_{\mathbf{G}}} w_1^R + \nabla_{v_2^L}^{g_{\mathbf{G}}} w_2^L - \nabla_{v_1^R}^{g_{\mathbf{G}}} w_2^L - \nabla_{v_2^L}^{g_{\mathbf{G}}} w_1^R \Big|_e, \kappa_i\right) \\ &\quad + g_{\mathbf{G}}\left(\nabla_{v_1^R}^{g_{\mathbf{G}}} w_1^R + \nabla_{v_2^L}^{g_{\mathbf{G}}} w_2^L - \nabla_{v_1^R}^{g_{\mathbf{G}}} w_2^L - \nabla_{v_2^L}^{g_{\mathbf{G}}} w_1^R \Big|_z, -dL_z \kappa_i\right) \\ &= -\frac{1}{2}g_{\mathbf{G}}([v_1, w_1] - [v_2, w_2], \kappa_i) - \frac{1}{2}g_{\mathbf{G}}([w_2, \kappa_i], v_1) - \frac{1}{2}g_{\mathbf{G}}([v_2, \kappa_i], w_1) \\ &\quad + \frac{1}{2}g_{\mathbf{G}}([v_1, w_1] - [v_2, w_2], \kappa_i) + \frac{1}{2}g_{\mathbf{G}}([w_2, \kappa_i], \text{Ad}(z^{-1})v_1) + \frac{1}{2}g_{\mathbf{G}}([v_2, \kappa_i], \text{Ad}(z^{-1})w_1) \\ &= -\frac{1}{2}g_{\mathbf{G}}([w_2, \kappa_i], v_1 - \text{Ad}(z^{-1})v_1) - \frac{1}{2}g_{\mathbf{G}}([v_2, \kappa_i], w_1 - \text{Ad}(z^{-1})w_1). \end{aligned}$$

Using this expression, we deduce the following.

Proposition 2.6. *Let $\widehat{V}, \widehat{W} \in T_p M$. Then, $h(\widehat{V}, \widehat{W}) = h_i(\widehat{V}, \widehat{W})\xi_i$ may be non-zero only when $V = (v_1, 0) \in \mathfrak{m}_1$ and $W = (0, w_2) \in \mathfrak{m}_2$. In such a case*

$$h_i(\widehat{V}, \widehat{W})_p = B([v_1, w_2], \kappa_i).$$

In particular, the mean curvature vector μ of M is identically zero, i.e., M is a minimal submanifold of N .

Proof. Let $V = (v_1, v_2)$, $W = (w_1, w_2) \in \mathfrak{p} \oplus \mathfrak{m}_1 \oplus \mathfrak{m}_2$. We discuss each relevant case separately:

- if $V, W \in \mathfrak{p}$, then $v_2 = -v_1 \in \mathfrak{c}$, $w_2 = -w_1 \in \mathfrak{c}$ and $\text{Ad}(z^{-1})|_{\mathfrak{c}} = \text{Id}_{\mathfrak{c}}$, whence

$$h_i(\widehat{V}, \widehat{W})_{(e,z)} = 0;$$

- if $V \in \mathfrak{p}$ and $W \in \mathfrak{m}_1$, then $v_2 = -v_1 \in \mathfrak{c}$, $w_1 \in \mathfrak{q}$ and $w_2 = 0$, whence

$$h_i(\widehat{V}, \widehat{W})_{(e,z)} = g_G([v_1, \kappa_i], w_1)_e = 0,$$

since $[v_1, \kappa_i] \in \mathfrak{c}$. The analogous conclusion holds when $V \in \mathfrak{p}$ and $W \in \mathfrak{m}_2$;

- if $V, W \in \mathfrak{m}_1$, then $v_1, w_1 \in \mathfrak{q}$ and $v_2 = 0 = w_2$, whence

$$h_i(\widehat{V}, \widehat{W})_{(e,z)} = 0.$$

Similarly, one has $h_i(\widehat{V}, \widehat{W})_{(e,z)} = 0$ when $V, W \in \mathfrak{m}_2$;

- if $V \in \mathfrak{m}_1$ and $W \in \mathfrak{m}_2$, then $v_1 \in \mathfrak{q}$, $v_2 = 0$ and $w_1 = 0$, $w_2 \in \mathfrak{q}$, whence

$$h_i(\widehat{V}, \widehat{W})_{(e,z)} = -g_G([w_2, \kappa_i], v_1)_e = B([w_2, \kappa_i], v_1) = B([v_1, w_2], \kappa_i).$$

□

2.3. The induced Bismut connection on M is Ricci flat and non-flat. We begin describing the restriction of the 2-forms $\iota_{\xi_i} H$ to $T_p M \times T_p M$.

Lemma 2.7. *For every $i = 1, \dots, k$, the 2-form $\iota_{\xi_i} H$ satisfies the following:*

- 1) $\iota_{\xi_i} H(\widehat{Y}, \cdot) = 0$, whenever $Y \in \mathfrak{p}$;
- 2) $\iota_{\xi_i} H|_{\widehat{\mathfrak{m}}_j|_p \times \widehat{\mathfrak{m}}_j|_p} = 0$, for $j = 1, 2$;
- 3) for $Y = (y_1, 0) \in \mathfrak{m}_1$ and $Z = (0, z_2) \in \mathfrak{m}_2$, $\iota_{\xi_i} H(\widehat{Y}, \widehat{Z}) = 2B([y_1, z_2], \kappa_i)$.

Proof. We can compute the expression of H at $p = (e, z)$ from equation (2.3). Using this expression together with Lemma 2.1, we have what follows.

- 1) Let $Y = (y, -y) \in \mathfrak{p}$. If $U = (u, -u) \in \mathfrak{p}$, then

$$H(\widehat{Y}, \widehat{U}, \xi_i) = -B([2y, 2u], \kappa_i) - B([2y, 2u], -\kappa_i) = 0,$$

while if $U = (u, 0) \in \mathfrak{m}_1$, then

$$H(\widehat{Y}, \widehat{U}, \xi_i) = -B([2y, u], \kappa_i) - B([2y, -u], -\kappa_i) = 0,$$

since $[y, u] \in \mathfrak{q}$ and $\kappa_i \in \mathfrak{c}$ are B -orthogonal. The same conclusion holds when $U \in \mathfrak{m}_2$.

2) Let $Y = (y_1, 0), Z = (z_1, 0) \in \mathfrak{m}_1$, then

$$H(\widehat{Y}, \widehat{Z}, \xi_i) = -B([y_1, z_1], \kappa_i) - B([-y_1, -z_1], -\kappa_i) = 0.$$

Similarly, if $Y = (0, y_2), Z = (0, z_2) \in \mathfrak{m}_2$, then

$$H(\widehat{Y}, \widehat{Z}, \xi_i) = -B([-y_2, -z_2], \kappa_i) - B([-y_2, -z_2], -\kappa_i) = 0.$$

3) If $Y = (y_1, 0) \in \mathfrak{m}_1$ and $Z = (0, z_2) \in \mathfrak{m}_2$, then

$$H(\widehat{Y}, \widehat{Z}, \xi_i) = -B([y_1, -z_2], \kappa_i) - B([-y_1, -z_2], -\kappa_i) = 2B([y_1, z_2], \kappa_i).$$

□

We now focus on the Ricci tensor of the Bismut connection $\overline{\nabla}$ on $M = (\mathbb{G} \times \mathbb{G})/C_{\text{diag}}$ defined by $(\overline{g}, \overline{H})$. From Proposition 2.3, we know that it has the following expression at $p = (e, z)$, for every $\widehat{Y}, \widehat{Z} \in T_p M$

$$(2.4) \quad \begin{aligned} \text{Ric}^{\overline{\nabla}}(\widehat{Y}, \widehat{Z}) &= -g\left(h(\widehat{Y}, e_j), h(\widehat{Z}, e_j)\right) + \frac{1}{4} H(\widehat{Y}, e_j, \xi_i) H(\widehat{Z}, e_j, \xi_i) \\ &\quad + \frac{1}{2} h_i(\widehat{Y}, e_j) H(\widehat{Z}, e_j, \xi_i) - \frac{1}{2} h_i(\widehat{Z}, e_j) H(\widehat{Y}, e_j, \xi_i), \end{aligned}$$

where (e_1, \dots, e_m) is a \overline{g} -orthonormal basis of $T_p M$.

Proposition 2.8. *The Bismut connection $\overline{\nabla}$ on M defined by the pair $(\overline{g}, \overline{H})$ is Ricci flat.*

Proof. We begin observing that $\text{Ric}^{\overline{\nabla}}(\widehat{Y}, \cdot)$ vanishes whenever $Y \in \mathfrak{p}$. Indeed, it follows from Proposition 2.6 that $h(\widehat{Y}, \cdot) = 0$. Moreover, we have $H(\widehat{Y}, e_j, \xi_i) = 0$ by 1) in Lemma 2.7.

Let us now focus on the symmetric part of $\text{Ric}^{\overline{\nabla}}$

$$\text{Ric}^{\overline{\nabla}, \text{Sym}}(\widehat{Y}, \widehat{Z}) = -g\left(h(\widehat{Y}, e_j), h(\widehat{Z}, e_j)\right) + \frac{1}{4} H(\widehat{Y}, e_j, \xi_i) H(\widehat{Z}, e_j, \xi_i).$$

From Proposition 2.6, we deduce that the summand

$$-g\left(h(\widehat{Y}, e_j), h(\widehat{Z}, e_j)\right) = -h_i(\widehat{Y}, e_j) h_i(\widehat{Z}, e_j)$$

is not zero if and only if either $Y, Z \in \mathfrak{m}_1$ and $e_j \in \widehat{\mathfrak{m}}_2|_p$ or $Y, Z \in \mathfrak{m}_2$ and $e_j \in \widehat{\mathfrak{m}}_1|_p$. We may choose the orthonormal basis (e_1, \dots, e_m) of $T_p M$ as follows: let (E_1, \dots, E_q) be a B -orthogonal basis of \mathfrak{q} such that $B(E_s, E_s) = -\frac{1}{2}$, $s = 1, \dots, q$, then

- e_1, \dots, e_k is a \overline{g} -orthonormal basis of $\widehat{\mathfrak{p}}|_p$;
- e_{k+1}, \dots, e_{k+q} is a \overline{g} -orthonormal basis of $\widehat{\mathfrak{m}}_1|_p$ with $e_{k+s} = \widehat{(E_s, 0)}$, for $s = 1, \dots, q$;
- $e_{k+q+1}, \dots, e_{k+2q}$ is a \overline{g} -orthonormal basis of $\widehat{\mathfrak{m}}_2|_p$ with $e_{k+q+s} = \widehat{(0, E_s)}$, for $s = 1, \dots, q$.

If $Y = (y_1, 0), Z = (z_1, 0) \in \mathfrak{m}_1$, we then have

$$h_i(\widehat{Y}, e_j) h_i(\widehat{Z}, e_j) = \sum_{s=1}^q h_i(\widehat{Y}, e_{k+q+s}) h_i(\widehat{Z}, e_{k+q+s}) = \sum_{s=1}^q B([y_1, E_s], \kappa_i) B([z_1, E_s], \kappa_i).$$

On the other hand, if $Y = (y_1, 0) \in \mathfrak{m}_1$, then we already know that $H(\widehat{Y}, e_j, \xi_i) = 0$ whenever $e_j \in \widehat{\mathfrak{p}}|_p$. Moreover, for $s = 1, \dots, q$, we have

$$H(\widehat{Y}, e_{k+s}, \xi_i) = 0,$$

by 2) in Lemma 2.7, and

$$H(\widehat{Y}, e_{k+q+s}, \xi_i) = 2B([y_1, E_s], \kappa_i),$$

by 3) in Lemma 2.7. Therefore, for all $Y = (y_1, 0), Z = (z_1, 0) \in \mathfrak{m}_1$, we have

$$\begin{aligned} \frac{1}{4} H(\widehat{Y}, e_j, \xi_i) H(\widehat{Z}, e_j, \xi_i) &= \frac{1}{4} \sum_{s=1}^q H(\widehat{Y}, e_{k+q+s}, \xi_i) H(\widehat{Z}, e_{k+q+s}, \xi_i) \\ &= \sum_{s=1}^q B([y_1, E_s], \kappa_i) B([z_1, E_s], \kappa_i) = h_i(\widehat{Y}, e_j) h_i(\widehat{Z}, e_j), \end{aligned}$$

whence it follows that $\text{Ric}^{\overline{\nabla}, \text{Sym}}(\widehat{Y}, \widehat{Z}) = 0$ for all $Y, Z \in \mathfrak{m}_1$. Similarly, if $Y = (0, y_2), Z = (0, z_2) \in \mathfrak{m}_2$, we obtain

$$h_i(\widehat{Y}, e_j) h_i(\widehat{Z}, e_j) = \sum_{s=1}^q B([y_2, E_s], \kappa_i) B([z_2, E_s], \kappa_i) = \frac{1}{4} H(\widehat{Y}, e_j, \xi_i) H(\widehat{Z}, e_j, \xi_i),$$

since $H(\widehat{Y}, e_{k+s}, \xi_i) = 2B([y_2, E_s], \kappa_i)$, for $s = 1, \dots, q$, and $H(\widehat{Y}, e_j, \xi_i) = 0$ otherwise. Thus, $\text{Ric}^{\overline{\nabla}, \text{Sym}}(\widehat{Y}, \widehat{Z}) = 0$ for all $Y, Z \in \mathfrak{m}_2$. We still have to examine the case where $Y \in \mathfrak{m}_1$ and $Z \in \mathfrak{m}_2$. Here, we have

$$\text{Ric}^{\overline{\nabla}, \text{Sym}}(\widehat{Y}, \widehat{Z}) = \frac{1}{4} H(\widehat{Y}, e_j, \xi_i) H(\widehat{Z}, e_j, \xi_i) = 0,$$

since $H(\widehat{Y}, e_j, \xi_i) = 0$ for $j = 1, \dots, k+q$ and $H(\widehat{Z}, e_j, \xi_i) = 0$ for $j = k+q+1, \dots, k+2q$. Therefore, the symmetric part of $\text{Ric}^{\overline{\nabla}}$ vanishes.

We now examine the skew-symmetric part of $\text{Ric}^{\overline{\nabla}}$

$$\text{Ric}^{\overline{\nabla}, \text{Skew}}(\widehat{Y}, \widehat{Z}) = \frac{1}{2} h_i(\widehat{Y}, e_j) H(\widehat{Z}, e_j, \xi_i) - \frac{1}{2} h_i(\widehat{Z}, e_j) H(\widehat{Y}, e_j, \xi_i).$$

The previous discussion shows that the summands might be non-zero only when $Y = (y_1, 0), Z = (z_1, 0) \in \mathfrak{m}_1$ and when $Y = (0, y_2), Z = (0, z_2) \in \mathfrak{m}_2$. In the first case, we have

$$\text{Ric}^{\overline{\nabla}, \text{Skew}}(\widehat{Y}, \widehat{Z}) = B([y_1, E_s], \kappa_i) B([z_1, E_s], \kappa_i) - B([z_1, E_s], \kappa_i) B([y_1, E_s], \kappa_i) = 0.$$

Similarly, $\text{Ric}^{\overline{\nabla}, \text{Skew}}(\widehat{Y}, \widehat{Z}) = 0$ also in the second case. Therefore $\text{Ric}^{\overline{\nabla}, \text{Skew}} = 0$. \square

To conclude the proof of the Main Theorem, we have to show that $\overline{\nabla}$ is not flat. From Section 2.2, we know that the curvature tensor of $\overline{\nabla}$ is given by

$$(2.5) \quad \begin{aligned} R^{\overline{\nabla}}(\widehat{X}, \widehat{Y}, \widehat{Z}, \widehat{U}) &= \left(h_i(\widehat{Y}, \widehat{Z}) + \frac{1}{2} H(\widehat{Y}, \widehat{Z}, \xi_i) \right) \left(h_i(\widehat{X}, \widehat{U}) + \frac{1}{2} H(\widehat{X}, \widehat{U}, \xi_i) \right) \\ &\quad - \left(h_i(\widehat{X}, \widehat{Z}) + \frac{1}{2} H(\widehat{X}, \widehat{Z}, \xi_i) \right) \left(h_i(\widehat{Y}, \widehat{U}) + \frac{1}{2} H(\widehat{Y}, \widehat{U}, \xi_i) \right), \end{aligned}$$

for all $\widehat{X}, \widehat{Y}, \widehat{Z}, \widehat{U} \in T_p M$. With similar computations as in the proof of Proposition 2.8, we can show the following.

Proposition 2.9. *Let $X = (x_1, 0)$, $Y = (y_1, 0) \in \mathfrak{m}_1$ and $Z = (0, z_2)$, $U = (0, u_2) \in \mathfrak{m}_2$, then*

$$R^{\overline{\nabla}}(\widehat{X}, \widehat{Y}, \widehat{Z}, \widehat{U}) = 4B([y_1, z_2], \kappa_i)B([x_1, u_2], \kappa_i) - 4B([x_1, z_2], \kappa_i)B([y_1, u_2], \kappa_i).$$

Proof. Consider $Y = (y_1, 0) \in \mathfrak{m}_1$ and $Z = (0, z_2) \in \mathfrak{m}_2$. Using Proposition 2.6 and 3) in Lemma 2.7, we see that the first factor of the first summand in the RHS of (2.5) is

$$h_i(\widehat{Y}, \widehat{Z}) + \frac{1}{2}H(\widehat{Y}, \widehat{Z}, \xi_i) = B([y_1, z_2], \kappa_i) + \frac{1}{2}2B([y_1, z_2], \kappa_i) = 2B([y_1, z_2], \kappa_i).$$

Analogous computations hold for the second factor as well as for both factors of the second summand in (2.5). Our claim then follows. \square

As an immediate consequence, we obtain that $\overline{\nabla}$ is not flat. Indeed, since $[\mathfrak{q}, \mathfrak{q}] = \mathfrak{c} \neq \{0\}$, we can choose $x, y \in \mathfrak{q}$ so that $[x, y] \in \mathfrak{c}$ is not zero, and we have the following.

Corollary 2.10. *Let $x, y \in \mathfrak{q}$ such that $[x, y] \in \mathfrak{c}$ is not zero. Then, for $X = (x, 0)$, $Y = (y, 0) \in \mathfrak{m}_1$ and $Z = (0, x)$, $U = (0, y) \in \mathfrak{m}_2$, we have*

$$R^{\overline{\nabla}}(\widehat{X}, \widehat{Y}, \widehat{Z}, \widehat{U}) = -4(B([x, y], \kappa_i))^2 = -4\|[x, y]\|^2 \neq 0.$$

Using the previous results, we can also show the next.

Proposition 2.11. *The torsion \overline{H} is not parallel with respect to $\overline{\nabla}$.*

Proof. First, we recall that for every $\widehat{X}, \widehat{Y}, \widehat{Z}, \widehat{U} \in T_p M$, the following identities hold (see e.g. equations (3.20) and (3.21) in [IP]):

$$d\overline{H}(\widehat{X}, \widehat{Y}, \widehat{Z}, \widehat{U}) = \mathfrak{S}_{\widehat{X}, \widehat{Y}, \widehat{Z}} \left[(\overline{\nabla}_{\widehat{X}} \overline{H})(\widehat{Y}, \widehat{Z}, \widehat{U}) + 2\overline{g} \left(\overline{H}(\widehat{X}, \widehat{Y}), \overline{H}(\widehat{Z}, \widehat{U}) \right) \right] - (\overline{\nabla}_{\widehat{U}} \overline{H})(\widehat{X}, \widehat{Y}, \widehat{Z}),$$

and

$$\mathfrak{S}_{\widehat{X}, \widehat{Y}, \widehat{Z}} R^{\overline{\nabla}}(\widehat{X}, \widehat{Y}, \widehat{Z}, \widehat{U}) = d\overline{H}(\widehat{X}, \widehat{Y}, \widehat{Z}, \widehat{U}) + (\overline{\nabla}_{\widehat{U}} \overline{H})(\widehat{X}, \widehat{Y}, \widehat{Z}) - \mathfrak{S}_{\widehat{X}, \widehat{Y}, \widehat{Z}} \overline{g} \left(\overline{H}(\widehat{X}, \widehat{Y}), \overline{H}(\widehat{Z}, \widehat{U}) \right).$$

Now, if the closed 3-form \overline{H} is $\overline{\nabla}$ -parallel, then we must have

$$\mathfrak{S}_{\widehat{X}, \widehat{Y}, \widehat{Z}} R^{\overline{\nabla}}(\widehat{X}, \widehat{Y}, \widehat{Z}, \widehat{U}) = 0,$$

for all $\widehat{X}, \widehat{Y}, \widehat{Z}, \widehat{U} \in T_p M$. As in Corollary 2.10, we choose $X = (x, 0)$, $Y = (y, 0) \in \mathfrak{m}_1$ and $Z = (0, x)$, $U = (0, y) \in \mathfrak{m}_2$ with $[x, y] \neq 0$. Then,

$$R^{\overline{\nabla}}(\widehat{X}, \widehat{Y}, \widehat{Z}, \widehat{U}) = -4\|[x, y]\|^2,$$

while

$$R^{\overline{\nabla}}(\widehat{Y}, \widehat{Z}, \widehat{X}, \widehat{U}) = 0 = R^{\overline{\nabla}}(\widehat{Z}, \widehat{X}, \widehat{Y}, \widehat{U}),$$

as one can easily check using Proposition 2.6 together with Lemma 2.7. Therefore,

$$\mathfrak{S}_{\widehat{X}, \widehat{Y}, \widehat{Z}} R^{\overline{\nabla}}(\widehat{X}, \widehat{Y}, \widehat{Z}, \widehat{U}) = -4\|[x, y]\|^2 \neq 0,$$

a contradiction. \square

3. REMARKS ON THE GEOMETRY OF M

In this section, we give some information on the geometry of $M = (G \times G)/K_{\text{diag}}$, where K is a subgroup of G as in the Main Theorem. In particular, we prove Proposition 1.1.

We first note that the group $G \times G$ acts on $M' := G \times (G/K)$ transitively as follows

$$(g_1, g_2) \cdot (x, yK) = (g_1 x g_2^{-1}, g_2 y K),$$

realizing M' as the orbit through (e, K) , so that $M \cong M'$. However, this diffeomorphism is not isometric if we endow M with the metric \bar{g} and M' with the product of g_G and the standard metric on the symmetric space G/K .

It is well-known that $\pi_1(G)$ is finite and that any quotient space of G by a compact subgroup of maximal rank has finite fundamental group and vanishing odd Betti numbers (see e.g. [Bor]). Therefore, if we represent M as $M' = G \times (G/K)$, then we immediately see that $\pi_1(M)$ is finite. Moreover, it follows by Künneth Theorem that

$$b_3(M) = b_3(G).$$

The fact that $b_3(G) = \ell$ is also well-known (see e.g. [MT]). This shows the first part of Proposition 1.1.

In particular, when G is simple, we have $b_3(M) = 1$, and the claim 1) of Proposition 1.1 follows from the fact that a manifold carrying a BRF pair with non-trivial torsion form has non-trivial third Betti number.

Finally, we discuss the case when G is semisimple not simple, proving 2) of Proposition 1.1. Let \tilde{G} be the universal cover of G with projection π . As K has maximal rank, it is well known that $\tilde{K} := \pi^{-1}(K^o)$ is connected and \tilde{G}/\tilde{K} covers G/K . Moreover, when we split $\tilde{G} = \prod_{i=1}^{\ell} G_i$ into the product of simple factors G_i , then $\tilde{K} = \prod_{i=1}^{\ell} K_i$, where $K_i := \tilde{K} \cap G_i$. Also, (G_i, K_i) are symmetric pairs if (G, K) is (see e.g. [Hel]). This implies that the manifold M is finitely covered by a product $\prod_{i=1}^{\ell} \tilde{M}_i$, where \tilde{M}_i are given by quotient spaces $(G_i \times G_i)/(K_i)_{\text{diag}}$.

Acknowledgements. The authors were supported by GNSAGA of INdAM and by the project PRIN 2017 “Real and Complex Manifolds: Topology, Geometry and Holomorphic Dynamics”. The authors would like to thank Jeffrey Streets and Mario Garcia-Fernández for their comments and for stimulating discussions.

REFERENCES

- [Agr] I. Agricola. The Srní lectures on non-integrable geometries with torsion. *Arch. Math. (Brno)* **42**, 5–84, 2006.
- [AFF] I. Agricola, A. C. Ferreira, T. Friedrich. The classification of naturally reductive homogeneous spaces in dimensions ≤ 6 . *Differential Geom. Appl.* **39**, 59–92, 2015.
- [AF] I. Agricola, T. Friedrich. A note on flat metric connections with antisymmetric torsion. *Differential Geom. Appl.* **28**, 480–487, 2010.
- [AK] D. V. Alekseevsky, B. N. Kimelfeld. Structure of homogeneous Riemannian spaces with zero Ricci curvature. *Funktsional. Anal. i Priložen* **9**, 5–11, 1975.
- [Bis] J.-M. Bismut. A local index theorem for non-Kähler manifolds,. *Math. Ann.* **284**, 681–699, 1989.
- [Bor] A. Borel. Sur la cohomologie des espaces fibre principaux et des espaces homogenes de groupes de Lie compacts. *Annals of Math.* **57**, 115–207, 1953.

- [CFMP] C. G. Callan, D. Friedan, E. J. Martinec, M. J. Perry. Strings in background fields. *Nuclear Phys. B* **262** (4), 593–609, 1985.
- [Car] É. Cartan. Sur les variétés à connexion affine et la théorie de la relativité généralisée (deuxième partie). *Ann. Ec. Norm. Sup.* **42**, 17–88, 1925.
- [CS] É. Cartan, J. A. Schouten. On Riemannian manifolds admitting an absolute parallelism. *Proc. Amsterdam* **29**, 933–946, 1926.
- [CMS] R. Cleyton, A. Moroianu, U. Semmelmann Metric connections with parallel skew-symmetric torsion. *Adv. Math.* **378**, 2021
- [FI] T. Friedrich, S. Ivanov. Parallel spinors and connections with skew-symmetric torsion in string theory. *Asian J. Math.*, **6**, 303–335, 2002.
- [GF] M. Garcia-Fernández. Ricci flow, Killing spinors, and T-duality in generalized geometry. *Adv. Math.* **350**, 1059–1108, 2019.
- [GJS] M. Garcia-Fernández, J. Jordan, J. Streets. Non-Kähler Calabi-Yau geometry and pluriclosed flow. arXiv:/2106.13716, 2021.
- [GFS] M. Garcia-Fernández, J. Streets. Generalized Ricci Flow. *AMS University Lecture Series* **76**, 2021.
- [Gau] P. Gauduchon. Hermitian connections and Dirac operators. *Boll. Un. Mat. Ital. B* **11**, 257–288, 1997.
- [Hel] S. Helgason. Differential Geometry, Lie Groups, and Symmetric Spaces. Academic Press, Inc., 1978
- [IP] S. Ivanov, G. Papadopoulos. Vanishing theorems and string backgrounds. *Classical Quantum Gravity* **18**, 1089–1110, 2001.
- [Jim] J.A. Jiménez. Riemannian 4-symmetric spaces *Trans. A.M.S.* **306**, 715–73, 1988.
- [MT] M. Mimura, H. Toda. Topology of Lie Groups, I and II. *Transl. Math. Monographs, A.M.S.* **91**, 1991.
- [OSW] T. Oliynyk, V. Suneeta, E. Woolgar. A gradient flow for worldsheet nonlinear sigma models. *Nuclear Phys. B* **739**, 441–458, 2006.
- [PR] F. Podestà, A. Raffero. Bismut Ricci flat manifolds with symmetries. arXiv:2202.00417, 2022.
- [Str] J. Streets. Regularity and expanding entropy for connection Ricci flow. *J. Geom. Phys.* **58**, 900–912, 2008.
- [Str2] J. Streets. Generalized geometry, T-duality, and renormalization group flow. *J. Geom. Phys.* **114**, 506–522, 2017.
- [Str3] J. Streets. Pluriclosed flow and the geometrization of complex surfaces. In: Chen J., Lu P., Lu Z., Zhang Z. (eds) Geometric Analysis. Progress in Mathematics, vol. 333. Birkhäuser, Cham. 2020.
- [ST1] J. Streets, G. Tian. A parabolic flow of pluriclosed metrics. *Int. Math. Res. Not.* **16**, 3101–3133, 2010.
- [ST2] J. Streets, G. Tian. Generalized Kähler geometry and the pluriclosed flow. *Nuclear Phys. B* **858**, 366–376, 2012.
- [ST3] J. Streets, G. Tian. Regularity results for pluriclosed flow. *Geom. Topol.* **17**, 2389–2429, 2013.
- [TV] F. Tricerri, L. Vanhecke. Homogeneous structures on Riemannian manifolds. London Math. Soc. Lecture Notes Series, vol. 83, Cambridge Univ. Press, Cambridge, 1983.

DIPARTIMENTO DI MATEMATICA E INFORMATICA “U. DINI”, UNIVERSITÀ DEGLI STUDI DI FIRENZE,
 VIALE MORGAGNI 67/A, 50134 FIRENZE, ITALY,
Email address: `fabio.podesta@unifi.it`

DIPARTIMENTO DI MATEMATICA “G. PEANO”, UNIVERSITÀ DEGLI STUDI DI TORINO, VIA CARLO ALBERTO 10, 10123 TORINO, ITALY
Email address: `alberto.raffero@unito.it`