

Contrast-Enhanced Ultrasound (CEUS) in Non-Traumatic Abdominal Emergencies



Authors

Diletta Cozzi¹ , Simone Agostini¹, Elena Bertelli¹, Michele Galluzzo², Emanuela Papa³, Germano Scevoli³ , Margherita Trinci², Vittorio Miele¹

Affiliations

- 1 Azienda Ospedaliero Universitaria Careggi, Emergency Radiology, Firenze, Italy
- 2 Azienda Ospedaliera San Camillo Forlanini, Emergency Radiology, Roma, Italy
- 3 Ospedale Sandro Pertini, Interventional Radiology, Roma, Italy

Key words

ultrasound; CT, acute, abdomen

received 03.08.2020

accepted after revision 01.05.2021

Bibliography

Ultrasound Int Open 2021; 7: E76–E86

DOI 10.1055/ a-1347-5875

ISSN 2199-7152

© 2021. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

Correspondence

Diletta Cozzi
Emergency Radiology
Azienda Ospedaliero Universitaria Careggi
Largo Brambilla 3
50139 Firenze
Italy
dilettaozzi@gmail.com

ABSTRACT

Conventional ultrasound imaging (US) is the first-line investigation in acute non-traumatic abdominal emergencies, but sometimes it needs further examinations, such as computed tomography (CT), to reach a certain diagnosis. Contrast-enhanced ultrasound (CEUS), through injection of contrast medium, may provide the radiologist with additional information that could not be investigated with baseline US. It could help reach a diagnosis and rapidly determine the proper therapy in an emergency setting. The purpose of this review is to explain and illustrate the various possibilities and limitations of CEUS in acute non-traumatic abdominal diseases, in particular acute inflammation, parenchymal infarcts, and hemorrhages.

Manuscript

In an emergency setting, ultrasound (US) imaging is the first-line approach, and it has a fundamental role in detecting and diagnosing various acute abdominal pathologies [1–3]. The many applications of US imaging in the study of abdominal diseases make US examination a versatile modality. Nevertheless, standard US imaging has several limitations in characterizing anatomic structures and vascularized lesions. Contrast-enhanced ultrasound (CEUS) is a technique that uses specific contrast agents that can improve the characterization of anatomical structures through the visualization of small vessels and microcirculation [4–7].

CEUS contrast agents are microbubbles made of a gas core and a stabilized biological shell. The second generation of sonography contrast media is a solution prepared very quickly and it is imme-

diately administrable. CEUS has the benefit of a high temporal resolution and the operator can detect the contrast transit in the arterial, portal-venous, and late phase. The arterial phase starts 10 sec after injection and lasts 25 – 30 sec. The portal-venous phase starts immediately after the arterial one. Furthermore, these contrast agents have a short half-life, are excreted rapidly from the lungs, and are very well tolerated. In fact, they can be injected into patients with renal insufficiency, hypotension, shock, without further preliminary laboratory tests, and are also indicated in critical patients [4, 5, 8]. CEUS contrast media are safe with a very low incidence of side effects. There are no cardio-, nephro- or hepato-toxicities and life-threatening anaphylactic reactions in abdominal applications have been reported with a rate of 0.001 % in the study of O'Connor et al. [9]. In Europe, CEUS has only been approved for a

limited number of non-traumatic abdominal pathologies in adults, focusing especially on liver disease. Various studies have shown it to be an excellent imaging technique also for non-hepatic applications. A fundamental practical approach is described in the EFSUMB Guidelines [10–12]. CEUS application in children is currently still “off-label” except for a few indications including vesico-ureteral reflux, but it could be a very useful method in the non-ionizing imaging of young patients [6, 13–17].


Today, in the emergency setting, CEUS is used predominantly in the study of blunt or penetrating traumatic lesions, especially for the diagnosis and follow-up of the solid parenchymal organ (so-called “non-operative management”) [8, 18–22]. Often CEUS is used after a baseline US examination, in an attempt to avoid radiation exposure a computed tomography (CT) examination, especially in hemodynamically stable minor trauma. Furthermore, CEUS is a valid method that can help the radiologist make rapid decisions in case of an unstable major trauma that cannot be moved to the radiology emergency department and needs immediate surgical treatment [8, 19, 20, 23, 24].

The experience with CEUS in abdominal non-traumatic pathology is still limited. In fact, the purpose of this article is to review the possibilities and limitations of CEUS in the acute abdominal setting, focusing on various scenarios (inflammation, ischemia, and hemorrhage) and their related radiological images.

Intra-abdominal hemorrhage

Intra-abdominal hemorrhage can be caused by spontaneous bleeding, for example during anticoagulant therapy, spontaneous rup-

ture of a solid mass, a hepatocellular carcinoma, an aortic aneurysm, or a hemorrhage within cystic lesions (► Fig. 1) [5, 25]. It can be secondary also to diagnostic procedures or therapies such as biopsy or femoral artery puncture in interventional maneuvers. Baseline US can detect the collection of bloody in the peritoneal or extraperitoneal space. Performing CEUS exam helps to recognize the real site of bleeding, through the direct visualization of the contrast leakage (similar to in angiography or during CT exam after contrast injection) [26]. On CEUS, the operator sees active contrast extravasation directly from a vessel (in a continuous or pulsatile form), like in the case of a spontaneous rupture of an abdominal vessel. The extravasation is seen as enhancing blood passing through the arterial wall defect and adjacent pooling [27]. CEUS may have difficulties in differentiating active free bleeding from a pseudoaneurysm. In this case, the differential diagnosis must be made through angiography or CT (► Fig. 2) [26, 28–30]. In the case of a ruptured abdominal mass, CEUS examination can detect the contrast bleeding within the tumor or into the adjacent hematoma (► Fig. 3) [31]. Also, in the case of patients with a complicated abdominal cystic mass or a vesical hemorrhage, CEUS can depict the bleeding site and differentiate it from active contrast extravasation or endoluminal clots (for example, in the case of massive hematuria). However, CEUS exam cannot depict small, deep, and multiple hemorrhages, especially those located outside the scanned area. In these cases, CT with contrast media injection is the main radiological examination to be performed, because of its panoramic and direct visualization of the exact bleeding vessel [32]. Moreover, it has to be remembered that clinical evaluation is fundamental. For example, with both B-mode and CEUS it is not possible to differen-



► **Fig. 1** Hemorrhagic corpus luteum. A young girl 14 y.o. with pelvic pain arrives at the emergency department and the US examination shows the presence of an enlarged left ovary compared to the contralateral. The CEUS exam shows the presence of anechoic (internal) cystic formation with an enhancing wall and an interruption of the peripheral profile (white arrow), findings compatible with rupture of the corpus luteum. It is also associated with free fluid in the pelvis.

tiate between a spontaneous or traumatic splenic rupture, because the imaging findings are the same [33, 34].


Parenchymal ischemia and infarction

In the case of parenchymal ischemia and infarction, CEUS can easily demonstrate the non-vascularized area, especially in solid organs like the spleen, liver, and kidney (that are also effectively examined by ultrasonography examination) [33, 34]. Suspected hypoechoic areas due to hypoperfusion already detected on baseline US can be confirmed with CEUS, which effectively shows the presence of microcirculation alterations. In fact, the alterations in perfusion after the administration of the ultrasound contrast medium appear as clearly visible hypoperfused areas due to the high intrinsic lesion-to-parenchyma contrast of CEUS [5]. In these cases, CEUS can be used instead of CT to reach a correct diagnosis and evaluate the extent of the damage. It is a valid method to perform noninvasive follow-up especially if the infarction is treated conservatively. A recent review by Tedesco et al. shows the diagnostic approach of CEUS in infarctions of the kidney, spleen and pancreas. This affirms that CEUS can replace CT in specific situations [10, 35, 36]. Furthermore, the study by Setola et al. describes the important role of CEUS in confirming suspected renal infarction on baseline US (► Fig. 4) [37]. The same CEUS semeiotics can be seen in testicular or ovarian torsion and infarction, torsion of the spermatic cord and epididymis, as described in the study by Badea et al. (► Fig. 5, 6) [12, 38–41]. An advantage of CEUS in the pediatric population is


the lack of radiation exposure. CEUS has also been reported useful in evaluating intestinal ischemia and small bowel infarction [42, 43]. Hata et al. reported a sensitivity of 85–100 % and a specificity of 98–100 % in diagnosing intestinal infarction with CEUS, from various etiologies [44].

Abdominal inflammation


Similar to ischemia and infarction, in inflammation and phlogosis, there is a higher lesion-to-parenchyma contrast ratio in CEUS examination where the hyperemia corresponds to increased parenchymal microcirculation. CEUS in inflammatory disease has a key role in differentiating between an acute phlogosis and a real abscess or phlegmon or inflammation from a gangrenous evolution [10, 45, 46]. This principle is applied to any acute abdominal inflammatory disease from cholecystitis to appendicitis, pancreatitis, pyelonephritis, or diverticulitis. In particular, in acute cholecystitis, CEUS can very easily detect the evident mural enhancement and any possible gangrenous evolution. In the last case, the lack of mural enhancement is a sign of perforation indicating the need for immediate surgical treatment (► Fig. 7) [47, 48]. Similarly, in acute appendicitis, CEUS improves the detection rate of the disease achieving 98 % accuracy and 100 % sensitivity in the diagnosis, as reported by Incesu et al. [49]. This method can be used especially in the pediatric population after baseline US, avoiding a CT examination. CEUS has been shown to have sufficient accuracy to determine complications (such as perforated or phlegmonous




► **Fig. 2** Splenic pseudoaneurysm. Young male patient 18 y.o. arrives at the emergency department with upper-left quadrant pain. Baseline US (arrows in a–b) demonstrates the presence of at least two parenchymal anechoic lesions with peripheral vascular signal at Doppler exam. CEUS (d–e) and CT (c) confirm the presence of two vascular lesions within the splenic parenchyma referable to pseudoaneurysm. The diagnosis of certainty and treatment were then made with angiography and embolization of pseudoaneurysms (f).




► **Fig. 3** Spontaneous HCC rupture. Case of spontaneous bleeding from an unknown liver lesion in a patient without additional comorbidities, which turned out to be hepatocellular carcinoma. CT with contrast medium (**a–b–d–e**) highlights the bleeding lesion in the left lobe, then confirmed by the CEUS examination (arrow in **f**) where there is a subcapsular arterial blush indicative of bleeding in progress, which also extends into the adjacent peritoneal cavity. Angiography confirms the presence of active bleeding (**c**), embolized and no longer evident in angiographic (**h**) and CEUS controls (**g**).




► **Fig. 4** Kidney infarct. Right cortical mesorenal infarction with triangular morphology, poorly visualized at baseline US (**a**) but without enhancement at CEUS (white arrow - **b**) compared to the remaining vascularized parenchyma.




► **Fig. 5** Acute testicular infarct. Segmental ischemia of a testicle visualized as a diffusely hypoechoic area with poor Doppler representation (white arrow in **a–b–c**). The infarcted part of the testicle is not vascularized or enhanced on CEUS (**d**).



► **Fig. 6** Acute ovarian torsion. Young girl 10 y.o. comes to the emergency department complaining of constant pain during defecation. The first US examination (**a–b**) demonstrates the presence of a coarse cystic mass within the right ovary, with slightly thickened walls with poor peripheral vascularization. CEUS (**d**) demonstrates the presence of a cystic lesion with poorly enhanced walls and with solid endoluminal parietal projections. The patient completed the diagnostic procedure with MRI (axial **c**, sagittal **e** and coronal **f** T2-weighted scans) which confirmed the presence of the adnexal lesion (arrow). Note the perfect agreement between the ultrasound images, CEUS, and MRI. The lesion turned out to be a cystic teratoma.




► **Fig. 7** Acute complicated cholecystitis. On the baseline US images (**a** and **c**) there are recognizable signs of cholecystitis with a gallbladder filled with sludge and microcalculi (arrow). CEUS highlights the evident enhancement of the gallbladder wall, which appears very thickened (**c** and **d**), without showing signs of complication such as an abscess or perforation.




► **Fig. 8** Acute diverticulitis. In **a**, the intestinal wall is visualized an axial scan, highlighting a hypoechoic exophytic lesion suggestive of a diverticulum. After administration of US contrast medium **b**, the evident enhancement of the inflamed intestinal wall, in the arterial phase (arrow).

and gangrenous appendicitis) [13, 49]. Another cause of abdominal pain that can mimic appendicitis is epiploic appendagitis. Also, in this case, CEUS may help the diagnosis by highlighting rounded enhancement in the abdominal fatty tissue within a necrotic center,

which is a typical finding in the disease [50]. In the case of acute pancreatitis, CEUS could be used to define the area of higher vascularization within the parenchyma. This technique is not used in routine daily work, first of all, because patients with acute pancre-



► **Fig. 9** Abscess in diverticulitis. Pain in the left iliac fossa in a man without further comorbidity. In the clinical suspicion of an abscess, a CEUS (a–b) examination was performed directly: it shows a nonhomogeneous constantly anechogenic fluid collection with marked peripheral arterial enhancement, typical of an abscess. The finding was confirmed by CT (c–d) which also makes small air bubbles stand out better, in the context of a peri-diverticular abscess.




► **Fig. 10** Abscess in Crohn's disease. Similarly to ► **Fig. 9**, this is a case of a 37-year-old male patient with Crohn's disease with an intestinal abscess tightly attached to the abdominal wall (arrow).


atitis have pain and it is not easy to examine the whole pancreas on US [10, 51]. In this case, CT exam performed in the emergency department is the gold standard and helps to grade the process [52].

In the case of acute gastrointestinal inflammation caused by diverticulitis, appendicitis, or inflammatory bowel disease (IBD – like Crohn disease), CEUS can improve the detection of their complications, such as abscess and phlegmon (► **Fig. 8, 9**) [53–55]. Abscess appears as a non-enhancing rounded lesion with hyperemic borders, while phlegmons are mass-like with a non-homogeneous pat-

tern of enhancement (► **Fig. 10**) [53, 56]. IBD is a lifelong chronic disease and patients are often young. Currently, US and CEUS are similar to CT and magnetic resonance imaging (MRI) in detecting active disease and follow-up of intestinal lesions [57, 58]. Detection and evaluation of vascular flow in the bowel wall is an important factor in the diagnosis and management of IBD. On baseline US, it may be difficult to distinguish the necrotic liquefied center of an abscess, but after contrast injection, the presence of non-enhanced necrotic central nucleus and a peripheral enhanced wall



► **Fig. 11** Renal abscess. CT **a** and US (**b–c**) signs of a renal abscess in the upper right pole in an elderly patient (arrow). The CEUS study (**c**) confirms the presence of an abscess with an anechoic necrotic center and peripheral enhancement, as demonstrated by the CT examination. The basic US scan **b** does not allow complete visualization of the pathology.



► **Fig. 12** Acute pyelonephritis. The basic US image in **a** highlights a blurred hypoechoic area in the right kidney. After the administration of contrast medium, the CEUS examination **b** confirms the presence of a triangular area with reduced enhancement compared to the remaining parenchyma: the finding refers to an area of pyelonephritis (arrows). It is also associated with perirenal free fluid collection. The images in **c** and **d** are images from a follow up CEUS study after twenty days of antibiotic therapy.

occasionally with internal septa with perifocal hyperemia can be easily demonstrated, especially during the arterial phase (► **Fig. 11**).

Similarly, in patients with acute pyelonephritis, the inflammatory areas within the cortical renal spaces are better depicted with CEUS than baseline US. These areas appear as rounded or wedge-shaped hypoechoic foci within the normally enhancing renal parenchyma, with marginal hyperemia (► **Fig. 12**) [62]. The presence of some perfusion can differentiate acute renal infection from infarction. In the latter scenario, the sonographic contrast signal is absent [63, 64].

Limits of CEUS

In an emergency setting, CEUS exam has several limitations, primarily related to the intrinsic limitations of US, i. e., intestinal movement, obesity, presence of free intraperitoneal air, and the inability of the patient to maintain breath-hold. Moreover, it does not have panoramic view, unlike CT which allows the complete visualization of the whole abdomen. CEUS may not diagnose deep or localized bleeding in a poorly studied abdominal parenchyma, such as the lower pole of the spleen or in the case of retroperitoneal collections. In these cases, an abdominal CT scan should be performed urgently. Despite these limitations, CEUS also has several advantages as explained in this review, including its rapid availability in an emergency context, patients who are bedridden, in intensive care units, or who are difficult to transport to the radiology department for a CT scan.

In conclusion, use of CEUS in the acute non-traumatic setting is not yet widespread, but it should be, especially because it is a simple and safe imaging modality in the study of inflammatory, ischemic, and hemorrhagic pathologies of the abdomen. Furthermore, CEUS improves the accuracy of baseline US imaging, while limiting the use of CT and related radiation exposure, especially in younger patients.

Funding

The authors declare that they did not receive any funding for the publication of this article.

Ethical Approval

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Formal consent is not required for this type of study.

This article does not contain any studies with animals by any of the authors.

Conflict of Interest

The authors declare that they have no conflict of interest related to the publication of this article.

References

- [1] Stoker J, Van Randen A, Laméris W et al. Imaging patients with acute abdominal pain. *Radiology* 2009; 253: 31–46
- [2] Choe J, Wortman JR, Micheals A et al. Beyond appendicitis: Ultrasound findings of acute bowel pathology. *Emerg Radiol* 2019; 26: 307–317
- [3] Van Randen A, Laméris W, Van Es HW et al. A comparison of the accuracy of ultrasound and computed tomography in common diagnoses causing acute abdominal pain. *Eur Radiol* 2011; 21: 1535–1545
- [4] Hunt D, Romero J. Contrast-enhanced ultrasound. *Magn Reson Imaging Clin N Am* 2017; 25: 725–736
- [5] Farina R, Catalano O, Stavolo C et al. Emergency radiology. *Radiol Med* 2015; 120: 73–84
- [6] Dietrich CF, Averkiou M, Nielsen MB et al. How to perform contrast-enhanced ultrasound (CEUS). *Ultras Int Open* 2017; 3: E2–E15
- [7] Emmanuel AL, Meijer RI, van Poelgeest E et al. Contrast-enhanced ultrasound for quantification of tissue perfusion in humans. *Microcirculation* 2019, doi:10.1111/micc.12588
- [8] Piccolo CL, Trinci M, Pinto A et al. Role of contrast-enhanced ultrasound (CEUS) in the diagnosis and management of traumatic splenic injuries. *J Ultrasound* 2018; 21: 315–327
- [9] O'Connor JP, Aboagye EO, Adams JE et al. Imaging biomarker roadmap for cancer studies. *Nat Rev Clin Oncol* 2017; 14: 169–186
- [10] Tedesco G, Sarno A, Rizzo G et al. Clinical use of contrast-enhanced ultrasonography beyond the liver: A focus on renal, splenic and pancreatic applications. *Ultrasonography* 2019; 38: 278–288
- [11] Claudon M, Dietrich CF, Choi BI et al. Guidelines and good clinical practice recommendations for contrast enhanced ultrasound (CEUS) in the liver – update 2012; a WFUMB-EFSUMB initiative cooperation with representatives of AFSUMB, AIUM, ASUM, FLAUS and ICUS. *Ultrasound Med Biol* 2013; 39: 187–210
- [12] Sidhu PS, Cantisani V, Dietrich CF et al. The EFSUMB guidelines and recommendations for the clinical practice of contrast-enhanced ultrasound (CEUS) in non-hepatic applications: update 2017 (short version). *Ultraschall Med* 2018; 154–180
- [13] Cogley JR, O'Connor SC, Houshyar R et al. Emergent pediatric US: What every radiologist should know. *Radiographics* 2012; 32: 651–665
- [14] Schreiber-Dietrich DG, Cui XW, Piscaglia F et al. Contrast-enhanced ultrasound in pediatric patients: A real challenge. *Z Gastroenterol* 2014; 52: 1178–1184
- [15] Menichini G, Sessa B, Trinci M et al. Accuracy of Contrast-Enhanced Ultrasound (CEUS) in the identification and characterization of traumatic solid organ lesions in children: A retrospective comparison with baseline-US and CE-MDCT. *Radiol Med* 2015; 120: 989–1001
- [16] Sidhu PS, Cantisani V, Deganello A et al. Role of contrast-enhanced ultrasound (CEUS) in paediatric practice: An EFSUMB position statement. *Ultraschall in Med* 2017; 38: 33–43
- [17] Trinci M, Piccolo CL, Ferrari R et al. Contrast-enhanced ultrasound (CEUS) in pediatric blunt abdominal trauma. *J Ultrasound* 2019; 22: 27–40
- [18] Pinto F, Valentino M, Romanini L et al. The role of CEUS in the assessment of hemodynamically stable patients with blunt abdominal trauma. *Radiol Med* 2015; 120: 3–11
- [19] Miele V, Piccolo CL, Galluzzo M et al. Contrast-enhanced ultrasound (CEUS) in blunt abdominal trauma. *Br J Radiol* 2016; 89: 20150823
- [20] Miele V, Piccolo CL, Sessa B et al. Comparison between MRI and CEUS in the follow-up of patients with blunt abdominal trauma managed conservatively. *Radiol Med* 2016; 121: 27–37

- [21] Tagliati C, Argalia G, Polonara G et al. Contrast-enhanced ultrasound in delayed splenic vascular injury and active extravasation diagnosis. *Radiol Med* 2019; 124: 170–175. doi:10.1007/s11547-018-0961-9
- [22] Tagliati C, Argalia G, Graziani B et al. Contrast-enhanced ultrasound in the evaluation of splenic injury healing time and grade. *Radiol Med* 2019; 124: 163–169. doi:10.1007/s11547-018-0954-8
- [23] Pinto F, Miele V, Scaglione M et al. The use of contrast-enhanced ultrasound in blunt abdominal trauma: advantages and limitations. *Acta Radiol* 2014; 55: 776–784
- [24] Sessa B, Trinci M, Ianniello S et al. Blunt abdominal trauma: role of contrast-enhanced ultrasound (CEUS) in the detection and staging of abdominal traumatic lesions compared to US and CE-MDCT. *Radiol Med* 2015; 120: 180–189
- [25] Mortelet KJ, Cantisani V, Brown DL et al. Spontaneous intraperitoneal hemorrhage: Imaging features. *Radiol Clin North Am* 2003; 41: 1183–1201
- [26] Catalano O, Sandomenico F, Mattace Raso M et al. Real-time, contrast-enhanced sonography: A new tool for detecting active bleeding. *J Trauma* 2005; 59: 933–939
- [27] Martegani A, Aiani L, Borghi C. The use of contrast-enhanced ultrasound in large vessels. *Eur Radiol* 2004; 14: P73–P86
- [28] Faccioli N, Foti G, Casagrande G et al. CEUS versus CT Angiography in the follow-up of abdominal aortic endoprostheses: Diagnostic accuracy and activity-based cost analysis. *Radiol Med* 2018; 123: 904–909
- [29] Corvino A, Sandomenico F, Setola SV et al. Added value of contrast-enhanced ultrasound (CEUS) with Sonovue® in the diagnosis of inferior epigastric artery pseudoaneurysm: report of a case and review of literature. *J Ultrasound* 2019; 22: 485–489. doi:10.1007/s40477-019-00398-x
- [30] Trinci M, Giangregorio C, Calabrese G et al. A rare case of non-traumatic intrasplenic pseudoaneurysms in a patient with acute T-cell lymphoblastic leukemia [published online ahead of print, 2019 Aug 1]. *J Ultrasound* 2019, 10.1007/s40477-019-00401-5. doi: doi:10.1007/s40477-019-00401-5
- [31] Sugihara T, Koda M, Okamoto T et al. Two patterns of contrast-enhanced ultrasonography with Sonazoid in spontaneous rupture of hepatocellular carcinoma: A report of four cases. *J Med Ultrason* 2001; 45: 319–323
- [32] Wells ML, Hansel SL, Bruining DH et al. CT for evaluation of acute gastrointestinal bleeding. *Radiographics* 2018; 38: 1089–1107
- [33] Rosling M, Trenker C, Neessee A et al. Spontaneous and traumatic splenic rupture: Retrospective clinica, B-Mode and CEUS analysis in 62 patients. *Ultrasound Int Open* 2018; 4: E30–E34
- [34] Dietrich CF, Averkiou M, Nielsen MB et al. How to perform Contrast-Enhanced Ultrasound (CEUS). *Ultrasound Int Open* 2018; 4: E2–E15
- [35] Valentino M, Bertolotto M, Derchi L et al. The role of contrast enhanced ultrasound in acute scrotal diseases. *Eur Radiol* 2011; 21: 1831–1840
- [36] Cantisani V, Bertolotto M, Weskott HP et al. Growing indications for CEUS: The kidney, testis, lymph nodes, thyroid, prostate, and small bowel. *Eur J Radiol* 2015; 84: 1675–1684. DOI: 10.1016/j.ejrad.2015.05.008
- [37] Setola SV, Catalano O, Sandomenico F et al. Contrast-enhanced sonography of the kidney. *Abdom Imaging* 2007; 32: 21–28
- [38] Von Herbay A, Schick D, Horger M et al. Low-MI-sonography with the contrast-agent SonoVue in the diagnosis of infarction of the spleen, kidney, liver and pancreas. *Ultraschall in Med* 2006; 27: 445–450
- [39] Badea R, Lucan C, Suci M et al. Contrast enhanced harmonic ultrasonography for the evaluation of acute scrotal pathology. *Pictorial essay. Med Ultrason* 2016; 18: 110–115
- [40] Moschouris H, Stamatou K, Lampropoulou E et al. Imaging of the acute scrotum: is there a place for contrast-enhanced ultrasonography? *Int Braz J Urol* 2009; 35: 692–705
- [41] Trinci M, Ianniello S, Galluzzo M et al. A rare case of accessory spleen torsion in a child diagnosed by ultrasound (US) and contrast-enhanced ultrasound (CEUS). *J Ultrasound* 2019; 22: 99–102
- [42] Giannetti A, Biscontri M, Randisi P et al. Contrast-enhanced sonography in the diagnosis of acute mesenteric ischemia: A case report. *J Clin Ultrasound* 2010; 38: 156–160
- [43] Kanzaki T, Hata J, Imamura H et al. Contrast-enhanced ultrasonography with Sonazoid for the evaluation of bowel ischemia. *J Med. Ultrasonics* 2012; 39: 161–167
- [44] Hata J, Kamada T, Haruma K et al. Evaluation of bowel ischemia with contrast-enhanced US: initial experience. *Radiology* 2005; 236: 712–715
- [45] Calabrese E, Catalano O, Nunziata A et al. Bedside contrast-enhanced sonography (CEUS) of the critically ill patient. *J Ultrasound Med* 2014; 33: 1685–1693
- [46] Adamietz B, Wenkel E, Uder M et al. Contrast enhanced sonography of the gallbladder: A tool in the diagnosis of cholecystitis? *Eur J Radiol* 2007; 61: 262–266
- [47] Sangrini E, Pecorelli A, Pettinari I et al. Contrast-enhanced ultrasonography to diagnose complicated acute cholecystitis. *Intern Emerg Med* 2015, doi:10.1007/s11739-015-1263-z
- [48] Yashima YTT, Nakahara S, Ishida K et al. Contrast-enhanced ultrasonographic image of gangrenous gallbladder. *J Med Ultrason* 2011; 38: 239–241
- [49] Incesu L, Yazicioglu AK, Selcuk MB et al. Contrast-enhanced power Doppler US in the diagnosis of acute appendicitis. *Eur J Radiol* 2004; 50: 201–209
- [50] Gorg C, Egbring J, Bert T. Contrast-enhanced ultrasound of epiploic appendagitis. *Ultraschall in Med* 2009; 30: 163–167
- [51] Golea A, Badea R, Socaciu M et al. Quantitative analysis of tissue perfusion using contrast-enhanced transabdominal ultrasound (CEUS) in the evaluation of the severity of acute pancreatitis. *Med Ultrason* 2010; 12: 198–204
- [52] Addeo G, Beccani D, Cozzi D et al. Groove pancreatitis: A challenging imaging diagnosis. *Gland Surg* 2019; 8: S178–S187
- [53] Ripollès T, Martínez-Pérez MJ, Paredes JM et al. Contrast-enhanced ultrasound in the differentiation between phlegmon and abscess in Crohn's disease and other abdominal conditions. *Eu J Radiol* 2013; 82: e525–e531
- [54] Buckley O, Geoghegan T, O'Riordan DS et al. Computed tomography in the imaging of colonic diverticulitis. *Clin Radiol* 2004; 59: 977–983
- [55] Sessa B, Galluzzo M, Ianniello S et al. Acute perforated diverticulitis: Assessment with multidetector computed tomography. *Seminars Ultrasound CT MRI* 2016; 37: 37–48. doi: 10.1053/j.sult.2015.10.003
- [56] Esteban JM, Aleixandre A, Hurtado M] et al. Contrast-enhanced power Doppler ultrasound in the diagnosis and follow up of inflammatory abdominal masses in Crohn's disease. *Eur J Gastroenterol Hepatol* 2003; 15: 253–259
- [57] Medellin A, Merrill C, Wilson SR. Role of contrast-enhanced ultrasound in evaluation of the bowel. *Abdom Radiol* 2017, doi:10.1007/s00261-017-1399-6
- [58] Cozzi D, Moroni C, Addeo G et al. Radiological patterns of lung involvement in inflammatory bowel disease. *Gastroenterol Res Pract* 2018, doi:10.1155/2018/5697846
- [59] Girlich C, Schacherer D, Lamby P et al. Innovations in contrast enhanced high resolution ultrasound improve sonographic imaging of the intestine. *Clin Hemor Microcirc* 2010; 45: 207–215

- [60] Catalano O, Sandomenico F, Mattace Raso M et al. Low mechanical index contrast-enhanced sonographic findings of pyogenic hepatic abscesses. *AJR* 2004; 182: 447–450
- [61] Kim KW, Choi BI, Park Sh et al. Pyogenic hepatic abscesses: distinctive features for hypovascular hepatic malignancy on contrast-enhanced ultrasound with SH U 508A: early experience. *Ultrasound Med Biol* 2004; 30: 725–733
- [62] Mitterberger M, Pinggera GM, Colleselli D et al. Acute pyelonephritis: comparison of diagnosis with computed tomography and contrast-enhanced ultrasonography. *BJU Int* 2008; 101: 341–344
- [63] Fontanilla T, Minaya J, Cortès C et al. Acute complicate pyelonephritis: contrast enhanced ultrasound. *Abdom Imaging* 2012; 37: 639–346
- [64] Bertelli E, Mercatelli L, Savi E et al. Surgical margin follow-up after nephron-sparing surgery: the possible role of CEUS.. *J Ultrasound* 2019, 10.1007/s40477-019-00413-1. doi: doi:10.1007/s40477-019-00413-1