

BRAIN COMMUNICATIONS

LETTER TO THE EDITOR

Failure of the glymphatic system by increases of jugular resistance as possible link between asthma and dementia

 Pasquale Gallina,^{1,2} Francesco Lolli,^{2,3} Duccio Cianti,¹ Francesco Perri² and Bernardino Porfirio³

¹Department of Neurosciences, Psychology, Drug Research and Child Health, University of Florence, 50134 Florence, Italy

²Careggi University Hospital, 50134 Florence, Italy

³Department of Clinical and Experimental Biomedical Sciences 'Mario Serio', University of Florence, 50134 Florence, Italy

Correspondence to: Pasquale Gallina, MD
Neurosurgery Unit, CTO Hospital
1, Largo Piero Palagi
50139 Florence, Italy
E-mail: pasquale.gallina@unifi.it

Nair *et al.*¹ provided evidence that relationships between diffusion-weighted magnetic resonance indices of neurodegeneration, CSF biomarkers and decline in cognitive function were consistently stronger in patients with asthma relative to controls. However, the mechanism underlying the association between pulmonary dysfunction and cognitive decline has remained unexplained.

To elaborate on this issue, it would be interesting to have information about a possible quantitative difference in the occurrence of perivascular spaces in the brain of asthmatic patients compared with controls; perivascular spaces are an imaging marker of glymphatic system failure. Perivascular spaces are part of the glymphatic system that is a brain-wide fluid transport pathway involved in the clearance of waste solutes (for the anatomy, physiology, pathology, imaging and clinical relevance of the glymphatic system, see Rasmussen *et al.*² and references within). Subarachnoid CSF enters into arteriolar perivascular spaces via a process known as perivascular space pumping that is driven by arterial wall motion. CSF exchanges with interstitial fluid, providing clearance of interstitial solutes. The fluid then leaves the brain mainly via venous perivascular spaces. CSF transport is facilitated by the expression of aquaporin-4 water channels on the perivascular spaces endfeet of astrocytes. If CSF flow is impaired, perivascular space flushing is less effective³ and CSF accumulates and stagnates inside these spaces that become visible at imaging. Derangement of the glymphatic system follows, interstitial waste molecules accumulate and neurodegeneration occurs.

Detection of a higher number of perivascular spaces in the asthmatic patients of the Rosenkranz group^{1,4} compared with controls might support dysfunction of the glymphatic system as responsible for the neurodegeneration in these patients.

We previously hypothesized that a disorder of CSF circulation may involve the upstream failure of extracranial haemodynamics.⁵ Indeed, on the arterial side, impairment of arterial perivascular space flow due to hypertension has been demonstrated,⁶ while on the venous side, impairment of venous perivascular space efflux has been reported in normal pressure hydrocephalus⁷ and presumed as a cause of dementia in patients with severe liver⁸ and pulmonary dysfunction.⁹ In particular, in patients with poor pulmonary function, dementia was hypothesized to be the ultimate phenomenon of a cascade of events involving the reduction of CSF intracranial outflow by increases in jugular resistances, stasis and accumulation of CSF in the interstitium and venous perivascular spaces and glymphatic failure.⁹

Such hypothesis, which would provide an explanation of the findings observed by Rosenkranz's group,^{1,4} might be assessed by demonstrating the association between dementia and chronic pulmonary heart disease in asthma. Moreover, *in vivo*¹⁰ and *post-mortem* studies¹¹ can investigate findings of glymphatic pathology in cognitively deteriorated asthmatic patients.

Finally, analysis of the duration of venous hypertension, and its burden in these patients, might indicate the threshold of onset of the neurodegenerative process and help

Received October 16, 2023. Revised October 16, 2023. Accepted February 09, 2024. Advance access publication February 13, 2024

© The Author(s) 2024. Published by Oxford University Press on behalf of the Guarantors of Brain.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

individuate the timing for possible CSF shunting.^{12,13} This procedure would aim to ameliorate glymphatic functioning by decongesting interstitium and venous perivascular spaces, thus preventing/limiting neurodegeneration and ultimately dementia.

Funding

This correspondence did not receive funding.

Competing interests

The authors report no competing interests.

Data availability

Data sharing is not applicable to this article as no new data were created or analysed.

References

1. Nair AK, Van Hulle CA, Bendlin BB, *et al.* Impact of asthma on the brain: Evidence from diffusion MRI, CSF biomarkers and cognitive decline. *Brain Commun.* 2023;5:fcad180.
2. Rasmussen MK, Mestre H, Nedergaard M. The glymphatic pathway in neurological disorders. *Lancet Neurol.* 2018;17:1016-1024.
3. Blair GW Jr, Thrippleton MJ, Shi Y, *et al.* Intracranial hemodynamic relationships in patients with cerebral small vessel disease. *Neurology.* 2020;94:e2258-e2269.
4. Rosenkranz MA, Dean DC III, Bendlin BB, *et al.* Neuroimaging and biomarker evidence of neurodegeneration in asthma. *J Allergy Clin Immunol.* 2022;149:589-598. e6.
5. Gallina P, Nicoletti C, Scollato A, Lolli F. The “glymphatic-lymphatic system pathology” and a new categorization of neurodegenerative disorders. *Front Neurosci.* 2021;15:669681.
6. Mestre H, Tithof J, Du T, *et al.* Flow of cerebrospinal fluid is driven by arterial pulsations and is reduced in hypertension. *Nat Commun.* 2018;9:4878.
7. Ringstad G, Vatnehol SAS, Eide PK. Glymphatic MRI in idiopathic normal pressure hydrocephalus. *Brain.* 2017;140:2691-2705.
8. Gallina P, Gallo O, Nicoletti C, Romanelli RG. A hydrodynamic hypothesis for the pathogenesis of glymphatic system impairment in hepatic encephalopathy. *J Hepatol.* 2019;71:228-229.
9. Gallina P, Boschi A, Angelini L, Lastrucci G. The possible link between poor pulmonary function and dementia. *Chest.* 2020;158:833-834.
10. Gomolka RS, Hablitz LM, Mestre H, *et al.* Loss of aquaporin-4 results in glymphatic system dysfunction via brain-wide interstitial fluid stagnation. *eLife.* 2023;12:e82232.
11. Boespflug EL, Simon MJ, Emmalyn L, *et al.* Targeted assessment of enlargement of the perivascular space in Alzheimer’s disease and vascular dementia subtypes implicates astroglial involvement specific to Alzheimer’s disease. *J Alzheimers Dis.* 2018;66:1587-1597.
12. Scollato A, Gallina P, Di Lorenzo N. Cerebrospinal fluid diversion in patients with enlarged Virchow–Robin spaces without ventriculomegaly. *Acta Neurol Scand.* 2016;133:75-80.
13. Scollato A, Lolli F, Lastrucci G, *et al.* Case report: A multiple sclerosis patient with imaging features of glymphatic failure benefitted from CSF flow shunting. *Front Neurosci.* 2022;16:863117.