

Article

Profound Transformations of Mediterranean Wetlands Compared to the Past: Changes in the Vegetation of the Fucecchio Marsh (Central Italy)

Lorenzo Lastrucci ¹  and Daniele Viciani ^{2,*} ¹ Botanical Collections, Natural History Museum of the University of Florence, University Museum System, Via G. La Pira 4, 50121 Florence, Italy; lorenzo.lastrucci@unifi.it² Department of Biology, University of Florence, Via G. La Pira 4, 50121 Florence, Italy

* Correspondence: daniele.viciani@unifi.it

Abstract

Although wetlands are key habitats for biodiversity conservation, they are also among the most threatened ecosystems in the world. They are mainly affected by human pressures and threats, even when they are included in protected areas. The Padule di Fucecchio area is one of the largest and most significant inland marshes in Italy. It is also a wetland of international importance, as defined by the Ramsar Convention. However, studies of the plant communities it contains are surprisingly scarce and out of date. To address this issue, a phytosociological survey of aquatic and marshy vegetation was conducted. This analysis provided an unparalleled census of the area's current aquatic and marsh vegetation. Eight different plant community types were reported in the former category and twenty-six in the latter, many of which were previously unknown in this territory. One of these is entirely novel and is described here for the first time. However, a comparison with previous data revealed that significant changes to the vegetation structure have occurred in recent decades. The hydrophyte communities have almost completely disappeared and many of the most sensitive plants in the most sensitive marsh communities have become rarer or disappeared. They have mostly been replaced by more resilient native plants and invasive alien species.

Keywords: aquatic and palustrine habitats; biological invasions; conservation; freshwater ecosystems; phytosociology; syntaxonom; vegetation



Academic Editors: Eusebio Cano Carmona and Kenneth R. Young

Received: 10 August 2025

Revised: 14 October 2025

Accepted: 16 October 2025

Published: 21 October 2025

Citation: Lastrucci, L.; Viciani, D. Profound Transformations of Mediterranean Wetlands Compared to the Past: Changes in the Vegetation of the Fucecchio Marsh (Central Italy). *Land* **2025**, *14*, 2096. <https://doi.org/10.3390/land14102096>

Correction Statement: This article has been republished with a minor change. The change does not affect the scientific content of the article and further details are available within the backmatter of the website version of this article.

Copyright: © 2025 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Wetlands are ecosystems of significant ecological, cultural, and economic value, playing a crucial role both locally and globally. They are vital for conserving biodiversity, as they are home to a wide variety of animal and plant species, some of which are endemic or endangered [1–4]. They also provide multiple ecological services, such as water purification, regulating surface water flow, filtering nutrients, protecting against flooding, acting as carbon sinks, and many others [3,5–9]. However, they are among the world's most threatened habitats, primarily due to human pressures and threats, such as agricultural and urban land transformation, overexploitation, water pollution, flow modification, habitat destruction or degradation, and invasion by alien species [1,3,8,10–12], not to mention that they are also subject to negative global impacts, such as climate change [13,14]. This worrying general trend has increased rapidly in recent years, even among wetlands included in protected areas [15–17]. Consequently, it is not surprising that freshwater ecosystems in

Europe and Italy also include many threatened habitats with an unfavourable conservation status [18–23]. Furthermore, general trends in the Mediterranean basin are exacerbated by the fact that many habitats not considered important for conservation in Central Europe become extremely rare and threatened in Southern European Mediterranean countries. However, these habitats are not included in the European Habitats Directive for the conservation of biodiversity [24] or on any other protection lists [25–29].

The Padule di Fucecchio area is one of the largest and most significant inland marshes in Italy. It is also one of the wetlands of international importance listed in the Ramsar Convention. It is home to several protected natural areas of European and local interest due to its rich natural environment and the unique features of its hydrogeology and landscape. This wetland is a crucial habitat for numerous bird species, including migratory waterfowl, and plays a vital role in maintaining regional biodiversity by providing essential breeding and feeding grounds for birds, as well as hosting a variety of invertebrates, amphibians, and plants [30]. Bird populations are regularly monitored [31], and there are also numerous studies of the area's flora, despite these being rather dated. Several specimens collected in the area from the 19th century to the present day are deposited in Tuscan herbariums [32–34]. Conversely, studies on plant communities are surprisingly scarce and out of date. To address this issue, a phytosociological survey focusing on aquatic and marshy vegetation was conducted in recent years. This work presents the results of these investigations, aiming to describe the current vegetation, compare the data with previous findings, and outline the changes that have occurred in the landscape.

2. Materials and Methods

2.1. Study Area

2.1.1. Topographic Location and Hydrological Features

The Padule di Fucecchio marsh, which covers an area of around 2000 hectares, is located in northern-central Tuscany, spanning the provinces of Florence and Pistoia and extending minimally into the provinces of Lucca and Pisa. Its centre coordinates are 43.800086° N, 10.799955° E (Figure 1).




Figure 1. The location of the study area in Italy (red dot) and its limits (red circle).

It encompasses the natural floodplain formed by streams and ditches originating from the Apennines to the north, the Montalbano Hills to the east, and the Cerbaie Hills to the west. Its only outlet is the Usciana canal, which flows into the Arno River. Located at an average altitude of around 13 m a.s.l., the marsh sits 5 m lower than the Arno floods, which can reach 18 m a.s.l. at the confluence with the Usciana canal. Therefore, the flow

towards the Arno River is hindered by the large quantity of sediments deposited by the major watercourse along its banks since the beginning of the Quaternary [30,32,35]. This territory has experienced numerous geological and climatic events from the Upper Pliocene to the present day. These events have determined changes in its geomorphology and the distribution of plant communities [36,37]. Human exploitation of the area's natural resources (fishing, agriculture, harvesting marsh herbs, etc.) since ancient times has been no less relevant in shaping the landscape [30]. From this perspective, the most profound alterations are those resulting from historical reclamation. This process began in the Renaissance period with the Medici family and was resumed in the second half of the 18th century by the Lorraine family. It continued until quite recent times [38]. Currently, a dense network of canals crosses the marsh area, while the permanent lake area depicted in ancient maps (e.g., Figure 2), has long since disappeared. Only during the rainy seasons, temporary lakes and ponds are present, some of which are used for hunting waterfowl [30]. The management of the water level currently relies primarily on hydraulic manoeuvres regulated by a Land Reclamation Consortium ("Consorzio di Bonifica del Padule di Fucecchio"). The effects of these manoeuvres on natural conservation, in terms of both flooding and drying times and the quality of inflowing water, have often been considered inadequate [39].



Figure 2. On the left, a historical topographic map of the Valdarno di Sotto plain, showing the Padule di Fucecchio marshes. The map was drawn up between 1770 and 1780, based on the original from 1747 [40]. On the right, the current aerial photograph of the same area.

2.1.2. Vegetated Landscape and Presence of Protected Areas

The Padule di Fucecchio is extensively characterised by reed beds covering large surface areas, in mosaic with wet meadows and shrublands (Figure 2). Concentric bands or a mosaic of meadows, poplar groves, and arable land develop around the marsh area [30]. This wetland is home to a protected area belonging to the Natura 2000 network, which was established by the Habitats Directive to promote the conservation of biodiversity across Europe [24]. This area (Natura 2000 code IT5130007) is a Special Area of Conservation and a Special Protection Area. The Padule di Fucecchio also contains other protected areas of provincial and local interest [30]. The Bosco di Chiusi (approximately 320 hectares), which is another biotope of high naturalistic value (Natura 2000 code IT5140010), is the last

remaining lowland forest in Valdinievole. It is located on the eastern side of the Padule di Fucecchio [23].

2.1.3. Climate and Bioclimate, Geological Substrates

With an average annual rainfall of about 1000 mm, approximately three months of moderate drought in summer, and an average annual temperature of 14.8 °C, the climate is usually classified as meso/sub-Mediterranean [41,42]. According to the classification of Thornthwaite and Mather [43], the climate belongs to the “second mesothermic, sub-oceanic” type, with a concentration of thermal efficiency in summer [42]. According to Pesaresi et al. [44], the study area has a temperate oceanic-sub-Mediterranean bioclimate.

In terms of geological substrates, the whole area is formed by alluvial sands, muds, and silts derived from Pliocene-Quaternary deposits [45].

2.2. Data Set and Data Analysis

The data set comprises 155 original georeferenced relevés carried out in plant communities dominated by aquatic and palustrine species. The data were collected in spring and summer of 2022 and 2023 using the classical phytosociological method by Braun-Blanquet [46,47] and its updates [48–51]. After converting the original Braun-Blanquet cover-abundance scale into the Van der Maarel ordinal scale [52], aquatic and palustrine coenoses were separated based on the growth forms of the dominant species [53]. This resulted in 16 relevés of aquatic plant communities and 139 relevés of palustrine ones. The resulting matrices (16 relevés × 23 species for aquatic plant communities and 139 relevés × 131 species for palustrine plant communities) were analysed using cluster analysis in the R environment [54]. This analysis used the Bray–Curtis dissimilarity vegdist function from the vegan package [55] and the Ward method from the hclust function in the stats package [54].

The names of plant species follow the Portal to the Flora of Italy [56]. The full names, including the authors, are provided in the tables. In the text, the names are given without the authors for brevity. We classified the plant communities according to the syntaxonomic nomenclature adopted in Vegetation of Europe by Mucina et al. [57], as well as subsequent updates [58]. However, in certain instances, the interpretation of the Italian Vegetation Prodrôme [59] was preferred, as detailed in the description paragraph. The syntaxonomical nomenclature is in accordance with the 4th edition of the ICPN [60]. All tables relating to vegetation types attributed to syntaxa that have already been described have been published as supplementary material, while the table relating to a new syntaxon that is described here has been published in the main body of the article. A comprehensive syntaxonomic scheme is provided in Appendix A, which includes the complete names of higher-rank syntaxa not cited in the text for brevity. All other syntaxa cited in the text but not included in the syntaxonomic scheme are also cited in full, including the authors and date.

3. Results and Discussion

The dendrograms resulting from the cluster analysis of the aquatic and palustrine plant communities are shown in Figures 3 and 4. These figures illustrate the correspondence between relevés and plant communities.

The classification provided by the cluster analysis was followed entirely for the aquatic communities and almost entirely for the palustrine ones; in fact, the position indicated by the dendrogram was taken into account for 136 out of 139 relevés. This is because, in accordance with the phytosociological procedure, greater importance is given to floristic composition in terms of both dominance and bioindication in order to discriminate between

phytocoenoses. The case of the community dominated by *Butomus umbellatus* can be regarded as an example, since it is clustered together with *Bolboschoenus glaucus* coenoses due to the minor presence of *B. glaucus* in the relevé, but is nevertheless considered autonomous. In a small number of other cases, the cluster analysis can differentiate some relevés based on floristic heterogeneity. However, these should be classified in the same plant community type, as species dominance and ecology are more indicative of this classification. The analysis identified 34 different plant community types, of which 8 were aquatic and 26 were palustrine. These can be classified into eight syntaxonomic classes (see Appendix A). Each community type is described and commented on in the following paragraphs.




Figure 3. Dendrogram resulting from the cluster analysis of aquatic plant communities. The numbers corresponding to the community type names are placed below the corresponding relevé numbers of the heading of Table S2. 1: *Persicarietum amphibiae*; 2: *Callitriche stagnalis* communities; 3: *Potamo pectinati*-*Myriophylletum spicati*; 4: *Potamogetonum denso-nodosi*; 5: *Potamo crispi*-*Ranunculetum trichophylli*; 6: *Callitriche brutia* communities; 7: *Lemnetum minuto-gibbae*; 8: *Callitriche platycarpa* communities.




Figure 4. Dendrogram resulting from the cluster analysis of palustrine plant communities. The numbers corresponding to the community type names are placed below the corresponding relevé numbers (see in the headings of Table 1 and Tables S3–S12 the line “Relevé number in cluster dendrogram”). 1: *Polypogono viridis*-*Paspaleetum distichi*; 2: *Caricetum ripariae*; 3: *Schoenoplectetum lacustris*; 4: *Bolboschoenetum glauci*; 5: *Butometum umbellati*; 6: *Cyperetum longi*; 7: *Frangulo alni*-*Salicetum cinereae*; 8: *Leucojum aestivum* communities; 9: *Juncus bulbosus*-*Gratiola officinalis* communities; 10: *Ranunculo flammulae*-*Juncetum bulbosi*; 11: *Eleocharitetum palustris*; 12: *Oenanthe aquatica*-*Rorippetum amphibiae*; 13: *Veronica peregrina* communities; 14: *Phragmitetum australis*; 15: *Bidens frondosa* communities; 16: *Phalaridetum arundinaceae*; 17: *Caricetum vesicariae*; 18: *Typhetum latifoliae*; 19: *Typhetum angustifoliae*; 20: *Glycerio*-*Sparganietum neglecti*; 21: *Cyperetum micheliano-diformis*; 22: *Echinochloo cruris-galli*-*Sporobolletum schoenoidis*; 23: *Phalarido arundinaceae*-*Bolboschoenetum laticarpi*; 24: *Portulaca oleracea* communities; 25: *Polygono lapathifolii*-*Xanthietum italici*; 26: *Echinochloo*-*Polygonetum lapathifolii*.

3.1. Aquatic Plant Communities

Lemnetum minuto-gibbae Liberman Cruz, Pedrotti et Venanzoni 1988 (Table S2, rel. 1).

In the study area, *Lemna minuta* grows in small monophytic stands and seems to be currently rather rare. This community can be attributed to *Lemnetum minuto-gibbae* association, described for Lake Titicaca in Bolivia [61] and reported in Central Italy for the lakes Trasimeno [62], Porta [63], and Massaciuccoli [64]. *Lemna minuta* is a species with a high invasive potential, due to its plasticity in respect to chemical water features and its high vegetative propagation capacity [65]. According to the Portal to the Flora of Italy [56], this alien species is present in many Italian regions and shows an invasive behaviour. It can often replace native species such as *L. minor*.

Persicarietum amphibiae Pohjala 1933 nom. mut. Prunier et al. 2019 (Table S2, rels. 2–5).

In the Padule di Fucecchio, communities dominated by the aquatic ecophene of *Persicaria amphibia* are currently the only hydrophytic communities with relatively large surface areas. These communities are only present in marsh areas where ponds (clearings in the reed beds used for hunting) have relatively clear water (Figure 5).



Figure 5. A flooded pond in April, partly covered by *Persicarietum amphibiae* vegetation.

P. amphibia-dominated stands are more or less dense, poor in species, and generally colonise ponds, canals, and ditches with still or slowly flowing, shallow, meso-eutrophic to eutrophic waters, subject to regular drying out (in this case, *P. amphibia* develops the terrestrial ecophene) on very clayey substrates [66]. In the past, these coenoses were often attributed to *Polygono-Potamogeton* *natantis* Soó 1964 association, but ecological studies have demonstrated that habitat requirements of communities dominated by *Potamogeton natans* and by *Persicaria amphibia* are different and support to split them into separate associations [67]. This vegetation type was already reported for Tuscany [68], though with different syntaxonomic invalid names (e.g., *Polygonetum natantis* Soó 1927, *nomen nudum*). Other authors consider *P. amphibia*-dominated communities referable to *Potamo natantis-Polygonetum natantis* Knapp et Stoffers 1962 association [69], but according to Prunier et al. [66], the valid priority name is *Polygonetum amphibium* “associaatio” Pohjala 1933, which they propose to update and change in *Persicarietum amphibiae*.

Potamo pectinati-Myriophylletum spicati Rivas Goday 1964 corr. Conesa 1990 (Table S2, rels. 6 and 7).

Small, sporadic, and species-poor stands dominated by *Myriophyllum spicatum* were found along the Pescia di Pescia stream. They can be attributed to the *Potamo pectinati-Myriophylletum spicati* association (in the past the invalid synonym *Myriophylletum spicati* Soó 1927 nom. nud. was often used), which has a broad ecology, but is more common in streams with eutrophic waters [70]. This vegetation type is considered an indicator of poor water quality [71], tolerating turbidity, drying, and human disturbance [69,72]. It has also been found in other Tuscan marshes, lakes, and ponds [64,73].

Potamogetonetenum denso-nodosi O. de Bolòs 1957 (Table S2, rels. 8 and 9).

Small stands dominated by *Potamogeton nodosus* were found in one of the northern peripheral canals of the marsh. Although *Potamogeton nodosus* communities are sometimes considered to be typical of running waters [74], their presence in standing waters is also rather common [75]. These communities, once probably more frequent in the Padule di Fucecchio canals, can be attributed to an impoverished aspect of the *Potamogetonetenum denso-nodosi* O. de Bolòs 1957 association, included in the *Potamogetonion* alliance, which has been reported both in standing and in flowing waters [62].

Callitriche platycarpa community (Table S2, rel. 10).

A small stand dominated by *Callitriche platycarpa* was found in a temporary pool within a mixed woodland area, on the border between the Padule di Fucecchio and the Bosco di Chiusi protected areas. This species is rather rare in Italy, and in Tuscany it is mainly known on the basis of old records [76]. Although a single community has recently been reported for the Tuscan Maremma [75], the phytosociological data available for Italy are very scarce. Felzines [77] reported the association *Veronico beccabungae-Callitrichetum platycarpae* Grube ex Felzines 2016 for France, indicated as *Veronico beccabungae-Callitrichetum platycarpae* Mériaux 1978 by Prunier et al. [66]. It belongs to the *Ranunculion fluitantis* alliance and indicates the reophile character of this community. However, this species can grow in different habitats [78]. As we found this coenosis in standing water, we provisionally attribute it to *Ranunculion aquatilis* alliance.

Callitriche stagnalis communities (Table S2, rels. 11 and 12).

In the same environment where we found the *Callitriche platycarpa* community, two small stands totally dominated by *C. stagnalis* occurred in shaded small ponds, subject to almost total drying up in the summer season and probably disturbed by trampling. *C. stagnalis* is typical of regions with hot summers and tolerates emergence and shading. It lives in shallow, meso-eutrophic, slightly acidic or neutral clear waters [66]. This species forms communities both in still and in running waters, showing a rather broad ecological behaviour. Consequently, from a syntaxonomic point of view, the classification of this vegetation type is various and not univocal. In running waters, the association *Callitrichetum stagnalis* Kaiser 1926 (also indicated in the past with the synonym *Callitrichetum stagnalis* Segal 1964) was sometimes reported in Europe and in Italy [66,79], while for still waters other types of associations, such as *Ranunculo ophioglossifolii-Callitrichetum stagnalis* Brullo, Scelsi et Spampinato 2001 [75,80] have been described. Frequently, however, both because of their floristic poverty and the poor ecological characterisation, communities dominated by this species are generically classified as phytocoenon or coenosis [73,81,82]. In our opinion, even for the monospecific stands surveyed in the study area, it is preferable to follow this type of interpretation, attributing them to the *Ranunculion aquatilis* alliance.

Callitriche brutia communities (Table S2, rels. 13 and 14).

Callitriche brutia is a Mediterranean–Atlantic species that can participate with different vegetation types. It can be found in aquatic communities of *Ranunculion aquatilis* alliance [75,83], or palustrine coenoses attributed to *Oenanthon* (today = *Eleocharito palustris-*

Sagittarion sagittifoliae) alliance [84], or to *Littorelletea* class [75,84], or, more frequently, when it dominates in Mediterranean temporary wet habitats, to *Isoëto-Nanojuncetea* class [85–89]. In the study area, *C. brutia*-dominated communities have been sporadically found in a peripheral position with respect to *Persicarietum amphibiae* communities and in contact with marsh vegetation (Figure 6), in environments that dry out earlier and are subject to greater fluctuations in water levels, so we refer them provisionally to the alliance *Ranunculion aquatilis*.



Figure 6. A small area of a flooded pond in April, mostly covered by *Callitriche brutia* communities.

Potamo crispus-*Ranunculetum trichophylli* Imchenetzky 1926 (Table S2, rels. 15 and 16).

Ranunculus trichophyllus is one of the more common hydrophytes in Tuscany and can be found in several habitat types, both in flowing and in standing waters [26,73]. However, in the study area, *R. trichophyllus*-dominated communities are very rare. Small and paucispecific stands were only found in one of the northern peripheral canals of the marsh. These coenoses can be attributed to the association *Potamo crispus*-*Ranunculetum trichophylli* (in the past, the invalid synonym *Ranunculetum trichophylli* Soó 1927 was used); typical of oligo-mesotrophic to eutrophic waters and tolerating high levels of turbidity, this vegetation type tolerates strong water level variations and can live for a short time even in complete emersion [69].

3.2. Palustrine Plant Communities

Echinochloa crus-galli-*Sporobolus schoenoides* Paradis and Lorenzoni 1994 nom. mut. nov. (Table S3).

The communities dominated by *Sporobolus schoenoides* grow in drying depressions at the edge of, or within, helophytic vegetation, such as *Bolboschoenus laticarpus* coenoses, or in the clearings resulting from the drying up of ponds and lakes (Figure 7).

S. schoenoides participates in various communities of ephemeral environments, gravitating mainly in coenoses of the *Isoëto-Nanojuncetea* class [90], but also in syntaxa indicating halophilous conditions, so much so that the association *Heleochoetum schoenoides* Topa 1939 is often placed by some authors in the class *Crypsietea aculeatae* Vicherek 1973 [91,92]. Paradis [93] analysed different types of community, which he initially differentiated from

Heleochloetum schoenoidis in several subassociations. These were subsequently formalised as new associations [94]. The comparison of these associations is used to evaluate the communities of the Padule di Fucecchio. Despite the presence of *Cyperus michelianus* in the Tuscan relevés, it is not possible to refer them to the association *Crypsio schoenoidis-Cyperetum micheliani* Martínez-Parras, Lorca, Esteban et Mesa 1988, which is always dominated by *C. michelianus* [95]. Furthermore, in the study area, the communities dominated by *C. michelianus* (see Table 1, sub *Cyperetum micheliano-diformis*) do not host *S. schoenoides* among the species and grow in rather different ecological situations, on more humid soils, and in topologically distant sites from those dominated by *S. schoenoides*. The absence of *Heliotropium supinum* from Tuscany, where it was reported based on ancient findings not recently confirmed [56], does not even allow the attribution of our communities to *Heliotropio supini-Heleochloetum schoenoidis* Rivas Goday 1955, recently reported by Tomaselli et al. [88] for Puglia and otherwise synonymized with *Crypsio schoenoidis-Cyperetum micheliani* by Rivas-Martínez et al. [96]. The communities of the Padule di Fucecchio can instead be attributed to the association *Echinochloo cruris-galli-Crypsidetum schoenoidis* (whose name should be changed to *Echinochloo cruris-galli-Sporoboletum schoenoidis*), an association that grows in rather eutrophic and compact silty-clayey soils, in which the floristic composition is enriched with species linked to disturbed humid environments and constituting a sort of link towards the communities of the *Bidentetea* class [93]. As regards the classification at the higher levels of the association, originally placed by the authors in the class *Isoëto-Nanojuncetea* and in the alliance *Heleochloion* Br.-Bl. 1952, it should be pointed out that currently Mucina et al. [57] consider *Heleochloion schoenoidis* Br.-Bl. ex Rivas Goday 1956 a typically halophilous alliance belonging to the class *Crypsietalia aculeatae*. Since we believe that maintaining the original classification at the class level is appropriate, as was performed by Tomaselli et al. [88]; for example, we attribute the association to the *Verbenion supinae* Slavnić 1951 alliance. This alliance traditionally included many coenoses that gravitated around the *Heleochloion* Br.-Bl. 1952 alliance. In fact, some authors considered *Verbenion supinae* and *Heleochloion* to be synonyms [89].




Figure 7. (A) A dried-up pond in August, the bottom of which is home to *Echinochloo cruris-galli-Sporoboletum schoenoidis* communities. (B) A *Sporobolus schoenoides* plant is shown in detail.

Table 1. *Cont.*[illegible]

Ranunculo flammulae-Juncetum bulbosi Oberdorfer 1957 (Table S4, rels. 1 and 2).

In a partially shaded site, in contact with aquatic *Callitriche platycarpa* vegetation, where the water level decreases, small communities dominated by *Juncus bulbosus* and *Ranunculus flammula* were detected. *Juncus bulbosus* has been indicated as a typical or frequent species in a large number of vegetation types [80,84,90,97], but when it is associated with *Ranunculus flammula* these communities can be attributed to *Ranunculo flammulae-Juncetum bulbosi*, an association which includes stands of low-growing perennial amphibious herbs occurring in the littoral zones of lakes, ponds, pools, and ditches along forest roads, periodically flooded by oligotrophic to mesotrophic, rarely dystrophic, water [98]. This vegetation type has been recorded in several western and central European countries [98–101] and can be attributed to *Littorellion uniflorae* alliance.

Juncus bulbosus-Gratiola officinalis communities (Table S4, rels. 3 and 4).

In sites ecologically similar and near to *Ranunculo flammulae-Juncetum bulbosi* communities, small stands dominated by *Juncus bulbosus* and with relevant cover values of *Gratiola officinalis* have been detected. *G. officinalis* typically grows in areas with wet soils for a fairly long period in spring, drying up in summer [75,84]. Some associations characterised by *G. officinalis* have been described (e.g., *Alismo lanceolatae-Gratioletum officinalis* Biondi and Bagella 2005) but our relevés are paucispecific and not dominated by *G. officinalis*. Therefore, this vegetation type could be interpreted as a situation in which the *Ranunculo flammulae-Juncetum bulbosi* association is replaced by these communities, in which *G. officinalis*, being less ecologically demanding than *R. flammula*, is advantaged by conditions of greater drying out and disturbance. These communities can always be attributed to the *Littorellion uniflorae* alliance.

Cyperetum micheliani-diformis ass. nova (Table 1, Holotypus: rel. 4).

In areas that flood during the winter and dry out during the summer, at the edges of reed beds and in clearings free of marsh vegetation, as well as along paths between reed beds, a vegetation type that is generally widespread throughout the study area grows, consisting of a mixture of native and alien species (Figure 8).



Figure 8. An aspect of *Cyperus*-dominated communities (*Cyperetum micheliani-diformis*).

These communities typically have two layers. The lower one is often dominated by *Cyperus michelianus*, with the presence, sometimes with high cover, of *Portulaca oleracea*, as well as the alien plants *Lindernia dubia*, *Amaranthus emarginatus*, and *Eclipta prostrata*. The

taller layer is often dominated by another alien species, *Cyperus difformis*, and is frequently accompanied by *Xanthium orientale* (which sometimes has high cover or is dominant), *Cyperus odoratus*, and *Echinochloa crus-galli*. In terms of their structure, physiognomy, and, to some extent, their floristic composition, these communities have many similarities to the *Cyperetum micheliano-glomerati* association described by Pellizzari [102] for the eastern stretch of the Po River. However, this last vegetation type is characterised by two species of *Cyperus* absent in our relevés (*C. glomeratus* and *C. squarrosus*, the latter of which is yet to be found in Tuscany), as well as by *Amaranthus tuberculatus*. Compared to *Cyperetum micheliano-glomerati*, in the relevés of the study area higher cover values are provided by *Cyperus difformis*, a species that infests rice crops and is characteristic of the *Oryzo-Cyperetum difformis* association W.Koch 1954 [102,103], and partly by *Xanthium orientale*, which sometimes becomes dominant. *X. orientale* participates in different communities linked to the disturbed environments of humid areas, gravitating towards *Chenopodium rubri* (e.g., *Polygono lapathifolii-Xanthietum italici* Pirola and Rossetti 1974) or *Bidention* (e.g., *Xanthio-Polygonetum persicariae* O.Bolós 1957) alliances. In our relevés, *Amaranthus emarginatus* is also present, while it is missing from the relevés by Pellizzari [102]. On average, the communities of the Padule di Fucecchio are much poorer in species, with an average of seven species per relevé compared to over eleven for the Po River. Given the differences highlighted, we propose establishing a new association named *Cyperetum micheliano-difformis*, which in the study area plays an ecological role similar to that of the *Cyperetum micheliano-glomerati* association in the Po River area. The characteristic species of the *Cyperetum micheliano-difformis* association are *Cyperus michelianus*, *C. difformis*, *Xanthium orientale*, and *Amaranthus emarginatus*. Similarly to how Pellizzari [102] classified the association he described, the floristic composition allows us to place *Cyperetum micheliano-difformis* in the *Bidention* alliance. This is consistent with the ecology of the communities, which develop on muddy emergent soils in eutrophic environments subject to significant human disturbance. In our relevés, some stands dominated by *Cyperus odoratus* can also be noted, an aspect which is more closely linked to the Po River communities described by Pellizzari [102], together with some other relevés in which the frequency and/or cover values of alien species is lower and in which *Cyperus michelianus* forms dense carpets with high cover values. This species generally characterises the coenoses of the *Isoëto-Nanojuncetea* class. Such situations may suggest that, as hypothesised for other coenoses in which *C. michelianus* coexists with other alien species [104], the described association may constitute communities that have evolved into coenoses with a strong presence of alien species from an original *Cyperetum michelianus* Horvatic 1931 or similar syntaxa, favoured over time by increased disturbance and water eutrophication.

Bidens frondosa communities (Table S5, rels. 1–4).

This vegetation type is dominated by the alien and invasive American neophyte *Bidens frondosa*, with a constant and often huge presence of another invasive neophyte, *Symphyotrichum squamatus*. These communities are found near reed beds, ponds, and canals, in slightly higher areas, where water retention is relatively less persistent, and the soil tends to dry more quickly. Similar coenoses have been referred by some authors [105] to *Polygonetum hydropiperis* Passarge 1965, the valid name of *Polygono hydropiperis-Bidentetum* (Koch 1926) Lohmeyer in Tüxen 1950, or to *Leersio-Bidentetum tripartitae* (Poli et J. Tüxen 1960) Zaliberova et al. 2000 *bidentetosum frondosae* (Felzines and Loiseau 2005), or to *Cypero eragrostidi-Bidentetum frondosae* [106]. Regardless, in all these cases, the paucispecific communities we detected lack some of the other characteristic or differential species of the above-mentioned associations (i.e., *Polygonum hydropiper*, *Leersia oryzoides*, *Cyperus eragrostis*), and we therefore prefer not to attribute them at the level of association, but only to *Bidention* alliance.

Veronica peregrina communities (Table S5, rels. 5–7).

Veronica peregrina is an American neophyte that in the study area sometimes forms communities in which it is dominant, especially along the wet banks of ditches and canals. This species, already reported for the Fucecchio marshland by Roma-Marzio et al. [107], is considered as a casual alien species in Tuscany, even if this status should be changed to naturalised according to our relevés, showing rather dense and not ephemeral stands in the study area. This species forms communities belonging to several types of syntaxa, from *Nanocyperion*, e.g., *Filaginello uliginosae-Veronicetum peregrinae* Molero Brion. et Romo 1988 [108], to the *Polygono-Poetea* class [109]. On the other hand, according to Julve [110], the species is characteristic of the *Bidention* alliance. Our relevés, paucispecific and formed in addition to *V. peregrina* by a few other species from the neighbouring coenoses (*Phragmites australis*, *Persicaria lapathifolia*, *Phalaris arundinacea*), do not allow an in-depth syntaxonomic analysis on the community's attribution at higher levels. From an ecological point of view, however, the coenoses develop at the limit of the vegetation of tall helophytes, in wet and eutrophic soils, disturbed by human activities (mowing, trampling, etc.) and by water level fluctuations, suggesting a provisional placement in the *Bidention* alliance.

Portulaca oleracea communities (Table S5, rels. 8 and 9).

In the same desiccated areas where *Echinochloo-Sporobolium* or *Cyperetum michelianii-difformis* are present, *Portulaca oleracea* sometimes becomes dominant for small stretches (Figure 9), and gives rise to paucispecific communities that can be considered an impoverished transition aspect between the most hygrophilous coenoses (*Bidention* alliance) and the driest and ruderal ones of the classes *Chenopodietea* Br.-Bl. in Br.-Bl. et al. 1952 or *Stellarietea mediae* Tüxen et al. ex von Rochow 1951, in which *P. oleracea* is often included [111,112]. This vegetation type, difficult to attribute to higher level syntaxa, can be provisionally referred to *Bidention* alliance.



Figure 9. A dried-up pond in August, with *Portulaca oleracea*-dominated communities growing on the bottom.

Polygono lapathifolii-Xanthietum italici Pirola et Rossetti 1974 (Table S6, rels. 1–3).

In the study area, in the late summer–autumn period, on the desiccating and desiccated sandy-silty, eutrophic substrates emerging from the dried water bodies, large surfaces are covered by annual communities dominated by *Xanthium orientale*, an invasive alien species widespread in Italy, accompanied by *Persicaria lapathifolia*, *Echinochloa crus-galli*, *Stachys palustris*, and a few other plants (Figure 10). This pioneer vegetation type is common alongside rivers, ponds, and lakes [68,113,114], and can be attributed to *Polygono lapathifolii*-*Xanthietum italici* association. Previously, *X. italicum*/*X. orientale* subsp. *italicum* was considered a distinct taxon, but is now considered a synonym of *X. orientale*, see Tomasello [115]. Since these communities form later in the growing season than *Cyperetum micheliani-diformis*, depending on when the survey is carried out, it is sometimes possible to detect transitional features, where species from the earliest communities and those in progress coexist (e.g., rels. 21 and 22 in Table 1).




Figure 10. (A) A flooded canal in early May. (B) The same canal in August, now dry, with *Polygono lapathifolii*-*Xanthietum italici* communities growing on the riverbed. (C) A drying pond in early July, with a belt dominated by *Bolboschoenus laticarpus* in the foreground. (D) Another pond in August, with *Polygono lapathifolii*-*Xanthietum italici* communities growing on the bottom.

Echinochloo-Polygonetum lapathifolii Soó et Csurös 1947 (Table S6, rels. 4–8).

In the areas bordering reed beds, canals, and *Bolboschoenus* coenoses, on moist soils that have emerged for longer periods, communities dominated by *Echinochloa crus-galli* and/or *Persicaria lapathifolia* can be found. They can be probably attributed to *Echinochloo-Polygonetum lapathifolii*, an association growing on soils rich in nutrients and with high pH [116]. These communities are rather similar to *Polygono-Xanthietum italici* from an ecological viewpoint, but can be distinguished from a floristic one, and have already been reported for other Tuscan and European wetlands [26,117,118]. Along the course of some large French rivers (Loire, Allier), for moist sandy areas that dry out progressively from mid-spring, Felzines and Loiseau [119] described an association named *Persicario lapathifoliae-Echinochloetum cruris-gallii*, whose relevés seem to be very similar to ours, though some floristic differences can be noted (e.g., the presence in French communities of *Panicum capillare*, *Leersia oryzoides*, *Eragrostis pectinacea*, and others). The problem of

a correct syntaxonomic attribution of Tuscan communities at the association level must be probably studied more in depth with a dedicated work, so they can be provisionally referred to *Chenopodion rubri* alliance.

Phragmitetum australis Savič 1926 nom. corr. (Table S7, rels. 1–10).

Phragmites australis forms dense species-poor stands along submerged and emergent shores of lakes, swamps, pools, ponds, riverbanks, and canals [120,121]. These communities largely characterise the plant landscape of the Padule di Fucecchio, and can be attributed to the *Phragmitetum australis* association, which is quite common in Italy [120,121] and Tuscany [26,75,122]. In several large wetlands of Central Italy, these coenoses seem to be affected by the die-back syndrome and have shown a serious decline [27]. Despite that, no signs of die-back (e.g., the clumping habitus) were observed in the study area. This is probably due to the fact that in the Padule di Fucecchio strong seasonal water fluctuations, naturally and even artificially regulated, reduce the risk of permanent submergence, which seems to be an unfavourable ecological condition for *Phragmitetum* [27].

Typhetum latifoliae Nowiński 1930 (Table S7, rels. 11–13).

The communities dominated by *Typha latifolia* can be attributed to *Typhetum latifoliae*, an association very common in Italy, which grows in several different habitats, such as ponds, shores of lakes, banks of slow-running streams, deltas, swamps, canals, and ditches [120], also tolerating the summer drying [123]. These communities are often very poor in species or even monospecific [124,125], a feature confirmed also in the study area, where this vegetation type is not very frequent and consists mainly of linear formations along peripheral ditches and canals.

Typhetum angustifoliae Pignatti 1953 (Table S7, rel. 14).

The communities dominated by *Typha angustifolia* can be attributed to *Typhetum angustifoliae*, an association typical of several wetland types which grows in mesotrophic to eutrophic waters, often monospecific or species-poor [120]. Though the association is rather common in Italy [120], and in Tuscany [64,122], in the Padule di Fucecchio it was found to be rather rare: we surveyed only one *T. angustifolia*-dominated community, in the widening of a ditch (probably of artificial origin), in a peripheral area of the marsh, between cultivated fields.

Schoenoplectetum lacustris Chouard 1924 nom. mut. nov. Lastrucci et al. 2023 (Table S7, rels. 15–19).

The communities dominated by *Schoenoplectus lacustris* can be attributed to the *Schoenoplectetum lacustris* association, which has a strong pioneer feature, and is reported for several habitat types, such as shores of mesotrophic to eutrophic lakes, ponds, or canals; it usually grows in deeper and longer-lasting water than other types of reed vegetation [22,120,126,127], and this is also confirmed in the study area.

Phalarido arundinaceae-Bolboschoenetum laticarpi Passarge 1999 corr. Krumbiegel 2006 (Table S7, rels. 20–29).

Bolboschoenus laticarpus, contrarily to *B. glaucus*, is also present in central Europe and lives only in freshwater habitats (Hroudová et al., 2009). In the study area, the communities in which it dominates are widespread, probably as much as *Bolboschoenetum glauci* or more. These dense and paucispecific coenoses can be probably attributed to *Phalarido arundinaceae-Bolboschoenetum laticarpi*, an association floristically rather heterogeneous living in floodplains of large rivers [128]. *Phalaris arundinacea* is not present in our relevé but is widespread in the area, while another differential species, *Xanthium orientale*, is frequent. To our knowledge, this vegetation type has not been previously reported for Italy, but it is likely not rare according to the updated distribution of the species [129].

Cyperetum longi Micevski 1957 (Table S8, rel. 1).

In the peripheral parts of the marsh, along the secondary ditches between meadows and arable land, small stands dominated by *Cyperus longus* are sporadically present. These communities grow in shallow water environments, which often dry out rather early, and can be attributed to *Cyperetum longi*, an association distributed in southern Europe and the Mediterranean [68,130,131], referable to *Magnocaricion elatae* alliance.

Caricetum ripariae Máthé et Kovács 1959 (Table S8, rels. 2–9).

This association lives in meso- and eutrophic wetlands that have emerged since the end of spring [120,121]. It was often found to be frequent in rather disturbed habitats [26,64]. In the study area, the association is quite widespread along the edges of canals, around reed beds, and in wet depressions. These communities are characterised by a large dominance of *Carex riparia* and by the presence of other species typical of upper syntaxonomic units, such as *Phragmites australis*, *Bolboschoenus glaucus*, and *B. laticarpus*, *Convolvulus sepium*, *Lythrum salicaria*. They can be attributed to *Magnocaricion gracilis* alliance.

Caricetum vesicariae Chouard 1924 (Table S8, rels. 10–12).

Along the lower part of the ditch banks, in humid depressions and meadows, some communities dominated by *Carex vesicaria* were sporadically detected. They are referable to *Caricetum vesicariae*, a typical association of mesotrophic to eutrophic habitats, permanently flooded or saturated with water for most of the year [120,132]. *Caricetum vesicariae* was already known for lowland and montane wetlands in Tuscany [122,133] and can be attributed to *Magnocaricion gracilis* alliance.

Phalaridetum arundinaceae Libbert 1931 (Table S8, rels. 13–19).

Phalaris arundinacea-dominated communities occur in complexes of marsh vegetation in lowland river floodplains, in shallow depressions, in the littoral zones of water bodies and ditches with still or flowing water, generally eutrophic and base-rich, and in other similar environments [68,82,123]. In the study area, these communities have been detected along canals and ditches, where soils remain more or less constantly wet, and are not directly affected or disturbed by currents. For these reasons, they can be attributed to *Phalaridetum arundinaceae*, and not to *Rorippo-Phalaridetum arundinaceae* Kopecký 1961, an association typical of less humid and more disturbed sites, with longer exposure to conditions of complete emersion [69,123]. In any case, even if it has not been detected by us, the presence of *Rorippo-Phalaridetum arundinaceae* in the area is possible.

Leucojum aestivum communities (Table S8, rels. 20 and 21).

Leucojum aestivum can participate with different vegetation types, and can be found both in the understory of floodplain hygrophilous woods, dominated, for example, by *Fraxinus angustifolia* subsp. *oxycarpa*, *Salix alba*, etc. [23,134,135], and in herbaceous hygrophilous communities of *Magnocaricetalia* [117,136]. These communities, surveyed mostly for their rarity and their conservation interest, were present at the edge of the Padule, in a clearing of a partially shaded forest site, of which they can probably be considered an herbaceous *sinusia*. They can be provisionally attributed to *Magnocaricion gracilis* alliance.

Glycerio-Sparganietum neglecti Koch 1926 (Table S9, rels. 1–3).

Some communities dominated by *Sparganium erectum* L. sensu lato were found in the peripheral canals of the Padule di Fucecchio. According to Landucci et al. [120,121], they can be attributed to *Glycerio-Sparganietum neglecti*, which should be considered as a “macro-association” including other associations used in the past, such as *Sparganietum erecti* Roll 1938 and *Sparganietum microcarpi* (Weber 1976) Passarge 1978, due to the frequent uncertain taxonomic identification of the species in this group [137]. These communities usually grow in environments considered intermediate between *Phragmition* and *Glycerio-Sparganion* ecological conditions, so their attribution to the alliance level is not always univocal [120,121,128]. In the study area these canals generally have strong fluctuations of water levels, justifying the attribution of *Glycerio-Sparganietum* to the alliance *Glycerio-Sparganion*.

Eleochariteum palustris Savič 1926 (Table S10, rels. 1–3).

These communities are dominated by *Eleocharis palustris* and can be attributed to the *Eleochariteum palustris* association. They show a typical pioneer behaviour, often developing in the wet soils emerging during the dry season [82]. In our relevés, we only found *E. palustris* subsp. *palustris*, but in the area, the presence of other subspecies is possible [138]. This association, typically poor in species, is rather common in Italy [75,120,122,139], but in the Padule di Fucecchio it is not so widespread. It was sometimes found along some canals or in the muddy areas surrounding the ponds (clearings for hunting) open in the reed beds.

Oenanthe aquatica-*Rorippa amphibia* Lohmeyer 1950 (Table S10, rels. 4–6).

Communities dominated by *Rorippa amphibia* generally occur in lowlands on lacustrine and fluvial sediment accumulations, and grow in marshes, ponds, pools, canals, and lentic sections of rivers, where in summer the water table often drops far enough to expose the bottom [123]. This vegetation type can be attributed to *Oenanthe aquatica*-*Rorippa amphibia*, an association sporadically present in the Padule di Fucecchio (Figure 11) and in other Tuscan wetlands [75].



Figure 11. An aspect of *Oenanthe aquatica*-*Rorippa amphibia* communities.

Bolboschoenetum glauci Grechushkina, Sorokin et Golub 2011 (Table S10, rels. 7–13).

After taxonomic and distributional revisions of the genus *Bolboschoenus* by Hroudová et al. [140,141] for Europe and by Di Natale et al. [129] for Italy, the knowledge on the distribution of this complex of species (*B. maritimus* sensu lato) and on their dominated vegetation types has been significantly updated [128,142]. *B. glaucus* does not occur in inland central Europe, can grow both in freshwater and saline habitats, and is mainly distributed in southern Europe, showing a thermophilic character [128,129]. In the study area, it forms dense, paucispecific, and rather widespread communities in which it dominates, which can be attributed to the *Bolboschoenetum glauci* association already indicated in other Tuscan wetlands [75,127].

Butometum umbellati Philippi 1973 (Table S10, rel. 14).

In the study area, *Butomus umbellatus* is sporadically present in some vegetation types (e.g., in *Schoenoplectetum lacustris* coenoses), but rarely it forms communities in which it dominates. We could perform only one relevé of this type that can be attributed to

Butometum umbellati association. In Italy and in Tuscany this vegetation type is generally sporadic [68,132,143], and in the Padule di Fucecchio it is probably much less widespread than in the past.

Polypogono viridis-Paspaleto distichi Br.-Bl. in Br.-Bl., Gajewski, Wraber et Walas 1936 nom. invers. et mut. nov. (Table S11, rels. 1–10).

These communities are characterised by the dominance of the invasive alien species *Paspalum distichum* [144] and in the past have generally been attributed to the *Paspalo distichi-Polypogonetum viridis* association. This vegetation type has a mostly Mediterranean distribution and often results in being very poor in species [64,81,113]. Our relevés also confirm this feature, showing a strong dominance of *P. distichum*. In the study area, the species is found in the disturbed banks of the canals and around the ponds (clearings for hunting) in the reed beds. It forms dense mono- or paucispecific carpets in contact with hygrophilous herbaceous or helophytic communities. In only a few cases that can be interpreted as transition aspects (e.g., rel. 10 in Table S11), some stands can be co-dominated by other species, such as *Cyperus* sp.pl. From a nomenclatural viewpoint, the association name requires reversal, since *P. distichum* always dominates in both the original description [145] and the aforementioned works (including our relevés). We therefore propose changing it to *Polypogono viridis-Paspaleto distichi* (*Polypogon viridis* is the accepted name by several floras of *Agrostis viridis* and *Agrostis semiverticillata*). Furthermore, since the *Paspalo distichi-Agrostion semiverticillatae* alliance is based on the *Paspalo distichi-Polypogonetum viridis* association [146], the alliance name could also be changed to *Polypogono viridis-Paspalion distichi* Br.-Bl. in Br.-Bl. et al. 1952 nom. invers. et mut. nov. Regarding attribution to higher-level taxa, Mucina et al. [57] refer the *Paspalo distichi-Agrostion semiverticillatae* alliance to the *Paspalo-Heleochoetalia* Br.-Bl. ex Rivas Goday 1956 order, of the *Bidentetea* class. This class includes summer annual pioneer lacustrine bank vegetation. However, both *Paspalum distichum* and *Polypogon viridis* are perennial species. Therefore, we prefer to attribute the *Polypogono viridis-Paspaleto distichi* association to the *Holoschoenetalia* order of the *Molinio-Arrhenatheretea* class, in accordance with the Italian Prodrome [59].

Frangulo alni-Salicetum cinereae Graebner et Hueck 1931 (Table S12, rels. 1–4).

Salix cinerea-dominated communities are typical of marshes and grow in conditions of prolonged submersion. They are an infrequent example of the palustrine edapho-hygrophilous series in the temperate bioclimatic region [82]. These communities appear as species-poor, dense shrublands, generally in contact with plant communities of *Phragmito-Magnocaricetea* class, or directly bordering water bodies with or without aquatic vegetation. In Padule di Fucecchio, they are uncommon and can be considered a relic because their ecological niche has been occupied by *Amorpha fruticosa*, an alien shrub that is widespread in the marsh and has been favoured directly and indirectly by human disturbance. *A. fruticosa* is also present in our relevés and sometimes dominates. This vegetation type and the problems posed by *A. fruticosa* were already known for other Tuscan wetlands, both near and far from the study area [26,133]. As to the syntaxonomic attribution, in the past these communities have been attributed either to *Salicetum cinereae* Zólyomi 1931 or to *Frangulo alni-Salicetum cinereae* associations which, according to Douda [147], are synonyms of *Salicetum pentandro-auritae* Passarge 1957. However, the floristic description of *Salicetum pentandro-auritae* appears to differ significantly from that of *S. cinerea* communities described under previous names. Furthermore, *Frangulo alni-Salicetum cinereae* has also been used in recent times [135], prompting us to provisionally maintain this name.

3.3. Past and Present Vegetation

As mentioned in the introduction, studies in the literature on the vegetation of the Padule di Fucecchio are particularly scarce. Furthermore, the level of detail with which this

work analyses the current vegetation has never been achieved before, making comparison with previous data difficult. Some phytosociological information, dated and at varying degrees of detail, on neighbouring areas [133,148,149], or on the very small provincial reserves within the Padule [150,151], is available, but most of the information originates from floristic studies, which provide general qualitative descriptions of the vegetation. Apart from some data relating to the Padule di Fucecchio contained in ancient Tuscan floras [152–154] or in specific works [155], the first comprehensive work on the flora in the study area is considered to be that by Nannizzi [156]. In this work, the plant landscape is also described in broad terms, although the direct observations date back to 1932–1933. Furthermore, Nannizzi's floristic list complicates matters as he also reports species found by previous authors without distinguishing between them. Consequently, if he does not mention them in the vegetation section, it is difficult to ascertain whether the taxon was discovered by him or only by others prior to his research.

Thirty years after Nannizzi's publication, Lorenzoni and Chiesura Lorenzoni [157] reported the presence and diffusion of some syntaxa in a preliminary note on ongoing research. However, the full study was never subsequently published. They mentioned: (1) *Lemnetum minoris* (Rubel 1930) Pignatti 1953, as very common; (2) *Potameto-Utricularietum* Br.-Bl. 1951, dominated by *Utricularia vulgaris* accompanied by various species of *Potamogeton*, as frequent in shallow waters; (3) *Potameto-Vallisnerietum* Br.-Bl. 1931, dominated by *Myriophyllum spicatum*, *Potamogeton natans*, *Vallisneria spiralis*, together with *Potamogeton crispus*, *Spirodela polyrrhiza*, as not very frequent; (4) *Myriophylleto-Nupharetum* Koch 1926, as the most common and widespread association, where *Nymphoides peltata*, *Myriophyllum verticillatum*, etc., predominate; and (5) three associations of *Phragmitetea*, i.e., *Typhetum angustifoliae* (Allorge 1921) Pign. 1953, with both *Typha angustifolia* and *T. latifolia*, *Phragmitetum communis* (Allorge 1921) Pign. 1953 and *Magnocaricetum elatae* Koch 1926, without reporting their frequency. Regardless of the names they used for the associations, which are incorrect today, it is clear that the vegetation situation was very different from the current one. Lorenzoni and Chiesura Lorenzoni [157] state that, overall, the vegetation at the time of their study corresponds to that of 1932–1933. However, the reciprocal relationship between the communities has changed. Reclamation work involving the almost total elimination of marshy areas and the stabilisation of canal banks determined a clear separation between aquatic and terrestrial hygrophilous vegetation. This allowed anthropised vegetation containing alien species (e.g., *Paspalum distichum*, *Amorpha fruticosa*, and *Bidens frondosa*), which were already present in 1932–1933, to become more widespread by 1968.

Subsequently Garbari [32] published a work on the flora and vegetation of Padule, presenting data collected between 1975 and 1977. This study also reported Nannizzi's species and observed qualitative and quantitative changes in the flora. Regarding the vegetation, Garbari [32] only conducted nine relevés, six of which were related to aquatic and marsh vegetation. These plots are large (80–400 m²), rather heterogeneous, and not in line with current trends of smaller dimensions and more homogeneous communities. However, they provide an overview of the vegetated landscape. His survey includes a reed bed with *Phragmites australis*; a community with *Typha latifolia* and several other marsh and aquatic species (*Potamogeton* sp. pl.); a community of hydrophytes with *Lemna gibba*, *L. minor* and *Wolffia arrhiza*; a community with *Glyceria fluitans* and *Azolla filiculoides* (sub *A. caroliniana*, see Lastrucci et al. [158]) with *Nymphaea alba* and *Hydrocharis morsus-ranae*; a sedge bed with *Carex elata* and others *Carex*; and a community dominated by *Oenanthe aquatica*, with several other aquatic and marsh species. Garbari [32] concludes that, at the time of his study, the aquatic and marsh vegetation of the Padule di Fucecchio was dominated everywhere by two main aspects. One of these was attributable to *Phragmitetea*, with *Phragmitetum communis* having replaced *Carex elata* communities in most of the territory,

although these were still present in residual strips. Another aspect was the vegetation in ditches and canals, which was dominated by *Typha*, *Oenanthe*, *Sparganium*, *Alisma*, and *Sagittaria*, among others.

The observed plant landscape is fundamentally different, particularly with regard to the aquatic vegetation, but also with regard to the marsh communities. Almost all of the aquatic vegetation has disappeared from the Padule di Fucecchio, with only a few small patches remaining, mainly in ditches and canals with clearer water, or in unusual habitats, such as small puddles along paths in the nearby forests. The most emblematic hydrophytes (*Nymphaea alba*, *Hydrocharis morsus-ranae*, *Utricularia* spp., various *Potamogeton* species, etc.) have apparently disappeared. Marsh vegetation is still dominated by *Phragmites australis*, but several marsh plants seem to have disappeared completely (e.g., *Sagittaria sagittifolia*) while others have decreased in number and spread (e.g., *Oenanthe aquatica*). Some species that play an important role today, such as *Xanthium orientale* and many other alien species (e.g., *Eclipta prostrata*, *Lindernia dubia*, etc.), were not even included in the floristic list by Garbari [32].

Why has the plant landscape changed so much? It is probably the result of a combination of factors. Following the Second World War, agricultural changes occurred, with a transition from an agricultural economy (the production of vegetable material for covering flasks and demijohns, as well as straw for chairs, was important in the area) and fishing, to a manufacturing-based economy. This led to highly polluted waters in the marsh. There was also the accidental arrival of several alien plant and animal species, particularly the Louisiana crayfish (*Procambarus clarkii*) and nutria (*Myocastor coypus*) which spread to the area in the late 1990s [39]. From an avifaunistic perspective, the crayfish has been a valuable resource for many marsh birds of conservation interest. However, as has already been observed in other wetlands, the increasing presence of these invasive species has led to the decline and subsequent destruction of numerous aquatic and marshy plant communities [159,160]. Conversely, the avifaunal value of the area has remained substantially unchanged over time and, in some cases, has even improved [161]. The application to the whole area of conservation rules designed specifically for birdlife, which hindered the owners from continuing (at least partly) a traditional management of the territory, was a practice which had in any case supported a certain biodiversity (both zoological and botanical) over time.

4. Conclusions

This study has enabled us to census the current aquatic and marsh vegetation of the Padule di Fucecchio in unprecedented detail. We ascertained the presence of eight aquatic plant communities and twenty-six palustrine ones. It also reports the presence of community types that were previously unknown in the area. One of these types (*Cyperetum micheliano-difformis*) is completely new and is described here for the first time. However, comparison with previous data has also enabled us to reasonably conclude that crucial changes to the vegetation structure have occurred in recent decades: hydrophyte communities have almost completely disappeared, and many sensitive marsh plant species have become rare or disappeared, being mostly replaced by more resilient native and invasive species. In addition to human disturbance and water pollution, the new vegetation arrangements were probably favoured by the arrival of alien fauna species that have had a profoundly negative impact on the state and dynamics of the vegetation. For these reasons, aquatic and marsh vegetation types should be considered, like other types of plant communities of conservation interest, semi-natural coenoses and should be actively managed. However, it should be kept in mind that management that favours the protection of certain ecosystem components (e.g., birdlife) may not be adequate for the conservation

of others (e.g., aquatic and marsh plants and vegetation). In a territory such as the Padule di Fucecchio, where it is possible to do so, areas with differentiated management should be created in agreement with local stakeholders, with the aim of conserving (and sometimes restoring) the various natural emergencies, both floristic and faunal.

Ultimately, this study confirms the findings of several scholars that the acquisition of taxonomic and syntaxonomic knowledge through herbarium and field surveys conducted by taxonomists remains fundamental to the management and conservation of nature [162–164].

Supplementary Materials: The following supporting information can be downloaded at <https://www.mdpi.com/article/10.3390/land14102096/s1>, Supplementary Tables, divided in: Table S1: General Legend; Table S2: Aquatic plant communities; Table S3: Palustrine plant communities of *Echinochloo cruris-galli-Sporoboletum schoenoidis* association; Table S4: Palustrine plant communities of *Littorellion uniflorae* alliance; Table S5: Palustrine plant communities of *Bidention tripartitae* alliance; Table S6: Palustrine plant communities of *Chenopodion rubri* alliance; Table S7: Palustrine plant communities of *Phragmition australis* alliance; Table S8: Palustrine plant communities of *Magnocaricion elatae* and *Magnocaricion gracilis* alliances; Table S9: Palustrine plant communities of *Glycerio-Sparganietum neglecti* association; Table S10: Palustrine plant communities of *Eleocharito palustris-Sagittarion sagittifoliae* alliance; Table S11: Palustrine plant communities of *Polypogono viridis-Paspaleum distichi* association; Table S12: Palustrine plant communities of *Frangulo alni-Salicetum cinereae* association.

Author Contributions: Conceptualization, investigation, data curation, methodology, formal analysis, validation, writing—original draft preparation, writing—review and editing, L.L. and D.V. All authors have read and agreed to the published version of the manuscript.

Funding: Daniele Viciani acknowledges the support of the National Biodiversity Future Center to University of Florence (Italy), funded by the Italian Ministry of University and Research, PNRR, Missione 4 Componente 2, Investimento 1.4, Project CN00000033.

Data Availability Statement: The original contributions presented in this study are included in the article/Supplementary Material. Further inquiries can be directed to the corresponding author.

Acknowledgments: We would like to thank Lorella Dell’Olmo for preparing Figures 1–4. We would also like to thank our friends Varaldo Macchi and Giovanni Riccioni, who have in-depth knowledge of the Fucecchio marshes. They accompanied us on several excursions and provided us with valuable information. Ultimately, we would like to thank the anonymous reviewers and editors for their comments and suggestions, which helped to improve the manuscript.

Conflicts of Interest: The authors declare no conflicts of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

Appendix A

Syntaxonomic Scheme

Lemnetea O. de Bolòs et Masclans 1955

 Lemnetalia minoris O. de Bolòs et Masclans 1955

 Lemnion minoris O. de Bolòs et Masclans 1955

Lemnetum minuto-gibbae Liberman Cruz, Pedrotti et Venanzoni 1988

Potamogetonetea Klika in Klika et Novák 1941

 Potamogetonetalia Koch 1926

 Potamogetonion Libbert 1931

Persicarietum amphibiae Pohjala 1933 nom. mut. Prunier et al. 2019

Potamo pectinati-Myriophylletum spicati Rivas Goday 1964 corr.

 Conesa 1990

Potamogetonetum denso-nodosi O. de Bolòs 1957

- Callitricho hamulatae-Ranunculetales aquatilis Passarge ex Theurillat in Theurillat et al. 2015
- Ranunculon aquatilis* Passarge ex Theurillat in Theurillat et al. 2015
- Callitriche platycarpa* community
- Callitrichetum stagnalis* Kaiser 1926
- Callitriche brutia* communities
- Potamo crispus*-*Ranunculetum trichophylli* Imchenetzky 1926
- Isoëto-Nanojuncetea Br.-Bl. et Tx. in Br.-Bl. et al. 1952
- Nanocyperetalia Klika 1935
- Verbenion supinae* Slavnić 1951
- Echinochloo cruris-galli-Sporobolus schoenoidis* Paradis & Lorenzoni 1994 nom. mut. nov.
- Littorelletea uniflorae Br.-Bl. et Tx. ex Westhoff, Dijk & Passchier 1946
- Littorelletalia uniflorae Koch ex Tx. 1937
- Littorellion uniflorae Koch ex Klika 1935
- Ranunculo flammulae-Juncetum bulbosi* Oberdorfer 1957
- Juncus bulbosus-Gratiola officinalis* communities
- Bidentetea Tx., Lohmeyer et Preising ex von Rochow 1951
- Bidentetalia Br.-Bl. et Tx. ex Klika et Hadač 1944
- Bidention tripartitae* Nordhagen ex Klika et Hadač 1944
- Cyperetum micheliano-difformis* ass. nova
- Bidens frondosa* communities
- Veronica peregrina* communities
- Portulaca oleracea* communities
- Chenopodium rubri* (Tx. in Poli et J. Tx. 1960) Hilbig et Jage 1972
- Polygonum lapathifolii-Xanthium italicum* Pirola et Rossetti 1974
- Echinochloo-Polygonetum lapathifolii* Soó et Csurös 1947
- Phragmito-Magnocaricetea Klika in Klika et Novák 1941
- Phragmitetalia Koch 1926
- Phragmiton australis* Koch 1926 nom. corr.
- Phragmitetum australis* Savič 1926 nom. corr.
- Typhetum latifoliae* Nowiński 1930
- Typhetum angustifoliae* Pignatti 1953
- Schoenoplectetum lacustris* Chouard 1924 nom. mut. Lastrucci et al. 2023
- Phalarido arundinaceae-Bolboschoenetum laticarpi* Passarge 1999 corr. Krumbiegel 2006
- Magnocaricetalia Pignatti 1953
- Magnocaricion elatae* Koch 1926
- Cyperetum longi* Micevski 1957
- Magnocaricion gracilis* Géhu 1961
- Caricetum ripariae* Máthé et Kovács 1959
- Caricetum vesicariae* Chouard 1924
- Phalaridetum arundinaceae* Libbert 1931
- Leucojum aestivum* communities
- Nasturtio-Glycerietalia Pignatti 1953
- Glycerio-Sparganion* Br.-Bl. et Sissingh in Boer 1942
- Glycerio-Sparganietum neglecti* Koch 1926
- Oenanthetalia aquaticae Hejný ex Balátová-Tuláčková et al. 1993
- Eleocharito palustris-Sagittarion sagittifoliae* Passarge 1964

Eleocharitetum palustris Savič 1926
Oenantho aquaticae-Rorippetum amphibiae Lohmeyer 1950
Bolboschoenetum glauci Grechushkina, Sorokin & Golub 2011
Butometum umbellati Philippi 1973
Molinio-Arrhenatheretea Tx. 1937
Holoschoenetalia Br.-Bl. ex Tchou 1948
Polypogono viridis-Paspalion distichi Br.-Bl. in Br.-Bl., Roussine & Nègre 1952 nom. invers. et mut. nov.
Polypogono viridis-Paspaleum distichi Br.-Bl. in Br.-Bl., Gajewski, Wraber & Walas 1936 nom. invers. et mut. nov.
Franguletea Doing ex Westhoff in Westhoff et Den Held 1969
Salicetalia auritae Doing 1962
Salicion cinereae T. Müller et Görs ex Passarge 1961
Frangulo alni-Salicetum cinereae Graebner et Hueck 1931

References

1. Dudgeon, D.; Arthington, A.H.; Gessner, M.O.; Kawabata, Z.; Knowler, D.J.; Lévêque, C.; Naiman, R.J.; Prieur-Richard, A.-H.; Soto, D.; Stiassny, M.L.J.; et al. Freshwater biodiversity: Importance, threats, status and conservation challenges. *Biol. Rev.* **2006**, *81*, 163–182. [\[CrossRef\]](#)
2. Acreman, M.; Hughes, K.A.; Arthington, A.H.; Tickner, D.; Dueñas, M.A. Protected areas and freshwater biodiversity: A novel systematic review distils eight lessons for effective conservation. *Conserv. Lett.* **2020**, *13*, e12684. [\[CrossRef\]](#)
3. Albert, J.S.; Destouni, G.; Duke-Sylvester, S.M.; Magurran, A.E.; Oberdorff, T.; Reis, R.E.; Winemiller, K.O.; Ripple, W.J. Scientists' warning to humanity on the freshwater biodiversity crisis. *Ambio* **2021**, *50*, 85–94. [\[CrossRef\]](#)
4. Fois, M.; Cuena-Lombraña, A.; Artufel, M.; Attard, V.; Cambria, S.; Farrugia, K.; Marignani, M.; Minissale, P.; Pace, A.; Sciandrello, S.; et al. Plant distribution and conservation in mediterranean islands' lentic wetlands: There is more than aquatic species. *Biodivers. Conserv.* **2024**, *33*, 1373–1392. [\[CrossRef\]](#)
5. De Groot, R.; Brander, L.; Van Der Ploeg, S.; Costanza, R.; Bernard, F.; Braat, L.; Christie, M.; Crossman, N.; Ghermandi, A.; Hein, L.; et al. Global estimates of the value of ecosystems and their services in monetary units. *Ecosyst. Serv.* **2012**, *1*, 50–61. [\[CrossRef\]](#)
6. Zhang, L.L.; Yin, J.X.; Jiang, Y.Z.; Wang, H. Relationship between the hydrological conditions and the distribution of vegetation communities within the Poyang lake national nature reserve, China. *Ecol. Inform.* **2012**, *11*, 65–75. [\[CrossRef\]](#)
7. Mitsch, W.J.; Bernal, B.; Hernandez, M.E. Ecosystem services of wetlands. *Int. J. Biodivers. Sci. Ecosyst. Serv. Manag.* **2015**, *11*, 1–4. [\[CrossRef\]](#)
8. Gardner, R.C.; Barchiesi, S.; Beltrame, C.; Finlayson, C.M.; Galewski, T.; Harrison, I.; Paganini, M.; Perennou, C.; Pritchard, D.E.; Rosenqvist, A.; et al. *State of the World's Wetlands and Their Services to People: A Compilation of Recent Analyses*; Ramsar Briefing Note No. 7; Ramsar Convention Secretariat: Gland, Switzerland, 2015.
9. Li, L.; Xu, H.; Zhang, Q.; Zhan, Z.; Liang, X.; Xing, J. Estimation methods of wetland carbon sink and factors influencing wetland carbon cycle: A review. *Carbon Res.* **2024**, *3*, 50. [\[CrossRef\]](#)
10. Hrivnák, R.; Kochjarová, J.; Oľahel'ová, H.; Paľove-Balang, P.; Slezák, M.; Slezák, P. Environmental drivers of macrophyte species richness in artificial and natural aquatic water bodies—comparative approach from two central European regions. *Ann. Limnol. Int. J. Limnol.* **2014**, *50*, 269–278. [\[CrossRef\]](#)
11. Fois, M.; Cuena-Lombraña, A.; Bacchetta, G. Knowledge gaps and challenges for conservation of Mediterranean wetlands: Evidence from a comprehensive inventory and literature analysis for Sardinia. *Aquat. Conserv. Mar. Freshw. Ecosyst.* **2021**, *31*, 2621–2631. [\[CrossRef\]](#)
12. Ballut-Dajud, G.A.; Sandoval Herazo, L.C.; Fernández-Lambert, G.; Marín-Muñoz, J.L.; López Méndez, M.C.; Betanzo-Torres, E.A. Factors affecting wetland loss: A review. *Land* **2022**, *11*, 434. [\[CrossRef\]](#)
13. Xi, Y.; Peng, S.; Ciaia, P.; Chen, Y. Future impacts of climate change on inland Ramsar wetlands. *Nat. Clim. Chang.* **2021**, *11*, 45–51. [\[CrossRef\]](#)
14. Leberger, R.; Geijzendorffer, I.R.; Gaget, E.; Gwelami, A.; Galewski, T.; Pereira, H.M.; Guerra, C.A. Mediterranean wetland conservation in the context of climate and land cover change. *Reg. Environ. Change* **2020**, *20*, 67. [\[CrossRef\]](#)
15. Reid, A.J.; Carlson, A.K.; Creed, I.F.; Eliason, E.J.; Gell, P.A.; Johnson, P.T.; Kidd, K.A.; MacCormack, T.J.; Olden, J.D.; Ormerod, S.J.; et al. Emerging threats and persistent conservation challenges for freshwater biodiversity. *Biol. Rev.* **2019**, *94*, 849–873. [\[CrossRef\]](#) [\[PubMed\]](#)

16. Fluët-Chouinard, E.; Stocker, B.D.; Zhang, Z.; Malhotra, A.; Melton, J.R.; Poulter, B.; Kaplan, J.O.; Goldewijk, K.K.; Siebert, S.; Minayeva, T.; et al. Extensive global wetland loss over the past three centuries. *Nature* **2023**, *614*, 281–286. [\[CrossRef\]](#)
17. Xu, T.; Weng, B.; Yan, D.; Wang, K.; Li, X.; Bi, W.; Li, M.; Cheng, X.; Liu, Y. Wetlands of international importance: Status, threats and future protection. *Int. J. Environ. Res. Public Health* **2019**, *16*, 1818. [\[CrossRef\]](#)
18. Janssen, J.A.M.; Rodwell, J.S.; Garcia Criado, M.; Gubbay, S.; Haynes, T.; Nieto, A.; Sanders, N.; Landucci, F.; Loidi, J.; Symank, A.; et al. *European Red List of Habitats*; Publications Office of European Union: Luxembourg, 2016; ISBN 978-92-79-61588-7.
19. Zivkovic, L.; Biondi, E.; Pesaresi, S.; Lasen, C.; Spampinato, G.; Angelini, P. The third report on the conservation status of habitats (Directive 92/43/EEC) in Italy: Processes, methodologies, results and comments. *Plant Sociol.* **2017**, *54*, 51–64.
20. Gigante, D.; Acosta, A.T.R.; Agrillo, E.; Armiraglio, S.; Assini, S.; Attorre, F.; Bagella, S.; Buffa, G.; Casella, L.; Giancola, C.; et al. Habitat conservation in Italy: The state of the art in the light of the first European Red List of Terrestrial and Freshwater Habitats. *Rend. Lincei. Sci. Fis. Nat.* **2018**, *29*, 251–265. [\[CrossRef\]](#)
21. Lazzaro, L.; Bolpagni, R.; Buffa, G.; Gentili, R.; Lonati, M.; Stinca, A.; Acosta, A.T.R.; Adorni, M.; Aleffi, M.; Allegrezza, M.; et al. Impact of invasive alien plants on native plant communities and Natura 2000 Habitats: State of the art, gap analysis and perspectives in Italy. *J. Environ. Manag.* **2020**, *274*, 111140. [\[CrossRef\]](#) [\[PubMed\]](#)
22. Viciani, D.; Vidali, M.; Gigante, D.; Bolpagni, R.; Villani, M.; Acosta, A.T.R.; Adorni, M.; Aleffi, M.; Allegrezza, M.; Angiolini, C.; et al. A first checklist of the alien-dominated vegetation in Italy. *Plant Sociol.* **2020**, *57*, 29–54. [\[CrossRef\]](#)
23. Gennai, M.; Gabellini, A.; Viciani, D.; Venanzoni, R.; Dell’Olmo, L.; Giunti, M.; Lucchesi, F.; Monacci, F.; Mugnai, M.; Foggi, B. The floodplain woods of Tuscany: Towards a phytosociological synthesis. *Plant Sociol.* **2021**, *58*, 1–28. [\[CrossRef\]](#)
24. European Commission. Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora. *Off. J. Eur. Union* **1992**, *206*, 50.
25. Benavent-González, A.; Lumbreras, A.; Molina, J.A. Plant communities as a tool for setting priorities in biodiversity conservation: A novel approach to Iberian aquatic vegetation. *Biodivers. Conserv.* **2014**, *23*, 2135–2154. [\[CrossRef\]](#)
26. Lastrucci, L.; Bonari, G.; Angiolini, C.; Casini, F.; Giallonardo, T.; Gigante, D.; Landi, M.; Landucci, F.; Venanzoni, R.; Viciani, D. Vegetation of Lakes Chiusi and Montepulciano (Siena, central Italy): Updated knowledge and new discoveries. *Plant Sociol.* **2014**, *51*, 29–55.
27. Lastrucci, L.; Cerri, M.; Coppi, A.; Dell’Olmo, L.; Ferranti, F.; Ferri, V.; Filipponi, F.; Foggi, B.; Galardini, R.; Reale, L.; et al. Spatial landscape patterns and trends of declining reed-beds in peninsular Italy. *Plant Biosyst.* **2019**, *153*, 427–435. [\[CrossRef\]](#)
28. Angiolini, C.; Viciani, D.; Bonari, G.; Lastrucci, L. Habitat conservation prioritization: A floristic approach applied to a Mediterranean wetland network. *Plant Biosyst.* **2017**, *151*, 598–612. [\[CrossRef\]](#)
29. Casavecchia, S.; Allegrezza, M.; Angiolini, C.; Biondi, E.; Bonini, F.; Del Vico, E.; Fanfarillo, E.; Foggi, B.; Gigante, D.; Gianguzzi, L.; et al. Proposals for improvement of Annex I of Directive 92/43/EEC: Central Italy. *Plant Sociol.* **2021**, *58*, 99–118. [\[CrossRef\]](#)
30. Bartolini, A. La Riserva Naturale del Padule di Fucecchio. Dieci anni di gestione (1996–2006). *Quad. Padule Fucecchio* **2007**, *4*, 1–165.
31. Puglisi, L.; Arcamone, E.; Franchini, M.; Giunchi, D.; Meschini, E.; Sacchetti, A.; Vanni, L.; Vezzani, A. *Atlante Degli Uccelli Nidificanti e Svernanti in Toscana. 2. Distribuzione, Abbondanza e Conservazione*; Centro Ornitologico Toscano: Tuscany, Italy; ETS Edizioni: Pisa, Italy, 2024.
32. Garbari, F. Indagine Floristica e Vegetazionale sul Padule di Fucecchio. In *Progetto Pilota per la Salvaguardia e la Valorizzazione del Padule di Fucecchio*; Consorzio di Bonifica del Padule di Fucecchio, Ed.; Arti Grafiche Giorgi & Gambi: Firenze, Italy, 1980; pp. 217–263.
33. Tomei, P.E.; Guazzi, E.; Kugler, P.C. *Le Zone Umide Della Toscana: Indagine Sulle Componenti Floristiche e Vegetazionali*; Regione Toscana: Firenze, Italy, 2001; p. 167.
34. Lastrucci, L.; Selvi, F.; Coppi, A.; Viciani, D. Ricerche botaniche del Museo di Storia Naturale dell’Università di Firenze sulle aree umide della Toscana. *Not. Della Soc. Bot. Ital.* **2024**, *6*, 112–113.
35. Padule di Fucecchio. 2025. Available online: https://it.wikipedia.org/wiki/Padule_di_Fucecchio (accessed on 15 May 2025).
36. Chetoni, R. Geologia: Idrologia ed Evoluzione Paleogeografica del Padule di Fucecchio. In *Il Padule di Fucecchio e il Lago di Sibolla. Natura e Storia*; Dani, F.R., Ed.; Centro Ricerca, Documentazione e Promozione Padule di Fucecchio: Castelmartini, Italy, 1999; pp. 41–51.
37. Corsi, R. Flora e Vegetazione NELL’AREA del Padule di Fucecchio. In *Il Padule di Fucecchio e il Lago di Sibolla. Natura e Storia*; Dani, F.R., Ed.; Centro Ricerca, Documentazione e Promozione Padule di Fucecchio: Castelmartini, Italy, 1999; pp. 53–74.
38. Malvolti, A. Il Padule Nella Storia. In *Il Padule di Fucecchio e il Lago di Sibolla. Natura e Storia*; Dani, F.R., Ed.; Centro Ricerca, Documentazione e Promozione Padule di Fucecchio: Castelmartini, Italy, 1999; pp. 21–40.
39. Bartolini, A.; Nocita, A.; Bartali, S.; Pini, G.; Vigiani, V.; Franzese, M.; Vezzani, A. *Studio per la Salvaguardia Della Biodiversità del S.I.R. Padule di Fucecchio*; Centro Ricerca, Documentazione e Promozione Padule di Fucecchio: Castelmartini, Italy, 2004; p. 68.

40. Regione Toscana. CASTORE—Cartografia Storica Regionale. Carta Topografica Della Pianura del Valdarno di Sotto con i Confini delle Respettive Comunità Dedotta Dall’Originale Autentico Eseguito Nell’Anno 1747. 2025. Available online: https://www502.regione.toscana.it/cartografia_storica_regionale_gi/cartografia_storica_regionale_smartviewer.html?img=11191 (accessed on 15 April 2025).
41. Stazione Meteorologica di Larciano Castelmartini. Available online: https://it.wikipedia.org/wiki/Stazione_meteorologica_di_Larciano_Castelmartini (accessed on 15 May 2025).
42. Bigi, L.; Rustici, L. *Regime Idrico dei Suoli e Tipi Climatici in Toscana*; Dipartimento dell’Agricoltura e delle Foreste: Tuscany/Firenze, Italy, 1984; p. 130.
43. Thornthwaite, C.W.; Mather, J.R. Instruction and tables for computing potential evapotranspiration and the water balance. *Lab. Climatol.* **1957**, *10*, 1–311.
44. Pesaresi, S.; Biondi, E.; Casavecchia, S. Bioclimates of Italy. *J. Maps* **2017**, *13*, 955–960. [\[CrossRef\]](#)
45. Carmignani, L.; Conti, P.; Cornamusini, G.; Pirro, A. Geological map of Tuscany (Italy). *J. Maps* **2013**, *9*, 487–497. [\[CrossRef\]](#)
46. Braun-Blanquet, J. *Plant Sociology: The Study of Plant Communities*; McGraw-Hill: New York, NY, USA, 1932.
47. Braun-Blanquet, J. *Pflanzensoziologie: Grundzüge der Vegetationskunde*, 3rd ed.; Springer: Wien, Austria, 1964; pp. 1–330.
48. Dengler, J.; Berg, C.; Jansen, F. New ideas for modern phytosociological monographs. *Ann. Bot.* **2005**, *5*, 195–213.
49. Dengler, J.; Chytrý, M.; Ewald, J. Phytosociology. In *Encyclopedia of Ecology*; Jorgensen, S.E., Fath, B.D., Eds.; Elsevier: Oxford, UK, 2008; pp. 2767–2779.
50. Biondi, E. Phytosociology today: Methodological and conceptual evolution. *Plant Biosyst.* **2011**, *145* (Suppl. S1), 19–29. [\[CrossRef\]](#)
51. Pott, R. Phytosociology: A modern geobotanical method. *Plant Biosyst.* **2011**, *145* (Suppl. S1), 9–18. [\[CrossRef\]](#)
52. Van der Maarel, E. Transformation of cover-abundance values in phytosociology and its effects on community similarity. *Vegetatio* **1979**, *39*, 97–114.
53. Arrigoni, P.V. A classification of plant growth forms applicable to the floras and vegetation types of Italy. *Webbia* **1996**, *50*, 193–203. [\[CrossRef\]](#)
54. R Core Team. *R: A Language and Environment for Statistical Computing*; R Foundation for Statistical Computing: Vienna, Austria, 2024; Available online: <https://www.R-project.org/> (accessed on 15 January 2025).
55. Oksanen, J.; Blanchet, F.G.; Friendly, M.; Kindt, R.; Legendre, P.; McGlinn, D.; Ouellette, M.-H.; Cunha, E.R.; Smith, T.; Stier, A.; et al. *Vegan: Community Ecology Package*, version 2.6-6.1.; Comprehensive R Archive Network: Vienna, Austria, 2024. Available online: <https://CRAN.R-project.org/package=vegan> (accessed on 15 January 2025).
56. Portale della Flora d’Italia—Portal to the Flora of Italy 2024.3. 2025. Available online: <http://dryades.units.it/floritaly> (accessed on 12 February 2025).
57. Mucina, L.; Bültmann, H.; Dierßen, K.; Theurillat, J.P.; Raus, T.; Čarni, A.; Šumberová, K.; Willner, W.; Dengler, J.; García, R.G.; et al. Vegetation of Europe: Hierarchical floristic classification system of vascular plant, bryophyte, lichen, and algal communities. *Appl. Veg. Sci.* **2016**, *19* (Suppl. S1), 3–264. [\[CrossRef\]](#)
58. Chytrý, M.; Řezníčková, M.; Novotný, P.; Holubová, D.; Preislerová, Z.; Attorre, F.; Biurrun, I.; Blažek, P.; Bonari, G.; Borovyk, D.; et al. FloraVeg.EU—An online database of European vegetation, habitats and flora. *Appl. Veg. Sci.* **2024**, *27*, e12798. [\[CrossRef\]](#)
59. Biondi, E.; Blasi, C. *Prodromo della Vegetazione Italiana*. 2015. Available online: <http://www.prodromo-vegetazione-italia.org/> (accessed on 12 February 2025).
60. Theurillat, J.P.; Willner, W.; Fernandez-Gonzalez, F.; Bultmann, H.; Čarni, A.; Gigante, D.; Mucina, L.; Weber, H. International Code of Phytosociological Nomenclature. *Appl. Veg. Sci.* **2021**, *24*, e12491. [\[CrossRef\]](#)
61. Liberman Cruz, M.; Pedrotti, F.; Venanzoni, R. Le associazioni della classe Lemnatea del lago Titicaca (Bolivia). *Hidrobiology* **1988**, *2*, 377–388.
62. Landucci, F.; Gigante, D.; Venanzoni, R. An application of the Cocktail method for the classification of the hydrophytic vegetation at Lake Trasimeno (Central Italy). *Fitosociologia* **2011**, *48*, 3–22.
63. Lastrucci, L.; Valentini, E.; Dell’Olmo, L.; Vietina, B.; Foggi, B. Hygrophilous vegetation and habitats of conservation interest in the area of the Lake Porta (Tuscany, Central Italy). *Atti Soc. Tosc. Sci. Nat. Mem. Serie B* **2016**, *122*, 131–146.
64. Lastrucci, L.; Dell’Olmo, L.; Foggi, B.; Massi, L.; Nuccio, C.; Vicenti, C.; Viciani, D. Contribution to the knowledge of the vegetation of the Lake Massaciuccoli (northern Tuscany, Italy). *Plant Sociol.* **2017**, *54*, 67–87. [\[CrossRef\]](#)
65. Bramley, J.L.; Reeve, J.T.; Dussart, B.J. The Distribution of an Alien Duckweed, *Lemna minuta*, Within the British Isles: Identification, Dispersal and Niche Constraints. In *Plant Invasions: General Aspects and Special Problems*; Pyšek, P., Prach, K., Rejmánek, M., Wade, M., Eds.; SPB Academic Publishing: Amsterdam, The Netherlands, 1995; pp. 181–185.
66. Prunier, P.; Greulich, F.; Béguin, C.; Boissezon, A.; Delarze, R.; Hegg, O.; Klötlzi, F.; Pantke, R.; Steffen, J.; Steiger, P.; et al. Phytosuisse: Il Riferimento Fitosociologico per la Svizzera. 2019; ISSN 2624-6171. Available online: [www.infoflora.ch/it/ambienti/phytosuisse-\(prunier-et-al.\).html](http://www.infoflora.ch/it/ambienti/phytosuisse-(prunier-et-al.).html) (accessed on 10 April 2025).
67. Kłosowski, S.; Tomaszewicz, H. Habitat requirements of *Polygonetum natantis* Soó 1927 and *Potamogetonetum natantis* Soó 1927 phytocenoses in north-eastern Poland. *Acta Soc. Bot. Pol.* **1986**, *55*, 141–157. [\[CrossRef\]](#)

68. Lastrucci, L.; Landi, M.; Angiolini, C. Vegetation analysis on wetlands in a Tuscan agricultural landscape (central Italy). *Biologia* **2010**, *65*, 54–68. [\[CrossRef\]](#)
69. Šumberová, K. Vegetace Vodních Rostlin Zakořeněných ve dně (Potametea). Vegetation of Aquatic Plants Rooted in the Bottom. In *Vegetace České Republiky 3. Vodní a Mokřadní Vegetace. Vegetation of the Czech Republic 3. Aquatic and Wetland Vegetation*; Chytrý, M., Ed.; Academia: Prague, Czech Republic, 2011; pp. 100–247.
70. Loidi, J.A.; Biurrun, I.G.; Herrera, M.G. La vegetación del centro-septentrional de España. *Itinera Geobot.* **1997**, *9*, 161–618.
71. Ceschin, S.; Zuccarello, V.; Caneva, G. Role of macrophyte communities as bioindicators of water quality Application on the Tiber River basin (Italy). *Plant Biosyst.* **2010**, *144*, 528–536. [\[CrossRef\]](#)
72. Zervas, D.; Tsiripidis, I.; Bergmeier, E.; Tsiaoussi, V. A phytosociological survey of aquatic vegetation in the main freshwater lakes of Greece. *Veg. Classific. Surv.* **2020**, *1*, 53–75. [\[CrossRef\]](#)
73. Viciani, D.; Angiolini, C.; Bonari, G.; Bottacci, A.; Dell’Olmo, L.; Gonnelli, V.; Zoccola, A.; Lastrucci, L. Contribution to the knowledge of aquatic vegetation of montane and submontane areas of Northern Apennines (Italy). *Plant Sociol.* **2022**, *59*, 25–35. [\[CrossRef\]](#)
74. Hrivnák, R. Aquatic plant communities in the catchment area of the Ipel’ river in Slovakia and Hungary. Part II. Class Potametea. *Thaiszia* **2002**, *12*, 137–160.
75. Lastrucci, L.; Selvi, F.; Bajona, E.; Sforzi, A.; Siccardi, E.; Viciani, D. Advancing Knowledge of Wetland Vegetation for Plant Diversity Conservation: The Case of Small Lakes, Ponds, and Pools in Maremma (Southern Tuscany, Central Italy). *Land* **2025**, *14*, 218. [\[CrossRef\]](#)
76. Lastrucci, L.; Saiani, D.; Mugnai, A.; Ferretti, G.; Viciani, D. Distribution novelties of the genus *Callitriche* (Plantaginaceae) in Italy from the study of the Herbarium Centrale Italicum collections. *Mediterr. Bot.* **2024**, *45*, e87474. [\[CrossRef\]](#)
77. Felzines, J.C. Contribution au prodrome des végétations de France: Les Potametea Klika in Klika & V. Novak 1941. *Doc. Phytosoc.* **2016**, *3*, 218–437.
78. Passarge, H. Mitteleuropäische Potamogetonetea, I. *Phytocoenologia* **1992**, *20*, 489–527. [\[CrossRef\]](#)
79. Ceschin, S.; Salerno, G. La vegetazione del basso corso del Fiume Tevere e dei suoi affluenti (Lazio; Italia). *Fitosociologia* **2008**, *45*, 39–74.
80. Brullo, S.; Scelsi, F.; Spampinato, F. *La Vegetazione dell’Aspromonte*; Laruffa Editore: Reggio Calabria, Italy, 2001.
81. Baldoni, M.; Biondi, E. La vegetazione del medio e basso corso del fiume Esino (Marche-Italia centrale). *Stud. Bot.* **1993**, *11*, 209–257.
82. Venanzoni, R.; Gigante, D. Contributo alla conoscenza della vegetazione degli ambienti umidi dell’Umbria (Italia). *Fitosociologia* **2000**, *37*, 13–63.
83. Lastrucci, L.; Foggi, B.; Selvi, F.; Becattini, R. Contributo alla conoscenza della vegetazione e della flora delle aree umide nel comprensorio di Capalbio (Provincia di Grosseto, Italia centrale). *Arch. Geobot.* **2007**, *10*, 1–30.
84. Gigante, D.; Maneli, F.; Venanzoni, R. Mediterranean temporary wet systems in inland Central Italy: Ecological and phytosociological features. *Plant Sociol.* **2013**, *50*, 93–112.
85. Bagella, S.; Caria, M.C.; Farris, E.; Filigheddu, R. Phytosociological analysis in Sardinian Mediterranean temporary wet habitats. *Fitosociologia* **2009**, *46*, 11–26.
86. Tomaselli, V.; Beccarisi, L.; Brullo, S.; Cambria, S.; Forte, L.; Minissale, P.; Veronico, G. Phytosociological research on temporary ponds in Apulia (southern Italy). *Mediterr. Bot.* **2020**, *41*, 15–41. [\[CrossRef\]](#)
87. Tomaselli, V.; Beccarisi, L.; Cambria, S.; Forte, L.; Minissale, P.; Sciandrello, S.; Veronico, G.; Brullo, S. Validation of associations for the temporary ponds of the class *Isoeto-Nanojuncetea* in Puglia (southern Italy). *Mediterr. Bot.* **2022**, *43*, e80627. [\[CrossRef\]](#)
88. Brullo, S.; Brullo, C.; Sciandrello, S.; Tavilla, G.; Cambria, S.; Tomaselli, V.; Ilardi, V.; Giusso del Galdo, G.; Minissale, P. The plant communities of the class *Isoeto-Nanojuncetea* in Sicily. *Plants* **2022**, *11*, 1214. [\[CrossRef\]](#)
89. Pedrotti, F.; Russo, G.; Murrja, E. Flora, plant communities and vegetation mapping of the lake d’Otri (Gargano, Apulia, Italy). *Contrib. Bot.* **2022**, *57*, 53–62. [\[CrossRef\]](#)
90. Brullo, S.; Minissale, P. Considerazioni syntassonomiche sulla classe *Isoeto-Nanojuncetea*. *Itinera Geobot.* **1998**, *11*, 263–290.
91. Eliáš, P.; Dítě, D.; Grulich, V.; Sádovský, M. Distribution and communities of *Crypsis aculeata* and *Heleochoa schoenoides* in Slovakia. *Hacquetia* **2008**, *7*, 5–20. [\[CrossRef\]](#)
92. Eliáš, P.; Dítě, D.; Dítě, Z. Halophytic Vegetation in the Pannonian Basin: Origin, Syntaxonomy, Threat, and Conservation. In *Handbook of Halophytes*; Grigore, M.N., Ed.; Springer: Cham, Switzerland, 2020. [\[CrossRef\]](#)
93. Paradis, G. Observations synécologiques sur des stations corses de trois thérophytes fini-estivales: *Crypsis aculeata*, *Crypsis schoenoides* et *Chenopodium chenopodioides*. *Monde Plantes* **1992**, *444*, 1–21.
94. Paradis, G.; Lorenzoni, C. Etude phytosociologique de communautés thérophytiques hygro-nitrophiles estivo-autumnales de la Corse (grpts. à *Crypsis aculeata*, *C. schoenoides*, *Glinus lotoides* et *Chenopodium chenopodioides*). Nouvelles propositions syntaxoniques. 2e contribution. *Monde Plantes* **1994**, *449*, 19–26.
95. Martínez-Parras, J.M.; Lorca, M.P.; Esteban, C.B.; Mesa, J.M. Algunas comunidades vegetales higrófilas e higrónitrófilas estivo-autumnales de la provincia de Granada. *Acta Bot. Barcinon.* **1988**, *37*, 271–279.

96. Rivas-Martinez, S.; Fernandez-Gonzalez, F.; Loidi, J.; Lous, M.; Penas, A. Syntaxonomical Checklist of vascular plant communities of Spain and Portugal to association level. *Itinera Geobot.* **2001**, *14*, 5–341.
97. Dawson, F.H.; Szoszkiewicz, K. Relationships of some ecological factors with the associations of vegetation in British rivers. *Hydrobiologia* **1999**, *415*, 117–122. [[CrossRef](#)]
98. Šumberová, K.; Navrátilová, J.; Čtvrtlíková, M.; Hájek, M.; Bauer, P. Vegetace Oligotrofních Vod (*Littorelletea uniflorae*). Vegetation of Oligotrophic Water Bodies. In *Vegetace České Republiky 3. Vodní a Mokřadní Vegetace. Vegetation of the Czech Republic 3. Aquatic and Wetland Vegetation*; Chytrý, M., Ed.; Academia: Prague, Czech Republic, 2011; pp. 268–308.
99. Ninot, J.M.; Carreras, J.; Carrillo, E.; Vigo, J. Syntaxonomic conspectus of the vegetation of Catalonia and Andorra. I: Hygrophilous herbaceous communities. *Acta Bot. Barc.* **2000**, *46*, 191–237.
100. Szankowski, M.; Kłosowski, S. Habitat variability of the *Littorelletea uniflorae* plant communities in Polish Lobelia lakes. *Hydrobiologia* **2006**, *570*, 117–126. [[CrossRef](#)]
101. Hrivnák, R.; Kochjarová, J.; Oľahel'ová, H. Vegetation of the aquatic and marshland habitats in the Orava region, including the first records of *Potametum alpini*, *Potametum zizii* and *Ranunculo-Juncetum bulbosi* in the territory of Slovakia. *Biologia* **2011**, *66*, 626–637.
102. Pellizzari, M. *Cyperus*-dominated vegetation in the eastern Po River. *Plant Sociol.* **2020**, *57*, 1–16. [[CrossRef](#)]
103. Piccoli, F.; Gerdol, R. Rice-field weed communities in Ferrara Province (northern Italy). *Aquatic Bot.* **1981**, *10*, 317–328. [[CrossRef](#)]
104. Melečková, Z.; Dítě, D.; Eliáš, P.; Schmidt, D. *Cyperus glomeratus* L. rediscovered in Slovakia. *Hacquetia* **2016**, *15*, 93–100. [[CrossRef](#)]
105. Šumberová, K.; Lososová, Z. Vegetace Jednoletých Nitrofilních Vlhkomilných Bylin (*Bidentetea tripartitae*). Vegetation of Annual Nitrophilous Wetland Herbs. In *Vegetace České Republiky 3. Vodní a Mokřadní Vegetace. Vegetation of the Czech Republic 3. Aquatic and Wetland Vegetation*; Chytrý, M., Ed.; Academia: Prague, Czech Republic, 2011; pp. 347–384.
106. Amigo, J. Los herbazales terofíticos higronitrófilos en el noroeste de la Península Ibérica (*Bidentetea tripartitae* Tüxen; Lohmeyer & Preising ex von Rochow 1951). *Lazaroa* **2006**, *27*, 43–58.
107. Roma-Marzio, F.; Lastrucci, L.; Guzzon, F.; Ardenghi, N.M.G.; Peruzzi, L.; Mossini, S. Nuove segnalazioni floristiche italiane 5. Flora vascolare (28–46). *Not. Soc. Bot. Ital.* **2018**, *2*, 205–210.
108. Maiorca, G.; Crisafulli, A.; Puntillo, D.; Signorino, G.; Spampinato, G. Wetland vegetation of the Tarsia Lake Regional Nature Reserve (Calabria, southern Italy). *Mediterr. Bot.* **2020**, *41*, 67–84.
109. Passarge, H. *Pflanzenengesellschaften Norostdeutschland*; J. Cramer: Berlin/Stuttgart, Germany, 1996.
110. Julve, P. Les Base de Données Botaniques. Baseveg. Baseflor. Programme Catminat. 2025. Available online: <https://www.tela-botanica.org/ressources/donnees/telechargements/> (accessed on 15 March 2025).
111. Chytrý, M.; Tichý, L. *Diagnostic, Constant and Dominant Species of Vegetation Classes and Alliances of the Czech Republic: A Statistical Revision*; Masaryk University: Brno, Czech Republic, 2003; Volume 108, pp. 1–231.
112. Lososová, Z. Portulacetum Oleraceae Felföldy 1942. In *Vegetace České Republiky. 2. Ruderalní, Plevelová, Skalní a Sut'ová Vegetace. Vegetation of the Czech Republic 2. Ruderal, Weed, Rock and Scree Vegetation*; Chytrý, M., Ed.; Academia: Prague, Czech Republic, 2009; pp. 198–200.
113. Biondi, E.; Baldoni, M. La vegetazione del fiume Marecchia (Italia Centrale). *Biogeographia* **1994**, *17*, 51–87. [[CrossRef](#)]
114. Biondi, E.; Vagge, I.; Baldoni, M.; Taffetani, F. Biodiversità fitocenotica e paesaggistica dei fiumi dell'Italia centro-settentrionale: Aspetti fitosociologici e sinfitosociologici. *Studi Trent. Sci. Nat. Acta Biol.* **2004**, *80*, 13–21.
115. Tomasello, S. How many names for a beloved genus?—Coalescent-based species delimitation in *Xanthium* L. (Ambrosiinae, Asteraceae). *Mol. Phylogenetics Evol.* **2018**, *127*, 135–145. [[CrossRef](#)] [[PubMed](#)]
116. Geisselbrecht-Taferner, L.; Mucina, L. *Bidentetea Tripartiti*. In *Pflanzenengesellschaften Österreich. Teil 1*; Mucina, L., Grabherr, G., Ellmauer, T., Eds.; Gustav Fischer Verlag: Jena/Stuttgart, Germany; New York, NY, USA, 1993; pp. 90–109.
117. Sanda, V.; Öllerer, K.; Burescu, P. *Fitocenozele din România*; Ars Docendi: București, Romania, 2008; p. 570.
118. Neacșu, A.; Arsene, G.G. Pioneer vegetation on water shores (*Bidenti-Polygonetum hydropiperi* Lohmeyer in R. Tüxen 1950 and *Echinochloo-Polygonetum lapathifolii* Soó et Csűrös 1974) in the Romanian Banat. *Res. J. Agric. Sci.* **2017**, *49*, 122–135.
119. Felzines, J.C.; Loiseau, J.E. Les groupements fluviaux des *Bidentetea* de la Loire moyenne, du bas Allier et de la Dordogne moyenne. Modifications apportées à la synsystème de la classe des *Bidentetea*. *Bull. Soc. Bot. Cent. Ouest* **2005**, *36*, 159–204.
120. Landucci, F.; Gigante, D.; Venanzoni, R.; Chytrý, M. Wetland vegetation of the class *Phragmito-Magno-Caricetea* in central Italy. *Phytocoenologia* **2013**, *43*, 67–100. [[CrossRef](#)]
121. Landucci, F.; Šumberová, K.; Tichý, L.; Hennekens, S.; Aunina, L.; Biță-Nicolae, C.; Borsukevych, L.; Bobrov, A.; Čarni, A.; De Bie, E.; et al. Classification of the European marsh vegetation (*Phragmito-Magnocaricetea*) to the association level. *Appl. Veg. Sci.* **2020**, *23*, 297–316. [[CrossRef](#)]
122. Lastrucci, L.; Angiolini, C.; Bonari, G.; Bottacci, A.; Gonnelli, V.; Zoccola, A.; Mugnai, M.; Viciani, D. Contribution to the knowledge of marsh vegetation of montane and submontane areas of Northern Apennines (Italy). *Plant Sociol.* **2023**, *60*, 25–36. [[CrossRef](#)]

123. Šumberová, K.; Hájková, P.; Chytrý, M.; Hroudová, Z.; Sádlo, J.; Hájek, M.; Hrivnák, R.; Navrátilová, J.; Hanáková, P.; Ekrt, L.; et al. Vegetace Rákosin a Vysokých Ostřic (*Phragmito-Magno-Caricetea*). Marsh Vegetation. In *Vegetace České Republiky 3. Vodní a Mokřadní Vegetace. Vegetation of the Czech Republic 3. Aquatic and Wetland Vegetation*; Chytrý, M., Ed.; Academia: Prague, Czech Republic, 2011; pp. 385–580.
124. Hrivnák, R. The plant communities of Phragmitetalia in the catchment area of the Ipel'river (Slovakia and Hungary), 2, Tall-sedge dominated wetlands (*Magnocaricion elatae*). *Biol. Bratisl.* **2004**, *59*, 457–476.
125. Fernez, T.; Causse, G. *Synopsis Phytosociologique des Groupements Végétaux d'Île-de-France, version 1*; Conservatoire Botanique National du Bassin Parisien: Paris, France, 2015; p. 89.
126. Poldini, L. *La Vegetazione del Carso Isontino e Triestino*; Lint: Trieste, Italy, 1989.
127. Lastrucci, L.; Ferretti, G.; Mantarano, N.; Foggi, B. Vegetation and habitat of conservation interest of the lake Acquato (Grosseto—Italy). *Plant Sociol.* **2019**, *56*, 19–30.
128. Hroudová, Z.; Hrivnák, R.; Chytrý, M. Classification of inland *Bolboschoenus*-dominated vegetation in Central Europe. *Phytocoenologia* **2009**, *39*, 205–215. [\[CrossRef\]](#)
129. Di Natale, S.; Lastrucci, L.; Hroudova, Z.; Viciani, D. A review of *Bolboschoenus* species (Cyperaceae) in Italy based on herbarium data. *Plant Biosyst.* **2022**, *156*, 261–270. [\[CrossRef\]](#)
130. Horvat, I.; Glavač, V.; Ellenberg, H. *Vegetation Südosteuropas*; G. Fischer Verlag: Stuttgart, Germany, 1974.
131. Devillers, P.; Devillers-Terschuren, J. *Application and Development of the Palearctic Habitat Classification in the Course of the Setting up of the Emerald Project, Malta*; Council of Europe: Strasbourg, France, 2001.
132. Buchwald, R. Vegetazione e odonotofauna negli ambienti acquatici dell'Italia Centrale. *Braun Blanquetia* **1994**, *11*, 1–77.
133. Lastrucci, L.; Viciani, D.; Nuccio, C.; Melillo, C. Indagine vegetazionale su alcuni laghi di origine artificiale limitrofi al Padule di Fucecchio (Toscana, Italia Centrale). *Ann. Mus. Civ. Rovereto* **2008**, *23*, 169–203.
134. Hahn, I.; Gergely, A.; Barabás, S. Changes in the active floodplain vegetation of the Szigetköz. *Ann. Bot.* **2011**, *1*, 1–8.
135. Poldini, L.; Vidali, M.; Castello, M.; Sburlino, G. A novel insight into the remnants of hygrophilous forests and scrubs of the Po Plain biogeographical transition area (Northern Italy). *Plant Sociol.* **2020**, *57*, 17–69. [\[CrossRef\]](#)
136. Postolache, G. Diversitatea cenotaxonomică a vegetației Republicii Moldova. *Rev. Bot.* **2022**, *25*, 24–43.
137. Lastrucci, L.; Gambirasio, V.; Prosser, F.; Viciani, D. First record of *Sparganium oocarpum* in Italy and new regional distribution data for *S. erectum* species complex. *Plant Biosyst.* **2024**, *158*, 595–600. [\[CrossRef\]](#)
138. Lastrucci, L.; Lazzaro, L.; Lunardi, L.; Fiorini, G.; Viciani, D. Morphometric surveys on *Eleocharis palustris* (L.) Roem. et Schult. (Cyperaceae). The contribution of herbarium investigations to the delimitation of its subspecies and their distribution in Italy. *Ann. Bot.* **2020**, *10*, 97–105. [\[CrossRef\]](#)
139. Angiolini, C.; Viciani, D.; Bonari, G.; Zoccola, A.; Bottacci, A.; Ciampelli, P.; Gonnelli, V.; Lastrucci, L. Environmental drivers of plant assemblages: Are there differences between palustrine and lacustrine wetlands? A case study from the northern Apennines (Italy). *Knowl. Manag. Aquat. Ecosyst.* **2019**, *420*, 34. [\[CrossRef\]](#)
140. Hroudová, Z.; Zákravský, P.; Frantík, T. Ecological differentiation of Central European *Bolboschoenus* taxa and their relationship to plant communities. *Folia Geobot.* **1999**, *34*, 77–96. [\[CrossRef\]](#)
141. Hroudová, Z.; Zákravský, P.; Ducháček, M.; Marhold, K. Taxonomy, distribution and ecology of *Bolboschoenus* in Europe. *Ann. Bot. Fenn.* **2007**, *44*, 81–102.
142. Grechushkina, N.A.; Sorokin, A.N.; Golub, V.B. The plant communities with domination of *Phragmites australis* and *Bolboschoenus glaucus* in the territory of Russian coast of the Azov Sea. *Nauchnyye Soobshchestva* **2011**, *20*, 105–115.
143. Prosser, F.; Sarzo, A. Flora e vegetazione di fossi nel settore trentino del fondovalle dell'Adige (Trentino-Italia settentrionale). *Ann. Mus. Civ. Rovereto* **2003**, *18*, 89–144.
144. Celesti-Grapow, L.; Alessandrini, A.; Arrigoni, P.V.; Banfi, E.; Bernardo, L.; Bovio, M.; Brundu, G.; Cagiotti, M.R.; Camarda, I.; Carli, E.; et al. Inventory of the non-native flora of Italy. *Plant Biosyst.* **2009**, *143*, 386–430. [\[CrossRef\]](#)
145. Braun-Blanquet, J.; Gajewski, W.; Wraber, M.; Walas, J. Classe des *Rudereto-Secalinetales*. Groupements messicoles; culturaux et nitrophiles-rudérales du cercle de végétation méditerranéen. *Prodr. Group. Vég. Montp.* **1936**, *3*, 1–37.
146. Braun-Blanquet, J.; Roussine, N.; Nègre, R. *Les Groupements Végétaux de la France Méditerranéenne*; Centre National de la Recherche Scientifique: Montpellier, France, 1952.
147. Douda, J. Salicetum Pentandro-Auritae Passarge 1957. In *Vegetace České Republiky. 4. Lesní a Křovinná Vegetace. Vegetation of the Czech Republic 4. Forest and Shrub Vegetation*; Chytrý, M., Ed.; Academia: Prague, Czech Republic, 2013; pp. 180–182.
148. Tomei, P.E.; Cenni, M. Il Bosco di Chiusi e la Paduletta di Ramone (PT): Note floristiche e vegetazionali. *Quad. Mus. Stor. Nat. Livorno* **1986**, *7*, 55–79.
149. Arrigoni, P.V. Documenti per la carta della vegetazione delle Cerbaie (Toscana settentrionale). *Parlatorea* **1997**, *2*, 39–71.
150. Corsi, R. La Flora e la Vegetazione. In *Lungo le Rotte Migratorie. Progetti di Ricerca Sulla Vegetazione, L'Avifauna e le Specie Aliene*; Venturato, E., Petrini, R., Eds.; Centro Ricerca, Documentazione e Promozione Padule di Fucecchio: Castelmartini, Italy, 2001; pp. 23–120.

151. Franzese, M. *Studio delle Comunità Vegetazionali ed Aggiornamento Cartografico della Vegetazione*; Centro di Ricerca, Documentazione e Promozione del Padule di Fucecchio: Larciano, Italy, 2004; p. 43.
152. Caruel, T. *Prodromo della Flora Toscana*; Le Monnier: Firenze, Italy, 1860–1864; p. 767.
153. Sandri, G.; Fantozzi, P. Contribuzione alla Flora della Val di Nievole. *Nuovo Giorn. Bot. Ital.* **1895**, *2*, 129–180.
154. Baroni, E. *Supplemento Generale al “Prodromo della Flora Toscana di T. Caruel”*; Società Botanica Italiana: Firenze, Italy, 1897; p. 633.
155. Pampanini, R. Gli esponenti più rimarchevoli e più rari della Flora toscana nel censimento dei Monumenti Naturali d’Italia. *Nuovo Giorn. Bot. Ital.* **1925**, *32*, 5–35.
156. Nannizzi, A. Ricerche idrobiologiche sul padule di Fucecchio. II. Fanerogame. *Boll. Pesca Piscicoltura. Idrobiol.* **1938**, *14*, 161–179.
157. Lorenzoni, G.G.; Chiesura Lorenzoni, F. Osservazioni sull’evoluzione della vegetazione del Padule di Fucecchio (Toscana). *Nuovo Giorn. Bot. Ital.* **1968**, *101*, 423–424.
158. Lastrucci, L.; Fiorini, G.; Lunardi, L.; Viciani, D. Herbarium survey on the genus *Azolla* (Salviniaceae) in Italy: Distributive and taxonomic implications. *Plant Biosyst.* **2019**, *153*, 710–719. [\[CrossRef\]](#)
159. Rodríguez, C.F.; Bécares, E.; Fernández-Aláez, M.; Fernández-Aláez, C. Loss of diversity and degradation of wetlands as a result of introducing exotic crayfish. *Biol. Invasions* **2005**, *7*, 75–85. [\[CrossRef\]](#)
160. Prigioni, C.; Balestrieri, A.; Remonti, L. Food habits of the coypu, *Myocastor coypus*, and its impact on aquatic vegetation in a freshwater habitat of NW Italy. *Folia Zool.* **2005**, *54*, 269–277.
161. Zarri, E. C.U.A. (Censimento Uccelli Acquatici e Svernanti) Padule di Fucecchio 2025. *Natura in Toscana*. 2025. Available online: <https://www.naturaintoscana.it/c-u-a-padule-di-fucecchio-2025/> (accessed on 15 May 2025).
162. Löbl, I.; Klausnitzer, B.; Hartmann, M.; Krell, F.T. The silent extinction of species and taxonomists—An appeal to science policymakers and legislators. *Diversity* **2023**, *15*, 1053. [\[CrossRef\]](#)
163. Eckert, I.; Bruneau, A.; Metsger, D.A.; Joly, S.; Dickinson, T.A.; Pollock, L.J. Herbarium collections remain essential in the age of community science. *Nat. Commun.* **2024**, *15*, 7586. [\[CrossRef\]](#) [\[PubMed\]](#)
164. Liu, D. World view. *Nature* **2024**, *633*, 741. Available online: <https://www.nature.com/articles/d41586-024-03072-3> (accessed on 7 November 2024). [\[CrossRef\]](#)

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

Table S1: General Legend of Table 1 and Tables S2-S12

In the headings of Table 1 and Tables S1-S11 the cover-abundance values of the species are given according to the Braun-Blanquet scale and correspond to the following ranges of cover:

Braun-Blanquet scale value Range of cover in the relevé

5	76-100%
4	51-75%
3	26-50%
2	6-25%
1	1-5%
+	<1%, several individuals
r	<0.5%, 1-2 small individuals

Table S2. Aquatic plant communities

Relevé number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16				
Area m ²	2	8	8	10	10	2	2	1	2	1	1	1	2	1	1	1				
Total cover (%)	100	100	100	85	95	70	60	100	100	75	60	60	85	95	100	60				
Latitude ° N	43,78162	43,78583	43,78583	43,78162	43,78162	43,82	43,82	43,85208	43,78995	43,80882	43,80847	43,80847	43,78107	43,78107	43,84561	43,84561				
Longitude ° E	10,81045	10,80778	10,80778	10,81045	10,81045	10,77778	10,77778	10,76774	10,79974	10,82106	10,82021	10,82021	10,81163	10,81163	10,79649	10,79649				
<i>Lemnetum minuto-gibbae</i>																				
Lemna minuta Kunth	5																			
<i>Persicarietum amphibiae</i>																				
Persicaria amphibia (L.) Delarbre aquatic ecophene	4				5	5	5	1												
<i>Potamo pectinati-Myriophylletum spicati</i>																				
Myriophyllum spicatum L.	4						4													
<i>Potamogetonetum denso-nodosi</i>																				
Potamogeton nodosus Poir.	5								5											
<i>Callitriche platycarpa community</i>																				
Callitriche platycarpa Kütz.											4									
<i>Callitriche stagnalis communities</i>																				
Callitriche stagnalis Scop.												4	4							
<i>Callitriche brutia communities</i>																				
Callitriche brutia Petagna subsp. brutia				+										5	5					
<i>Potamo crispi-Ranunculetum trichophylli</i>																				
Ranunculus trichophyllus Chaix															5	3				
<i>Other species</i>																				
Paspalum distichum L.								+		+										
Eleocharis palustris (L.) Roem. & Schult. subsp. palustris													1	1	r	2	2			
Echinochloa crus-galli (L.) P.Beauv. subsp. crus-galli	3	1															+			
Bolboschoenus glaucus (Lam.) S.G.Sm.	1	+																		
Bolboschoenus maritimus (L.) Palla s.l.															1	1				
Lythrum salicaria L.													2	+						
Phragmites australis (Cav.) Trin. ex Steud.									+	+										
Persicaria lapathifolia (L.) Delarbre										1										
Ranunculus repens L.											1									
Sparganium erectum L.										1										
Ranunculus flammula L.												+								
Alisma plantago-aquatica L.														+						
Convolvulus sepium L.																+				
Gratiola officinalis L.											+									
Juncus bulbosus L.											+									

Table S3. Palustrine plant communities of *Echinochloo cruris-galli*-*Sporoboletum schoenoidis* association

[illegible]

Table S4. Palustrine plant communities of *Littorellion uniflorae* alliance

Relevé number	1	2	3	4
Relevé number in cluster dendrogram	117	118	119	120
Area m ²	8	4	4	4
Total cover (%)	75	65	70	80
Latitude ° N	43.80882	43.80876	43.80998	43.80998
Longitude ° E	10.82106	10.82085	10.82573	10.82573
<i>Ranunculo flammulae-Juncetum bulbosi</i>				
Ranunculus flammula L.	1	4		
Juncus bulbosus L.	2	2	4	3
<i>Juncus bulbosus - Gratiola officinalis comm.</i>				
Gratiola officinalis L.	1		2	3
<i>Other species of Littorellion and upper units</i>				
Callitriche platycarpa Kütz.	2			
<i>Other species</i>				
Ajuga reptans L.			+	2
Leucjum aestivum L. subsp. aestivum	1	+		
Lycopus europaeus L.	1		1	
Alisma lanceolatum With.			+	
Alisma plantago-aquatica L.	+			
Carex acutiformis Ehrh.	2			
Carex spicata Huds. subsp. spicata				2
Carex vesicaria L.	2			
Leucanthemum sp.			+	
Myosotis scorpioides L. / M. laxa Lehm. subsp. cespitosa (Schultz) Hyl. ex Nordh.				+
Potentilla reptans L.			+	
Ranunculus repens L.	2			
Rumex conglomeratus Murray	+			
Sphagnum sp.		1		
Symphyotrichum squamatum (Spreng.) G.L.Nesom			+	

Table S5. Palustrine plant communities of *Bidenton tripartitae* alliance

Relevé number	1	2	3	4	5	6	7	8	9
Relevé number in cluster dendrogram	129	114	110	113	125	123	124	90	91
Area m ²	20	10	4	4	2	2	1	4	4
Total cover (%)	100	100	100	100	95	90	85	70	75
Latitude ° N	43,78923	43,78014	43,78101	43,78014	43,78973	43,79054	43,79069	43,82516	43,82589
Longitude ° E	10,79813	10,80701	10,80775	10,80701	10,79782	10,80131	10,80168	10,79771	10,79754
<i>Bidens frondosa</i> comm.									
<i>Bidens frondosa</i> L.	5	4	5	5					
<i>Symphytotrichum squamatum</i> (Spreng.) G.L.Nesom	+	3	2	1					
<i>Convolvulus sepium</i> L.	2	+	2	4					
<i>Veronica peregrina</i> comm.									
<i>Veronica peregrina</i> L.					4	5	5		
<i>Portulaca oleracea</i> comm.									
<i>Portulaca oleracea</i> L.								4	4
<i>Sporobolus schoenoides</i> (L.) P.M.Peterson								+	1
<i>Other species of Bidenton and upper units</i>									
<i>Persicaria lapathifolia</i> (L.) Delarbre	+		+		+	2			
<i>Amaranthus blitum</i> var. <i>emarginatus</i> (Moq. ex Uline & W.L.Bray) Lambinon				+				+	
<i>Echinochloa crus-galli</i> (L.) P.Beauv.		+						1	
<i>Bidens connata</i> Muhl. ex Willd.	2								
<i>Oxybasis chenopodioides</i> (L.) S.Fuentes, Uotila & Borsch								1	
<i>Xanthium orientale</i> L.		1							
<i>Dysphania ambrosioides</i> (L.) Mosyakin & Clemants				+					
<i>Eclipta prostrata</i> (L.) L.				+					
<i>Other species</i>									
<i>Phragmites australis</i> (Cav.) Trin. ex Steud.		1	1	1		2	1		
<i>Amorpha fruticosa</i> L.		+	+			+			
<i>Lycopus europaeus</i> L.	1	+							
<i>Lythrum salicaria</i> L.					+	+			
<i>Phalaris arundinacea</i> L.	+						1		
<i>Plantago major</i> L.		+			1				
<i>Persicaria amphibia</i> (L.) Delarbre terrestrial ecophene							1		
<i>Ranunculus repens</i> L.					1				
<i>Rorippa prostrata</i> (J.P.Bergeret) Schinz & Thell.						1			
<i>Rumex conglomeratus</i> Murray	1								
<i>Agrostis stolonifera</i> L.	+								
<i>Anthemis</i> sp.					+				
<i>Bolboschoenus maritimus</i> (L.) Palla s.l. (probably <i>B. laticarpus</i> or <i>B. glaucus</i>)	+								
<i>Cardamine apennina</i> Lihová & Marhold					+				
<i>Veronica anagallis-aquatica</i> L. subsp. <i>anagallis-aquatica</i>					+				

Table S6. Palustrine plant communities of *Chenopodion rubri* alliance

Relevé number	1	2	3	4	5	6	7	8
Relevé number in cluster dendrogram	136	45	46	3	4	112	77	78
Area m ²	10	40	40	8	8	20	20	20
Total cover (%)	100	90	100	100	100	100	100	100
Latitude ° N	43.80137	43.79357	43.79357	43.78583	43.78583	43.78014	43.815	43.815
Longitude ° E	10.78718	10.79662	10.79662	10.80778	10.80778	10.80701	10.80917	10.80917

Polygono lapathifolii-Xanthietum italici

Xanthium orientale L.

Persicaria lapathifolia (L.) Delarbre

5	5	5	1	+	1	2
1	2	3	4	+		
		3	1	5	5	5

Echinochloo-Polygonetum lapathifolii

Echinochloa crus-galli (L.) P.Beauv.

+ 2

Other species of Chenopodion rubri and upper units

Amaranthus blitum var. emarginatus (Moq. ex Uline & W.L.Bray) Lambinon

r + 3 1 + 1

Portulaca oleracea L.

1 1 + +

Cyperus odoratus L.

1 +

Abutilon theophrasti Medik.

1

Amaranthus retroflexus L.

+ +

Bidens frondosa L.

1

Bidens tripartita L.

1

Eclipta prostrata (L.) L.

+

Other species

Lythrum salicaria L.

+ + 1 1 +

Bolboschoenus glaucus (Lam.) S.G.Sm.

+ +

Convolvulus sepium L.

1 +

Cyperus michelianus (L.) Delile

r +

Stachys palustris L.

3 1

Amorpha fruticosa L.

+

Bolboschoenus laticarpus Marhold, Hroudová, Ducháček & Zákr.

+

Bolboschoenus maritimus (L.) Palla s.l. (probably B. lat +

Chaerophyllum temulum L.

2

Erigeron sp.

+

Epilobium tetragonum L.

+

Ludwigia palustris (L.) Elliott

2

Lycopus europaeus L.

+

Paspalum distichum L.

+

Plantago major L.

+

Populus nigra L.

+

Solanum nigrum L.

+

Sonchus asper (L.) Hill

+

Veronica anagalloides Guss. subsp. anagalloides

+

Table S7. Palustrine plant communities of *Phragmiton australis* alliance

Relevé number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29
Relevé number in cluster dendrogram	84	18	85	14	57	79	86	23	6	42	37	17	103	109	101	105	61	35	56	111	72	59	63	60	68	62	73	74	75
Area m ²	50	20	50	20	5	50	50	40	4	50	10	20	20	20	20	20	10	10	20	4	20	20	20	20	20	10	20	2	20
Total cover (%)	100	100	100	100	100	100	100	100	100	100	100	100	100	60	100	100	100	100	80	100	100	100	100	100	100	100	100	100	100
Latitude ° N	43.82823	43.82	43.82459	43.77389	43.78139	43.815	43.82606	43.81417	43.78583	43.7939	43.78333	43.81528	43.77672	43.79454	43.7861	43.80049	43.77804	43.79972	43.77364	43.78014	43.77861	43.77804	43.77804	43.77804	43.77804	43.77804	43.77804	43.77804	43.77804
Longitude ° E	10.79712	10.77778	10.79151	10.81028	10.80583	10.80917	10.79895	10.79194	10.80778	10.79446	10.82333	10.76806	10.81172	10.81877	10.8127	10.78918	10.80863	10.82028	10.81634	10.80701	10.80833	10.80863	10.80891	10.80863	10.80974	10.80891	10.80835	10.80669	10.80669
<i>Phragmitetum australis</i> Phragmites australis (Cav.) Trin. ex Steud.	4	5	4	5	5	5	5	5	5	5			+	1															
<i>Typhetum latifoliae</i> Typha latifolia L.											5	5	5																
<i>Typhetum angustifoliae</i> Typha angustifolia L.														4															
<i>Schoenoplectetum lacustris</i> Schoenoplectus lacustris (L.) Palla																5	5	5	5	5									
<i>Phalarido arundinaceae-Bolboschoenetum laticarpi</i> Bolboschoenus laticarpus Marhold, Hroudová, Ducháček & Zákr.											1						2			4	3	5	5	5	5	4	4	5	5
<i>Other species of Phragmiton australis and upper units</i> Lythrum salicaria L. Bolboschoenus glaucus (Lam.) S.G.Sm. Phalaris arundinacea L. Carex riparia Curtis Cyperus longus L. Bolboschoenus maritimus (L.) Palla s.l. (probably B. laticarpus or B. glaucus) Lysimachia vulgaris L. Oenanthe aquatica (L.) Poir. Sparganium erectum L. s.l. Teucrium scordium L.		+	1	1		2	+		1	+		+	+	+		+					+	+	1	+	+				

Other species

[illegible]

[illegible]

[illegible]

Table S9. Palustrine plant communities of *Glycerio-Sparganietum neglecti* association

Relevé number	1	2	3
Relevé number in cluster dendrogram	39	34	38
Area m ²	10	10	10
Total cover (%)	100	100	100
Latitude ° N	43.7942	43.7994	43.7833
Longitude ° E	10.8267	10.8192	10.8247
<i>Glycerio-Sparganietum neglecti</i>			
Sparganium erectum L. s.l.	5	5	5
<i>Other species of Glycerio-Sparganion and upper units</i>			
Lythrum salicaria L.	1	+	+
Phalaris arundinacea L.	1		
Bolboschoenus laticarpus Marhold, Hroudová, Ducháček & Zákr.			+
Carex otrubae Podp.	+		
<i>Other species</i>			
Convolvulus sepium L.	2	+	
Agrostis stolonifera L.	1	+	
Rumex conglomeratus Murray	1		
Rubus ulmifolius Schott			+

[illegible]

Glyceria maxima (Hartm.) Holmb. subsp. maxima
 Stachys palustris L.

1

1

Other species

Paspalum distichum L.

+

3

1

2

+

1

Convolvulus sepium L.

+

3

+

+

2

Xanthium orientale L.

+

2

1

+

1

Agrostis stolonifera L.

1

2

Cyperus eragrostis Lam.

+

+

Persicaria amphibia (L.) Delarbre aquatic ecophene

+

1

Poa trivialis L.

1

1

Amorpha fruticosa L.

1

Elymus repens (L.) Gould subsp. repens

1

Persicaria amphibia (L.) Delarbre terrestrial ecophene

+

Plantago major L.

+

Ranunculus trichophyllus Chaix

1

Rumex conglomeratus Murray

+

Table S11. Palustrine plant communities of *Polypogono viridis-Paspaletum distichi* association

Relevé number	1	2	3	4	5	6	7	8	9	10
Relevé number in cluster dendrogram	106	107	92	87	137	5	1	2	16	52
Area m ²	4	4	4	4	4	4	4	4	4	10
Total cover (%)	100	100	100	100	80	100	100	100	100	100
Latitude ° N	43.78553	43.78553	43.82589	43.82475	43.85208	43.78583	43.78583	43.78583	43.78583	43.76992
Longitude ° E	10.81947	10.81947	10.79754	10.7978	10.76774	10.80778	10.80778	10.80778	10.80778	10.81013
<i>Polypogono viridis-Paspaletum distichi</i>										
<i>Paspalum distichum</i> L.	5	5	5	5	5	5	5	5	5	3
<i>Other species of Polypogono viridis-Paspalion distichi and upper units</i>										
<i>Trifolium repens</i> L.		+								
<i>Other species</i>										
<i>Echinochloa crus-galli</i> (L.) P.Beauv.			1	+	+					
<i>Bolboschoenus glaucus</i> (Lam.) S.G.Sm.							+	1		
<i>Cyperus rotundus</i> L.	1	2								
<i>Digitaria sanguinalis</i> (L.) Scop.	1	1								
<i>Erigeron canadensis</i> L.	1	+								
<i>Hypochaeris glabra</i> L.	+	1								
<i>Oxalis</i> sp.	+	+								
<i>Persicaria lapathifolia</i> (L.) Delarbre				+	+					
<i>Setaria italica</i> (L.) P.Beauv. subsp. <i>viridis</i> (L.) Thell.	+	+								
<i>Amaranthus blitum</i> var. <i>emarginatus</i> (Moq. ex Uline & W.L.Bray) Lambinon						1				
<i>Atriplex prostrata</i> Boucher ex DC.										+
<i>Bidens frondosa</i> L.										1
<i>Convolvulus sepium</i> L.										1
<i>Cyperus eragrostis</i> Lam.										3
<i>Daucus carota</i> L.	+									
<i>Eclipta prostrata</i> (L.) L.										+
<i>Erigeron bonariensis</i> L.		+								
<i>Persicaria amphibia</i> (L.) Delarbre terrestrial ecophene							1			
<i>Phragmites australis</i> (Cav.) Trin. ex Steud.										1
<i>Portulaca oleracea</i> L.						1				
<i>Senecio vulgaris</i> L.	+									
<i>Urtica dioica</i> L.					+					
<i>Verbena officinalis</i> L.	1									
<i>Veronica persica</i> Poir.	+									
<i>Xanthium orientale</i> L.						+				

Table S12. Palustrine plant communities of *Frangulo alni-Salicetum cinereae* association

Relevé number	1	2	3	4
Relevé number in cluster dendrogram	48	49	99	138
Area m ²	100	100	50	50
Total cover (%)	100	100	100	100
Latitude ° N	43.7924	43.7924	43.8201	43.78995
Longitude ° E	10.7954	10.7954	10.805	10.79974

Frangulo alni-Salicetum cinereae

Salix cinerea L.	2	4	5	5
Amorpha fruticosa L.	4	2	1	

Other species

Urtica dioica L.			2	
Chaerophyllum sp.				1
Aster sp.			+	
Atriplex prostrata Boucher ex DC.			+	
Convolvulus sepium L.				+
Lycopus europaeus L.			+	
Phalaris arundinacea L.			+	
Phragmites australis (Cav.) Trin. ex Steud.				+