

Effectiveness of low-carbohydrate diets for long-term weight loss in obese individuals: A meta-analysis of randomized controlled trials

Giovanni Antonio Silverii MD¹  | Claudia Cosentino MD² |
 Federica Santagiuliana RD³ | Francesco Rotella MD⁴ | Federica Benvenuti MD⁴ |
 Edoardo Mannucci MD¹  | Barbara Cresci MD³ 

¹Biomedical Sciences Department, Florence University, Florence

²Diabetology Unit, Pistoia San Jacopo Hospital, Pistoia, Italy

³Diabetology, Careggi Hospital, Florence, Italy

⁴Psychiatry, Careggi Hospital, Florence, Italy

Correspondence

Giovanni Antonio Silverii, MD, Experimental and Clinical Biomedical Sciences “Mario Serio” Department, University of Florence, Largo Brambilla 3, 50134, Florence, Italy.
 Email: giovanniantonio.silverii@unifi.it

Funding information

This research was performed as a part of the institutional activity of the unit, with no specific funding. Open Access Funding provided by Università degli Studi di Firenze within the CRUI-CARE Agreement.

Abstract

Aim: To assess whether low-carbohydrate (LC) diets are associated with differences in weight loss and well-being in people with obesity, and their cardiovascular and renal safety.

Materials and Methods: A meta-analysis of randomized controlled trials longer than 3 months, retrieved through an extensive search on MedLine and Embase databases, comparing weight loss with LC and control diets in people with body mass index (BMI) greater than 30 kg/m², was conducted.

Results: We retrieved 25 trials. Compared with controls, LC diets were associated with significant reduction of body weight at 3-4 (MD -2.59 [-3.93, -1.25] kg) and 6-8 months (MD -2.64 [-4.32, -0.95]), but no difference at 10-14 and 18-30 months, and significantly greater BMI reduction at 3-4 months (-1.66 [-2.70, -0.61] kg/m²), but not at other time points. Because only four trials reported data on renal function and psychological variables, renal safety and impact on well-being could not be assessed. Differences in fasting plasma glucose at any time point were not statistically significant. No significant differences in total or LDL cholesterol or blood pressure were found in the long term, whereas a long-term reduction of triglycerides (23.26 [-45.53, -0.98] mg/dl at 18-30 months), and increase of HDL cholesterol (MD 4.94 [0.30, 9.57] mg/dl at 18-30 months), were observed.

Conclusion: LC diets are associated with greater short-term weight loss than non-carbohydrate-restricted diets and a longer term favourable effect on cardiovascular risk factors. Further evidence on long-term efficacy and renal safety is needed before LC diets can be recommended as the preferred diets in obese people.

KEYWORDS

low-carbohydrate diets, meta-analysis, obesity, weight loss

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Diabetes, Obesity and Metabolism* published by John Wiley & Sons Ltd.

1 | INTRODUCTION

Obesity is associated with an increased risk of type 2 diabetes, cardiovascular disease, malignancies, and mortality.¹⁻⁴ The management of obesity is aimed at improving overall health, rather than merely at weight loss.⁵ Although intentional weight loss with some interventions has been associated with reduced mortality,⁶ no specific treatment for obesity has been proven to increase life expectancy in clinical trials, and unintentional weight loss may even be associated with increased mortality.⁷ Epidemiological studies reporting the effects of weight loss (including both unintentional and intentional) are inconclusive.⁸ Nevertheless, weight loss is expected to reduce the burden of obesity-associated morbidity and mortality⁹; therefore, a reasonable and slowly progressive weight reduction is advised, usually accounting for 5%-10% of initial weight.¹⁰ However, more ambitious targets may be advisable in those who are at higher risk of cardiovascular and metabolic complications.¹¹ Dietary modifications, together with an increase in physical activity and reduction of inactivity, are the first-line therapy for weight loss, encompassing modifications in caloric intake, eating habits, and nutrient composition.¹² The ideal diet is defined as being safe, healthy, nutritionally adequate, culturally acceptable, and economically affordable, and it should ensure long-term compliance and effectiveness.¹³ Most guidelines suggest a 600 kcal daily energy deficit and a reduction in fat intake.¹⁴⁻¹⁶ In particular, the so-called 'Mediterranean Diet', which consists in reducing the intake of fat, especially if saturated, and refined sugar, while increasing the consumption of vegetables and raw carbohydrates, is often considered the preferred approach, because long-term epidemiological data show lower overall mortality and morbidity in those who adhere to such a regimen.^{17,18}

More recently, some scientific societies have stressed the feasibility and effectiveness of low-carbohydrate (LC) diets for the treatment of obesity.^{5,19} The definition of LC diets is heterogeneous, with different degrees of carbohydrate restriction.²⁰⁻²³ Modern LC diets usually rely on proteins, rather than fats, to ensure energy intake, aiming at preserving muscle mass and limiting the negative impact of lipid metabolism, often implying the use of expensive protein supplements.²⁴ Because a greater protein dietary intake could be associated with a faster decline of glomerular filtration in the long term,²⁵ although this issue is controversial,²⁶ a dietary protein overload in obese individuals who are already at risk for renal diseases may raise concerns for renal safety.

The evidence on the effects of LC diets in the treatment of obesity has been summarized in some meta-analyses.²⁷⁻³³ However, those meta-analyses often include both obese and non-obese cases,^{27,29,31-33} and in some cases include observational studies together with clinical trials.²⁸ In addition, the results of available meta-analyses are usually driven by short- and very short-term trials, without separate analyses for longer term studies,³⁰ and they provide no specific information on renal safety in the longer term.^{27,28,30,31}

The primary aim of our meta-analysis is to assess the specific effect of carbohydrate restriction in the treatment of obesity; thus we explored differences between carbohydrate-restricted diets and non-

carbohydrate-restricted diets concerning weight loss and renal safety in obese individuals. The secondary aim is the exploration of possible effects of specific carbohydrate restriction on blood pressure, lipid profile, and blood glucose, together with its effects on the perceived quality of life and adherence to the prescribed diet.

2 | METHODS

This meta-analysis is reported following the criteria of the PRISMA statement.³⁴ The review protocol was submitted for registration to the PROSPERO website (#268453; <https://www.crd.york.ac.uk/PROSPERO/>).

2.1 | Search strategy and selection criteria

A systematic search on PubMed, Cochrane, clinicaltrials.gov and Embase databases was performed, collecting all randomized clinical trials written in English and performed on humans up to 1 November 2021. The full search string is reported in Table S1. Further studies were manually searched in references from retrieved papers.

Studies were included if they fulfilled the following criteria: randomized controlled trials; comparison of a LC diet with a non-carbohydrate-restricted diet (see below for definitions); apart from diet composition, no difference in treatment protocol between the two arms; duration of the trial of at least 12 weeks; end-of-study body weight, or body mass index (BMI), reported for both treatment arms; and studies enrolling only individuals with a BMI more than 30 kg/m², or separate analyses of subgroups of cases with a BMI more than 30 kg/m² in trials with wider inclusion criteria.

Studies were eligible for inclusion if they combined a dietary intervention with another non-pharmacological intervention type (e.g. prescribed exercise/physical activity, cognitive-behaviour therapy, psychological support), if this was equivalent across dietary intervention arms.

The diets were defined as follows, according to available nutrition guidelines^{35,36}:

1. non-carbohydrate-restricted diets: 45%-60% of total calories from carbohydrates;
2. mild LC diets: 26%-45% of total calories from carbohydrates; and
3. very LC diets: less than 26% of total calories from carbohydrates and/or less than 130 g of carbohydrates daily.

2.2 | Endpoints

The principal endpoints were the differences in mean BMI expressed as kg/m² between all LC and balanced diets after 3-4, 6-8, 10-14 and 18-30 months, and the difference in mean body weight between all LC and balanced carbohydrate diets after 3-4, 6-8, 10-14 and 18-30 months.

The secondary endpoints were the difference in mean, total, HDL and LDL cholesterol, and systolic blood pressure between all LC diets

and balanced carbohydrate diets after 3-4, 6-8, 10-14 and 18-30 months, and the difference in quality of life and adherence to prescribed diet between all LC diets and balanced carbohydrate diets after 3-4, 6-8, 10-14 and 18-30 months, and at the endpoint.

2.3 | Data collection

Titles and abstracts were screened independently by two authors, and potentially relevant articles were retrieved in full text format. For all published trials, results reported in published papers and supplements were used as the primary source of information; when the required information on protocol or outcomes was not available in the main publication, secondary publications were used for retrieval of the missing information; whenever necessary, an attempt at retrieval of the missing information was performed consulting the clinicaltrials.gov registry. The identification of relevant abstracts, the selection of studies, and data extraction were performed independently by two of the authors (GAS and BC), and conflicts were resolved by a third investigator (EM). The risk of bias was assessed using the features proposed by the Cochrane Collaboration³⁷ by two of the authors (FB and CC), and conflicts were resolved through discussion with a third investigator (EM); reporting bias was assessed for each main outcome.

2.4 | Statistical analyses

Mantel-Haenszel odds ratio (MH-OR) with 95% confidence interval (95% CI) and between-group difference in means (weighted mean difference [MD]) with 95% CI were calculated, on an intention-to-treat basis, for dichotomic and continuous outcomes, respectively, using the Wald-type confidence interval methods calculator. Heterogeneity was assessed by using I^2 statistics, using DerSimonian and Laird variance estimator. A random-effects model was applied as the primary analysis. Funnel plots for HbA1c levels were examined to estimate possible publication/disclosure bias. All analyses were performed using Review Manager 5.3.5 (The Cochrane Collaboration, 2014). The GRADE methodology³⁷ was used to assess the quality of the body of retrieved evidence, using the GRADE pro-GDT software (GRADEpro Guideline Development Tool; McMaster University, 2015). A sensitivity post hoc analysis was performed comparing weight loss in different treatment arms at 3-4 and 10-14 months, selecting only trials for which both the 3-4 and the 10-14 month follow-up results were available. A post hoc subgroup analysis was performed, dividing trials in which the protein content in the intervention group was below or above 30% of total daily calories.

3 | RESULTS

3.1 | Trial characteristics

Figure S1 reports the trial flow summary. Of the 7850 items, after removing duplicates, 886 were selected for retrieval of

the full text. Of those, 25 trials, overall enrolling 1233 cases on LC diets and 1209 cases on balanced diets, fulfilled the inclusion criteria.

The main characteristics of included trials are reported in Table 1. Out of 26 studies, 19 excluded individuals with kidney disease, and 13 excluded those with previous cardiovascular disease. Nine studies excluded individuals with diabetes, whereas only five included those affected by diabetes, while four included cases with or without diabetes; seven studies did not provide information on this issue. The risk of bias is reported in Figures S2 and S3.

3.2 | Weight loss

All the included trials, except for two,^{39,40} reported body weight or BMI data only at some time points; the analysis for body weight was therefore performed on 20, 12, 10 and three trials at 3-4, 6-8, 10-14 and 18-30 months, respectively (Figure 1). LC diets were associated with a significantly higher reduction of body weight at 3-4 (MD -2.59 [$-3.93, -1.25$] kg, $P = .0001$) and 6-8 months (MD -2.64 [$-4.32, -0.95$] kg, $P = .002$) with respect to balanced diets, with no heterogeneity ($I^2 = 0$). The difference in reduction of body weight between the two arms was no longer significant at 10-14 months (-2.30 [$-5.00, +0.41$] kg, $I^2 = 19$) and it totally disappeared at 18-30 months (MD $+0.89$ [$-2.32, +4.10$] kg, $I^2 = 0$). No publication bias was found (Figure S4).

We performed an additional analysis including only those studies providing data on body weight both at 3-4 and 10-14 months. Seven studies were available. LC diets were associated with a significantly greater reduction of body weight at 3-4 months (MD -2.72 [$-4.64, -0.80$], $P = .005$, $I^2 = 0$), which was no longer significant at 10-14 months (MD -2.50 [$-5.28, 0.27$], $I^2 = 8$) (Figure S6).

We also performed a post hoc subgroup analysis to explore the effect of protein dietary content on weight loss at any time point. No difference in weight loss was observed between trials in which the protein content of the intervention group was below or above 30% of total calorie intake (Figure S7).

Data on BMI were available for five, eight, six and one trial at 3-4, 6-8, 10-14 and 18-30 months, respectively. LC diets were associated with a significant reduction of BMI at 3-4 months (-1.66 [$-2.70, -0.61$] kg/m², $P = .002$), but not at other time points (Figure 2). No publication bias was found (Figure S5).

No significant difference was observed in BMI between trials in which the protein content of the intervention group was below or above 30% of total calorie intake (Figure S8).

3.3 | Renal function


Only two studies^{41,42} reported serum creatinine at endpoint, showing lower values in the LC values (MD -0.12 [$-0.17, -0.07$] mg/dl, $I^2 = 0\%$), which was already present at baseline (MD -0.10 [$-0.15,$

TABLE 1 Baseline characteristics of the included studies

Study	D	CKD Excl	CVD Excl	DM (%)	F (%)	Country	N		Age (y)	CHO % (g)			Fat % (g)			Prot % (g)			Energy (kcal)			BMI		
							I	C		I	C	I	C	I	C	I	C	I	C	I	C	I	C	
Bales 2017 ⁵⁵	6	No	Yes	NR	100	United States	51	29	60.0	40	55	30	30	30	15	15	(-500)	(-500)	37.5	38.3				
Bazzano 2014 ⁴¹	12	Yes	Yes	0	66	United States	75	73	46.9	(40)	55	-	30	-	-	-	-	-	35.4	35.4				
Brehm 2009 ⁵⁶	12	Yes	Yes	100	58	United States	26	27	56.5	(20)	55	-	30	-	15	-	-	-	36	36				
Cornier 2006 ⁵⁷	3	Yes	Yes	NR	100	United States	14	13	53.1	45	60	40	20	34	17	(-400)	-	-	34	35				
Dalle Grave 2013 ⁴²	12	No	No	NR	58	Italy	43	45	46.7	46	63	20	-	34	-	1350	-	-	45.8	45.4				
Daly 2006 ⁵⁷	3	Yes	No	100	52	United Kingdom	51	51	58.9	(70)	55	-	-	-	-	-	-	-	35	37				
De Luis 2009 ⁵⁹	3	No	Yes	NR	72	Spain	52	66	45.6	38	52	36	27	26	20	1507	1500	35.2	35.9					
De Luis 2015 ⁶⁰	9	Yes	Yes	NR	71	Spain	110	101	50.5	33	53	33	27	34	20	1050	1093	35.5	36.8					
Ebbeling, 2007 ²⁰	12	Yes	Yes	0	80	United States	36	37	27.5	40	55	35	25	25	25	-	-	37.2	36.6					
Foster 2003 ⁶¹	12	Yes	Yes	0	68	United States	33	30	44.1	(50)	60	-	25	-	15	-	1500	33.9	34.4					
Foster 2010 ³⁹	24	No	No	0	68	United States	153	154	45.6	(50)	55	-	30	-	15	-	1500	36.1	36.1					
Goday 2016 ⁴³	4	Yes	No	100	65	Spain	45	44	54.6	(50)	55	-	30	-	15	700	-	33	33					
Goldstein 2011 ²¹	12	No	No	100	48	Israel	26	26	56	(50)	60	-	20	-	20	-	1350	33	33					
Iqbal 2010 ⁴⁰	24	Yes	No	100	10	United States	70	74	60	32	50	-	30	-	-	(-500)	38	37						
Kerksick 2010 ^{61,62}	3	Yes	Yes	0	100	United States	43	65	34.9	20	55	30	15	50	30	1200	1200	36	35					
Kitabchi 2013 ⁶²	6	Yes	No	0	100	United States	14	18	35.7	40	55	30	30	30	15	1800	1800	41.3	37					
Morris 2020 ⁶³	3	Yes	Yes	100	55	United Kingdom	21	12	67	25	55	-	-	(60)	-	900	-	34.8	36.4					
Perticone 2019 ²²	12	Yes	Yes	NR	43	Italy	28	28	46.9	20	57	55	27	25	13	1000	(-500)	40.5	38.8					
Porter Starr 2016 ^{64,65}	6	Yes	No	NR	79	United States	41	26	68.2	40	55	30	30	30	15	(-500)	(-500)	36.4	37.2					
Racette 1995 ⁶⁵	3	Yes	Yes	0	100	United States	13	10	39	25	60	50	15	25	15	-	-	33.2	34.5					
Samaha 2003 ^{66,67}	6	Yes	No	40	18	United States	64	68	53	(30)	-	-	30	-	-	(-500)	42.9	42.9						
Stentz 2016 ⁶⁷	6	Yes	No	0	79	United States	12	13	42.1	40	55	30	30	30	15	1800	-	40.5	37.4					
Tonstad 2014 ⁶⁸	4	No	No	20	77	United States	82	91	48.4	(120)	55	-	-	-	-	-	-	36.6	36.3					
Yancy 2004 ³⁸	6	Yes	Yes	0	79	United States	59	60	44.9	(120)	-	-	30	-	-	(-750)	34.6	34						
Yancy 2015 ⁴⁴	11	Yes	No	23	27	United States	53	49	55	(120)	-	-	30	-	-	(-500)	36	36						

Note: Carbohydrates, fats and proteins are expressed in % of daily total intake, or grams when in brackets. Energy is expressed in kcal; energy is in brackets when expressed as the difference between daily recommended intake and prescribed calorie intake.

Abbreviations: BMI, body mass index; C, control; CHO, carbohydrate; CKD, chronic kidney disease; CVD, cardiovascular disease; D, duration (expressed as months); DM, diabetes mellitus; Excl, excluded; F, females; I, intervention; N, number; NR, not reported; Prot, proteins.



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting for weight (reporting bias)
- (G) Selective reporting for renal function (reporting bias)
- (H) Other bias

FIGURE 1 Difference in body weight (expressed as kg) at A, 3-4, B, 6-8, C, 10-14, and D, 18-30 months between low-carbohydrate (carb) and balanced carb diets. Risk of bias legend: A = random sequence generation (selection bias); B = allocation concealment (selection bias); C = blinding of participants and personnel (performance bias); D = blinding of outcome assessment (detection bias); E = incomplete outcome data (attrition bias); F = selective reporting for weight (reporting bias); G = selective reporting for renal function (reporting bias); H = other bias. “+” = low risk; “?” = unknown risk; “-” = high risk. CI, confidence interval; MD, mean difference; N, number




FIGURE 2 Difference in body mass index (BMI, expressed as kg/m²) at A, 3-4, B, 6-8, C, 10-14, and D, 18-30 months between low-carbohydrate (carb) and balanced carb diets. Risk of bias legend: A = random sequence generation (selection bias); B = allocation concealment (selection bias); C = blinding of participants and personnel (performance bias); D = blinding of outcome assessment (detection bias); E = incomplete outcome data (attrition bias); F = selective reporting for weight (reporting bias); G = selective reporting for renal function (reporting bias); H = other bias. "+" = low risk; "?" = unknown risk; "-" = high risk. CI, confidence interval; MD, mean difference; N, number

-0.05] mg/dl). One study²² reported the Chronic Kidney Disease Epidemiology Collaboration-calculated estimated glomerular filtration

rate (eGFR) (MD +4.00 [-2.39, 10.39] ml/min), whereas another study⁴³ reported no significant difference from baseline in both

TABLE 2 Psychological variables

Study	Domain	Scale	Subscale	Baseline		3-4 mo		6-8 mo		10-14 mo		18-30 mo	
				I	C	I	C	I	C	I	C	I	C
Dalle Grave 2013 ⁴²	Anxiety	BAI				7.5	6.4	9.2	7.6	10	5.9		
	Depression	BDI			9	11	8	9	10	9			
	Body uneasiness	BUT			56	53	35	38	39	35			
	Binge eating	BES			7.5	8.1	5.6	5.5	6.9	5.8			
Foster 2010 ³⁹	Craving	FCI	Sweets	2.6	2.5	2.3	2.2	2.3	2.2	2.4	2.5	2.5	2.2
			High-fats	1.9	1.9	2.0	1.8	2.1	2.0	2.1	1.8	2.0	1.6
			Carb/starch	2.1	2.2	1.9	2.1	1.8	2.1	1.9	2.3	1.9	2.1
			Fast-food fats	2.6	2.6	2.3	2.4	2.4	2.5	2.4	2.6	2.5	2.3
	Food preference	FPQ	Complex carbs	5.5	5.5	5.0	5.2	4.9	4.5	4.7	5.1	4.5	5.1
			Sugar	6.0	5.9	5.8	5.7	5.6	5.7	5.6	5.6	5.4	5.6
			Proteins	5.9	5.8	5.8	5.6	5.8	5.3	5.6	5.4	5.6	5.1
	Appetite	Appetite rating change	Hunger			-7	0	-3	0	-1	0	-5	-2
			Bothered by hunger			-3	0	0	3	0	3	0	3
			Eat in reaction to food cues			-12	-4	-6	0	-3	0	-3	-4
			Thoughts about food			-9	-3	-4	-2	-3	0	-3	-3
Morris 2020 ⁶³	Diabetes-related distress	PAID			14.4	20.7	No significant difference between groups						
	Belief	BS			4.3	4.5	4.7	4.6					
	Motivation	MS			3.7	4.1	4.6	4.1					
Yancy 2015 ⁴⁴	Quality of life	IWQOL-lite	Total			72	No significant difference between groups						

Abbreviations: BAI, Beck anxiety inventory; BDI, Beck depression inventory; BES, binge-eating scale; BS, belief score; BUT, body uneasiness test; C, control; carb, carbohydrate; FCI, food craving index; I, intervention; IWQOL-lite, impact of weight on quality of life-lite questionnaire; MS, motivation score; PAID, problem areas in diabetes.

groups by Modification of Diet in Renal Disease-calculated eGFR, without showing any data.

3.4 | Glycaemic control

Differences in fasting plasma glucose between LC diets and control arms were not statistically significant at any time point (Table S2).

3.5 | Cardiovascular risk factors

LC diets were associated with a significant increase of HDL cholesterol at 10-14 (MD 2.38 [0.29, 4.47] mg/dl) and 18-30 months (MD 4.94 [0.30, 9.57] mg/dl), but not at 3-4 and 6-8 months (Table S2), whereas no difference in total or LDL cholesterol was found at any time point (Table S2). A reduction in triglycerides was observed at 3-4, 10-14 and 18-30 months (MD -1.78-20.63 [-35.37, -5.89], -27.09 [-38.29, -15.90] and -23.26 [-45.53, -0.98] mg/dl, respectively), but not at 6 months. No difference was found in blood pressure at

any time point, with the only exception of lower diastolic blood pressure at 3-4 and 6-8 months in LC diets (MD -3.22 [-5.90, -0.53] and -1.78 [-3.10, -0.45] mmHg, respectively; Table S2).

3.6 | Adherence to diet

Retention to studies was 72.3% for LC diets and 70.8% for control diets, with no significant difference between the two groups ($P = .39$). However, only 40 individuals on LC and 31 on control diets reported diet dissatisfaction as a reason for dropout ($P = .20$). Eight cases on the LC diet (vs. none in the control arms) dropped out for safety concerns, such as an increase in LDL or creatinine, or ketosis (MH-OR 3.44 [0.84, 14.02]).

3.7 | Psychological variables

Only four studies reported data on psychological variables; of those, one did not report outcome data for each arm.⁴⁴ Assessment

measures included scales for binge eating, food cravings and appetite, food preferences, anxiety, depression, obesity- and diabetes-related quality of life, beliefs, and motivation (Table 2). Because no single instrument was used in more than one study, no meta-analysis was performed. In all those studies, weight loss was associated with a significant score reduction, all within the normal range, of measures of anxiety, depression, binge eating and body uneasiness, with no differences between LC and control diets. Reported outcomes are summarized in Table 2.

3.8 | GRADE scoring of available evidence

GRADE scoring for principal endpoints is reported in Table S3. The overall quality of evidence was assessed as high for body weight at 3-4, 6-8 and 10-14 months and for BMI at 3-4 months; as moderate for weight at 18-30 months and for BMI at 6-8 and 10-14 months; and as low for BMI at 18-30 months and for renal function at the endpoint.

4 | DISCUSSION

LC diets are associated with a moderately greater weight loss than non-carbohydrate-restricted diets in the short term. This difference seems to disappear in the longer term, although the number of available studies is insufficient to draw a definitive conclusion after longer than 12 months. Consistent results are obtained when restricting the analysis to trials for which both short- and medium-term results are available. This is in line with previous findings in meta-analyses including obese and overweight individuals.³² A previous meta-analysis, which reported a significant weight loss at 12-14 months, also included non-obese overweight cases and only explored very LC diets.²⁹ A more recent pairwise meta-analysis with different trial inclusion criteria, which did not report any comparison in weight loss between LC and low-fat diets, highlighted a negative correlation between actual carbohydrate intake and weight loss at 6 and at 12 months,²⁷ without providing any longer term data. However, such an analytical approach could overestimate the therapeutic effect of prescribed carbohydrate consumption, because cases with a greater adherence to prescriptions of LC diets could be more prone to weight loss per se.

Adherence to prescribed regimens is a major limiting factor of the efficacy of dietary interventions in obesity.⁴⁵ In some trials, actual carbohydrate intake in the LC diet arm could have been different from that prescribed.^{21,40} On the other hand, long-term adherence may be lower when the prescribed regimen is very different from usual (spontaneous) dietary intake⁴⁶; in fact, traditional eating habits in many countries include the consumption of a relevant amount of carbohydrates. This could reduce the effectiveness of LC diets in comparison with balanced diets.

Reduction of carbohydrate intake can be obtained either by increasing the fat or protein content of the diet, or both. The

observed effects on weight loss could therefore depend on carbohydrate restriction or the increase in intake of another nutrient. In a post hoc subgroup analysis, protein intake did not appear to moderate weight loss at any time point; however, the limited number and size of available trials does not allow drawing definitive conclusions on this point, which deserves further specific investigation.

The authors of most available trials appear to agree on a possible issue of the renal safety of LC diets, because impaired renal function is usually among the exclusion criteria. Inexplicably, most of those studies did not report any results on renal function at the end of the study, except for only four trials; although no significant differences between treatment arms were detectable in those trials, these results could have been altered by publication bias or disclosure bias. Currently, the renal safety of LC diets remains unknown.

Another potential concern regarding LC diets is cardiovascular safety because the increase in fat intake could have adverse effects on lipid profile and other risk factors.⁴⁷ Increased ketogenesis determined by extreme carbohydrate restriction could theoretically reduce the risk of cardiovascular disease.⁴⁸ Conversely, observational studies suggest that carbohydrate-rich Mediterranean-style diets are associated with reduced cardiovascular morbidity and mortality,^{17,18} whereas a LC intake has been associated with a higher cardiovascular risk.⁴⁶ The duration and the size of samples enrolled in randomized trials comparing LC and balanced diets in the treatment of obesity is too small for assessing their effects on cardiovascular events. In addition, many of the available trials excluded individuals with established cardiovascular disease, who are at higher risk for cardiovascular events. However, data on cardiovascular risk factors were reassuring, with a reduction in triglycerides and an increase in HDL cholesterol, in line with previous reports.⁴⁹

LC diets did not appear to have an advantage over control diets in the reduction of fasting plasma glucose. This result is in line with that of a previous meta-analysis reporting a transient reduction of HbA1c, followed by a modest deterioration of HbA1c in the longer term when LC diets are applied to individuals with type 2 diabetes.⁵⁰ On the other hand, the large majority of subjects included in the present meta-analysis was not affected by diabetes; the dietary intervention is less probable to produce a relevant effect on fasting plasma glucose when baseline levels are within the normal range.

The improvement of quality of life and psychological well-being is one of the aims of the treatment of obesity.⁵¹ Despite this fact, most available trials did not explore these domains. Reported data show that, not surprisingly, weight loss per se improves psychological status,⁵²⁻⁵⁴ but they fail to highlight any relevant difference between LC and balanced diets. Further studies, enrolling larger samples, are needed to clarify this point.

Several limitations should be considered in the interpretation of the results of this meta-analysis. The definition of LC diets is heterogeneous across studies, with different degrees of carbohydrate restriction; despite this fact, the observed heterogeneity for the principal outcomes was very low. Most trials are comparatively small, limiting the precision of estimates of treatment effect. In addition, most

studies have a short follow-up, limiting the possibility of extending results to longer term treatment. Notably, most long-term trials were performed in the United States; their results could be only partly applicable to different cultural contexts, such as those of Mediterranean countries, where adherence to a LC diet could theoretically be more problematic. Furthermore, many trials show relevant methodological limitations, thus reducing the quality of evidence. For example, allocation and detection bias could have led to an overestimation or underestimation of the efficacy of LC diets in some trials. In addition, the use of medication for obesity or other conditions (such as diabetes, hypertension and hyperlipidaemia) was not considered among the outcomes; differences in medication use could therefore have interfered with the results. On the other hand, this meta-analysis has some strengths: the clear definition of the target population for the dietary intervention (i.e. obese subjects only) increases the reliability of results, which is strengthened by their low heterogeneity.

This systematic review and meta-analysis shows that, in comparison with non-carbohydrate-restricted diets, LC diets are associated with a greater short-term weight loss, with no clear differences in efficacy over the longer term. Data on cardiovascular risk factors are reassuring, while the renal safety of LC diets is undetermined. Further trials are needed to clarify the balance between the benefits and harms of this dietary approach, including more thorough reporting of potential detrimental effects (such as those on renal function) and a wider assessment of psychological well-being and quality of life. The exploration of further outcomes, such as cognitive decline and the design of larger scale trials on hard endpoints, such as major cardiovascular events, the incidence of diabetes and renal failure, would provide a more robust assessment of the clinical effects of specific carbohydrate restriction in the treatment of obesity.

AUTHOR CONTRIBUTIONS

GAS was involved in design, data collection, analysis and writing the manuscript. CC and BC were involved in design, data collection and manuscript revision. EM was involved in the design, analysis and writing the manuscript. FS, FB and FR were involved in data collection and manuscript revision. The manuscript was drafted, revised and approved by all the authors in accordance with ICJME standards for authorship. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

ACKNOWLEDGEMENTS

This research was performed as a part of the institutional activity of the unit, with no specific funding. Open Access Funding provided by Università degli Studi di Firenze within the CRUI-CARE Agreement.

CONFLICT OF INTEREST

The authors have no conflicts of interest to disclose.

COMPLIANCE WITH ETHICAL STANDARDS

This article does not contain any studies with human participants or animals performed by any of the authors.

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/dom.14709>.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ORCID

Giovanni Antonio Silverii  <https://orcid.org/0000-0002-6695-3213>

Edoardo Mannucci  <https://orcid.org/0000-0001-9759-9408>

Barbara Cresci  <https://orcid.org/0000-0002-8730-0748>

REFERENCES

- Riaz H, Khan MS, Siddiqi TJ, et al. Association between obesity and cardiovascular outcomes: a systematic review and meta-analysis of mendelian randomization studies. *JAMA Netw Open*. 2018;1:e183788. doi:10.1001/jamanetworkopen.2018.3788
- Arnold M, Pandeya N, Byrnes G, et al. Global burden of cancer attributable to high body-mass index in 2012: a population-based study. *Lancet Oncol*. 2015;16:36-46. doi:10.1016/S1470-2045(14)71123-4
- Wang J, Yang D-L, Chen Z-Z, Gou B-F. Associations of body mass index with cancer incidence among populations, genders, and menopausal status: A systematic review and meta-analysis. *Cancer Epidemiol*. 2016;42:1-8. doi:10.1016/j.canep.2016.02.010
- Renahan AG, Tyson M, Egger M, Heller RF, Zwahlen M. Body-mass index and incidence of cancer: a systematic review and meta-analysis of prospective observational studies. *Lancet*. 2008;371:569-578. doi:10.1016/S0140-6736(08)60269-X
- Yumuk V, Frühbeck G, Oppert JM, Woodward E, Toplak H. An EASO position statement on multidisciplinary obesity management in adults. *Obes Facts*. 2014;7:96-101. doi:10.1159/000362191
- Carlsson LMS, Sjöholm K, Jacobson P, et al. Life expectancy after bariatric surgery in the swedish obese subjects study. *N Engl J Med*. 2020;383:1535-1543. doi:10.1056/nejmoa2002449
- De Stefani F d C, Pietraroia PS, Fernandes-Silva MM, Faria-Neto J, Baena CP. Observational evidence for unintentional weight loss in all-cause mortality and major cardiovascular events: a systematic review and meta-analysis. *Sci Rep*. 2018;8:15447. doi:10.1038/s41598-018-33563-z
- Chen C, Ye Y, Zhang Y, Pan XF, Pan A. Weight change across adulthood in relation to all cause and cause specific mortality: prospective cohort study. *BMJ*. 2019;367. doi:10.1136/bmj.l5584
- Garvey WT, Mechanick JL, Brett EM, et al. American association of clinical endocrinologists and American college of endocrinology comprehensive clinical practice guidelines for medical care of patients with obesity. *Endocr Pract*. 2016;22:1-203. doi:10.4158/EP161365.GL
- Donnelly JE, Blair SN, Jakicic JM, et al. Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults. *Med Sci Sports Exerc*. 2009;41:459-471. doi:10.1249/MSS.0b013e3181949333
- Schutz D. European practical and patient-Centred guidelines for adult obesity management in primary care. *Obes Facts*. 2019;12:40-66. doi:10.1159/000496183
- Amerio ML, Angrisani L, Annuzzi G, et al. Standard Italiani per la Cura dell'Obesità SIO-ADI 2016-2017. *SIO (Società Ital Dell'Obesità)*. 2017;1:1-292.
- Koliaki C, Spinos T, Spinou M, Brinia ME, Mitsopoulou D, Katsilambros N. Defining the optimal dietary approach for safe, effective and sustainable weight loss in overweight and obese adults. *Healthcare (Basel)*. 2018;6(3):73-86. doi:10.3390/healthcare6030073

14. Raynor HA, Champagne CM. Position of the academy of nutrition and dietetics: interventions for the treatment of overweight and obesity in adults. *J Acad Nutr Diet*. 2016;116:129-147. doi:10.1016/j.jand.2015.10.031
15. Wharton S, Lau DCW, Vallis M, et al. Obesity in adults: a clinical practice guideline. *CMAJ*. 2020;192:E875-E891. doi:10.1503/cmaj.191707
16. Jensen MD, Ryan DH, Apovian CM, et al. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: A report of the American college of cardiology/American heart association task force on practice guidelines and the obesity society. *J Am Coll Cardiol*. 2014;63:2985-3023. doi:10.1016/j.jacc.2013.11.004
17. Fidanza F, Alberti A, Lanti M, Menotti A. Mediterranean Adequacy Index: correlation with 25-year mortality from coronary heart disease in the Seven Countries Study. *Nutr Metab Cardiovasc Dis*. 2004;14:254-258. doi:10.1016/S0939-4753(04)80052-8
18. Trichopoulou A, Costacou T, Bamia C, Trichopoulos D. Adherence to a Mediterranean diet and survival in a Greek population. *N Engl J Med*. 2003;348:2599-2608. doi:10.1056/NEJMoa025039
19. Caprio M, Infante M, Moriconi E, et al. Very-low-calorie ketogenic diet (VLCKD) in the management of metabolic diseases: systematic review and consensus statement from the Italian Society of Endocrinology (SIE). *J Endocrinol Invest*. 2019;42:1365-1386. doi:10.1007/s40618-019-01061-2
20. Ebbeling CB, Leidig MM, Feldman HA, Lovesky MM, Ludwig DS. Effects of a low-glycemic load vs low-fat diet in obese young adults: A randomized trial. *JAMA*. 2007;297:2092-2102. doi:10.1001/jama.297.19.2092
21. Goldstein T, Kark JD, Berry EM, Adler B, Ziv E, Raz I. The effect of a low carbohydrate energy-unrestricted diet on weight loss in obese type 2 diabetes patients - A randomized controlled trial. *e-SPEN*. 2011;6:e178-e186. doi:10.1016/j.eclnm.2011.04.003
22. Perticone M, Maio R, Sciacqua A, et al. Ketogenic diet-induced weight loss is associated with an increase in Vitamin D levels in obese adults. *Molecules*. 2019;24:1-12. doi:10.3390/molecules24132499
23. Landry MJ, Crimarco A, Gardner CD. Benefits of low carbohydrate diets: a settled question or still controversial? *Curr Obes Rep*. 2021;10:409-422. doi:10.1007/s13679-021-00451-z
24. Moon J, Koh G. Clinical evidence and mechanisms of high-protein diet-induced weight loss. *J Obes Metab Syndr*. 2020;29:166-173. doi:10.7570/jomes20028
25. Malhotra R, Lipworth L, Cavanaugh KL, et al. Protein intake and long-term change in glomerular filtration rate in the Jackson Heart Study. *J Ren Nutr*. 2018;28:245-250. doi:10.1053/j.jrn.2017.11.008
26. Oosterwijk MM, Groothof D, Navis G, Bakker SJL, Laverman GD. High-normal protein intake is not associated with faster renal function deterioration in patients with type 2 diabetes: a prospective analysis in the DIALECT Cohort. *Diabetes Care*. 2021;45:35-41. doi:10.2337/dc21-1211
27. Willems AEM, Sura-De Jong M, Van Beek AP, Nederhof E, Van Dijk G. Effects of macronutrient intake in obesity: a meta-analysis of low-carbohydrate and low-fat diets on markers of the metabolic syndrome. *Nutr Rev*. 2021;79:429-444. doi:10.1093/nutrit/nuaa044
28. Muscogiuri G, Ghoch E. European guidelines for obesity management in adults with a very low-calorie ketogenic diet: a systematic review and meta-analysis. *Obes Facts*. 2021;222-245. doi:10.1159/000515381
29. Nordmann AJ, Nordmann A, Briel M, et al. Effects of low-carbohydrate vs low-fat diets on weight loss and cardiovascular risk factors. *Arch Intern Med*. 2006;166:285. doi:10.1001/archinte.166.3.285
30. Collet J-P, Thiele H, Barbato E, et al. ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Eur Heart J*. 2020;2020:1-79. doi:10.1093/eurheartj/ehaa575
31. Bueno NB, De Melo ISV, De Oliveira SL, Da Rocha AT. Very-low-carbohydrate ketogenic diet v. low-fat diet for long-term weight loss: A meta-analysis of Randomised controlled trials. *Br J Nutr*. 2013;110:1178-1187. doi:10.1017/S0007114513000548
32. Johnston BC, Kanters S, Bandayrel K, et al. Comparison of weight loss among named diet programs in overweight and obese adults: a meta-analysis. *JAMA*. 2014;312:923-933. doi:10.1001/jama.2014.10397
33. Hashimoto Y, Fukuda T, Oyabu C, et al. Impact of low-carbohydrate diet on body composition: meta-analysis of randomized controlled studies. *Obes Rev*. 2016;17:499-509. doi:10.1111/obr.12405
34. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71. doi:10.1136/bmj.n71
35. Italian Society of human nutrition. *LARN: Nutrients and Energy Intake Standard Levels for the Italian Population: 4th revision*. Milan: SICS Editions; 2014.
36. Scientific Opinion on the essential composition of total diet replacements for weight control. *EFSA J*. 2015;13:3957. doi:10.2903/j.efsa.2015.3957
37. Higgins JPT, Sterne JAC, Savović J, et al. A revised tool for assessing risk of bias in randomized trials. In: Chandler J, McKenzie J, Boutron I, Welch V, eds. *Cochrane Methods. Cochrane Database of Systematic Reviews*; 2016. doi:10.1002/14651858.CD201601
38. Guyatt GH, Oxman AD, Schünemann HJ, Tugwell P, Knottnerus A. GRADE guidelines: a new series of articles in the Journal of Clinical Epidemiology. *J Clin Epidemiol*. 2011;64:380-382. doi:10.1016/j.jclinepi.2010.09.011
39. Foster GD, Wyatt HR, Hill JO, et al. Weight and metabolic outcomes after 2 years on a low-carbohydrate versus low-fat diet. *Obstet Gynecol Surv*. 2010;65:769-770. doi:10.1097/OGX.0b013e31821342ba
40. Iqbal N, Vetter ML, Moore RH, et al. Effects of a low-intensity intervention that prescribed a low-carbohydrate vs. a low-fat diet in obese, diabetic participants. *Obesity*. 2010;18:1733-1738. doi:10.1038/oby.2009.460
41. Bazzano LA, Hu T, Reynolds K, et al. Effects of low-carbohydrate and low-fat diets: a randomized trial. *Ann Intern Med*. 2014;161:309-318. doi:10.7326/M14-0180
42. Dalle Grave R, Calugi S, Gavasso I, El Ghoch M, Marchesini G. A randomized trial of energy-restricted high-protein versus high-carbohydrate, low-fat diet in morbid obesity. *Obesity*. 2013;21:1774-1781. doi:10.1002/oby.20320
43. Goday A, Bellido D, Sajoux I, et al. Short-Term safety, tolerability and efficacy of a very low-calorie-ketogenic diet interventional weight loss program versus hypocaloric diet in patients with type 2 diabetes mellitus. *Nutr Diabetes*. 2016;6:e230. doi:10.1038/nutd.2016.36
44. Yancy WS, Mayer SB, Coffman CJ, Voils CI. Effect of allowing choice of diet on weight loss. A randomized trial. *Ann Intern Med*. 2015;162:805-814. doi:10.7326/M14-2358.Effect
45. Monnier L, Schlienger J-L, Colette C, Bonnet F. The obesity treatment dilemma: Why dieting is both the answer and the problem? A mechanistic overview. *Diabetes Metab*. 2021;47:101192. doi:10.1016/j.diabet.2020.09.002
46. Barber TM, Hanson P, Kabisch S, Pfeiffer AFH, Weickert MO. The low-carbohydrate diet: short-term metabolic efficacy versus longer-term limitations. *Nutrients*. 2021;13:1-15. doi:10.3390/nu13041187
47. Hammad S, Pu S, Jones PJ. Current evidence supporting the link between dietary fatty acids and cardiovascular disease. *Lipids*. 2016;51:507-517. doi:10.1007/s11745-015-4113-x
48. Ferrannini E. Sodium-glucose co-transporters and their inhibition: clinical physiology. *Cell Metab*. 2017;26:27-38. doi:10.1016/j.cmet.2017.04.011
49. Chawla S, Radenkovic D. The effect of low-fat and low-carbohydrate diets on weight loss and lipid levels: a systematic review. *Nutrients*. 2020;12:1-21. doi:10.3390/nu12123774

50. Silverii GA, Botarelli L, Dicembrini I, et al. Low-carbohydrate diets and type 2 diabetes treatment: a meta-analysis of randomized controlled trials. *Acta Diabetol.* 2020;57:1375-1382. doi:10.1007/s00592-020-01568-8
51. Bischoff SC, Boirie Y, Cederholm T, et al. Towards a multidisciplinary approach to understand and manage obesity and related diseases. *Clin Nutr.* 2017;36:917-938. doi:10.1016/j.clnu.2016.11.007
52. Patsalos O, Keeler J, Schmidt U, Penninx BWJH, Young AH, Himmerich H. Diet, obesity, and depression: a systematic review. *J Pers Med.* 2021;1:1-19.
53. Blaine BE, Rodman J, Newman JM. Weight loss treatment and psychological well-being: a review and meta-analysis. *J Health Psychol.* 2007;12:66-82. doi:10.1177/1359105307071741
54. Fabricatore AN, Wadden TA, Higginbotham AJ, et al. Intentional weight loss and changes in symptoms of depression: a systematic review and meta-analysis. *Int J Obes (Lond).* 2011;35:1363-1376. doi:10.1038/ijo.2011.2
55. Bales CW, Starr KNP, Orenduff MC, et al. Influence of protein intake, race, and age on responses to a weight-reduction intervention in obese women. *Curr Dev Nutr.* 2017;1:1-10. doi:10.3945/cdn.117.000703
56. Brehm BJ, Lattin BL, Summer SS, et al. One-year comparison of a high-monounsaturated fat diet with a high-carbohydrate diet in type 2 diabetes. *Diabetes Care.* 2009;32:215-220. doi:10.2337/dc08-0687
57. Cornier MA, Donahoo WT, Pereira R, et al. Insulin sensitivity determines the effectiveness of dietary macronutrient composition on weight loss in obese women. *Obes Res.* 2005;13:703-709. doi:10.1038/oby.2005.79
58. Daly ME, Paisey R, Paisey R, et al. Short-term effects of severe dietary carbohydrate-restriction advice in Type 2 diabetes - a randomized controlled trial. *Diabet Med.* 2006;23:15-20. doi:10.1111/j.1464-5491.2005.01760.x
59. De Luis DA, Sagrado MG, Conde R, Aller R, Izaola O. The effects of two different hypocaloric diets on glucagon-like peptide 1 in obese adults, relation with insulin response after weight loss. *J Diabetes Complications.* 2009;23:239-243. doi:10.1016/j.jdiacomp.2007.12.006
60. De Luis DA, Aller R, Izaola O, Romero E. Effects of a high-protein/low-carbohydrate versus a standard hypocaloric diet on adipocytokine levels and cardiovascular risk factors during 9 months, role of rs6923761 gene variant of glucagon-like peptide 1 receptor. *J Endocrinol Invest.* 2015;38:1183-1189. doi:10.1007/s40618-015-0304-9
61. Foster GD, Wyatt HR, Hill JO, et al. A randomized trial of a low-carbohydrate diet for obesity. *N Engl J Med.* 2003;348:2082-2090. doi:10.1056/nejmoa022207
62. Kerksick CM, Wismann-bunn J, Fogt D, et al. Changes in weight loss, body composition and cardiovascular disease risk after altering macronutrient distributions during a regular exercise program in obese women. *Nutr J.* 2010;9:59. doi:10.1186/1475-2891-9-59
63. Kitabchi AE, McDaniel KA, Wan JY, et al. Effects of high-protein versus high-carbohydrate diets on markers of β -Cell function, oxidative stress, lipid peroxidation, proinflammatory cytokines, and adipokines in obese, premenopausal women without diabetes. *Diabetes Care.* 2013;36:1919-1925. doi:10.2337/dc12-1912
64. Morris E, Aveyard P, Dyson P, et al. A food-based, low-energy, low-carbohydrate diet for people with type 2 diabetes in primary care: A randomized controlled feasibility trial. *Diabetes Obes Metab.* 2020;22:512-520. doi:10.1111/dom.13915
65. Porter Starr KN, Pieper CF, Orenduff MC, et al. Improved function with enhanced protein intake per meal: a pilot study of weight reduction in frail, obese older adults. *J Gerontol A Biol Sci Med Sci.* 2016;71:1369-1375. doi:10.1093/gerona/glv210
66. Racette SB, Schoeller DA, Kushner RF, Neil KM, Herling-laffaldano K. Effects of aerobic exercise and dietary carbohydrate on energy expenditure and body composition during weight reduction in obese women. *Am J Clin Nutr.* 1995;61:486-494. doi:10.1093/ajcn/61.3.486
67. Samaha FF, Iqbal N, Seshadri P, et al. A low-carbohydrate as compared with a low-fat diet in severe obesity. *N Engl J Med.* 2003;348:2074-2081. doi:10.1056/NEJMoa022637
68. Stentz FB, Brewer A, Wan J, et al. Remission of pre-diabetes to normal glucose tolerance in obese adults with high protein versus high carbohydrate diet: randomized control trial. *BMJ Open Diabetes Res Care.* 2016. doi:10.1136/bmjdr-2016-000258
69. Tonstad S, Malik N, Haddad E. A high-fibre bean-rich diet versus a low-carbohydrate diet for obesity. *J Hum Nutr Diet.* 2014. doi:10.1111/jhn.12118
70. Yancy WS. A low-carbohydrate, ketogenic diet versus a low-fat diet to treat obesity and hyperlipidemia. *Ann Intern Med.* 2004;140:769-779.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Silverii GA, Cosentino C, Santagiuliana F, et al. Effectiveness of low-carbohydrate diets for long-term weight loss in obese individuals: A meta-analysis of randomized controlled trials. *Diabetes Obes Metab.* 2022; 24(8):1458-1468. doi:10.1111/dom.14709