



The maltreated eco-phenotype of eating disorders: A new diagnostic specifier? A systematic review of the evidence and comprehensive description

Eleonora Rossi ^{a,1}, Emanuele Cassioli ^{a,2}, Cristiano Dani ^{a,3}, Giorgia Marchesoni ^a, Alessio M. Monteleone ^{b,4}, Stephen A. Wonderlich ^{c,5}, Valdo Ricca ^a, Giovanni Castellini ^{a,*,6}

^a Psychiatry Unit, Department of Health Sciences, University of Florence, Florence, Italy

^b Department of Psychiatry, University of Campania L. Vanvitelli, Naples, Italy

^c Sanford Center for Biobehavioral Research, Fargo, ND, United States of America

ARTICLE INFO

Keywords:

Eating disorders
Anorexia nervosa
Bulimia nervosa
Binge-eating disorder
Childhood maltreatment
Early trauma
Childhood abuse
Maltreated eco-phenotype
Systematic review
Diagnostic specifier

ABSTRACT

This systematic review aimed to summarize the evidence on the existence of a distinct phenotypic expression of Eating Disorders (EDs) associated with childhood maltreatment (CM), the so-called maltreated eco-phenotype of EDs. PRISMA standards were followed. Articles providing data about the characteristics of individuals with an ED reporting CM were included. Relevant results were extracted and summarized. A quality assessment was performed. A total of 1207 records were identified and screened, and 97 articles published between 1994 and 2023 were included. Findings revealed distinct biological and clinical features in patients with EDs reporting CM, including neuroanatomical changes, altered stress responses, ghrelin levels, inflammation markers, and gut microbiota composition. Clinically, CM correlated with severer eating behaviors, higher psychiatric comorbidity, impulsivity, emotional dysregulation, and risky behaviors. Additionally, CM was associated with poorer treatment outcomes, especially in general psychopathology and psychiatric comorbidities. This review highlighted the need to move towards an etiologically informed nosography, recognizing CM not merely as a risk factor, but also as an etiologic agent shaping different eco-phenotypic variants of EDs.

1. Introduction

The concept of disease eco-phenotype encompasses the manifestation of pathologies beyond the immediate physical level to incorporate a constellation of factors, including socio-cultural dynamics, personal experiences, and traumatic events such as childhood maltreatment (CM) (Belsky and Pluess, 2013). This theoretical framework facilitates a deepened understanding of the interplay between genetic predisposition and the unique set of experiences of the individual in the development and course of disorders. Thus, it allows the differentiation of clinical manifestations of diseases based on developmental trajectories, with

implications in terms of personalized and targeted treatments. In the field of psychiatry, a particular focus has been posed on the role of CM in modifying the phenotypic presentation of mental illnesses (Teicher et al., 2022; Teicher and Samson, 2013).

CM is remarkably widespread, with a prevalence around 15% within the general population and exceeding 50% among patients with psychiatric disorders (Struck et al., 2020). Historically, research predominantly emphasized sexual and physical abuse as risk factors for psychopathology. However, literature has progressively broadened its perspective on CM, with a growing emphasis on the profound impact of emotional abuse and neglect (Teicher et al., 2022). Accordingly, the

* Corresponding author.

E-mail address: giovanni.castellini@unifi.it (G. Castellini).

¹ 0000-0003-4755-3879

² 0000-0003-3623-7096

³ 0000-0002-2705-5453

⁴ 0000-0002-6786-4458

⁵ 0000-0002-0659-153X

⁶ 0000-0003-1265-491X

<https://doi.org/10.1016/j.neubiorev.2024.105619>

Received 17 December 2023; Received in revised form 1 March 2024; Accepted 6 March 2024

Available online 9 March 2024

0149-7634/© 2024 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

World Health Organization (WHO) provided a comprehensive definition of CM, emphasizing the betrayal of a child by an individual in a position of responsibility, trust, or power as a core aspect of CM: "Child maltreatment comprises abuse and neglect that occurs to children under 18 years of age. It includes all types of physical and/or emotional ill-treatment, sexual abuse, neglect, negligence as well as commercial or other forms of exploitation, all of which result in actual or potential harm to the child's health, development or dignity within the context of relationships of responsibility, trust, or power." (World Health Organization, 2022). Extensive research over the years elucidated the role of all forms of CM as transdiagnostic risk factors for psychopathology given their influence on the developmental trajectory of individuals, encompassing biological, emotional processing, and self-concept dimensions (McLaughlin et al., 2016, 2020; Pollak, 2015; Teicher et al., 2016). Moving from the observation that CM not only increases the likelihood of developing psychopathology, but also activates molecular, neuroanatomical, and psychological modifications that distinguish maltreated and non-maltreated individuals, numerous studies provided descriptions of the distinct characteristics exhibited by patients with various mental disorders when a history of childhood trauma is present (Teicher et al., 2022; Teicher and Samson, 2013). Notably, Teicher and Samson (2013) extensively described the maltreated eco-phenotypes of mood, anxiety, and substance abuse disorders, highlighting key features such as earlier age at onset, heightened symptom severity, more pernicious disease course, increased risk of suicide, compromised quality of life, and higher prevalence of psychiatric comorbidities compared to individuals with the same diagnoses but no history of CM (Teicher and Samson, 2013). Moreover, neurobiological findings revealed specific alterations associated with CM, including reduced hippocampal volume, heightened amygdala reactivity, dysregulation of the hypothalamus-pituitary-adrenal (HPA) axis, and changes in inflammatory markers (Teicher and Samson, 2013). An increasing number of studies showed similar findings within the scope of other psychiatric disorders, including personality disorders and psychoses (Teicher et al., 2022; Euler et al., 2021; Thomas et al., 2019). This led to a proposed necessity to modify the current diagnostic nosology, preferably incorporating specifiers to refer to the CM-associated eco-phenotype of mental illnesses (Teicher et al., 2022).

Eating disorders (EDs) - encompassing Anorexia Nervosa (AN), Bulimia Nervosa (BN), Binge Eating Disorder (BED), and Other Specified Feeding or Eating Disorders (OSFED) - are among the psychiatric illnesses most strongly associated with the presence of childhood trauma, with prevalence rates reaching up to 59% (Molendijk et al., 2017). The current nosology describes these disorders as having a common psychopathological core characterized by an overemphasis on the importance of body shape and weight in determining one's self-esteem, and by pathological eating behaviors ranging from fasting, to binge eating, or purging (American Psychiatric Association, 2022; World Health Organization, 2019). Although the Diagnostic and Statistical Manual (DSM) of mental disorders acknowledges CM as a potential risk factor for the development of EDs, neither the DSM nor the International Classification of Diseases currently recognize its potential influence on the phenotypic presentation of these disorders through specific codes or diagnostic specifiers (American Psychiatric Association, 2022; World Health Organization, 2019). Despite this, in recent years, the study of trauma as a possible phenotypic modifier of EDs gained increasing interest and a great number of studies have been published on this subject.

The main systematic reviews or meta-analysis of recent literature in this field have each focused on some specific aspects of the topic. In particular, the meta-analysis by Molendijk et al. (2017) elucidated to what extent CM affects some of the defining clinical features of EDs, including age at onset, prevalence of psychiatric comorbidities, suicidal behaviors, binge-purge behaviors, and disease severity (Molendijk et al., 2017). Various reviews, on the other hand, focused on the possible psychopathological and/or biological mechanisms involved in the link between childhood trauma and the development/maintenance of EDs

(Chami et al., 2019; Marciello et al., 2020; Rabito-Alcón et al., 2021; Trottier and MacDonald, 2017; Cascino and Monteleone, 2024). For what concerns the impact of CM on treatment outcome, Day et al. (2023) offered an in-depth analysis of how CM and the presence of Post-Traumatic Stress Disorder (PTSD) can alter responses to standard treatments (Day et al., 2023), while Brewerton (2019) and Mitchell et al. (2021) provided an overview of the importance of a trauma-informed care and practice for patients with EDs (Brewerton, 2019; Mitchell et al., 2021). However, a systematic review of the literature is missing that provides a comprehensive description of the current knowledge about the specific characteristics of patients with EDs and a history of CM from an integrated psychopathological, clinical, biological, and treatment-informed perspective. Furthermore, in recent years, alongside the exponential growth of interest in this area, the development of statistical methodologies such as network analysis and structural equation modeling (SEM) allowed deepening the knowledge of the complex interactions between psychopathological variables involved in shaping disease phenotypes both cross-sectionally and longitudinally. However, there is currently no systematic review that is recent enough to provide an overview of studies that have adopted these innovative approaches.

Therefore, the objective of this study was to conduct an updated systematic review of the literature about the role of CM in modifying the phenotypic presentation of EDs from the multi-faceted perspective of biological, clinical, psychological, and psychopathological correlates, longitudinal characteristics, and treatment outcomes to provide a summary of the evidence of the existence of a maltreated eco-phenotype of EDs and a comprehensive description of its defining features.

2. Methods

The present systematic review was conducted and reported following the updated Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (Page et al., 2021).

2.1. Search strategy and selection criteria

Both the PubMed and ClinicalTrials.gov databases were searched for peer-reviewed articles on the topics of both EDs and CM. Additional field tags were embedded in the search string in order to exclude articles that were not written in English and reviews or meta-analyses. However, an initial search was performed without filtering out non-English language articles, in order to take note of the number of excluded articles based on language. For the ClinicalTrials.gov database, the search was limited to studies marked as "Completed" or "Terminated". The full search string is provided below.

(eating disorder OR anorexia nervosa OR bulimia nervosa OR binge eating disorder OR disordered eating) AND (childhood trauma OR childhood abuse OR childhood maltreatment OR early trauma OR early abuse OR early maltreatment OR maltreated ecophenotype OR traumatic ecophenotype OR maltreated subtype OR traumatic subtype OR complex ptsd OR complex post-traumatic stress disorder OR adverse childhood experiences OR maltreated OR adolescent trauma OR adolescent maltreatment OR adolescent sexual abuse OR adolescent emotional abuse OR adolescent physical abuse OR childhood neglect OR early neglect OR neglect) AND eng[la] NOT review[pt] NOT systematic review[pt] NOT meta-analysis[pt]

The final search was conducted on the 21st of July, 2023, and included all results from database inception. Studies were included according to the following inclusion criteria (screened in this order): published after January 1st 1994; written in English; provided data on human participants; provided original data; were not case reports or series; provided quantitative data; included patients with current clinical diagnoses of EDs according to DSM-IV (or later editions) or ICD-10 (or later editions); (American Psychiatric Association, 1994; World Health Organization, 1994) assessed CM using validated instruments or clinical interviews following acceptable definitions of CM; provided data or analyses related to individuals suffering from an ED describing

specific characteristics associated with CM. In cases where a study included aggregated samples (e.g., patients with a current ED diagnosis together with recovered patients, or patients with CM together with individuals who reported adverse experiences in adulthood), it was included only if it reported results on the subgroups of interest for the current systematic review, or alternatively, it was excluded. Articles were excluded if they were found to be duplicate reports of previously published studies.

2.2. Study selection, data extraction, and quality assessment

Any articles published before 1994 were excluded from the retrieved records. Titles and abstracts were screened independently by CD and GM. After the first screening phase, full reports of included records were retrieved and further assessed for eligibility by CD and GM. Finally, CD and GM conducted data extraction on eligible reports, retrieving the following information: country where patient recruitment took place, study design, diagnostic criteria used for assessing ED, sample size for each diagnostic subgroup (only for the samples where the analyses of interest were conducted), participants' gender, age range (underage, adults, or mixed sample), instrument used for assessing CM, and results relevant to the topic of the review. A result was considered relevant when it described the characteristics of patients with EDs that were specific to the subgroup of individuals reporting CM, as opposed to those without a history of CM (e.g., comparisons between patients with and without CM, or associations between levels of exposure to CM and another measure). Since the validity of negative results in scientific literature is considered to be quite low in the absence of an appropriate discussion on the achieved statistical power, and given that the latter is rarely reported, especially in observational studies (Hebert et al., 2002), negative results were extracted and reported only if they represented one of the main outcomes, for which the study was powered.

The methodological quality assessment of cross-sectional studies was conducted by CD and GM, using the Newcastle-Ottawa Scale (NOS) adaptation for cross-sectional studies (Modesti et al., 2016). The same instrument was used for longitudinal studies, with the two additional items from the original NOS related to prospective evaluations (i.e., adequate follow-up duration and presence of attrition bias).

In cases of disagreement among reviewers at any phase of the review process, the full-text report was retrieved and independently assessed by expert reviewers ER and EC, with discussions held until a consensus was reached. A consensus was reached for all studies. Furthermore, a random subset comprising 30% of the full-text reports was independently reviewed by ER and EC to ensure the quality of the process and data.

2.3. Narrative synthesis

Given the inherent heterogeneity of the data, a narrative synthesis approach was chosen to summarize and analyze the findings (Rodgers et al., 2009). The synthesis process involved identifying emerging themes within the included studies, which were grouped according to shared characteristics of the main findings. These groupings allowed for the organization of the narrative synthesis. Overall, three main types of findings related to the maltreated eco-phenotype were searched:

1. Biological correlates: results that describe differences in terms of organ anatomy and function, including endocrinological, neurobiological and immunological features.
2. Clinical, psychological, and psychopathological correlates: findings showing different ED characteristics, in terms of etiopathogenetic, diagnostic, psychopathological and behavioral features.
3. Longitudinal characteristics and treatment outcomes: results from studies that highlight a distinct disease trajectory following the diagnosis of ED, in terms of varying prognosis and response to the different treatments offered.

For each main theme, a coherent and comprehensive account of all key findings from each study was summarized and reported.

3. Results

The complete flow diagram of the study selection process is reported in Fig. 1. The initial search identified a total of 1207 records, of which 1053 were screened and 97 were finally included in the review. The characteristics and main findings of the included studies are reported in Supplementary Table 1, 2, and 3. Two reports referred to the same sample, therefore they were summarized together (Vieira et al., 2017, 2018). Most of the selected studies were conducted in Europe ($n_{Italy} = 38$, $n_{France} = 4$, $n_{Spain} = 4$, $n_{Germany} = 3$, $n_{Belgium} = 1$, $n_{Denmark} = 1$, $n_{Norway} = 1$, $n_{Poland} = 1$, $n_{Portugal} = 1$), while the remaining were conducted in the United States ($n = 14$), Canada ($n = 12$), the United Kingdom ($n = 12$), Israel ($n = 3$), Japan ($n = 1$) and South Korea ($n = 1$) (Supplementary Table 1, 2, and 3). Regarding the study design, the vast majority of included reports were cross-sectional in nature ($n = 81$, including one with a retrospective chart review methodology), whereas 15 were longitudinal (of which 2 were retrospective) and 1 was a randomized controlled trial (RCT) (Supplementary Table 1, 2 and 3). More than half of the included studies used the DSM-IV criteria for diagnosing EDs ($n = 54$), while 38 studies used the DSM-5 criteria, and 5 studies used the ICD-10 criteria (Supplementary Table 1, 2, and 3).

In terms of methodological quality, the median score for cross-sectional studies was 7, with an interquartile range of 7–8, out of a possible maximum of 10. For longitudinal studies, the median score was 10, with an interquartile range of 9–11, out of a maximum of 12. These scores suggest that, despite the inherent limitations of observational studies, the overall quality was commendable. Most studies demonstrated acceptable sample sizes and response rates, utilized validated instruments for the assessment of CM, and, in the case of longitudinal studies, ensured adequate follow-up durations and minimized the risk of attrition bias.

3.1. The maltreated eco-phenotype of EDs: biological correlates

3.1.1. Neuroanatomy

Neuroimaging studies performed in patients with AN and BN showed preliminary evidence of specific modifications induced by CM at a neuroanatomical level (Cascino et al., 2023a, 2023b; Molina-Ruiz et al., 2022; Monteleone et al., 2019a). Maltreated patients exhibited reduced gray matter volume in the right paracentral lobule and left inferior temporal gyrus, reduced white matter integrity in corpus callosum, internal capsule, posterior thalamic radiation, longitudinal fasciculus, and corona radiata as compared to non-maltreated counterparts (Monteleone et al., 2019a). A negative correlation between gray and white matter changes and the levels of emotional and physical neglect was observed (Monteleone et al., 2019a). Furthermore, maltreated patients showed lower cortical thickness values in the left superior frontal, right caudal middle frontal, and right superior parietal gyri (Cascino et al., 2023a) and significantly lower local gyrification index in the left middle temporal gyrus (Cascino et al., 2023b). Finally, Molina-Ruiz et al. (2022) showed that volumes of the putamen (both hemispheres) were negatively correlated with self-reported early traumatization scores in combined analysis of ED and control subjects, although the result did not reach statistical significance in each diagnostic subgroup (Molina-Ruiz et al., 2022).

3.1.2. Endogenous stress response system

Several studies showed that early trauma impacts the hypothalamus-pituitary-adrenal (HPA) axis and sympathetic nervous system activity in patients with AN and/or BN (Castellini et al., 2012; Díaz-Marsá et al., 2007; Lelli et al., 2019; Meneguzzo et al., 2022; Monteleone et al., 2015, 2020a, 2021, 2018a, 2018b). Patients with CM had lower 24-h urinary free cortisol levels, (Meneguzzo et al., 2022) lower morning plasmatic




Fig. 1. Preferred reporting items for systematic reviews and meta-analysis (PRISMA) flow diagram.

cortisol levels (Lelli et al., 2019) and lower *saliva cortisol awakening response (CAR)* (Monteleone et al., 2015, 2020a, 2018a) than non-maltreated patients, with a significant negative correlation between the number of traumas and CAR (Monteleone et al., 2018a). As a further demonstration of dampened basal activity of the endogenous stress response system in patients with EDs reporting early trauma (Monteleone et al., 2020), showed that maltreated patients had decreased morning secretion of *salivary alpha-amylase*, (Monteleone et al., 2020a) a marker of the sympathetic nervous system activity. Furthermore Diaz-Marsa et al. (2007), showed that in patients with AN-bp and BN cortisol suppression after the administration of 0.25 mg dexamethasone was significantly correlated with the presence of childhood traumatic events (Díaz-Marsá et al., 2007), suggesting the presence of an increased sensitization of the hypothalamic-pituitary glucocorticoid receptors in this group. In contrast with these results, the study performed by Castellini et al. (2012) did not show a direct effect of childhood abuse (sexual or physical) on plasmatic cortisol levels (Castellini et al., 2012). However, it underlined a positive association between plasmatic cortisol levels and sexual desire in physically or sexually abused patients, suggesting that these women would experience an automatic fear response determining significant increase in cortisol levels when perceiving sexual desire. For what concerns the effect of psychotherapy in terms of modifying cortisol levels, Lelli et al. (2019)

showed that hypocortisolism persisted after CBT-E in patients with AN and BN reporting CM (Lelli et al., 2019).

Two studies evaluated the biological and emotional response to psychosocial stress exposure (Trier Social Stress Test – TSST) of maltreated patients with EDs as compared to those without CM (Monteleone et al., 2021, 2018b). Contrasting results were found in patients with AN. Indeed, one study showed that maltreated women exhibited a blunted cortisol response and reduced anxiety increase after TSST as compared to non-maltreated patients (Monteleone et al., 2018b), whereas another one did not confirm this result (Monteleone et al., 2021). In particular, the latter study showed that both maltreated patients with AN and BN had heightened emotional reactivity, lower levels of hunger and more severe body dissatisfaction after TSST. Furthermore, higher cortisol production was observed in patients with AN, regardless of the presence of CM, and in patients with BN and emotional CM. Emotional and not physical CM was the most significant determinant of the experimental differences observed in this study.

3.1.3. Inflammation markers and ghrelin

Early trauma was also demonstrated to impact on inflammation markers (Bou Khalil et al., 2022; Rodríguez-Quiroga et al., 2021) and ghrelin (Rossi et al., 2021). In particular, Bou Khalil et al. (2022) showed that maltreated patients with AN had a higher

neutrophil-to-lymphocyte ratio (Bou Khalil et al., 2022), whereas Rodríguez-Quiroga et al. (2021) demonstrated that CM was associated with higher levels of **tumor necrosis factor-alpha (TNF- α)** in a mixed sample of patients with EDs (Rodríguez-Quiroga et al., 2021). Furthermore, Rossi et al. (2021) showed in a mixed sample of patients with EDs including AN, BN, and BED, that those reporting CM had higher **ghrelin** levels, and early traumatic experiences moderated the association between ghrelin and overeating-related pathological behaviors (Rossi et al., 2021). The association between childhood sexual abuse and binge eating was mediated by elevated ghrelin, suggesting a possible involvement of this orexigenic molecule in maintaining pathological overeating-related behaviors in traumatized patients.

3.1.4. Microbiota-gut-brain axis

Only one study is available regarding the impact of CM on microbiota-gut-brain axis (Castellini et al., 2023a), showing that CM predicted lower **fecal levels of propionic and butyric acid** and higher levels of trait anxiety in a mixed sample of individuals with AN, BN and BED. A serial mediation model showed an indirect effect of CM on higher ED-specific psychopathology through the mediation of impaired levels of butyric acid, which in turn determined increased anxiety. PERMANOVA analysis showed that patients in the cluster characterized by CM, increased anxiety, binge eating behaviors, and reduced butyric acid showed different **microbiome composition**.

3.1.5. Genetics, monoaminergic, and glucocorticoid systems

Several studies evaluated the interactions between CM and genetic factors in determining neurobiological abnormalities and behavioral tendencies in patients with EDs (Groleau et al., 2014, 2012a; Steiger et al., 2007, 2008, 2013). Patients who were carriers of the low-function allele of the dopamine D2 receptor (DRD2) Taq1A polymorphism showed higher rates of sensation seeking, only in the presence of a history of childhood sexual abuse (Groleau et al., 2012a). However, in patients with a bulimia-spectrum disorder, no significant difference was found between those with early trauma (sexual or physical) and those without it in terms of DRD2 gene promoter methylation (Groleau et al., 2014). On the other side, with a classification based on a biallelic model of serotonin transporter promoter region (5-HTTLPR) polymorphism, multiple regression analyses indicated significant proportions of variance in stimulus seeking and insecure attachment to be explained by abuse \times genotype interaction effects, with greater psychopathology always occurring in S-allele carriers who had been abused (Steiger et al., 2007). Furthermore, patients who were carriers of the S-allele showed higher levels of dissociative behavior, only in the presence of CM (Steiger et al., 2008). Furthermore, the presence of early sexual abuse was associated with reduced serotonin transporter density on platelets (Steiger et al., 2004). No effects related to childhood abuse were observed regarding methylation of the glucocorticoid receptor gene (NR3C1) promoter (Steiger et al., 2013).

3.2. The maltreated eco-phenotype of EDs: clinical, psychological, and psychopathological correlates

3.2.1. ED-symptoms

Castellini et al. (2018) and Lelli et al. (2019) found that patients with AN or BN and a history of childhood abuse had a lower **age at disease onset** (Lelli et al., 2019; Castellini et al., 2018). Furthermore, according to Castellini et al. (2018) abused patients had a **history of overweight during childhood** more frequently (Castellini et al., 2018).

Several studies underlined that patients with **bingeing/purging subtypes of EDs**, namely AN-bp or BN, reported CM more frequently than those with AN-r (Díaz-Marsá et al., 2007; Carretero-García et al., 2012; Carter et al., 2006; Kjaersdam Tellús et al., 2021; Longo et al., 2019; Longo and Marzola, 2020; Rienecke et al., 2022; Speranza et al., 2003; Van Gerko et al., 2005). Furthermore, among patients with AN-r, a higher rate of sexual abuse was observed when a previous history of BN

was reported (Santonastaso and Zanetti, 2006).

An association between CM and greater severity of **pathological eating behaviors** was observed, specifically binge-eating (Castellini et al., 2018; Van Gerko et al., 2005; Groth et al., 2020; Waller et al., 2001; Wonderlich et al., 2007), purging (Castellini et al., 2018; Rienecke et al., 2022; Van Gerko et al., 2005; Groth et al., 2020; Waller et al., 2001; Wonderlich et al., 2007; Cascino et al., 2021), and restricting behaviors (Rienecke et al., 2022). Furthermore, among patients with BN or BED, night eating syndrome with binge eating was significantly associated with the presence of higher levels of physical neglect (Latzer et al., 2020).

Most of the studies underlined significant associations across the different ED diagnoses between childhood trauma exposure and increased severity of various dimensions of **ED-specific psychopathology** including drive for thinness and/or shape, weight, and eating concerns (Meneguzzo et al., 2022, 2021; Monteleone et al., 2021, 2020b; Carter et al., 2006; Van Gerko et al., 2005; Wonderlich et al., 2007; Latzer et al., 2020; Groleau et al., 2012b; Guillaume et al., 2016; Kong and Bernstein, 2009; Racine and Wildes, 2015; Tasca et al., 2013). In particular, Guillaume et al. (2016) underlined an association between the number of traumas experienced and an increased severity of ED symptoms (Guillaume et al., 2016), whereas Meneguzzo et al. (2021) reported a more severe clinical impact in case of repetition of adulthood trauma exposure after childhood abuse (Meneguzzo et al., 2021). Aligning with these results, according to Pugh et al. (2018) the relative power of the ED “voice” was related to childhood emotional abuse, partly through the mediation of dissociation (Pugh et al., 2018). However, some studies did not confirm these results and showed no significant differences between patients with AN or BN reporting childhood sexual and/or physical abuse or not in terms of severity of ED psychopathology (Lelli et al., 2019; Castellini et al., 2018; Calugi et al., 2018; Favaro et al., 1998).

Furthermore, an association between the presence of early traumatic experiences and **negative body image/body dissatisfaction was observed** (Monteleone et al., 2021; Cascino et al., 2021; Kong and Bernstein, 2009; Muehlenkamp et al., 2011). In particular, Sweetingham et al. (2008) showed that having been teased about appearance by peers and having been verbally bullied by family members were both associated with higher body dissatisfaction, and shame mediated the relationship between these variables (Sweetingham and Waller, 2008).

3.2.2. Psychiatric comorbidities, psychosocial functioning, and quality of life

CM was associated with greater **general psychopathology** (Meneguzzo et al., 2022, 2021; Castellini et al., 2018; Carter et al., 2006; Latzer et al., 2020; Muehlenkamp et al., 2011), worse **psychosocial functioning** (Guillaume et al., 2016; Meneguzzo et al., 2021) and poorer **quality of life** (Meneguzzo et al., 2021; Monteleone et al., 2023). In particular, Guillaume et al. (2016) underlined a crude dose-effect between the number of traumas experienced and an increased severity of daily functioning impairment (Guillaume et al., 2016).

For what concerns psychiatric comorbidities, CM predicted an increased likelihood of **mood disorders** (Wonderlich et al., 2007) including depression (Meneguzzo et al., 2022; Carter et al., 2006; Longo and Marzola, 2020; Waller et al., 2001; Latzer et al., 2020; Groleau et al., 2012b; Kong and Bernstein, 2009; Richardson et al., 2008) and bipolar disorder type II (Castellini et al., 2018). Leraas et al. (2018) through a latent profile analysis showed that CM was associated with a “stable depressed” profile of patients with BN characterized by low affective lability and high affect intensity (Leraas et al., 2018). Furthermore, CM was associated with increased **obsessive-compulsive symptoms** (Castellini et al., 2018; Carter et al., 2006; Kong and Bernstein, 2009; Lockwood et al., 2004), **anger/hostility** (Wonderlich et al., 2007; Strangio et al., 2017), and **anxiety** (Meneguzzo et al., 2022; Castellini et al., 2023a; Carter et al., 2006; Wonderlich et al., 2007; Cascino et al., 2021; Richardson et al., 2008). In particular, Stein et al. (2013)

underlined that childhood sexual abuse was associated with increased fear of life (Stein et al., 2013). Moreover, patients with CM reported increased **post-traumatic stress symptoms** (Cascino et al., 2021; Meneguzzo et al., 2021; Brewerton et al., 2021; Castellini et al., 2020a; Hazzard et al., 2021; Steiger et al., 2001). Specifically, several studies underlined an association between CM and **dissociation** (Castellini et al., 2018; Longo and Marzola, 2020; Waller et al., 2001; Muehlenkamp et al., 2011; Longo et al., 2023; Nagata et al., 1999), including somatoform dissociation (Longo et al., 2023) and depersonalization (Castellini et al., 2018). Finally, Dalgleish et al. (2003) showed that self-reported parental abuse predicted a tendency to produce over general memories when exposed to negative emotional cues (Dalgleish et al., 2003).

For what concerns the area of **personality disorders**, an association was observed between a history of CM and comorbid borderline personality disorder psychopathology in patients with AN or BN (Steiger et al., 2007; Wonderlich et al., 2007; Spiegel et al., 2022), particularly in the presence of polytrauma (Uttinger et al., 2016). Moreover, Rodríguez-Quiroga 2021 showed that a history of CM was associated with higher delirious, narcissistic, and paranoid personality traits in a mixed sample of patients with EDs (Rodríguez-Quiroga et al., 2021). Finally, Grilo & Masheb (2002) showed that in a sample of patients with BED, among the different types of CM only emotional abuse predicted the presence of a personality disorder (Grilo and Masheb, 2002). In particular, patients with cluster C personality disorders, especially avoidant personality disorder, were significantly more likely to have clinically meaningful levels of emotional abuse (Grilo and Masheb, 2002).

3.2.3. Impulsivity, emotional dysregulation, and interoceptive awareness

The presence of any childhood trauma predicted greater **impulsivity** and impulsive behaviors (Castellini et al., 2012, 2018; Monteleone et al., 2021, 2020b; Steiger et al., 2007; Kong and Bernstein, 2009; Strangio et al., 2017; Corstorphine et al., 2007; Dodd et al., 2022; Favaro et al., 2004; Schaefer et al., 2021). Furthermore, increasing evidence showed a significant association between CM and **emotional dysregulation** (Racine and Wildes, 2015; Strangio et al., 2017; Schaefer et al., 2021; Cassioli et al., 2022). In particular, Racine and Wildes showed that of the three forms of childhood abuse (physical, sexual and emotional), emotional abuse was the most strongly related to emotion dysregulation in patients with AN, both in restricting and binge-eating/purging subtypes (Racine and Wildes, 2015). Moreover, Monteleone et al. (2021) provided an experimental evidence of the heightened emotional reactivity of patients reporting CM when exposed to social stress through the adoption of the TSST (Monteleone et al., 2021). Finally, a severer deficit in **interoceptive awareness**, an indirect measure of emotion regulation, was underlined among patients reporting CM (Monteleone et al., 2021, 2020b; Kong and Bernstein, 2009).

3.2.4. Self-harm behaviors, suicidality, and substance abuse

Several studies underlined an association between CM and higher rates of **non-suicidal self-injury (NSSI)** in patients with AN, BN or BED (Vieira et al., 2017, 2018; Wonderlich et al., 2007; Favaro et al., 1998, 2008; Muehlenkamp et al., 2011; Steiger et al., 2001; Nagata et al., 1999; Corstorphine et al., 2007; Dohm et al., 2002; Favaro and Santonastaso, 1997, 1998, 1999, 2000; Fennig and Hadas, 2010). In particular, Favaro & Santonastaso (2000) underlined that impulsive self-injury in patients with AN was predicted by childhood sexual abuse and anxiety, in contrast with compulsive self-injury which was predicted by obsessionality and age (Favaro and Santonastaso, 2000). Furthermore, many studies found that patients with AN and/or BN with a history of CM showed a higher risk of **suicidal ideation and suicidal attempts** (Vieira et al., 2018; Favaro et al., 1998; Nagata et al., 1999; Favaro et al., 2008; Fennig and Hadas, 2010; Arnold et al., 2023; Mayes et al., 2014; Nickel et al., 2006; Smith et al., 2016).

For what concerns **alcohol and substance abuse**, many studies underlined an association between these risky behaviors and CM in

samples of patients with different kinds of EDs, including AN, BN and BED (Castellini et al., 2018; Richardson et al., 2008; Corstorphine et al., 2007; Schaefer et al., 2021; Dohm et al., 2002).

3.2.5. Sexuality

The relationship between CM and sexuality has been investigated both in terms of sexual dysfunctions (e.g. problems in sexual desire, arousal, orgasm, and satisfaction) and of dysregulated sexuality/hypersexuality. Although CM did not show a direct effect in determining severer sexual dysfunctions before treatment in patients with AN or BN (Castellini et al., 2012; Cassioli et al., 2020), a paradoxical association between sexual desire and increased hematic cortisol levels was observed in patients sexually or physically abused, suggesting that in this population sexual activity might elicit an automatic fear response (see also 3.2.2) (Castellini et al., 2012). Furthermore, patients with AN or BN reporting childhood sexual abuse did not show a significant improvement in sexual functioning after treatment (Castellini et al., 2013). For what concerns the issue of hypersexual behaviors, Castellini et al. (2020) showed that childhood trauma was associated with **dysregulated sexuality** (Castellini et al., 2020a). In particular, in the subgroup of traumatized patients hypersexuality was associated with emotion dysregulation and psychopathology. Finally, Walsh and Burns (2000) demonstrated that patients with both AN and history of sexual abuse had higher response times for sexual maturation words than for control-related words, and greater cognitive interference from sexual maturation words than the non-abused patients with AN (Walsh and Burns, 2000).

3.2.6. Psychological constructs

Several psychological constructs were associated with childhood trauma exposure. In particular, feelings of ineffectiveness (Monteleone et al., 2019, 2020ba, 2021, 2018b; Cascino et al., 2021; Groleau et al., 2012b; Kong and Bernstein, 2009), social insecurity (Monteleone et al., 2019a, 2021, 2018b), perfectionism (Kong and Bernstein, 2009; Monteleone et al., 2020b), asceticism (Monteleone et al., 2021, 2020b), the presence of a more external locus of control (Waller, 1998), self-criticism (Speranza et al., 2003), low self-esteem (Carter et al., 2006; Latzer et al., 2020; Muehlenkamp et al., 2011), interpersonal problems (Carter et al., 2006; Dodd et al., 2022), interpersonal distrust (Monteleone et al., 2021), negative self-directed style and self-discrepancy (i.e., the difference between one's actual self and who one believes they ought to be) (Borg et al., 2022), early maladaptive schemas (e.g. disconnection and rejection) (Meneguzzo et al., 2021) and core beliefs of abandonment, functional dependency, defectiveness/shame, emotional deprivation or inhibition, failure to achieve, mistrust/abuse, social isolation, unrelenting standards, and vulnerability to harm (Waller et al., 2001).

3.2.7. Complex relationships between psychopathological dimensions

Rodgers et al. (2019) through a Bayesian network approach showed that symptoms importance and **network structure of ED psychopathology** differed between individuals with various ED diagnosis with and without a history of CM (Rodgers et al., 2019). Specifically, in patients with CM a 4-symptom pathway emerged, leading from overvaluation of weight and shape, and ending in overeating, through the mediation of loss of control and depressed mood, which emerged as the more important driving symptoms. On the other side, the ED symptom network in patients without CM was organized around the overinvestment in weight and shape and the resulting efforts to control or alter these aspects through dieting and excessive exercise. In the latter group, the driving symptoms were overeating and overvaluation of weight and shape.

The network analysis approach identified the core role of **emotional abuse** among the different kinds of CM in the phenomenology of EDs. In particular, two network analysis studies underlined that all types of CM were connected to psychopathology through the mediation of emotional

abuse in samples of patients with AN restricting type or with binge-purging symptoms (i.e., with AN purging type or BN), (Monteleone et al., 2019b) and with BN or BED (Monteleone et al., 2022). Interoceptive awareness was involved in the associations between emotional abuse and ED-core symptoms in all the diagnostic groups (Monteleone et al., 2019b, 2022) and a pivotal role in this association was also played by ineffectiveness in the binge-purge group (Monteleone et al., 2019b, 2022) and by impulsivity in patients with BED (Monteleone et al., 2022). Aligning with these results, a network analysis study performed in underaged patients with AN, BN, or OSFED showed that the links between physical and sexual abuse, psychopathology, and life satisfaction took place through the emotional abuse node (Monteleone et al., 2023). Furthermore, emotional neglect was connected to life satisfaction both directly and via emotional abuse, feeling disliked by others, and feelings of inferiority and worthlessness before treatment (Monteleone et al., 2023).

In terms of possible *mediators between CM and ED-specific psychopathology*, many other psychopathological and psychological variables were demonstrated to be involved in patients with different ED diagnosis through mediation analysis, path analysis, or structural equation models, and included emotion dysregulation (Racine and Wildes, 2015), attachment anxiety and avoidance (Tasca et al., 2013), feelings of ineffectiveness (Groleau et al., 2012b), affective instability (Groleau et al., 2012b), depression (Kong and Bernstein, 2009), early maladaptive schemas (e.g. disconnection and rejection) (Meneguzzo et al., 2021), and negative core beliefs (Waller et al., 2001). In particular, Waller et al. (2001) showed that depression mediated the association between childhood sexual abuse, core beliefs of abandonment and mistrust/abuse and frequency of bingeing, whereas both dissociation and depression mediated the association between core beliefs of defectiveness/shame and frequency of vomiting (Waller et al., 2001).

Several studies were performed evaluating the *mediators between CM and psychiatric comorbidities/risky behaviors* in patients with EDs. In particular, path analysis showed that emotion dysregulation and impulsivity mediated the relationship between CM and the presence of problematic alcohol/drug use in patients with BN (Schaefer et al., 2021). In terms of predicting NSSI, structural equation modeling analyses showed that the latent variable related to childhood abuse was associated to NSSI through the latent variables related to general psychopathology and low self-esteem, which in turn predicted dissociative symptoms and body dissatisfaction, respectively (Muehlenkamp, 2011). Furthermore, simple mediation analysis in the context of path analysis in a mixed sample of patients with EDs indicated that childhood sexual abuse was associated with the presence of two or more stressful life events in the 12 months preceding the first ED symptoms, which in turn were associated with the presence of non-suicidal self-injury (Vieira et al., 2018). Furthermore, childhood physical abuse had an indirect effect on non-suicidal self-injury through higher levels of negative self-evaluation, substance use and suicide attempts (Vieira et al., 2018). Finally, Dodd et al. (2022) showed that among women with BN childhood trauma indirectly predicted NSSI through atypical cognitions such as odd thinking, unusual perceptual experiences, and quasi-psychotic thinking.

3.3. The maltreated eco-phenotype of EDs: longitudinal characteristics and treatment outcomes

Several studies investigated whether a history of CM was associated with a distinct longitudinal outcome (e.g. drop-out, diagnostic cross-over, hospitalization), and different response to treatment, according to different therapeutic approaches.

First of all, CM was associated with increased likelihood of *drop-out* in patients with AN or BN, both in patients treated with CBT (Castellini et al., 2018; Mahon et al., 2001a, 2001b) or interpersonal psychotherapy approaches (Mahon et al., 2001a, 2001b). In particular, Castellini et al. (2018) showed that subjects with both childhood abuse and neglect had

the greatest probability of drop-out (Castellini et al., 2018). Furthermore, Carter et al. (2006) evaluated the impact of CM on treatment outcome of an intensive group therapy in patients with AN and showed that patients with AN-bp and childhood sexual abuse terminated treatment earlier as compared to other patients (Carter et al., 2006). However, an overall difference on the rate or timing of premature termination of treatment between patients reporting or not childhood sexual abuse was not observed (Carter et al., 2006).

Regarding the longitudinal trend of *ED symptoms*, several studies are available involving patients with AN or BN treated with CBT. In particular, Cassioli et al. (2022) showed that as the severity of CM increased, the longitudinal trend of ED-psychopathology worsened in patients with AN, and higher baseline levels of emotional dysregulation mediated this effect (Cassioli et al., 2022). Accordingly, Lelli et al. (2019) showed that ED psychopathology did not improve after treatment in patients with BN reporting childhood sexual and/or physical abuse (Lelli et al., 2019). In contrast with these results, Calugi et al. (2018) showed that patients with AN reporting childhood sexual abuse similarly improved in terms of BMI and ED-specific psychopathology as compared to those not reporting it (Calugi et al., 2018), and the same result regarding ED-psychopathology was found in sexually and/or physically abused patients with AN by Lelli et al. (2019) or with both AN or BN by Castellini et al. (2018) (Lelli et al., 2019; Castellini et al., 2018). Moreover, Castellini et al. (2018) and Lelli et al. (2019) showed that traumatized and non-traumatized patients did not differ in terms of remission rates from the ED (Lelli et al., 2019; Castellini et al., 2018). However, Castellini et al. (2018) showed that patients reporting childhood trauma underwent diagnostic cross-over and were hospitalized more frequently (Castellini et al., 2018). In terms of outcomes regarding various eating behaviors, Rienecke et al., 2022 showed that among patients with various ED diagnoses treated with individual and group psychotherapy including trauma specific interventions, those reporting higher adverse childhood experiences had higher binge eating scores at follow-up, but not purging or restrictive behaviors (Rienecke et al., 2022). Concerning patients with BED, Hazzard et al. (2021) demonstrated that lifetime PTSD and childhood abuse were associated with higher binge-eating frequency at follow-up both in patients treated with Integrative Cognitive-Affective Therapy for BED and with CBT administered through guided self-help (Hazzard et al., 2021). The association between childhood abuse and binge-eating frequency at follow-up was moderated by lifetime PTSD, such that childhood abuse was associated with higher binge-eating frequency only in the presence of a history of PTSD (Hazzard et al., 2021).

The negative impact of CM on the longitudinal trend of *general psychopathology and psychiatric comorbidities* was underlined by several studies adopting both CBT and psychodynamic approaches. In particular, for what concerns CBT, Lelli et al. (2019) and Castellini et al. (2018) showed that patients with AN or BN reporting CM did not improve in terms of general psychopathology at follow-up (Lelli et al., 2019; Castellini et al., 2018). Furthermore, Castellini et al. (2018) showed that at follow-up patients with childhood abuse reported greater rates of use of antidepressants and had higher levels of psychiatric comorbidity, specifically a mood disorder comorbidity or obsessive-compulsive disorder (Castellini et al., 2018). In studies of psychodynamic therapy, Strangio et al. (2017) showed that in a mixed sample of patients with various ED diagnosis, those with a history of CM reported a significant decrease only in psychoticism, whereas those not reporting CM had a significant decrease in somatization, obsessive-compulsive symptoms, general psychopathology, phobic anxiety, interoceptive deficits, and dissociation (Strangio et al., 2017). In contrast with these results, Calugi et al. (2018) showed that patients with AN and a history of sexual abuse similarly improved after intensive CBT-E in terms of work, social functioning and general psychopathology as compared to those without a history of childhood sexual abuse (Calugi et al., 2018).

Finally, patients with AN and BN and a history of CM treated with

CBT did not improve in terms of *hypocortisolemia* (Lelli et al., 2019), nor in *sexual functioning*, unlike those without CM (Castellini et al., 2013). Moreover, Castellini et al. (2020) evaluated the impact of CM on the longitudinal trajectory of *amenorrhea* in patients with AN treated with CBT-E, and time-to-event analysis demonstrated that a history of CM predicted an earlier resumption of menses (Castellini et al., 2020b).

For what concerns the *mediators of treatment response* in traumatized patients, Kopland et al. (2023) demonstrated in a sample of patients with AN, BN or OSFED treated with compassion-focused therapy that improvements in ED psychopathology were predictive of enhancements in self-compassion, but only when CM levels were low. Conversely, in the presence of higher CM levels, an inverse relationship between these variables was observed and the reduction in ED symptoms was predicted by improvements in self-compassion (Kopland et al., 2023). Furthermore, Cassioli et al. (2022) demonstrated the pivotal role of emotional dysregulation in the maintenance of higher levels of ED psychopathology in patients treated with CBT-E in the presence of early trauma (Cassioli et al., 2022). Finally, Monteleone et al. (2023) evaluated the *modification of the relationship between variables after treatment* through the comparison of the network structure of psychopathology before and after a CBT-oriented multimodal treatment in underaged patients with AN, BN or OSFED (Monteleone et al., 2023). This study showed that prior to treatment, emotional neglect was connected to life satisfaction both directly and via emotional abuse, feeling disliked by others, and feelings of inferiority and worthlessness (as previously mentioned, see paragraph 3.2.7). Following discharge, only the negative association between emotional neglect and excellent life

conditions persisted (Monteleone et al., 2023).

4. Discussion

The present systematic review provided an accurate description of the evidence about the existence of a maltreated eco-phenotype of EDs and a comprehensive picture of its defining features from a biological, psychopathological, clinical, and treatment response perspective, which are graphically represented in Fig. 2.

Regarding the biological correlates, the majority of available studies concerned neuroanatomical modifications and alterations in the endogenous stress response system. Specifically, neuroimaging studies revealed variations in both gray and white matter, along with reduced cortical thickness and a diminished gyrification index in specific brain regions among individuals with EDs and a history of childhood trauma (see Supplementary Table 1), supporting the hypothesis of the role of CM in influencing the course of brain development (Teicher et al., 2016; Hanson et al., 2010; Kelly et al., 2013). As for the endogenous stress response system, results seem to suggest that CM may determine a functional exhaustion of the HPA axis. Indeed, several studies showed lower urinary and plasmatic cortisol levels, and lower saliva CAR in individuals reporting CM (see Supplementary Table 1). Furthermore, Lelli et al. (2019) showed that these alterations persisted after treatment (Lelli et al., 2019), matching with what was demonstrated in other populations regarding the long-lasting effects of early trauma on the HPA axis (Schär et al., 2022; Van Voorhees and Scarpa, 2004). Moreover, the studies evaluating the emotional and biological reactivity to an




Fig. 2. Graphical summary of main results.

experimental psychosocial stressor (TSST) showed that CM predicted a different emotional reactivity and a diverse response of the HPA axis in this type of context (Monteleone et al., 2021, 2018b), aligning with the hypothesis that childhood traumatic experiences lead to increased interpersonal vulnerability and changes in threat-related social and emotional processing (McLaughlin et al., 2020; Pollak, 2015). Although studies related to other biological correlates of CM in patients with EDs are few, the literature seemed to indicate an association with modifications in inflammation markers, ghrelin levels, fecal short chain fatty acids, and microbiome composition (see [Supplementary Table 1](#)), supporting the role of CM as a factor that impacts the entire organism, including the immune system and the gut-brain axis (Danese et al., 2017; Rohr et al., 2023; Zhang et al., 2022). Moreover, CM has been shown to influence the expression of phenotypic characteristics associated with the presence of specific genes among patients with EDs (see [Supplementary Table 1](#)), including DRD2 and 5-HTTLPR, aligning with the hypothesis of CM as an epigenetic modifier (Pollak, 2015; Parade et al., 2021; Thumfart et al., 2022).

CM was also associated with specific clinical features, including an earlier age at onset, the bingeing/purging subtypes of EDs, and greater severity of pathological eating behaviors (see [Supplementary Table 2](#)). As for the psychopathological aspect, early trauma showed a clear impact on determining greater severity of general psychopathology, more psychiatric comorbidity, including personality disorders, worse psychosocial functioning and quality of life, higher levels of impulsivity, emotional dysregulation, deficits in interoceptive awareness, dysregulated sexuality, risky behaviors such as non-suicidal self-injury, suicidality, and alcohol and substance abuse (see [Supplementary Table 2](#)). Various studies have also highlighted that specific psychological constructs were more represented in patients with a history of CM, attaining both the interpersonal area, such as social insecurity, and the intrapsychic area, such as increased self-criticism and core beliefs of defectiveness/shame (see [Supplementary Table 2](#)).

Data on the severity of ED-specific psychopathology exhibited some inconsistencies, with most studies showing a positive association between CM and worse ED psychopathology, including a poorer body image, but there were also studies that did not confirm this association (see [Supplementary Table 2](#)). Regarding this topic, studies adopting innovative statistical methods allowing an accurate evaluation of the relationship between variables highlighted that the complex associations between psychopathological dimensions and the network structure of ED psychopathology differed in patients reporting CM (see [Supplementary Table 2](#)). In light of this evidence, it appears that the maltreated eco-phenotype of EDs should not be exclusively observed from the quantitative perspective of severity, particularly concerning ED-specific psychopathology. Instead, it should be considered from the viewpoint of a possible distinct psychopathological structure of the disorder.

Regarding the longitudinal outcomes, the literature showed that the maltreated eco-phenotype of EDs is characterized by a worse response to currently proposed treatments, particularly in terms of higher dropout rates, diagnostic cross-over, hospitalization rates, long-term maintenance of higher levels of general psychopathology, psychiatric comorbidity, and sexual dysfunctions (see [Supplementary Table 3](#)). Furthermore, specific psychopathological dimensions were demonstrated to be involved in the maintenance of ED psychopathology in the presence of CM, including emotional dysregulation and low self-compassion (see [Supplementary Table 3](#)). However, data on the course of ED-specific psychopathology over time were somewhat discordant. In particular, some studies adopting the CBT approach showed that the longitudinal trajectory of ED psychopathology was not affected by the presence of CM (Lelli et al., 2019; Castellini et al., 2018; Calugi et al., 2018), while Cassioli et al. (2022) found that as trauma severity increased, treatment response worsened also in the area of ED-specific symptoms (Cassioli et al., 2022). It should be noted that the studies that did not show an impact of CM on the longitudinal course of ED psychopathology did not consider the severity of the experienced

trauma. Furthermore, they neglected the area of emotional abuse in the definition of childhood trauma, considering exclusively that of sexual and/or physical abuse. This is a fundamental fact to consider, given that several studies seemed to suggest that emotional abuse represents the core element of CM involved in modifying the phenotypic characteristics of EDs (Monteleone et al., 2023, 2019b, 2022). Another apparently controversial result was that reported by Castellini et al. (2020) regarding the fact that CM predicted an earlier menses resumption in amenorrheic patients with AN (Castellini et al., 2020b). However, it should be noted that other factors associated with the presence of early trauma and weight recovery also predicted menses resumption, including diagnostic cross-over, making it possible to speculate that in patients with CM menses resumption should not be necessarily viewed as a marker of recovery from the ED, but rather of a transition to a new phase of the disease (Castellini et al., 2020b).

The results of this review underlined that the correlates of CM did not show significant differences across the various ED diagnosis. Furthermore, many of the characteristics of the maltreated eco-phenotype of EDs highlighted in this review were found to be similar to those observed in other psychiatric disorders, including the presence of trauma-related neuroanatomical alterations, modifications of the HPA axis, a higher prevalence of comorbidity, emotional dysregulation, and risky behaviors (Teicher et al., 2022; Teicher and Samson, 2013). Moreover, changes in social information processing as well as high levels of self-criticism and low self-compassion have been consistently acknowledged as transdiagnostic factors conferring heightened risk for multiple types of psychopathology in individuals reporting childhood traumatic experiences (McLaughlin et al., 2020; Lassri and Gewirtz-Meydan, 2022; Sachs-Ericsson et al., 2006; Soffer et al., 2008). Similarly, several studies showed an association of CM not just with a severer clinical presentation (Teicher et al., 2022; Teicher and Samson, 2013), but also with a qualitative modification of the symptomatic expression and network structure of the core psychopathological features of other mental disorders (Medeiros et al., 2020; Wrobel et al., 2023), and with worse responses to standard treatments (Euler et al., 2021; Thomas et al., 2019; Nanni et al., 2012; Schückher et al., 2019). The overlapping of the characteristics of the maltreated eco-phenotype of EDs with those of the maltreated eco-phenotypes of other mental disorders from a biological and clinical point of view, and in terms of longitudinal course, underlined the role of early trauma as a factor which is transdiagnostically associated with specific modifications of the phenotypic expression of mental disorders. These findings align with the hypothesis that CM triggers alterations that are intrinsically tied to the traumatic experience itself, regardless of whether psychopathology subsequently develops or not, and irrespective of the specific disorder the individual may eventually develop (McLaughlin et al., 2020; Bege-mann et al., 2023), which will then depend on the interaction with other risk and resilience factors including genetic and biological components (Teicher and Samson, 2013).

Beyond identifying shared characteristics with the maltreated eco-phenotypes of other psychiatric conditions, the findings of this systematic review allowed to underline distinct features inherent to the maltreated eco-phenotype of EDs. These distinguishing traits seem to arise from the interaction between the biological and psychopathological changes induced by early trauma and the specific psychopathological and clinical features of EDs. For instance, within this context, low self-esteem, shame, and the early maladaptive schemas typical of post-traumatic psychopathology manifested in terms of greater body dissatisfaction (Muehlenkamp et al., 2011; Sweetingham and Waller, 2008) and higher tendency to overvalue the importance of body shapes and weight (Meneguzzo et al., 2021). Furthermore, binge/purge eating behaviors were the behavioral correlate of negative core beliefs (e.g., defectiveness and shame) (Waller et al., 2001). Trauma-related psychopathological dimensions, including emotional dysregulation, affective instability, and the tendency to lose control, were expressed in this populations in terms of an exacerbation of ED-specific psychopathology

(Groleau et al., 2012b; Racine and Wildes, 2015; Cassioli et al., 2022), whereas dissociation predicted binge-eating and purging symptoms (Waller et al., 2001). Feelings of ineffectiveness phenotypically manifested in terms of more severe ED symptoms (Groleau et al., 2012b), specifically greater body dissatisfaction and bulimia (Monteleone et al., 2019b, 2022), whereas depressive symptoms were linked to greater drive for thinness (Kong and Bernstein, 2009), body dissatisfaction (Kong and Bernstein, 2009), overeating (Rodgers et al., 2019), and bingeing/vomiting behaviors (Waller et al., 2001). Finally, difficulties in interoceptive awareness correlated with drive for thinness (Monteleone et al., 2019b), bulimic symptoms (Monteleone et al., 2019b, 2022), and body dissatisfaction (Monteleone et al., 2022). It is also noteworthy that in the area of sexuality aspects of post-traumatic and ED-specific psychopathology intertwined in a unique way (Castellini et al., 2012, 2013, 2020a). Furthermore, these findings highlighted a specific interplay between trauma-induced biological changes and those resulting from malnutrition and disordered eating behaviors (see [Supplementary Table 1](#)). A notable example of this was the elevation of ghrelin levels (Rossi et al., 2021), which increase in response to both stress and purging behaviors, as well as the alterations in the endogenous stress-response system, particularly within the HPA axis, which have been observed in response to both early trauma and underweight conditions. Furthermore, the pattern of menstrual alterations found in these patients is typical of the maltreated eco-phenotype of EDs and not of what was observed in other areas of mental health (Castellini et al., 2020b). Finally, the evidence related to changes in the gut microbiota seems to suggest that they may be attributable not only to the effects of early trauma but also to those of disordered eating (Castellini et al., 2023a). Regarding the neuroanatomical aspects, studies in this field are still preliminary. However, it is intriguing to note that the brain areas implicated in trauma-related modifications in patients with EDs have been shown to be involved with functions such as higher-order visual perception, emotional regulation processes, self-awareness, brain reward systems, the recognition and interpretation of body image-related information, the processing of stimuli from one's own body, and taste processing (Cascino et al., 2023a, 2023b; Monteleone et al., 2019a). Future studies will need to clarify to what extent the neuroanatomical alterations observed in patients with EDs are specific to this population as opposed to others, and whether they may be involved in promoting the development of ED-specific psychopathology as opposed to other types of psychopathology in individuals exposed to CM. Thoroughly understanding the interaction between post-traumatic psychopathological and biological features and ED-specific expressions as they manifest in the maltreated eco-phenotype of EDs is crucial to ensure the identification and treatment of both aspects appropriately. Focusing solely on the non-specific aspects, which are common to all maltreated eco-phenotypes, would indeed cause to overlook those post-traumatic manifestations that are expressed through the specific features of the disorder.

Overall, this review indicates that the wounds of childhood trauma can create both biological and relational vulnerability, potentially triggering psychopathological imbalances which may not merely manifest as a generic ED, but rather as a distinct disease eco-phenotype, characterized by unique psychopathological and biological expressions, and by a specific longitudinal trajectory. Therefore, this findings support the hypothesis proposed by Teicher et al. regarding the need to modify the approach to mental disorders from a clinical, research, and nosographic perspective, moving towards considering early trauma as a fundamental phenotypic and, consequently, diagnostic modifier (Teicher et al., 2022; Teicher and Samson, 2013). Specifically, Teicher et al. recommended a moderate revision of the DSM consisting in the addition of the new diagnostic specifier “with maltreatment history” or “with early life stress” to the primary DSM diagnosis when a history of early trauma is present as a first step to acknowledge the critical role of CM as a phenotypic modifier of psychiatric disorders and to move towards the development of an etiologically informed classification system (Teicher et al., 2022;

Teicher and Samson, 2013). Adopting this perspective in EDs would allow for the identification of the specific needs of patients reporting CM, improving clinical management. Furthermore, it would improve research in this field, allowing to stratify patient groups according to the presence or not of CM and therefore making it more targeted. Finally, it would enable the adoption of trauma-focused treatments when a post-traumatic etiopathogenesis is recognized. Recently, Rossi et al. (2024) demonstrated that the add-on of Eye Movement Desensitization and Reprocessing (EMDR), a treatment originally developed for PTSD, to CBT-E in patients with AN and a history of CM improved treatment response, not just in terms of ED-specific psychopathology and weight recovery, but also of reduction of general psychopathology and dissociative symptoms (Rossi et al., 2024). Furthermore, the RCT performed by Trottier et al. (2022) proved the benefits of an integrated ED-PTSD approach comprehending cognitive processing therapy for PTSD (Resick et al., 2017) and CBT-E interventions in terms of improvements in post-traumatic symptoms and anxiety in patients with EDs and comorbid PTSD as compared to standard CBT (Trottier et al., 2022). Furthermore, potential innovative pharmacological interventions targeting biological maintenance mechanisms, such as alterations in the stress-response system or gut microbiota, could be studied in order to improve the prognosis of patients reporting a history of CM. A trauma-informed nosography would be particularly important in the field of EDs, given the poor response to standard treatments (Atwood and Friedman, 2020; Halmi, 2013; Miskovic-Wheatley et al., 2023), the burden of these diseases (Castellini et al., 2023b; Van Hoeken and Hoek, 2020), and the consequent need to find new therapeutic strategies that move towards personalized approaches.

Despite the large number of studies included in this systematic review, it is necessary to underline the limitations of the literature on this subject. First, all the included studies evaluated female populations, with only a very small minority including male subjects. Moreover, only a few studies considered underage populations, or patients with BED, and studies related to the biological correlates of the maltreated eco-phenotype are still scarce. There was a general lack of reporting and consideration of sociodemographic factors in the studies reviewed, underscoring the need for further research in this area. Considering that the age of first exposure appears to modulate the impact of CM on mental health (Teicher et al., 2016; Kaplow and Widom, 2007; Dunn et al., 2023), this remains an unresolved issue in the field of the maltreated eco-phenotype in EDs due to the lack of data. Most of the studies included were cross-sectional, and the longitudinal ones were still mostly observational in design. However, while this generally means that the evidence reported should be considered of low quality, it does not mean that they cannot provide valuable insights, and it is important to make some considerations. In the field of CM correlates, especially in a single diagnostic area like EDs, it would be unreasonable to expect studies such as RCTs, beyond studies on treatment outcomes. Therefore, in this context, high-quality evidence would typically mean that the study design minimizes bias, using appropriate statistical methods controlling for potential confounding variables, and using recognized diagnostic criteria and psychometric tools. As indicated by the quality assessment, these points were present in most of the included studies. Moreover, many of the findings were confirmed by multiple studies, which strengthens the level of evidence. On the contrary, results reported by a single study or those on which there was no agreement in different reports should be considered more preliminary, pending future data on the subject. Regarding longitudinal studies on treatment outcomes, it is advisable that future RCTs in the field of EDs consider the presence or absence of CM as a moderator of outcome. It would also be essential to expand research with longitudinal studies to more thoroughly investigate the psychopathological dimensions that are associated with the maintenance of symptoms over time in the presence of a history of trauma, and to identify increasingly specific targets for structuring innovative treatments. Finally, many studies, especially the less recent ones, considered only childhood sexual/physical abuse,

neglecting the importance of emotional abuse and neglect. However, more recent studies have overcome this limitation. It might also be relevant to investigate the impact of other types of early trauma, such as bereavement. Adverse childhood events different from CM could lead to a distinct traumatized eco-phenotype with characteristics that are more or less overlapping with those of the maltreated one, which was the subject of this review.

The present study shows various strengths, particularly the investigation of a wide range of correlates of childhood trauma through a very inclusive research string, which allowed the inclusion of a large number of studies and thus a comprehensive description of the maltreated eco-phenotype of EDs from various perspectives. In addition, the screening of papers was extensively discussed among multiple researchers, allowing for accurate verification of the extrapolated data. However, it also has some weaknesses, notably that the research was confined to two databases (PubMed and ClinicalTrials.gov), and only reports published in English were considered. Moreover, negative results that did not represent the primary outcome of the study were excluded, as they did not provide data for which an adequate power calculation had been performed. While this represents a strength in terms of statistical accuracy, it also may introduce a bias related to giving more relevance to positive results.

In conclusion, this systematic review provided, for the first time, an accurate and comprehensive description of the maltreated eco-phenotype of EDs. It highlighted the importance of moving towards an etiologically informed nosography that considers the role of early trauma not just as a risk factor for the development of these disorders, but also as an etiologic agent capable of producing eco-phenotypic variants of EDs from the multifaceted perspectives of biology, psychopathology, and longitudinal course. Recognizing the importance of CM as a phenotypic modifier would improve the clinical approach to EDs, facilitate and empower research in this field, and guide the development of new therapeutic strategies that consider the specific needs and psychopathological features of individuals reporting a history of childhood trauma.

Funding sources

None.

Author statement

Conceptualization: E.R., E.C. and G.C.; Data curation: E.R., E.C., C.D. and G.M.; Investigation: E.R., E.C., C.D. and G.M.; Methodology: E.R. and E.C.; Project administration: E.R. and E.C.; Resources: E.R. and E.C.; Supervision: E.R. and E.C.; Validation: E.R., E.C., A.M.M., S.A.W., V.R. and G.C.; Visualization: E.R. and E.C.; Writing – original draft: E.R. and E.C.; Writing - review & editing: E.R., E.C., C.D., G.M., A.M.M., S.A.W., V.R. and G.C.

Data availability

Data will be made available on request.

Acknowledgments

None.

Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at [doi:10.1016/j.neubiorev.2024.105619](https://doi.org/10.1016/j.neubiorev.2024.105619).

References

- American Psychiatric Association. The Diagnostic and Statistical Manual of Mental Disorders (4th Edition). (American Psychiatric Association, Washington, DC, 1994).
- American Psychiatric Association, 2022. Diagnostic and Statistical Manual of Mental Disorders. American Psychiatric Association Publishing. <https://doi.org/10.1176/appi.books.9780890425787>.
- Arnold, S., Correll, C.U., Jaite, C., 2023. Frequency and correlates of lifetime suicidal ideation and suicide attempts among consecutively hospitalized youth with anorexia nervosa and bulimia nervosa: results from a retrospective chart review. *Bord. Personal. Disord. Emot. Dysregulation* 10, 10.
- Atwood, M.E., Friedman, A., 2020. A systematic review of enhanced cognitive behavioral therapy (CBT-E) for eating disorders. *Int. J. Eat. Disord.* 53, 311–330.
- Begemann, M.J.H., et al., 2023. Childhood trauma is associated with reduced frontal gray matter volume: a large transdiagnostic structural MRI study. *Psychol. Med.* 53, 741–749.
- Belsky, J., Pluess, M., 2013. Beyond risk, resilience, and dysregulation: phenotypic plasticity and human development. *Dev. Psychopathol.* 25, 1243–1261.
- Borg, S.L., et al., 2022. Relationships between childhood abuse and eating pathology among individuals with binge-eating disorder: examining the moderating roles of self-discrepancy and self-directed style. *Eat. Disord.* 30, 355–369.
- Bou Khalil, R., et al., 2022. Neutrophil-to-lymphocyte ratio (NLR) variations in relationship with childhood maltreatment in patients with anorexia nervosa: a retrospective cohort study. *Eat. Weight Disord. - Stud. Anorex. Bulim. Obes.* 27, 2201–2212.
- Brewerton, T.D., 2019. An overview of trauma-informed care and practice for eating disorders. *J. Aggress. Maltreatment Trauma* 28, 445–462.
- Brewerton, T.D., et al., 2021. Provisional posttraumatic stress disorder is associated with greater severity of eating disorder and comorbid symptoms in adolescents treated in residential care. *Eur. Eat. Disord. Rev.* 29, 910–923.
- Calugi, S., et al., 2018. Anorexia nervosa and childhood sexual abuse: treatment outcomes of intensive enhanced cognitive behavioural therapy. *Psychiatry Res* 262, 477–481.
- Carretero-García, A., et al., 2012. Repeated traumatic experiences in eating disorders and their association with eating symptoms. *Eat. Weight Disord. Stud. Anorex. Bulim. Obes.* 17, e267–e273.
- Carter, J.C., Bewell, C., Blackmore, E., Woodside, D.B., 2006. The impact of childhood sexual abuse in anorexia nervosa. *Child Abuse. Negl.* 30, 257–269.
- Cascino, G., et al., 2021. How is the history of early traumatic exposure associated with the psychopathological outcomes of COVID-19 related lockdown and subsequent re-opening in people with eating disorders? *Front. Psychiatry* 12, 789344.
- Cascino, G., et al., 2023a. Childhood maltreatment is associated with cortical thinning in people with eating disorders. *Eur. Arch. Psychiatry Clin. Neurosci.* 273, 459–466.
- Cascino, G., et al., 2023b. Association between childhood maltreatment and cortical folding in women with eating disorders. *Eur. J. Neurosci.* 58, 2868–2873.
- Cascino, G., Monteleone, A.M., 2024. Early traumatic experiences and the hypothalamus-pituitary-adrenal axis in people with eating disorders: A narrative review. *Psychoneuroendocrinology* 159, 106665.
- Cassoli, E., et al., 2020. Sexuality, embodiment and attachment style in anorexia nervosa. *Eat. Weight Disord. Stud. Anorex. Bulim. Obes.* 25, 1671–1680.
- Cassoli, E., et al., 2022. A 1-year follow-up study of the longitudinal interplay between emotion dysregulation and childhood trauma in the treatment of anorexia nervosa. *Int. J. Eat. Disord.* 55, 98–107.
- Castellini, G., et al., 2012. Childhood abuse, sexual function and cortisol levels in eating disorders. *Psychother. Psychosom.* 81, 380–382.
- Castellini, G., et al., 2013. Childhood sexual abuse moderates the relationship between sexual functioning and eating disorder psychopathology in Anorexia Nervosa and Bulimia Nervosa: A 1-year follow-up. *Study J. Sex. Med.* 10, 2190–2200.
- Castellini, G., et al., 2018. Different outcomes, psychopathological features, and comorbidities in patients with eating disorders reporting childhood abuse: a 3-year follow-up study. *Eur. Eat. Disord. Rev.* 26, 217–229.
- Castellini, G., et al., 2020a. The impact of COVID-19 epidemic on eating disorders: a longitudinal observation of pre versus post psychopathological features in a sample of patients with eating disorders and a group of healthy controls. *Int. J. Eat. Disord.* 53, 1855–1862.
- Castellini, G., et al., 2020b. Predictors of resumption of menses in Anorexia Nervosa: a 4-Year longitudinal study. *Psychosom. Med.* 82, 782–786.
- Castellini, G., et al., 2020a. Dysregulated sexuality in women with eating disorders: the role of childhood traumatic experiences. *J. Sex. Marital Ther.* 46, 793–806.
- Castellini, G., et al., 2023a. Gut microbiota metabolites mediate the interplay between childhood maltreatment and psychopathology in patients with eating disorders. *Sci. Rep.* 13, 11753.
- Castellini, G., et al., 2023b. Use and misuse of the emergency room by patients with eating disorders in a matched-cohort analysis: What can we learn from it? *Psychiatry Res* 328, 115427.
- Chami, R., Monteleone, A.M., Treasure, J., Monteleone, P., 2019. Stress hormones and eating disorders. *Mol. Cell. Endocrinol.* 497, 110349.
- Corstorphine, E., Waller, G., Lawson, R., Ganis, C., 2007. Trauma and multi-impulsivity in the eating disorders. *Eat. Behav.* 8, 23–30.
- Dalgleish, T., et al., 2003. Self-reported parental abuse relates to autobiographical memory style in patients with eating disorders. *Emotion* 3, 211–222.
- Danese, A., Lewis, J.S., 2017. Psychoneuroimmunology of early-life stress: the hidden wounds of childhood Trauma? *Neuropsychopharmacology* 42, 99–114.
- Day, S., Hay, P., Tannous, Wadad K., Fatt, S.J., Mitchison, D., 2023. A systematic review of the effect of PTSD and trauma on treatment outcomes for eating disorders. *Trauma Violence Abus.* 152483802311673. <https://doi.org/10.1177/15248380231167399>.

- Díaz-Marsá, M., et al., 2007. Findings with 0.25 mg dexamethasone suppression test in eating disorders: *Association with Childhood Trauma*. *CNS Spectr.* 12, 675–680.
- Dodd, D.R., Crosby, R.D., Cao, L., Gordon, K.H., Wonderlich, S.A., 2022. Borderline personality disorder symptoms as mediational mechanisms linking childhood trauma and nonsuicidal SELF-INJURY among women with bulimia nervosa. *Int. J. Eat. Disord.* 55, 372–381.
- Dohm, F.-A., et al., 2002. Self-harm and substance use in a community sample of Black and White women with binge eating disorder or bulimia nervosa. *Int. J. Eat. Disord.* 32, 389–400.
- Dunn, E., et al., 2023. Sensitive periods for the effect of child maltreatment on psychopathology symptoms in adolescence. *Complex Psychiatry* 9, 145–153.
- Euler, S., et al., 2021. Impact of childhood maltreatment in borderline personality disorder on treatment response to intensive. *Dialect. Behav. Ther. J. Personal. Disord.* 35, 428–446.
- Favaro, A., et al., 2004. The relationship between temperament and impulsive behaviors in eating disordered subjects. *Eat. Disord.* 13, 61–70.
- Favaro, A., et al., 2008. Self-injurious behavior and attempted suicide in purging bulimia nervosa: associations with psychiatric comorbidity. *J. Affect. Disord.* 105, 285–289.
- Favaro, A., Santonastaso, P., 1997. Suicidality in eating disorders: clinical and psychological correlates. *Acta Psychiatr. Scand.* 95, 508–514.
- Favaro, A., Santonastaso, P., 1998. Impulsive and compulsive self-injurious behavior in bulimia nervosa: prevalence and psychological correlates. *J. Nerv. Ment. Dis.* 186, 157–165.
- Favaro, A., Santonastaso, P., 1999. Different types of self-injurious behavior in bulimia nervosa. *Compr. Psychiatry* 40, 57–60.
- Favaro, A., Santonastaso, P., 2000. Self-injurious behavior in anorexia nervosa. *J. Nerv. Ment. Dis.* 188, 537–542.
- Favaro, A., Grave, R.D., Santonastaso, P., 1998. Impact of a history of physical and sexual abuse in eating disordered and asymptomatic subjects. *Acta Psychiatr. Scand.* 97, 358–363.
- Fennig, S., Hadas, A., 2010. Suicidal behavior and depression in adolescents with eating disorders. *Nord. J. Psychiatry* 64, 32–39.
- Grilo, C.M., Masheb, R.M., 2002. Childhood maltreatment and personality disorders in adult patients with binge eating disorder: maltreatment and personality disorders. *Acta Psychiatr. Scand.* 106, 183–188.
- Groleau, P., et al., 2012a. Dopamine-system genes, childhood abuse, and clinical manifestations in women with Bulimia-spectrum. *Disord. J. Psychiatr. Res.* 46, 1139–1145.
- Groleau, P., et al., 2012b. Childhood emotional abuse and eating symptoms in bulimic disorders: An examination of possible mediating variables: Childhood Emotional Abuse and Bulimic Disorders. *Int. J. Eat. Disord.* 45, 326–332.
- Groleau, P., et al., 2014. Methylation of the dopamine D2 receptor (DRD2) gene promoter in women with a bulimia-spectrum disorder: Associations with borderline personality disorder and exposure to childhood abuse. *J. Psychiatr. Res.* 48, 121–127.
- Groth, T., Hilsenroth, M., Boccio, D., Gold, J., 2020. Relationship between trauma history and eating disorders in adolescents. *J. Child Adolesc. Trauma* 13, 443–453.
- Guillaume, S., et al., 2016. Associations between adverse childhood experiences and clinical characteristics of eating disorders. *Sci. Rep.* 6, 35761.
- Halmi, K.A., 2013. Perplexities of treatment resistance in eating disorders. *BMC Psychiatry* 13, 292.
- Hanson, J.L., et al., 2010. Early stress is associated with alterations in the orbitofrontal cortex: a tensor-based morphometry investigation of brain structure and behavioral risk. *J. Neurosci.* 30, 7466–7472.
- Hazzard, V.M., et al., 2021. Treatment outcomes of psychotherapy for binge-eating disorder in a randomized controlled trial: Examining the roles of childhood abuse and post-traumatic stress disorder. *Eur. Eat. Disord. Rev.* 29, 611–621.
- Hebert, R.S., Wright, S.M., Dittus, R.S., Elasy, T.A., 2002. Prominent medical journals often provide insufficient information to assess the validity of studies with negative results. *J. Negat. Results Biomed.* 1, 1.
- Kaplow, J.B., Widom, C.S., 2007. Age of onset of child maltreatment predicts long-term mental health outcomes. *J. Abnorm. Psychol.* 116, 176–187.
- Kelly, P.A., et al., 2013. Cortical thickness, surface area, and gyrification abnormalities in children exposed to maltreatment: neural markers of vulnerability? *Biol. Psychiatry* 74, 845–852.
- Kjaersdam Tellús, G., Lauritsen, M.B., Rodrigo-Domingo, M., 2021. Prevalence of various traumatic events including sexual trauma in a clinical sample of patients with an eating disorder. *Front. Psychol.* 12, 687452.
- Kong, S., Bernstein, K., 2009. Childhood trauma as a predictor of eating psychopathology and its mediating variables in patients with eating disorders. *J. Clin. Nurs.* 18, 1897–1907.
- Kopland, M.C.G., Vrabel, K., Melsom, L., Hoffart, A., Johnson, S.U., 2023. Self-compassion in eating disorders and childhood trauma: A study of within-person effects in a randomized controlled trial. *Psychother. Res.* 33, 640–653.
- Lassri, D., Gewirtz-Meydan, A., 2022. Self-Compassion Moderates the Mediating Effect of Self-Criticism in the Link Between Childhood Maltreatment and Psychopathology. *J. Interpers. Violence* 37, NP21699–NP21722.
- Latzer, Y., et al., 2020. Childhood maltreatment in patients with binge eating disorder with and without night eating syndrome vs. control. *Psychiatry Res* 293, 113451.
- Lelli, L., Castellini, G., Cassioli, E., Monteleone, A.M., Ricca, V., 2019. Cortisol levels before and after cognitive behavioural therapy in patients with eating disorders reporting childhood abuse: a follow-up study. *Psychiatry Res* 275, 269–275.
- Leraas, B.C., et al., 2018. Affect-based profiles of bulimia nervosa: the utility and validity of indicators assessed in the natural environment. *Psychiatry Res* 259, 210–215.
- Lockwood, R., Lawson, R., Waller, G., 2004. Compulsive features in the eating disorders: a role for trauma? *J. Nerv. Ment. Dis.* 192, 247–249.
- Longo, P., Marzola, E., 2020. De Bacco, C., Demarchi, M. & Abbate-Daga, G. Young patients with anorexia nervosa: the contribution of post-traumatic stress disorder and traumatic events. *Medicina* 57, 2.
- Longo, P., Bertorello, A., Panero, M., Abbate-Daga, G., Marzola, E., 2019. Traumatic events and post-traumatic symptoms in anorexia nervosa. *Eur. J. Psychotraumatology* 10, 1682930.
- Longo, P., Marzola, E., Martini, M., Amodeo, L., Abbate-Daga, G., 2023. Anorexia nervosa and somatoform dissociation: a neglected body-centered perspective. *J. Trauma Dissociation* 24, 141–156.
- Mahon, J., Bradley, S.N., Harvey, P.K., Winston, A.P., Palmer, R.L., 2001a. Childhood trauma has dose-effect relationship with dropping out from psychotherapeutic treatment for bulimia nervosa: a replication. *Int. J. Eat. Disord.* 30, 138–148.
- Mahon, J., Winston, A.P., Palmer, R.L., Harvey, P.K., 2001b. Do broken relationships in childhood relate to bulimic women breaking off psychotherapy in adulthood? *Int. J. Eat. Disord.* 29, 139–149.
- Marciello, F., et al., 2020. Early traumatic experiences and eating disorders: a focus on the endogenous stress response system. *J. Psychopathol.* 26, 77–84.
- Mayes, S.D., et al., 2014. Correlates of suicide ideation and attempts in children and adolescents with eating disorders. *Eat. Disord.* 22, 352–366.
- McLaughlin, K.A., et al., 2016. Maltreatment exposure, brain structure, and fear conditioning in children and adolescents. *Neuropsychopharmacology* 41, 1956–1964.
- McLaughlin, K.A., Colich, N.L., Rodman, A.M., Weissman, D.G., 2020. Mechanisms linking childhood trauma exposure and psychopathology: a transdiagnostic model of risk and resilience. *BMC Med* 18, 96.
- Medeiros, G.C., et al., 2020. Childhood maltreatment and impact on clinical features of major depression in adults. *Psychiatry Res* 293, 113412.
- Meneguzzo, P., et al., 2021. Associations between trauma, early maladaptive schemas, personality traits, and clinical severity in eating disorder patients: a clinical presentation and mediation analysis. *Front. Psychol.* 12, 661924.
- Meneguzzo, P., et al., 2022. Urinary free cortisol and childhood maltreatments in eating disorder patients: new evidence for an ecophenotype subgroup. *Eur. Eat. Disord. Rev.* 30, 364–372.
- Miskovic-Wheatley, J., et al., 2023. Eating disorder outcomes: findings from a rapid review of over a decade of research. *J. Eat. Disord.* 11, 85.
- Mitchell, K.S., Sciolò, E.R., Galovski, T., Belfer, P.L., Cooper, Z., 2021. Posttraumatic stress disorder and eating disorders: maintaining mechanisms and treatment targets. *Eat. Disord.* 29, 292–306.
- Modesti, P.A., et al., 2016. Panethnic Differences in Blood Pressure in Europe: a systematic review and meta-analysis. *PLoS One* 11, e0147601.
- Molendijk, M.L., Hoek, H.W., Brewerton, T.D., Elzinga, B.M., 2017. Childhood maltreatment and eating disorder pathology: a systematic review and dose-response meta-analysis. *Psychol. Med.* 47, 1402–1416.
- Molina-Ruiz, R.M., et al., 2022. Striatal volumes as potential biomarkers in eating disorders: a pilot study. *Rev. Psiquiatr. Salud Ment.* 15, 65–73.
- Monteleone, A.M., et al., 2015. Childhood trauma and cortisol awakening response in symptomatic patients with anorexia nervosa and bulimia nervosa: childhood Trauma and HPA Axis in Eating Disorders. *Int. J. Eat. Disord.* 48, 615–621.
- Monteleone, A.M., et al., 2018a. Impaired cortisol awakening response in eating disorder women with childhood trauma exposure: evidence for a dose-dependent effect of the traumatic load. *Psychol. Med.* 48, 952–960.
- Monteleone, A.M., et al., 2018b. Deranged emotional and cortisol responses to a psychosocial stressor in anorexia nervosa women with childhood trauma exposure: evidence for a "maltreated ecophenotype". *J. Psychiatr. Res.* 104, 39–45.
- Monteleone, A.M., et al., 2019b. The association between childhood maltreatment and eating disorder psychopathology: a mixed-model investigation. *Eur. Psychiatry* 61, 111–118.
- Monteleone, A.M., et al., 2019a. The effects of childhood maltreatment on brain structure in adults with eating disorders. *World J. Biol. Psychiatry* 20, 301–309.
- Monteleone, A.M., et al., 2020b. Parental bonding, childhood maltreatment and eating disorder psychopathology: an investigation of their interactions. *Eat. Weight Disord. Stud. Anorex. Bulim. Obes.* 25, 577–589.
- Monteleone, A.M., et al., 2020a. Early traumatic experiences impair the functioning of both components of the endogenous stress response system in adult people with eating disorders. *Psychoneuroendocrinology* 115, 104644.
- Monteleone, A.M., et al., 2021. Emotional traumatic experiences significantly contribute to identify a maltreated ecophenotype sub-group in eating disorders: Experimental evidence. *Eur. Eat. Disord. Rev.* 29, 269–280.
- Monteleone, A.M., et al., 2022. The connection between childhood maltreatment and eating disorder psychopathology: a network analysis study in people with bulimia nervosa and with binge eating disorder. *Eat. Weight Disord. - Stud. Anorex. Bulim. Obes.* 27, 253–261.
- Monteleone, A.M., et al., 2023. Pathways between childhood maltreatment and life satisfaction in adolescents with eating disorders: a network analysis. *Eur. Eat. Disord. Rev.* 31, 724–733.
- Muehlenkamp, J.J., Claes, L., Smits, D., Peat, C.M., Vandereycken, W., 2011. Nonsuicidal self-injury in eating disordered patients: a test of a conceptual model. *Psychiatry Res* 188, 102–108.
- Nagata, T., Kiriike, N., Iketani, T., Kawarada, Y., Tanaka, H., 1999. History of childhood sexual or physical abuse in Japanese patients with eating disorders: relationship with dissociation and impulsive behaviours. *Psychol. Med.* 29, 935–942.
- Nanni, V., Uher, R., Danese, A., 2012. Childhood maltreatment predicts unfavorable course of illness and treatment outcome in depression: a meta-analysis. *Am. J. Psychiatry* 169, 141–151.

- Nickel, M.K., et al., 2006. Familial and sociopsychopathological risk factors for suicide attempt in bulimic and in depressed women: Prospective study. *Int. J. Eat. Disord.* 39, 410–417.
- Page, M.J., et al., 2021. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* n71. <https://doi.org/10.1136/bmj.n71>.
- Parade, S.H., et al., 2021. A systematic review of childhood maltreatment and DNA methylation: candidate gene and epigenome-wide approaches. *Transl. Psychiatry* 11, 134.
- Pollak, S.D., 2015. Multilevel developmental approaches to understanding the effects of child maltreatment: recent advances and future challenges. *Dev. Psychopathol.* 27, 1387–1397.
- Pugh, M., Waller, G., Esposito, M., 2018. Childhood trauma, dissociation, and the internal eating disorder 'voice'. *Child Abus. Negl.* 86, 197–205.
- Rabito-Alcón, M.F., Baile, J.I., Vanderlinden, J., 2021. Mediating factors between childhood traumatic experiences and eating disorders development: a systematic review. *Children* 8, 114.
- Racine, S.E., Wildes, J.E., 2015. Emotion dysregulation and anorexia nervosa: an exploration of the role of childhood abuse: Emotion Regulation. *Child. Abus., Anorex. Int. J. Eat. Disord.* 48, 55–58.
- Resick, P.A., Monson, C.M., Chard, K.M., 2017. Cognitive Processing Therapy for PTSD: A Comprehensive Manual. Guilford Press, New York.
- Richardson, J., et al., 2008. Relevance of the 5-HTTLPR polymorphism and childhood abuse to increased psychiatric comorbidity in women with bulimia-spectrum disorders. *J. Clin. Psychiatry* 69, 981–990.
- Rienecke, R.D., et al., 2022. Adverse childhood experiences among a treatment-seeking sample of adults with eating disorders. *Eur. Eat. Disord. Rev.* 30, 156–167.
- Rodgers, M., et al., 2009. Testing methodological guidance on the conduct of narrative synthesis in systematic reviews: effectiveness of interventions to promote smoke alarm ownership and function. *Evaluation* 15, 49–73.
- Rodgers, R.F., et al., 2019. Structural differences in eating disorder psychopathology after history of childhood abuse: Insights from a Bayesian network analysis. *J. Abnorm. Psychol.* 128, 795–805.
- Rodríguez-Quiroga, A., MacDowell, K.S., Leza, J.C., Carrasco, J.L., Díaz-Marsá, M., 2021. Childhood trauma determines different clinical and biological manifestations in patients with eating disorders. *Eat. Weight Disord. - Stud. Anorex. Bulim. Obes.* 26, 847–857.
- Rohr, J.C., et al., 2023. History of childhood physical abuse is associated with gut microbiota diversity among adult psychiatric inpatients. *J. Affect. Disord.* 331, 50–56.
- Rossi, E., et al., 2021. Ghrelin as a possible biomarker and maintaining factor in patients with eating disorders reporting childhood traumatic experiences. *Eur. Eat. Disord. Rev.* 29, 588–599.
- Rossi, E., et al., 2024. Eye movement desensitisation and reprocessing as add-on treatment to enhanced cognitive behaviour therapy for patients with anorexia nervosa reporting childhood maltreatment: A quasi-experimental multicenter study. *European eating disorders review : the journal of the Eating Disorders Association* 32 (2), 322–337. <https://doi.org/10.1002/erv.3044>.
- Sachs-Ericsson, N., Verona, E., Joiner, T., Preacher, K.J., 2006. Parental verbal abuse and the mediating role of self-criticism in adult internalizing disorders. *J. Affect. Disord.* 93, 71–78.
- Santonastaso, P., Zanetti, T., 2006. De Antoni, C., Tenconi, E. & Favaro, A. Anorexia nervosa patients with a prior history of bulimia nervosa. *Compr. Psychiatry* 47, 519–522.
- Schaefer, L.M., et al., 2021. Examining the roles of emotion dysregulation and impulsivity in the relationship between psychological trauma and substance abuse among women with bulimic-spectrum pathology. *Eat. Disord.* 29, 276–291.
- Schär, S., Mürmer-Lavanchy, I., Schmidt, S.J., Koenig, J., Kaess, M., 2022. Child maltreatment and hypothalamic-pituitary-adrenal axis functioning: A systematic review and meta-analysis. *Front. Neuroendocrinol.* 66, 100987.
- Schückler, F., Sellin, T., Engström, I., Berglund, K., 2019. History of childhood abuse is associated with less positive treatment outcomes in socially stable women with alcohol use disorder. *BMC Women's Health* 19, 159.
- Smith, C.E., et al., 2016. Is childhood trauma associated with lifetime suicide attempts in women with bulimia nervosa? *Eat. Weight Disord. Stud. Anorex. Bulim. Obes.* 21, 199–204.
- Soffer, N., Gilboa-Schechtman, E., Shahar, G., 2008. The relationship of childhood emotional abuse and neglect to depressive vulnerability and low self-efficacy. *Int. J. Cogn. Ther.* 1, 151–162.
- Speranza, M., et al., 2003. Depressive psychopathology and adverse childhood experiences in eating disorders. *Eur. Psychiatry* 18, 377–383.
- Spiegel, J., et al., 2022. Emotional abuse interacts with borderline personality in adolescent inpatients with binge-purging eating disorders. *Eat. Weight Disord. Stud. Anorex. Bulim. Obes.* 27, 131–138.
- Steiger, H., et al., 2001. Association of serotonin and cortisol indices with childhood abuse in bulimia nervosa. *Arch. Gen. Psychiatry* 58, 837.
- Steiger, H., et al., 2004. Serotonin function, personality-trait variations, and childhood abuse in women with bulimia-spectrum eating disorders. *J. Clin. Psychiatry* 65, 830–837.
- Steiger, H., et al., 2007. The 5HTTLPR polymorphism, prior maltreatment and dramatic-erratic personality manifestations in women with bulimic syndromes. *J. Psychiatry Neurosci.* JPN 32, 354–362.
- Steiger, H., et al., 2008. Dissocial behavior, the 5HTTLPR polymorphism, and maltreatment in women with bulimic syndromes. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* 147B, 128–130.
- Steiger, H., Labonté, B., Groleau, P., Turecki, G., Israel, M., 2013. Methylation of the glucocorticoid receptor gene promoter in bulimic women: Associations with borderline personality disorder, suicidality, and exposure to childhood abuse. *Int. J. Eat. Disord.* 46, 246–255.
- Stein, D., et al., 2013. Attitudes toward life and death and suicidality among inpatient female adolescents with eating. *Disord. J. Nerv. Ment. Dis.* 201, 1066–1071.
- Strangio, A.M., et al., 2017. The effect of abuse history on adolescent patients with feeding and eating disorders treated through psychodynamic therapy: comorbidities and outcome. *Front. Psychiatry* 8.
- Struck, N., et al., 2020. Childhood maltreatment and adult mental disorders – the prevalence of different types of maltreatment and associations with age of onset and severity of symptoms. *Psychiatry Res* 293, 113398.
- Sweetingham, R., Waller, G., 2008. Childhood experiences of being bullied and teased in the eating disorders. *Eur. Eat. Disord. Rev.* 16, 401–407.
- Tasca, G.A., et al., 2013. Attachment insecurity mediates the relationship between childhood trauma and eating disorder psychopathology in a clinical sample: a structural equation model. *Child Abus. Negl.* 37, 926–933.
- Teicher, M.H., Samson, J.A., 2013. Childhood maltreatment and psychopathology: a case for ecophenotypic variants as clinically and neurobiologically distinct subtypes. *Am. J. Psychiatry* 170, 1114–1133.
- Teicher, M.H., Samson, J.A., Anderson, C.M., Ohashi, K., 2016. The effects of childhood maltreatment on brain structure, function and connectivity. *Nat. Rev. Neurosci.* 17, 652–666.
- Teicher, M.H., Gordon, J.B., Nemeroff, C.B., 2022. Recognizing the importance of childhood maltreatment as a critical factor in psychiatric diagnoses, treatment, research, prevention, and education. *Mol. Psychiatry* 27, 1331–1338.
- Thomas, S., Höfler, M., Schäfer, I., Trautmann, S., 2019. Childhood maltreatment and treatment outcome in psychotic disorders: a systematic review and meta-analysis. *Acta Psychiatr. Scand.* 140, 295–312.
- Thumfart, K.M., Jawaid, A., Bright, K., Flachsmann, M., Mansuy, I.M., 2022. Epigenetics of childhood trauma: Long term sequelae and potential for treatment. *Neurosci. Biobehav. Rev.* 132, 1049–1066.
- Trottier, K., MacDonald, D.E., 2017. Update on Psychological Trauma, Other Severe Adverse Experiences and Eating Disorders: State of the Research and Future Research Directions. *Curr. Psychiatry Rep.* 19, 45.
- Trottier, K., Monson, C.M., Wonderlich, S.A., Crosby, R.D., 2022. Results of the first randomized controlled trial of integrated cognitive-behavioral therapy for eating disorders and posttraumatic stress disorder. *Psychological medicine* 52 (3), 587–596. <https://doi.org/10.1017/S0033291721004967>.
- Utzing, L.M., et al., 2016. A latent profile analysis of childhood trauma in women with bulimia nervosa: Associations with borderline personality disorder psychopathology: Classifying Trauma and BPD Psychopathology in BN. *Int. J. Eat. Disord.* 49, 689–694.
- Van Gerko, K., Hughes, M.L., Hamill, M., Waller, G., 2005. Reported childhood sexual abuse and eating-disordered cognitions and behaviors. *Child Abus. Negl.* 29, 375–382.
- Van Hoeken, D., Hoek, H.W., 2020. Review of the burden of eating disorders: mortality, disability, costs, quality of life, and family burden. *Curr. Opin. Psychiatry* 33, 521–527.
- Van Voorhees, E., Scarpa, A., 2004. The Effects of Child Maltreatment on the Hypothalamic-Pituitary-Adrenal Axis. *Trauma Violence Abus.* 5, 333–352.
- Vieira, A.I., et al., 2017. Putative Risk Factors for Non-Suicidal Self-Injury in Eating Disorders: Non-Suicidal Self-Injury in Eating Disorders. *Eur. Eat. Disord. Rev.* 25, 544–550.
- Vieira, A.I., et al., 2018. Eating disorders and non-suicidal self-injury: Structural equation modelling of a conceptual model. *Eur. Eat. Disord. Rev.* 26, 431–437.
- Waller, G., 1998. Perceived control in eating disorders: Relationship with reported sexual abuse. *Int. J. Eat. Disord.* 23, 213–216.
- Waller, G., et al., 2001. The Psychopathology of Bulimic Women Who Report Childhood Sexual Abuse: The Mediating Role of Core Belief. *J. Nerv. Ment. Dis.* 189, 700–708.
- Walsh, J., Burns, F., 2000. Sexual maturation and control issues among sexually abused and non-abused anorexia patients. *Br. J. Clin. Psychol.* 39, 307–310.
- Wonderlich, S.A., et al., 2007. The effects of childhood trauma on daily mood lability and comorbid psychopathology in bulimia nervosa. *J. Trauma. Stress* 20, 77–87.
- World Health Organization. ICD-10: International classification of diseases (10th revision). (<https://icd.who.int/browse10>) (1994).
- World Health Organization. ICD-11: International classification of diseases (11th revision). (<https://icd.who.int/>) (2019).
- World Health Organization. Child maltreatment. (<https://www.who.int/news-room/fact-sheets/detail/child-maltreatment>) (2022).
- Wrobel, A.L., et al., 2023. Childhood trauma and depressive symptoms in bipolar disorder: A network analysis. *Acta Psychiatr. Scand.* 147, 286–300.
- Zhang, Y., et al., 2022. Characteristics and mediating effect of gut microbiota with experience of childhood maltreatment in major depressive disorder. *Front. Neurosci.* 16, 926450.