

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/330557463>

Opportunistic alien catfish: unexpected findings in the diet of the alien species *Ictalurus punctatus* in Central Italy

Article in Archives of Polish Fisheries - January 2019

DOI: 10.2478/aopf-2018-0027

CITATIONS

0

READS

34

5 authors, including:



Phillip Haubrock
Senckenberg Research Institute

31 PUBLICATIONS 8 CITATIONS

[SEE PROFILE](#)



Martina Azzini
University of Florence

2 PUBLICATIONS 0 CITATIONS

[SEE PROFILE](#)



Ivan Fribbi
University of Florence

3 PUBLICATIONS 0 CITATIONS

[SEE PROFILE](#)



Alberto Francesco Inghilesi
University of Florence

38 PUBLICATIONS 134 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Diversity of Thaumaleidae in Europe [View project](#)



Lake Arreo Ecological State Recovery [View project](#)

Opportunistic alien catfish: unexpected findings in the diet of the alien species *Ictalurus punctatus* in Central Italy

Phillip J. Haubrock, Martina Azzini, Ivan Fribbi, Alberto F. Inghilesi, Elena Tricarico

Received – 25 July 2018/Accepted – 10 December 2018. Published online: 31 December 2018; ©Inland Fisheries Institute in Olsztyn, Poland
Citation: Haubrock P.J., Azzini M., Fribbi I., Inghilesi A.F., Tricarico E. 2018 – Opportunistic alien catfish: unexpected findings in the diet of the alien species *Ictalurus punctatus* in Central Italy – Fish. Aquat. Life 26: 239-242.

Abstract. Alien fish species are known for the negative impacts they can have on native species and their ability to shape recipient communities. One such case is the North American channel catfish, *Ictalurus punctatus* (Raf.), in the Arno River in central Italy. During river bank reconstruction, specimens of this species (n=40) were collected, and the stomach content was analysed. As a result, we highlight the opportunistic feeding behavior of this alien species that is responding rapidly to newly available resources and also poses a threat to native species.

Keywords: *Ictalurus punctatus*, feeding, opportunistic, impact

Introduction

Dietary analyses of invasive species is a common technique employed by biologists examining invasions, and such analyses have revealed much information about the biology and associated impact of invasive alien species (Kolar and Lodge 2002, Feyrer et al. 2003). Opportunistic fish species are known especially for their generally plastic diet and adaptability (Tyus and Nikirk 1990, García-Berthou 2002, Copp et al. 2009), and they often exhibit high variability among populations, years, and even seasons (Panek and Weis 2013, Purdom et al. 2015).

By nature, catfish are known to be opportunistic scavengers (Andrews et al. 1977, Tucker and Hargreaves 2004). In the Arno River in Florence, Italy, the abundant alien North American Channel catfish, *Ictalurus punctatus* (Raf.), has become a highly abundant species in biomass and number (Nocita 2007). In its native habitat, this catfish has been observed exhibiting seasonally variable, generally opportunistic predation (Busbee 1969, Ploeg and Tucker 1994), but information about it in Italy is rare or only anecdotal (Ligas 2007, Haubrock et al. 2017, 2018). In a highly anthropogenically affected ecosystem like the Arno River, alien aquatic species dominate in number and biomass (Nocita and Lenuzza 2016) and engage in interactions

P.J. Haubrock [✉]
Department of River Ecology and Conservation, Senckenberg
Research Institute and Natural History Museum Frankfurt,
Clamecystraße 12, 63571 Gelnhausen, Germany
e-mail: Phillip.Haubrock@hotmail.de

P.J. Haubrock, A.F. Inghilesi, E. Tricarico
Department of Biology, University of Florence, Via Romana 17,
50125 Florence, Italy

M. Azzini, I. Fribbi
Scienze delle Produzioni Agroalimentari e dell'Ambiente,
University of Florence, Piazzale delle Cascine 18, 50144 Florence,
Italy

A.F. Inghilesi
NEMO, Nature and Environment Management Operators s.r.l., Piazza
M. D'Azeglio 11, 50121 Florence, Italy

within the aquatic ecosystem and potentially with semi-aquatic organisms (Gelwick and McIntyre 2017).

During mechanical reconstruction work by the local Arno River Basin Authority on the vegetation zone along the Arno River in the city of Florence, Italy, plant material and gravel were discarded into the river (Fig. 1). In addition to its function as a natural barrier, this zone is characterized by large energy, nutrient, and biotic flow because of its proximity to aquatic (Cummins 1974, Sedel et al. 1974) and terrestrial ecosystems (Odum 1978). Eliminating this zone is likely to have negative a impact on both aquatic and terrestrial ecosystems (Odum 1978), while discarding plant material and gravel into the river might have an unforeseeable impact on the aquatic ecosystem.

To investigate how this abundant catfish species responds to such an external impact on the aquatic ecosystem, *Ictalurus punctatus* was sampled, and their diet was analysed. This type of study can help increase our understanding of alien fish behavior in general while also providing much needed information about the feeding behavior of alien catfish.

Materials and methods

Sampling

Fish were caught during construction work (March to June 2018) in the segment of the Arno River (43°45'49.9"N, 11°18'04.2"E) located in the city of Florence. *Ictalurus punctatus* was caught using standard fishing rods (2.20-3.90 m), 0.22-0.36 mm monofilament line, and size 2-26 fishing hooks baited with a variety of baits (maggots, worms, freshly cut liver, or bait-fish) that were deployed on the bottom and in the mid water column. The fish caught were placed on ice before being transported to the laboratory of the Department of Biology and the Natural History Museum La Specola in Florence.

Data and dietary analyses

Total fish length (TL) and weight (g) were measured (Norton et al. 1976), and the sex of the fish was determined. Prey items in the stomach of the catfish were identified to the lowest possible taxa under a standard stereo-microscope. Fragmented prey items were considered to be part of whole organisms, and they were counted as such. Only the numbers of the prey items were recorded. Consumed prey items were analysed as the frequency of occurrence ($\% = 100 \times A^I \times N^{-1}$) where A^I is the number of fish preying on species I, N is the total number of fish analysed (excluding those with empty stomachs), and Number is of food items (N%) following the formula

$$N^i = \frac{N_i}{\sum_{i1}^Q N_i},$$

where N_i is the number of food category i.

Results

During the study, both riverbanks were cleared of vegetation, some of which was discarded into the river resulting in increased nutrient input (Figure 1), which strongly increased turbidity. The specimens collected (21 male, 19 female) ranged from 13.5 to 64 cm in length (33.8 ± 2.2 cm) and weighed

Table 1

Dietary analysis results for *Ictalurus punctatus* (n=40) sampled from the Arno River

Prey	N%	F%
Aquatic plants	20.67	60
Terrestrial plants	1.68	7.5
Crustaceans	29.6	50
Aquatic molluscs	2.79	5
Hirudinea	0.56	2.5
Aquatic insects	8.94	32.5
Terrestrial insects	23.46	40
Small Cyprinidae	7.26	25
Large Cyprinidae	0.56	2.5
Fish larvae	1.68	7.5
Small mammals	0.56	2.5
Small birds	1.68	7.5
Testudines	0.56	2.5




Figure 1. North riverbank of the Arno River after the mechanical removal of vegetation.

between 20 and 3,500 g (538.5 ± 106.3 g). The results of the dietary analyses are listed in Table 1.

Discussion

Sampling predatory fish species, and especially opportunistic ones, to analyze their diets and potential diet shifts can provide important information about imbalanced trophic interactions or other disturbances (Kolar and Lodge 2002, Syväranta et al. 2010). Additionally, when performed for alien species, these analyses can reveal important information about the behavior and the impact of the species studied (Copp et al. 2009, Moris and Atkins 2009). In the case of the present study, plant material (aquatic and terrestrial), crustaceans (*Procambarus clarkii*, *Dikerogammarus villosus*), and terrestrial insects were the most dominantly consumed prey,

while the percentage shares of other prey categories were lower (Table 1). Nonetheless, the high percentages of terrestrial insects were based on items such as Scaraboidae, the larvae of which can be found in the soil, and other beetles (e.g., Coccinellidae and Curculionidae), which are items that are usually found in vegetated areas. The identification of multiple juvenile birds (Columbidae) and mammals (*Rattus norvegicus*) indicates opportunistic feeding behavior, which likely intensified because of the reconstruction works on the river banks that increased the availability of these prey. Indeed, comparing these observations with those from diet analyses of native populations showed the increased consumption of terrestrial prey items. Apart from these high quantities of terrestrial prey items, the dietary spectrum observed was similar to that identified among native populations (Bailey and Harrison 1948, Tyus and Nikirk 1990).

Lastly, the remains of an alien pond slider (*Trachemys scripta*) was noted in the stomach of *I. punctatus*. This is the first documented occurrence of an alien Ictaluridae feeding on Testudines outside its native range, and, at least to our knowledge, the first ever documentation of this behavior. This underscores the opportunistic nature of this species and thus, its potentially negative impact on recipient species communities (Cadi and Joly 2003, Fritz et al. 2005, Strayer 2010). In addition to the mechanical removal of the vegetation zone having had a negative impact on both aquatic and terrestrial ecosystem and their inhabitants, it can be concluded from the additional prey identified that *Ictalurus punctatus* exhibits opportunistic feeding behaviour when given the opportunity.

Acknowledgements. We would like to thank Paolo Agnelli, Fausto Barbagli, and Claudia Corti for the identification of highly decomposed material. Funding was provided by the Aquainvad-ED project (2020 Marie Skłodowska-Curie ITN-2014-ETN-642197).

Author contributions. P.J.H. and E.T. designed the study; M.A., I.F., and P.J.H. sampled the specimens for the study; M.A. and I.F. analysed the samples; P.J.H. and A.F.I. performed the analyses; P.J.H. and M.A. wrote the manuscript.

References

- Andrews W.H., Wilson C.R., Poelma P.L., Romero A. 1977 – Bacteriological survey of the channel catfish (*Ictalurus punctatus*) at the retail level – J. Food Sci. 42: 359-363.
- Bailey R.M., Harrison Jr H.M. 1948 – Food habits of the southern channel catfish (*Ictalurus lacustris punctatus*) in the Des Moines River, Iowa – Trans. Am. Fish. Soc. 75: 110-138.
- Busbee R.L. 1968 – Piscivorous activities of the channel catfish – The Progressive Fish-Culturist 30: 32-34.
- Cadi A., Joly P. 2003 – Competition for basking places between the endangered European pond turtle (*Emys orbicularis galloitalica*) and the introduced red-eared slider (*Trachemys scripta elegans*) – Can. J. Zool. 81: 1392-1398.
- Copp G.H., Robert Britton J., Cucherousset J., García-Berthou E., Kirk R., Peeler E., Stakėnas S. 2009 – Voracious invader or benign feline? A review of the environmental biology of European catfish *Silurus glanis* in its native and introduced ranges – Fish Fish. 10: 252-282.
- Cummins K.W. 1974 – Structure and function of stream ecosystems – BioScience 24: 631-64.
- Feyrer F., Herbold B., Matern S.A., Moyle P.B. 2003 – Dietary shifts in a stressed fish assemblage: consequences of a bivalve invasion in the San Francisco Estuary – Environ. Biol. Fish. 67: 277-288
- Fritz U., Fattizzo T., Guicking D., Tripepi S., Pennisi M.G., Lenk P., Wink M. 2005 – A new cryptic species of pond turtle from southern Italy, the hottest spot in the range of the genus *Emys* (Reptilia, Testudines, Emydidae) – Zool. Scr. 34: 351-371.
- García-Berthou E. 2002 – Ontogenetic diet shifts and interrupted piscivory in introduced largemouth bass (*Micropterus salmoides*) – Int. Rev. Hydrobiol. 87: 353-363.
- Gelwick F.P., McIntyre P.B. 2017 – Trophic Relations of Stream Fishes – In: Methods in Stream Ecology, Volume 1. Third Edition (Eds) F.R. Hauer, G.A. Lamberti, Academic Press: 457-479.
- Haubrock P.J., Johovic I., Balzani P., Inghilesi A.F., Nocita A., Tricarico E. 2017 – Assessing the impacts of the invasive Channel catfish *Ictalurus punctatus* in central Italy – In: 20th International Conference of Aquatic Species, October 22-26, 2017, Fort Lauderdale, Florida.
- Haubrock P.J., Balzani P., Johović I., Inghilesi A.F., Tricarico E. 2018 – The effects of two different preservation methods on morphological characteristics of the alien channel catfish *Ictalurus punctatus* (Rafinesque, 1818) in European freshwater – Croat. J. Fish. 76: 80-84.
- Kolar C.S., Lodge D.M. 2002 – Ecological predictions and risk assessment for alien fishes in North America – Science 298(5596): 1233-1236.
- Ligas A. 2007 – Population dynamics of the Channel Catfish, *Ictalurus punctatus* (Rafinesque, 1818), in the Ombrone river (Tuscany, Italy) – Atti Soc. tosc. Sci. nat., Mem., Serie B, 114: 57-62.
- Morris J.A., Akins J.L. 2009 – Feeding ecology of invasive lionfish (*Pterois volitans*) in the Bahamian archipelago – Environ. Biol. Fish. 86: 389-398.
- Nocita A. 2007 – La fauna ittica del bacino dell'Arno – Biologia Ambientale, 21: 97-105.
- Nocita A., Lenuzza A. 2016 – Le specie ittiche aliene in Toscana e il caso del Siluro europeo.
- Norton V.M., Davis K.B., Nishimura H. 1976 – A technique for sexing channel catfish – Trans. Am. Fish. Soc. 105: 460-462
- Odum E.P. 1978 – Ecological importance of the riparian zone. Strategies for Protection and Management of Floodplain Wetlands and Other Riparian Ecosystems. USDA Forest Serv. GTR-WO-12.
- Panek F.M., Weis J.S. 2013 – Diet of the Eastern Mudminnow (*Umbra pygmaea* DeKay) from two geographically distinct populations within the North American native range – Northeast. Nat. 20:37-48.
- Ploeg M.V.D., Tucker C.S. 1994 – Seasonal trends in flavor quality of channel catfish, *Ictalurus punctatus*, from commercial ponds in Mississippi – J. Appl. Aquacult. 3: 121-140.
- Purdom T., Schoofs A., Hodgson J., Maki T., Hodgson J. 2015 – A thirty-year diet record of largemouth bass (*Micropterus salmoides*) from a small north temperate lake – Bios 86: 20-30.
- Sedell J., Triska F.J., Hall J., Anderson N., Lyford J. 1974 – Sources and fates of organic inputs in coniferous forest streams – In: Integrated research in the coniferous forest biome (Ed.) R.H. Waring, AIBS Symp. Conif. Forest Ecosystems. Conif. Forest Biome Bull. No. 5: 57-59.
- Strayer D.L. 2010 – Alien species in fresh waters: ecological effects, interactions with other stressors, and prospects for the future – Freshw. Biol. 55 (Suppl. 1): 152-174.
- Syväranta J., Cucherousset J., Kopp D., Crivelli A., Céréghino R., Santoul F. 2010 – Dietary breadth and trophic position of introduced European catfish *Silurus glanis* in the River Tarn (Garonne River basin), southwest France – Aquat. Biol. 8: 137-144.
- Tucker C.S., Hargreaves J.A. 2004 – Biology and culture of channel catfish – Amsterdam, the Netherlands: Elsevier.
- Tyus H., Nikirk N. 1990 – Abundance, Growth, and Diet of Channel Catfish, *Ictalurus punctatus*, in the Green and Yampa Rivers, Colorado and Utah – Southw. Nat. 35: 188-198.