

Impact of shared facilities in advancing solid-state NMR research: 2025 edition[☆]

Robert W. Schurko^{a,b}, Chad M. Rienstra^c, Christopher P. Jaroniec^d, Alexandar L. Hansen^d, W. Trent Franks^d, David L. Bryce^e, Andreas Brinkmann^f, Victor Tersikh^f, Steven P. Brown^g, Dinu Iuga^g, Carine van Heijenoort^h, Franck Fayonⁱ, Sylvain Bertaini^j, Carlos Alfonso^k, Göran Karlsson^l, Gerhard Gröbner^m, Marek J. Potrzebowskiⁿ, Linda Cerofolini^{o,p,q}, Enrico Ravera^{o,p,q}, Marco Fragai^{o,p,q}, Moreno Lelli^{o,p,q}, Anne Lesage^r, Guido Pintacuda^r, Miquel Pons^s, Luís Mafra^t, José F. Schneider^u, Gustavo A. Monti^{v,w}, Rodolfo H. Acosta^{v,w}, Horacio M. Pastawski^{v,w}, Brijith Thomas^x, Yury G. Kolyagin^y, Vipin Agarwal^z, Guangjin Hou^{aa}, Feng Deng^{ab}, Kai Xue^{ac,ad}, Takanori Kigawa^{ae}, G.N. Manjunatha Reddy^{af,*} 

^a Department of Chemistry & Biochemistry, Florida State University, Tallahassee, FL, 32306, United States

^b National High Magnetic Field Laboratory (NHMFL), Tallahassee, FL, 32310, United States

^c Department of Biochemistry, University of Wisconsin-Madison, National Magnetic Resonance Facility at Madison (NMRFAM), Madison, WI, 53706, United States

^d Department of Chemistry and Biochemistry, Campus Chemical Instrument Center NMR Facility, and the National Gateway Ultrahigh Field NMR Center, The Ohio State University, Columbus, OH, 43210, United States

^e Department of Chemistry and Biomolecular Sciences, Centre for Catalysis Research and Innovation, and Nexus for Quantum Technologies, University of Ottawa, Ottawa, Ontario, K1N6N5, Canada

^f Metrology, National Research Council Canada, 1200 Montreal Road, M40, Ottawa, Ontario, K1A 0R6, Canada

^g Department of Physics, University of Warwick, Coventry, CV4 7AL, United Kingdom

^h Institut de Chimie des Substances Naturelles, Structural Chemistry and Biology team, CNRS, Université Paris-Saclay, 1 av. de la terrasse, Gif-sur-Yvette, 91190, France

ⁱ CEMHTI, CNRS, University of Orléans, 1D Avenue de la Recherche Scientifique, 45071, Orléans, Cedex 2, France

^j IM2NP, CNRS, Aix-Marseille Université, Univ. de Toulon, av. Escadrille Normandie Niemen, Marseille, 13397, France

^k COBRA, CNRS, Université de Rouen Normandie, 1 rue Tesnière, Mont-Saint-Aignan, 76821, France

^l Swedish NMR Centre at the University of Gothenburg, Gothenburg, Sweden

^m Swedish NMR Centre at Umeå University, Department of Chemistry, Umeå University, Umeå, Sweden

ⁿ Centre of Molecular and Macromolecular Studies, Polish Academy of Sciences, Sienkiewicza 112, Lodz, 90-363, Poland

^o Magnetic Resonance Centre (CERM) Università degli Studi di Firenze, Via Luigi Sacconi 6, Sesto Fiorentino, Firenze, 50019, Italy

^p Dipartimento di Chimica "Ugo Schiff," Università degli Studi di Firenze, Via della Lastruccia 3-13, Sesto Fiorentino, Firenze, 50019, Italy

^q Consorzio Interuniversitario Risonanze Magnetiche di Metalloproteine (CIRMMMP), Via Luigi Sacconi 6, Sesto Fiorentino, Firenze, 50019, Italy

^r High Field NMR Center (CRMN), CNRS/ENS Lyon/Université Claude Bernard Lyon 1, 5 rue de la Doua, Villeurbanne, 69100, France

^s Departament de Química Inorgànica i Orgànica, Universitat de Barcelona., Baldiri Reixac, 10-12, 08028, Barcelona, Spain

^t CICECO-Aveiro Institute of Materials, Department of Chemistry, University of Aveiro, Aveiro, 3810-193, Portugal

^u Instituto de Física de São Carlos – Universidade de São Paulo, Avenida Trabalhador São-carlense 400, São Carlos, SP, 13566-590, Brazil

^v Facultad de Matemática, Astronomía, Física y Computación, Universidad Nacional de Córdoba, Córdoba, 5016, Argentina

^w LANAIS-RMS, Instituto de Física Enrique Gaviola, CONICET, Córdoba, 5016, Argentina

^x Science Division, New York University Abu Dhabi, P.O. Box, Abu Dhabi, 129188, United Arab Emirates

^y Department of Chemistry, Lomonosov Moscow State University, Lenin Hills 1, Bld. 3, Moscow, 119992, Russia

^z Tata Institute of Fundamental Research Hyderabad, Hyderabad, Telangana, 500046, India

^{aa} State Key Laboratory of Catalysis, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian, 116023, China

^{ab} National Center for Magnetic Resonance in Wuhan, State Key Laboratory of Magnetic Resonance Spectroscopy and Imaging, Wuhan Institute of Physics and Mathematics, Innovation Academy for Precision Measurement Science and Technology, Chinese Academy of Sciences, Wuhan, 430071, China

^{ac} Centre of High Field NMR Spectroscopy and Imaging, Nanyang Technological University, 21 Nanyang Link, 63737, Singapore

^{ad} School of Physical and Mathematical Sciences, Nanyang Technological University, 21 Nanyang Link, 637371, Singapore

^{ae} NMR Operation Team, RIKEN Center for Integrative Medical Sciences, Yokohama, Kanagawa, 230-0045, Japan

This article is part of a special issue entitled: Advancements in SSNMR published in Solid State Nuclear Magnetic Resonance.

[☆] Given their role as Editor-in-Chief, David Bryce had no involvement in the peer-review of this article and has no access to information regarding its peer-review. Full responsibility for the editorial process for this article was delegated to another journal editor.

* Corresponding author. University of Lille, CNRS, Centrale Lille Institut, Univ. Artois, UMR 8181–UCCS– Unité de Catalyse et Chimie Du Solide, Lille, F-59000, France.

E-mail address: gnm.reddy@univ-lille.fr (G.N. Manjunatha Reddy).

<https://doi.org/10.1016/j.ssnmr.2025.102053>

Received 10 October 2025; Received in revised form 25 November 2025; Accepted 25 November 2025

Available online 28 November 2025

0926-2040/© 2025 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

A B S T R A C T

Shared research facilities (SRFs) offer researchers cost-effective access to advanced analytical instrumentation that individual laboratories may find challenging to acquire or maintain. By centralizing resources, SRFs support a diverse user community including students, early-career scientists, senior principal investigators, and industrial collaborators, while providing expert technical support and ensuring efficient use of infrastructure and funding. These facilities not only drive research productivity and foster interdisciplinary collaboration, but also serve as centers for training the next generation of scientists. In this article, SRFs that offer solid-state nuclear magnetic resonance (NMR) capabilities are discussed, highlighting representative examples, their accessibility, governance models, technical operations, application areas, and data-sharing practices. Usage data reveal that solid-state NMR-based SRFs strongly align with high-priority research goals, contributing to impactful projects across chemistry, life sciences, and materials science, as reflected in publication outcomes. The article also emphasizes that the collaborative networks among SRFs enhance knowledge exchange and resource coordination. Such coordinated inter-facility partnerships are expected to address emerging challenges, ultimately supporting sustainable infrastructure that meets the evolving needs of the solid-state NMR community.

1. Introduction

Solid-state NMR spectroscopy has become an indispensable characterization technique for elucidating the atomic- and molecular-level structures of a wide array of solid materials, particularly for systems that push the limits of conventional analytical techniques and/or where other methods provide ambiguous structural information. Emerging in the mid-20th century, NMR spectroscopy evolved into sophisticated approaches to probe nuclear spins in bulk matter, laying the foundation for today's advanced characterization spanning solids, liquids, and soft matter (e.g., polymers, gels, colloids, foams, biological materials). In the case of solid-state NMR, overcoming early challenges [1], such as addressing inherent sensitivity and resolution limitations, has driven transformative developments in the field. Notably, in the 1970s, techniques like Magic-Angle Spinning (MAS) and Cross-Polarization (CP) significantly improved spectral resolution and sensitivity [2,3]. The 1980–1990s saw continued progress, including the advent of high-power radiofrequency (RF) amplifiers, commercialization of CP/MAS probes, access to magnetic fields of >10 T, and 2D correlation experiments [4–10]. All of this paved the way for high-resolution solid-state NMR studies of polycrystalline materials, polymers, pharmaceuticals, glasses, and biological solids, including membrane proteins and amyloids. New pulse sequences involving dipolar decoupling and recoupling for internuclear distance measurements, and those for separation of the effects of isotropic and anisotropic interactions arising for half-integer quadrupolar nuclei [11,12], have opened further opportunities for the study of inorganic, hybrid, and complex biological materials. These advances have positioned solid-state NMR spectroscopy to fill information gaps left by X-ray diffraction (XRD), neutron diffraction/scattering, and/or electron microscopy (EM) techniques, especially for systems lacking long-range order or requiring detailed structural information on local atomic environments.

In the 21st century, solid-state NMR spectroscopy continues to advance to meet complex scientific demands for higher sensitivity, resolution, faster data collection, and higher throughput analyses [8]. Structural characterization of volume-limited samples is increasingly performed in their native or functional states—examples include biomacromolecules in complex cellular environments, protein–drug complexes to natural biopolymers including biomass and plant cell walls, catalytic supports, functional polymers, gels and electrolytes, coatings, energy materials, and semiconductor thin films used in optoelectronics, photonics, and energy conversion/storage devices [13–18]. Large-scale SRFs for solid-state NMR have played, and will continue to play, transformative roles across scientific disciplines and are important for advancing research in all aspects of chemistry, materials multidisciplinary, and life sciences. Progress in solid-state NMR-based structural analysis increasingly depends on access to state-of-the-art instrumentation and methods, including high magnetic fields (≥ 800 MHz) [19], robust pulse sequences and optimization of experimental parameters for

high-quality data acquisition [7,20], innovations in probe technology such as fast MAS systems ($\nu_{\text{rot}} = 30\text{--}160$ kHz) [21–23]. Additional capabilities such as MAS cryoprobes for room-temperature signal enhancement [24–26], and dynamic nuclear polarization (DNP) measurements at relatively low-temperatures [27–29], and those enabling a broad range of variable-temperature measurements [30,31], as well as those suitable for *in situ* experiments, and *in operando* studies [18, 32–35], have also become equally important. Solid-state NMR can act in a complementary manner with other structure determination techniques, which has led to the development of fields such as NMR crystallography (i.e., the combined use of solid-state NMR, XRD methods, and molecular modeling to predict, solve, and refine crystal structures) [36,37]. Alongside, computational prediction of NMR parameters, geometry optimization, and machine learning (ML)/artificial intelligence (AI)-driven structure determination are growing in popularity, which will be essential for NMR data interpretation and discovery acceleration [38–45].

The growing complexity and costs of these technologies, plus the expertise required for their optimal use, make national and regional shared solid-state NMR facilities strategically essential. Key idea of shared resources is to provide centralized access, expert support, and state-of-the-art infrastructure for high-end experimentation. In doing so, they create new opportunities in solid-state NMR for both experts and non-experts alike, fostering broader interdisciplinary collaboration.

This article showcases the growing influence of SRFs in pushing solid-state NMR capabilities and applications forward. It also illustrates how coordinated, multi-institutional networks and shared infrastructure enable transformative research that may be often challenging to pursue within the confines of individual research groups and laboratories.

2. Examples of shared NMR infrastructure and facilities

Several SRFs across the United States, Canada, the United Kingdom, Europe, South America, the Middle East, Africa, Australia, and Asia provide access to solid-state NMR infrastructure, offering complementary resources and domain-specific expertise. These centers house diverse instrumentation: solid-state and DNP-NMR spectrometers across a range of field strengths, custom probes, extreme-condition and *in situ/in operando* setups, and apparatus for DNP enhancements. Many of these NMR-based SRFs are integrated into broader research networks and paired with facilities for sample preparation, high-throughput data acquisition, and analysis. Some SRFs integrate complementary analytical methods such as magnetic resonance imaging (MRI), electron paramagnetic resonance (EPR), Fourier-transform ion cyclotron resonance (FT-ICR) mass spectrometry, synchrotron facilities, cryogenic electron microscopy (cryo-EM), computational modelling, data analysis software, enabling holistic and multimodal investigations (see Table 1). Although dozens of SRFs serve the global NMR community, the selected examples below illustrate diverse operational models, shared practices,

Table 1

Examples of SRFs include: National (N), International (IN), Collaborative (C), Industrial (ID), Coordinated Access (CA), Network for Advanced NMR (NAN), National Research facility (NRF) and Hybrid (HYB) modes. The HYB model extends access beyond NMR to other magnetic field-based techniques such as EPR, MRI, and FT-ICR mass spectrometry, while also integrating complementary capabilities including sample preparation, synchrotron characterization, microscopy, data analysis software, and modeling. PANACEA (Pan-European solid-state NMR Infrastructure for Chemistry-Enabling Access) is meta-infrastructure (MI) that coordinates access, expertise, and development efforts across national platforms. This federated model enables users to draw on complementary instrumentation and leading solid-state NMR expertise distributed across multiple centers, while harmonizing operational and training practices. Detailed hardware specifications are available on the individual facility websites or the discussion below. CCP-NC: Collaborative Computational Project for NMR Crystallography.

Shared research facility	Modes of access	Complementary techniques
PANACEA, meta-infrastructure, USA, EU	MI, CA	
National High Magnetic Field Laboratory (NHMFL or MagLab), Florida, USA	N, IN, ID, C, HYB, PANACEA	EPR, FT-ICR, MRI
National Magnetic Resonance Facility at Madison (NMRFAM), Wisconsin, WI, USA	N, IN, ID, C, HYB	Modeling, Data analysis software
Campus Chemical Instrument Center NMR Facility and National Gateway Ultrahigh Field NMR Center, Columbus, OH, USA	N, IN, ID, C, HYB, NAN	EPR, FT-ICR, Proteomics, Cryo-electron microscopy
Infranalytics, French meta-infrastructure	N, IN, CA, HYB, PANACEA	EPR, FT-ICR
The UK high-field solid-state NMR national research facility	N, IN, ID, C, CA, HYB, PANACEA	NMR crystallography (CCP-NC)
Government of Canada Ultrahigh-Field NMR Collaboration Platform, Canada	N, IN, ID, C, CA	
Swedish national NMR facility (SwedNMR)	N, IN, ID, C, CA, PANACEA	
Centro Risonanze Magnetiche (CERM), Firenze, Italy	N, IN, ID, C, CA, HYB, PANACEA	Synchrotron, Cryo-electron microscopy
Spanish National NMR Facilities, Barcelona, Madrid, and Bilbao, INSTRUCT	N, IN, ID, C, HYB	Synchrotron, Cryo-electron microscopy
Portuguese NMR Centre, University of Aveiro, Aveiro, Portugal	N, IN, ID, C, CA, PANACEA	
Shared NMR facility, PAS-Lodz, Poland	N, IN, C, ID	Computational modelling, X-ray diffraction/scattering
Core analytical facilities at NYU-AD, UAE and KAUST, SA	N, C, ID, CA	Microscopy, Electron diffraction
Multiple SRFs in China	N, IN, C, CA, HYB	EPR, FT-ICR, MRI
National Center for Magnetic Resonance in Wuhan	N, C, IN	MRI
Multiple SRFs in Russia	N, C, ID, HYB	EPR, FT-ICR, MRI, X-ray diffraction/scattering
Multiple NRFs in India	N, C, ID, CA	
Shared NMR facility, NTU, Singapore	N, IN, CA	
RIKEN Yokohama NMR Research Infrastructure, Kangawa, Japan	N, C, ID, CA	
Multiple SRFs Australia, New Zealand	C, ID, CA	
SRFs in Brazil	C, ID, CA	
Multiple SRFs in South Africa	C, ID, CA	
SRFs in Córdoba, Argentina	C, ID, CA	

opportunities and challenges, guiding principles, and common interests. The highlighted centers (Fig. 1: ordered geographically from left to right) are driving interdisciplinary research and innovation in NMR methodology and instrumentation.

Since the early 2000s, the growth of many SRFs—particularly national facilities has experienced a shift toward high magnetic fields (18.8–35 T) in conjunction with fast MAS (50–160 kHz) probes and custom-built probes for extreme conditions in order to broaden the scope of solid-state NMR applications. In addition, more than sixty DNP-NMR spectrometers spanning frequencies from 400 MHz/263 GHz to 900 MHz/593 GHz (some of which are accessible via SRFs) are accelerating structural studies of challenging solids. At the time of writing this article, several DNP-enabled NMR facilities are actively supporting the broader solid-state NMR community including 400 MHz/263 GHz systems (Fig. 1: geographically from left to right) in: Santa Barbara CA, Ames IA, Dayton OH, Princeton NJ (U. S. A.); Bruker (U. S. A.); Manchester and Cambridge (U. K.); Utrecht (Netherlands); Lyon, Grenoble, and Marseille (France); Bruker (France); Berlin, Rostock, Frankfurt and Darmstadt (Germany); Gothenburg, (Sweden); École Polytechnique Fédérale de Lausanne/EPFL and Eidgenössische Technische Hochschule Zürich/ETH Zurich (Switzerland); Aveiro (Portugal); Rehovot (Israel); Melbourne (Australia); Jeddah (Saudi Arabia); Wuhan (China); Kyoto and Tsukuba (Japan)—with new systems under installation in Virginia (USA); Bangalore (India); Shanghai (China). Facilities equipped with 600 MHz/395 GHz DNP systems include San Diego, CA, Tallahassee, FL, New York, NY, Dallas, TX, Columbus, OH, and Bruker Billerica, MA (U. S. A); Guelph, ON and Edmonton, AB (Canada); Nottingham (U. K.); Nijmegen (Netherlands); Göttingen, Aachen, and Jülich (Germany); ETH Zurich (Switzerland); Doha (Qatar); Abu Dhabi (U. A. E.); and Beijing (China), with further systems being acquired in Ames IA (U. S. A.), Hangzhou and Hefei (China). High-field DNP facilities include 800 MHz/527 GHz systems in Pacific Northwest National Laboratory/PNNL WS (U. S. A.); Utrecht (Netherlands); Lyon and Paris (France); Munich, Jülich and Berlin (Germany). In addition, EPFL (Switzerland) hosts a 900 MHz/593 GHz DNP system. Hardware specifications at SRFs may change due to upgrades including field strengths, consoles, probe types, DNP capabilities, and maintenance cycles, and the acquisition/replacement of specialized capabilities. Information on key hardware capabilities at each SRF alongside indicative output metrics (user numbers, allocated time, research topics, publication counts and data management plans) are presented in the discussion sections features by individual SRFs (*vide infra*), or directly obtained from the annual reports published in the dedicated websites of the SRFs. High sensitivity and resolution obtained in NMR spectra acquired at high fields and/or with DNP NMR capabilities are expected to facilitate rapid data acquisition and analysis, structural information obtained from which can be further refined through computational modeling, automation, and data-driven approaches such as NMR crystallography, structure prediction, and ML approaches. Therefore, advances in upgrading of the associated hardware from central- (CPU) to graphics processing unit (GPU), exascale interconnects (hybrid CPU-GPU architectures), GPU-dense nodes, and liquid-cooled systems, dramatically boost throughput for molecular modelling, and ML-driven simulations.

2.1. SRFs in Florida, FL, U. S. A. (Robert W. Schurko)

The National High Magnetic Field Laboratory (NHMFL or MagLab, for short) is an NSF-funded U.S. research facility spanning three institutions: the Florida State University (FSU in Tallahassee, FL; headquarters), University of Florida (UF, Gainesville, FL), and Los Alamos National Laboratory (LANL, Los Alamos, NM). The MagLab features a number of world-record magnets and high magnetic fields, and spans research areas such as physics, chemistry, materials science, engineering, biology, biochemistry, and medicine. There are seven user programs: (i) DC field (FSU), which features steady, continuous magnetic fields up to 45 T [46], including resistive and hybrid magnets; (ii) NMR/MRI (FSU), which has high-field solid-state NMR and MRI/S (including in vivo animal imaging) and probe development for solid-state NMR and MRI; (iii) Advanced Magnetic Resonance Imaging and Spectroscopy (AMRIS, UF), which has high-field solution NMR and



Fig. 1. Approximate locations of selected shared NMR facilities. Black dots indicate facilities with GHz-class spectrometers, and blue stars represent shared NMR facilities (300-900 MHz range). Red color ‘+’ symbols represent DNP NMR facilities. For accurate site details and institutional affiliations, please refer to the discussion in the following sections. (For interpretation of the references to color in this figure legend, the reader is referred to the Web version of this article.)

specialized NMR/MRI for biological and brain studies; (iv) EMR (FSU), which features high-frequency ESR techniques for studying unpaired electrons in materials (and working with the NMR/MRI division on DNP methods); (v) ICR (FT Ion Cyclotron Resonance Mass Spectrometry, FSU), which offers ultra-high mass resolution and accuracy; (vi) High B/T (UF), combining magnetic fields (up to ~ 15 T) with ultra-low temperatures (~ 0.4 mK) for quantum fluids/solids research; and (vii) Pulsed Field Facility (LANL), which uses ultra-high magnetic field pulses up to ~ 100 T for short-duration field experiments. The MagLab served 12,625 users from across the U.S. and around the world from 2018 to 2024, with ca. 50 % of these representing students and postdocs. On average, user counts from universities, government laboratories, and industry are 1390, 280, and 137, respectively, per annum. At the time of writing of this article, all user programs are open access with no user fees, and the only requirement is submission of a brief proposal that is reviewed by scientific peers.

The NMR/MRI and AMRIS user programs collectively serve ca. 550 users and offer $>17,000$ magnet days per year, with the majority of research focusing on chemistry, materials science, biology, and medicine. There are three flagship instruments, including (i) the 36 T series-connected hybrid (36T-SCH) [47], which is the highest field NMR magnet in the world (operating at 35.2 T/1.5 GHz for ^1H NMR); (ii) the 21.1 T/900 MHz ultra-widebore magnet (900 UWB) [48], which is used for in vivo MRI as well as solid-state NMR of quadrupolar nuclides; and (iii) the 395 GHz/14.1 T/600 MHz DNP NMR platform [49], which features a unique quasioptical microwave table that enables simultaneous MAS DNP NMR and Overhauser DNP (ODNP) NMR experiments. There are also 11 other NMR platforms with magnetic fields ranging from 7.1 T to 20.0 T. The NMR Technology Group in the NMR/MRI user program specializes in the construction of solid-state NMR probes with unique capabilities, including unique triple-resonance tuning configurations, low-frequency configurations for unresponsive low- γ nuclides, MAS rates >100 kHz, low-E coils for “lossy” biological samples, and special designs for GHz-class NMR platforms [50].

The large number of magnet days and high volume of users mandates that careful consideration be given to the number of support scientists and technicians, efficient scheduling, and magnet day usage, triaging of experiments, tracking of research output, and safety concerns. The NMR/MRI User Program is overseen by a Director, who reports not only

to the Director of the NHFML, but also to External Advisory and User Committees. Collectively, these partnerships govern future scientific directions, acquisition of equipment, and support of high-impact research projects, along with training and recruitment of users and education and mentorship activities for students. Staff scientists (referred to as Research Faculty or Research Associates at the MagLab), work directly with users on experiments, training of students, and dissemination of publications—they also conduct their own independent research projects that serve to improve the rapidity, efficiency, and quality of solid-state NMR experiments, which greatly benefits our users (especially non-experts in NMR). The user support of technical staff is also crucial, who ensure that all instruments are well-maintained, software is up to date, and new hardware and probes are consistently available—this latter aspect ensures that our users have access to the most cutting-edge NMR hardware available.

Many options are provided to current and potential users of the NMR/MRI facilities at the MagLab. For instance, users are offered the opportunity to work on site (this is especially valuable for students and/or initiation of projects) or to run their experiments remotely. Research faculty and technical staff work with directly with users to choose the most suitable magnets, spectrometers, probes, experiments, and pulse sequences, all with the goal of achieving meaningful scientific outcomes. Expert NMR users have the opportunity to work on methodological development and attend workshops at MagLab (e.g., high-field solid-state NMR, DNP, and probe/coil workshops), whereas non-experts are tutored and mentored in the application and interpretation of complex NMR data sets. With education and training in mind, we have recently launched the MagLab Summer School on Solid-State NMR Spectroscopy—our first full-length offering had 18 student participants from a variety of different background and educational levels (undergrad to postdoc) and featured lectures, problem sets, tours, workshops, and hands-on experimentation. In addition, in partnership with the Pan-European solid-state NMR Infrastructure for Chemistry-Enabling Access (PANACEA, which brings together seven national infrastructures from Europe and that at the MagLab), a workshop on DNP for European and American students and postdocs was recently offered at MagLab. Many of these ideas and access/training models are detailed in the following sections.

2.2. SRFs in Madison, WI, U. S. A. (Chad M. Rienstra)

The National Magnetic Resonance Facility at Madison (NMRFAM, Department of Biochemistry at the University of Wisconsin-Madison) is a national SRF supported by the National Institutes of Health (NIGMS) through the P41 Biomedical Technology Research Resource (BTRR) and R24 National and Regional Resource programs, along with user fees. The facility provides open access to students, researchers, and principal investigators, supporting research in biochemistry, structural biology, chemistry, materials science, software for data analysis, NMR methodology, and instrumentation. As a key partner in the Network for Advanced NMR (NAN), NMRFAM contributes to a nationwide effort to enhance NMR infrastructure and collaboration. Through this network, two new 1.1 GHz NMR spectrometers have been made available to the broader NMR community, one at NMRFAM dedicated to solid-state NMR, and the other at the Complex Carbohydrate Research Center (CCRC) configured for solution NMR.

Solid-state NMR capabilities have been substantially augmented at NMRFAM over the last five years through strategic expansion of high-field instrumentation and supporting infrastructure. Recent expansions include additions of several magnets (two 600 MHz, two 750 MHz, 900 MHz, and 1.1 GHz) with solid-state capabilities and a suite of probes with custom features. The 1.1 GHz and 900 MHz instruments are equipped with Bruker NEO consoles and feature an advanced suite of probes including BlackFox 1.6 mm HCN/HPC for biological samples, Phoenix 1.2 mm and 1.6 mm HFX for fast MAS applications, Phoenix 3.2 mm for low-gamma nuclei, and Bruker 0.7 mm HCN for ultrafast MAS. A custom-developed lock system for the 1.1 GHz spectrometer significantly enhances spectral resolution at ultra-high fields [51]. Probes for the 900 MHz system are compatible with the 1.1 GHz system, including BlackFox low-E field probes in 1.6 mm and 2.5 mm formats optimized for biological NMR applications. Moderate field instrumentation includes a 750 MHz wide spectrometer and three 600 MHz instruments dedicated to solids that provide additional capacity and serve as platforms for method development, staging and sample evaluation.

The facility maintains comprehensive rotor coverage from 0.7 mm to 5.0 mm, including 1.2 mm, 1.6 mm, 2.5 mm, and 3.2 mm sizes, optimized for different experimental requirements. All instruments emphasize compatibility with legacy Varian equipment, currently supported by PhoenixNMR and Resynant, ensuring continuity for established experimental protocols. The instrument suite enables detection of a broad range of nuclei including ^1H , ^{13}C , ^{15}N , ^{19}F , ^{31}P , and quadrupolar species, supporting advanced multidimensional NMR experiments across magnetic fields ranging from 600 MHz to 1.1 GHz. Technical innovations include advanced amplifier compensation systems [52], automated optimization software [53] and shimming protocols [54] that deliver improved homogeneity and spectral resolution. Comprehensive sensitivity studies across the magnetic field range have been conducted, providing quantitative performance metrics and optimization guidelines for experimental design [55]. Photoillumination probes enable control of pH within the rotor during solids measurements [56].

By integrating state-of-the-art instrumentation, automated data acquisition systems, advanced analysis software, and collaborative expertise, NMRFAM plays an important role in driving methodological innovation and expanding the scientific reach of solid-state NMR spectroscopy. The facility provides comprehensive user support including training programs, collaborative assistance in experimental design, data acquisition optimization, and analysis consultation. All instruments are accessible to the user community through the Network for Advanced NMR (NAN) and the NMRFAM website, supporting both local and external users from academic and industrial sectors.

2.3. SRFs in Columbus, OH, U. S. A. (Christopher P. Jaroniec, Alexandar L. Hansen, W. Trent Franks)

The solid-state NMR facilities at The Ohio State University (OSU,

Columbus, OH) are comprised of the Campus Chemical Instrument Center (CCIC) NMR Facility and the National Gateway Ultrahigh Field NMR Center with the first Bruker 1.2 GHz NMR spectrometer in the US, supported by a recent mid-scale research infrastructure award from the National Science Foundation, as its centerpiece. In addition to the ultrahigh field 1.2 GHz instrument, which has solid-state and solution NMR capabilities, the CCIC NMR facility staff operate and maintain an additional eight moderate-to-high field Bruker high-resolution NMR instruments including an 800 MHz solid-state NMR spectrometer obtained through a National Institutes of Health High End Instrumentation award, a 600 MHz/395 GHz DNP solid-state NMR spectrometer, a 800 MHz spectrometer with solids, solution and micro-imaging capabilities, as well as two 600 MHz, 700 MHz, 800 MHz and 850 MHz solution NMR spectrometers, most of which are equipped with cryoprobes. The solid-state NMR spectrometers are described in additional detail below, and collectively this comprehensive suite of state-of-the-art NMR instruments is available for use by academic researchers and industrial partners across the nation in diverse disciplines such as structural biology, materials science, and metabolomics [22,57–65].

The 1.2 GHz Bruker Avance Neo NMR spectrometer is equipped with a set of probes to support a wide range of scientific applications. For solid-state NMR research on materials this system offers a 3.2 mm HX probe for low-gamma nuclei (^{197}Au to ^{13}C ; 20–300 MHz) and a 1.9 mm HX probe for high-gamma nuclei (^{13}C to ^{31}P ; 300–484 MHz). The facility also offers premier fast- and ultra-fast MAS capabilities for biological samples with 1.3 mm and 0.7 mm HCN probes with MAS rates of up to 67 kHz and 111 kHz, respectively, which are ideal for advanced studies employing proton detection for enhanced sensitivity. The Bruker Avance III HD Ascend 600 MHz wide-bore NMR DNP solid-state NMR spectrometer is equipped with a 395 GHz gyrotron and a low-temperature MAS cabinet. The system offers several DNP and room-temperature probes for studying biological solids, powders, and micro-crystalline samples, including 3.2 mm HCN and HX low-temperature MAS probes for experiments carried out at 100 K as well as a 1.3 mm HCN room temperature probe. The Bruker Avance III HD Aeon 800 MHz wide-bore solid-state NMR spectrometer comes with an extensive collection of advanced probes, including 1.3 mm and 0.7 mm MAS probes capable of spinning speeds up to 67 kHz and 111 kHz, respectively, as well as 3.2 mm HXY and Efree HCN probes. A multi-purpose Bruker Avance Neo Ascend 800 MHz NMR instrument is equipped with a 4.0 mm gradient HR-MAS probe ideal for studies of biological solids such as membrane proteins and tissue samples, and can also accommodate the 0.7 mm, 1.3 mm and 3.2 mm Efree HCN probes.

While the facilities at Ohio State have a strong focus on solid-state NMR, they are also home to a world-class suite of instruments for solution-state NMR and are integrated with a comprehensive mass spectrometry core. For solution-state NMR studies, the 1.2 GHz spectrometer is equipped with a 3 mm TCI (HCN) Cryoprobe, which is ideally suited for resolution-limited studies of large proteins, nucleic acids, and complex mixtures found in metabolomics. To support high-throughput studies, the system is also equipped with a Chilled SampleCase for storing and automatically loading up to 24 samples at controlled temperatures and an Automated Tuning & Matching (ATM) module for streamlined operation. This is complemented by an 850 MHz NMR with a TCI cryoprobe and a 24-sample SampleCase for high-throughput biological studies, as well as a 700 MHz NMR with a TXO cryoprobe and a 510-sample SampleJet. The 700 MHz instrument offers the highest ^{13}C -detection sensitivity of all the solution instruments, providing unique capabilities for studying intrinsically disordered proteins (IDPs), natural products, and polysaccharides. The broader CCIC also includes a Mass Spectrometry and Proteomics (MSP) facility, which houses ten instruments and offers 26 distinct service options. This allows users access to complementary techniques such as Fourier Transform Ion Cyclotron Resonance (FT-ICR) mass spectrometry and targeted compound quantification, creating a hybrid environment for multifaceted materials and biomolecular characterization.

The NMR facilities at OSU support a large and diverse user base consisting of OSU internal users as well as users from external academic institutions, government and non-profit organizations, and industrial partners. Access is provided to both NMR experts and non-experts, with facility staff offering expert training, consultation on experimental design, data collection, and analysis to ensure users can achieve meaningful scientific outcomes. In the fiscal year 2025 alone, the CCIC NMR facility and National Gateway Ultrahigh Field NMR Center served over 200 users representing 25 institutions, and supported external research grants in funding resulting in over 30 publications.

2.4. SRFs in Canada (David L. Bryce, Andreas Brinkmann, Victor Terskikh)

The Government of Canada Ultrahigh-Field NMR Collaboration Platform is operated by the National Research Council Canada with support from Laboratories Canada, and a consortium of other Canadian Government Departments and Universities. Located in Ottawa, the centerpiece of the facility is a Bruker 900 MHz (21.1 T) instrument equipped with a wide array of commercial and custom-built solution-state and solid-state solenoid and magic-angle-spinning (MAS) probes, in particular for low-gamma and quadrupolar nuclei. Since its inception in 2005/06 as the National Ultrahigh-Field NMR Facility for Solids, this facility has received support from the Canada Foundation for Innovation, the Natural Sciences and Engineering Research Council of Canada, the National Research Council Canada, Laboratories Canada, the University of Ottawa, Bruker Biospin, and several other partners. Since its inauguration the facility has hosted visitors and collected experimental data for users in academia, government, and industry from across Canada and worldwide. In 2020, the spectrometer has been significantly upgraded and is now configured to alternate between solution- and solid-state NMR and features a recently acquired liquids cryoprobe for increased sensitivity. Access is provided through partnership agreements and on a per-use cost-recovery basis. As of 2025, well over 330 peer-reviewed publications have relied upon results obtained using the 900 MHz instrument. See, for example, refs [66–76].

In addition to this national platform, several universities in Canada operate local and regional NMR Core Facilities which are open to users and/or sample submission for a fee. Examples include the NMR center at the University of Alberta, the NMR Core Facility at the University of Ottawa, and Dalhousie University's Nuclear Magnetic Resonance Research Resource. In addition, 600 MHz/395 GHz DNP NMR facilities are available at University of Guelph, Guelph, and University of Alberta, Edmonton. There are currently no GHz-class magnets available for solid-state NMR in Canada.

2.5. SRFs in Warwick, U. K. (Steven P. Brown, Dinu Iuga)

The Engineering and Physical Sciences Research Council (EPSRC)- and Biotechnology and Biological Sciences Research Council (BBSRC)-funded UK high-field solid-state NMR national research facility (NRF) hosted at the University of Warwick has been providing access to researchers since 2010, initially at magnetic field of 850 MHz, expanding to include 1 GHz in 2020 and 1.2 GHz in 2025. The facility has been contributing to the UK's fundamental and applied research in academia and industry. The access model, which supports external (within the UK) users, fosters collaboration across disciplines and institutions. By combining cutting-edge technology with other areas of research such as the calculation of NMR parameters using NMR crystallography methods (benefitting from funding by EPSRC since 2010 for the collaborative computational project in NMR crystallography, CCP-NC), the NRF is uniquely positioned to enable cutting-edge research across materials, physical and life science applications, including batteries, catalysts, energy materials, pharmaceuticals, plant cell walls and protein interactions. The access range and impacts of the work carried out at the NRF are illustrated by a few representative examples in the areas

biological assemblies [77,78], pharmaceuticals [79,80], hydrogels [81], biomass characterization [82,83], metal organic frameworks [84,85], inorganic and hybrid materials [86–89], and NMR methodology [90, 91]. In this way, the NRF not only accelerates scientific discovery but also strengthens the UK's position in advanced characterization via shared NMR facilities. In addition, the Warwick NMR facility enables access to international users through the Pan-European NMR network (PANACEA, see below) facilitated by the EU. In the last 5 years (2020–2024), the UK high field NMR facility served over 400 researchers including PhD students, postdoctoral researchers, principal investigators, and industrial collaborators, yielding to over 100 publications. The NRF operates 24/7, 365 days a year, with all time (excluding a small amount of maintenance days and facility manager time) allocated to the user community.

2.6. SRFs in Europe

Several NMR-based SRFs across Europe offer access through both locally managed centers and the broader Pan-European NMR infrastructure: Germany, Switzerland, the Netherlands, France, Italy, and Spain have developed SRFs accessible to the national users and the EU users, some of which are presented in the below sections. In a separate article published as part of this special issue, Pan-European solid-state NMR Infrastructure for Chemistry-Enabling Access (PANACEA) project more elaboratively features the collaborative effort between these SFRs.

2.7. Multiple SRFs in France (Carine van Heijenoort, Franck Fayon, Sylvain Bertaina, Carlos Alfonso)

In France, the national research infrastructure “Infranalytics”, established in 2022 and led by the CNRS Chemistry Department, unites three major user facilities: NMR (solid- and solution-state, and imaging), EPR (spectroscopy and imaging), and high-resolution FT-ICR mass spectrometry, all operating at very high magnetic fields. This decentralized national SRF provides access to cutting-edge magnetic field-based analytical techniques through a network of thirteen sites across France, housing 23 high-field instruments: 9 NMR spectrometers (up to 28.2 T, ^1H = 1200 MHz), 8 EPR spectrometers (up to 9.4 T, 263 GHz), and 6 FT-ICR systems (up to 18.0 T). Operated by internationally recognized research teams with their expertise in various fields, these platforms deliver complementary insights into chemical composition, molecular organization, and dynamics across diverse samples. Specifically, NMR applications span healthcare, biological samples, environmental science, energy, catalysis, advanced materials, and agri-food innovation [22,92–102]. The Infranalytics SRF is open to national and international users, providing approximately 900 days of access per year for NMR, 480 days per year for ESR/EPR, and 420 days per year for FT-ICR measurements, over the range of available instruments. Beyond providing access, the Infranalytics SRF also coordinates national-level investments in advanced instrumentation. Interdisciplinary in nature, it serves the scientific needs of users who cannot independently acquire such sophisticated tools, ensuring access to world-class analytical capabilities. This coordinated framework enhances both the quality and efficiency of access by (i) directing users to the most suitable platform based on expert-reviewed project evaluations, (ii) strengthening complementarity among partner facilities to optimize resources and plan future equipment, (iii) harmonizing user training programmes, and (iv) coordinating investments in ultra-high-field technologies to extend access for French and international research teams, including those using standard magnetic fields.

Infranalytics experimental NMR facility (previously known as l'infrastructure de Recherche Résonance Magnétique Nucléaire à Très Haut Champ, IR-RMN-THC) today brings together a 1.2 GHz (Lille) and a 1 GHz (Lyon) spectrometer both equipped for solution, or solid-state applications with unique ultra-fast MAS capabilities at 1 GHz, two 950 MHz (Gif-sur Yvette, Grenoble), and one 900 MHz (Lille) for

biomolecule and biosolid applications with *in situ* high pressure or sample illumination capabilities at 950 MHz, one 850 MHz (Orléans) for solid-state NMR with unique high-temperature facilities, and two MAS DNP NMR 800 MHz systems (Paris, and Lyon) with polarizers for DNP-dissolution. A new 1 GHz spectrometer with solution and MAS cryoprobes (Bordeaux, 2026) is envisaged to join the range of facilities. Each spectrometer offers 100 days of access per year, with an exception to 1.2 GHz spectrometer, which is open to users 200 days per year. Over the last five years, Infranalytics' ultra-high-field NMR facility has provided users with more than 4890 experiment days, corresponding to the completion of 481 scientific projects (more than 96 projects and 980 experiment days per year on average). Across all research areas, this results in over 100 publications each year from users and affiliated research teams, spanning disciplines such as chemistry, biology, materials science, environmental science, and earth sciences.

2.8. Multiple SRFs in Sweden (Göran Karlsson, Gerhard Gröbner)

SwedNMR is a national and nation-wide research infrastructure that provides access to advanced NMR instrumentation and expert support across all areas of NMR spectroscopy in Sweden. SwedNMR supports a diverse range of applications from the atomic-level characterization of organic and pharmaceutical compounds, proteins, RNA, and DNA to studies of batteries, solar cells, and clinical samples [103–112]. The infrastructure operates through a distributed, node-based model organized into three thematic modules: bioNMR, materials NMR, and translational NMR. SwedNMR is supported by the Swedish Research Council and involves the following participating NMR partners: University of Gothenburg, Chalmers University of Technology, Lund University, Linköping University, Royal Institute of Technology (KTH), Stockholm University, Swedish University of Agricultural Sciences (SLU), Umeå University, and Uppsala University. The infrastructure includes three access nodes located in Gothenburg, Stockholm, and Umeå, providing researchers with on-site and remote access to high-field NMR systems, while five expert nodes offer technical consultation, methodological development, and user training. SwedNMR is open to users from all scientific backgrounds and levels of expertise, from academic groups and industrial partners to early-career researchers and students, thus promoting widespread adoption of NMR across disciplines. The main solid-state NMR centre is located in Umeå, together with the smaller Stockholm and Gothenburg nodes offers access to solid-state including cryo-MAS and DNP-NMR capabilities. Together, these sites offer broad access to advanced solid-state NMR, including cryogenic MAS and DNP-NMR. The flagship instrument is the 850 MHz solid-state NMR spectrometer in Umeå, equipped with a 0.7 mm HXY MAS probe, alongside a 600 MHz system featuring a 3.2 mm Cryo-MAS probe. Additional instrumentation includes an 800 MHz spectrometer with a 1.3 mm MAS probe (expected in 2026) and a 600 MHz HX MAS system in Stockholm, as well as 400 MHz systems with 3.2 mm HXY and HFX DNP capabilities in Gothenburg. Between 2022 and 2024, these facilities collectively supported 205 users across approximately 3600 access days. On average, SwedNMR annually serves 150 external users from Sweden, supporting 30 research projects per year. During 2020–2024, this usage has led to over 209 peer-reviewed publications and dissemination activities across a wide range of disciplines, including biomacromolecular research (40 % usage), catalysis and advanced materials (30–40 % usage), cellulose-based biopolymers, and pharmaceutical and other applications (20–30 % usage).

2.9. SRF in Łódź, Poland (Marek J. Potrzebowski)

The NMR Laboratory at the Centre of Molecular and Macromolecular Studies, Polish Academy of Sciences (CMMS PAS, Łódź) serves as a central hub, among others, for solid-state NMR spectroscopy in Poland. Funded through research grants from the National Science Centre (NCN), internal resources and industrial contracts, the laboratory

supports around 30 domestic users annually (2020–2025) and engages in scientific collaborations with several European countries including the Czech Republic, Croatia, France, and Portugal, as well as collaborative effort with research labs in the United States. The SRF at Łódź is equipped with four solid-state NMR spectrometers with fields ranging from 9.4 T to 14.1 T (AVANCE Neo 400, AVANCE III WB 400, AVANCE III 500, and AVANCE III 600), along with a diverse set of probeheads supporting rotor diameters from 4 mm to 1.3 mm, with dual-channel broadband and triple-resonance dedicated probes, enabling analysis of several NMR active nuclei crucial for studying biological samples, pharmaceuticals, organic and organic–inorganic hybrid materials. The NMR Center in Łódź offers tailored support for research projects ranging from routine analysis of powders with basic ^{13}C CP-MAS to high-resolution 2D correlation NMR experiments for spectral analysis in conjunction with advanced NMR crystallography, and solid-state NMR methods development [113–124]. For projects beyond the SRF's capabilities, researchers from Poland are encouraged to apply for access through the PANACEA network. Encouraged by the success of the local capability and international collaboration, Poland is currently working toward forming a national NMR infrastructure, obtained a favorable evaluation, making a step towards investments in the Polish Roadmap for a shared NMR Research Infrastructure.

2.10. SRF in Florence, Italy (Linda Cerofolini, Enrico Ravera, Marco Fragai, Moreno Lelli)

Italy's Magnetic Resonance Center (CERM–Centro Risonanza Magnetica) at the University of Florence provides access to 13 NMR spectrometers (9.4–28.2 T, 400–1200 MHz), spanning solution- and solid-state applications in structural biology, drug–protein interactions, catalysis, pharmaceuticals, and materials. Of these instruments, the 1.2 GHz, 850, 800, and 700 MHz, are equipped with solid-state NMR probes and a 600 MHz instrument is equipped with an HR-MAS probe. Access is offered through three routes: Instruct-ERIC (transnational, EU including supported Italian users), Instruct-ITALIA (national structural biology network, started its activity in early 2020), and PANACEA (2021 onwards, global solid-state NMR for chemistry, pharma, and materials). At the national level, CERM has been providing access to state-of-the-art NMR instrumentation since 1990 and transnational access since 1994, overall, with more than 30 years of experience in hosting users. Since 2010, CERM together with its associated Consortium for the Magnetic Resonance of Metalloproteins (CIRMMP) is also included in the “Italian Roadmap of Research Infrastructures of Pan-European Interest”. CERM is also the Italian node of the European Instruct-ERIC, the European research infrastructure consortium for integrated structural biology defined in the European Strategy Forum on Research Infrastructures (ESFRI) Roadmap and active since 2012. CERM is contributing to the Instruct-ITALIA infrastructure with its NMR or the EPR instrumentation, complementing facilities offering platforms and techniques which include 3D structural methods such as X-Ray crystallography and cryo-EM, Circular Dichroism, and other biophysical platforms. In parallel, CERM/CIRMMP is also the core centre of the Instruct-ITALIA network, an infrastructure to promote and foster an integrated approach in structural biology at the national level.

Regarding solid-state NMR, CERM offers probes ranging from 4 mm to 0.7 mm HCN (MAS speed up to 111 kHz) to support a wide range of scientific applications. The 1.2 GHz Bruker Avance Neo NMR spectrometer with 0.7 mm probe is particularly suited for advanced studies employing proton detection with enhanced sensitivity and resolution without the need for deuteration, and enabling detailed investigations of dynamics and interactions in the solid state. The Bruker Avance III 850 MHz wide-bore system with 3.2 mm CP MAS DVT $^{15}\text{N}/^{13}\text{C}/^1\text{H}$, 1.3 mm CP MAS $^1\text{H}^{19}\text{F}/\text{BB}/^{15}\text{N}$ and 0.7 mm CP MAS $^1\text{H}/^{13}\text{C}/^{15}\text{N}$ probe-heads for biomolecular applications. The Bruker Avance III 800 MHz standard-bore system with 3.2 mm CP MAS DVT E-free $^{15}\text{N}/^{13}\text{C}/^1\text{H}$, 1.3 mm 3γ $^1\text{H}^{19}\text{F}/\text{BB-X}/\text{BB-Y}$ and 1.3 mm CP MAS $^1\text{H}/^{13}\text{C}/^{15}\text{N}$ probes allow to

perform measurements on inorganic materials and biomaterials with high salt concentrations. The Bruker Avance NEO 700 MHz systems come with two CP MAS BB/BB/1H probe-heads (4- and 3.2 mm) which allow to record also solid-state NMR spectra on ^{27}Al and ^{11}B nuclei.

Specifically, the CERM/CIRMMP infrastructure provides state-of-the-art instrumentation and expertise to perform the most comprehensive array of experiments needed for the structural and dynamic characterization of biological macromolecules and their complexes, with well-known expertise in structural determination of protein, paramagnetic proteins (CERM contributed to the identification of structure of a paramagnetic metalloprotein (MMP12) uniquely through solid-state NMR restraints [125] and protein amyloids) [126,127], antibiotics [128,129], biomaterials [130], material sciences [131,132] and DNP-enhanced NMR [99,133–135] are also active fields of research. The research at CERM is devoted to integrated approaches, and not only in structural biology. CERM facilities are also enabled with other complementary techniques that are also included in the user access possibilities, such as EPR, protein production for in-cell NMR, computational services, and proprietary infrastructure dedicated to industrial research providing access to instrumentation and expertise. A Cryo-EM instrument and an X-ray diffraction platform are available at the facility jointly with the Chemical Department. Together, these facilities supported 205 users over approximately 3600 access days, resulting in more than 209 peer-reviewed publications and dissemination outputs. Specifically, for solid-state NMR, during the last 5 years (2020–2024) 42 users accessed the facility with more than 560 access days distributed in the areas of biomacromolecular and cellular systems, advanced materials, and pharmaceutical sciences. These contributions reflect diverse domains including biomacromolecular systems, advanced functional materials, cellulose-based biopolymers, and pharmaceutical sciences as well solid-state NMR and DNP-NMR methodology, demonstrating broad interdisciplinary impact.

2.11. SRF in Aveiro, Portugal (Luis Mafra)

The Portuguese NMR Centre, established in 2023–2024 at the University of Aveiro, is Portugal's largest NMR facility. It houses six spectrometers, including a 700 MHz Bruker Avance III HD for high-resolution studies of biomaterials and polymers, a 400 MHz Bruker Avance Neo DNP-NMR with cryogenic MAS probe, and four mid-field systems (300, 400, and two 500 MHz), and operates within both the national PTNMR network and the Pan-European PANACEA network. Managed by 10 permanent researchers, two technical staff, and several PhD students, the Centre benefits from funding from CICECO (Centre for Research in Ceramics and Composite Materials)–Aveiro Institute of Materials, which fosters strong interdisciplinary research. Operating through PANACEA's peer-review system (see below - PANACEA project), the Centre serves approximately 32 local users annually, supporting both academia and industry (e.g., Hovione, Bondalti). Services include a wide range of multinuclear 1D/2D MAS NMR methods. The research at this SRF also involving developing novel MAS-DNP methodologies to polarize gases in confined spaces and exploring additive manufacturing (3D printing) to produce custom NMR components [136]. The Centre has been a hub for methodological innovation, with expertise in developing advanced solid-state NMR techniques. Key contributions include enhancing the sensitivity and resolution for challenging quadrupolar nuclei using methods such as multiple-quantum (MQ) and satellite-transition (ST) MAS, and developing high-resolution ^1H -based combined rotation and multiple-pulse spectroscopy (CRAMPS) approaches at fast MAS, combined with ^1H -inverse detection and ^1H – ^1H double-quantum–single-quantum (DQ-SQ) recoupling. Experimental capabilities are complemented by computational approaches, including first-principles calculations and ab initio molecular dynamics. Research spans materials science and biophysics, supported by National (FCT) and European Research Council (ERC) grants. Applications include atomic-scale studies of CO_2

chemisorption and physisorption in nanoporous materials such as metal-organic frameworks (MOFs), covalent-organic frameworks (COFs), and amine-functionalized silicas (using DNP-enhanced $^{13}\text{C}/^{15}\text{N}$ NMR); investigations of battery electrolytes, *in situ* photo-electrochemical NMR, drug delivery systems, and pharmaceuticals, bioadhesives, biomass, studies of energy materials, glasses, heterogeneous catalyst acidity (e.g., ^{31}P probe molecules in zeolites); and luminescent nanothermometry [137–147]. Specifically, the biophysical research addresses protein–lipid interactions in membrane dynamics, integrated with complementary methods, while machine learning with neural network potentials is applied to model gas interactions and reactions under confinement.

2.12. Coordinated access between SRFs in EU and in U. S. A. (Anne Lesage, Guido Pintacuda)

The PANACEA project is a European distributed infrastructure designed to facilitate access to advanced solid-state NMR instrumentation for the broader chemistry community. Funded by the EU Horizon 2020 INFRAIA-02-2020 (*Integrating Activities for Starting Communities*), PANACEA has been operating since 2021, and brings together eight NMR-based SRFs operating as national infrastructures across Europe as well as a strategic partnership with the US-based National High-Magnetic Field Laboratory [148]. It includes the High-Field NMR Center in Lyon (CRMN, France), the Extreme Condition and Materials: High Temperature and Irradiation Laboratory in Orléans (CEHMTI, France), the High-Field Solid-State NMR National Research Facility (NRF) in Warwick (UK), the Danish Center for Ultrahigh-Field NMR Spectroscopy in Aarhus (Denmark), the Magnetic Resonance Center in Florence (CERM/CIRMMP, Italy), the Solid-State NMR Facility for Advanced Materials Science in Radboud (the Netherlands), the Aveiro Institute of Materials, in Aveiro (CICECO, Portugal), and the Swedish NMR Centre in Gothenburg (Sweden).

Unlike the above-mentioned centralized facilities, PANACEA acts as a meta-infrastructure, coordinating access, expertise, and development efforts across national platforms. This federated model allows users to benefit from complementary instrumentation and forefront expertise in solid-state NMR spread across multiple centers, while aligning operational and training practices. The consortium currently offers access to 30 NMR spectrometers (with proton frequencies ranging from 100 to 1500 MHz), fully equipped for the most advanced solid-state NMR applications, including 6 DNP systems, enabling a wide range of applications from pharmaceuticals and biomolecular systems to fine chemicals, cosmetics, food, materials, fuel, polymers, and clean energy tries [148]. As of mid-2025, more than 200 projects have been supported through over 1100 trans-national access days, with an average project length of 5–6 days. The user activities represent between 10 and 17% of the available time at each participating infrastructure, preserving capacities for national users and local in-house research programs. To support this transnational access, PANACEA has implemented common user interfaces, streamlined proposal review processes, and provided training resources to assist non-expert users in adopting solid-state NMR techniques. Beyond access, PANACEA promotes methodological and technological innovation through joint research activities involving four key partners: Bruker Biospin, Mestrelab, Weizmann Institute of Science, and École Polytechnique Fédérale de Lausanne (EPFL). These collaborations have enhanced instrument performances, software usability, and analytical workflows, impacting areas as probe technology, remote access tools, metadata standards, and NMR crystallography. In turn, these joint research activities improve the quality of the trans-national access to the infrastructures for the users. By leveraging and connecting national capabilities, PANACEA exemplifies a next-generation model of infrastructure coordination, serving as a backbone for European solid-state NMR while fostering global partnerships and enhancing the reach, efficiency, and impact of shared scientific resources.

2.13. SRFs in Barcelona, Madrid, and Bilbao, Spain (Miquel Pons)

Spain's high-field NMR centers are organized as a distributed national research infrastructure under the Infraestructura Científico-Técnica Singular (ICTS) program, with three service nodes located in Barcelona, Madrid, and Bilbao. These centers provide competitive access to researchers through evaluation by an independent external committee. The facilities feature NMR spectrometers in the 18.8 T–23.5 T (800 MHz–1 GHz) range. The GHz instruments in Barcelona and Bilbao employ high-temperature superconducting magnets, eliminating the need for liquid helium cooling below 2 K. Each service node offers distinct expertise to the users: the Barcelona center focuses on intrinsically disordered proteins and pharmaceutical biologics; the Madrid node is advancing biosolids NMR with a MAS cryoprobe and is soon to install a fast MAS probe on its 800 MHz instrument; and the Bilbao node has strengths in glycobiology and metabolomics [26,149–156]. All three centers are actively involved in integrative structural biology efforts, in close collaboration with complementary large-scale facilities such as the ALBA synchrotron located within 30 km proximity to the Barcelona node, and cryo-EM hubs co-located near each NMR site. The three high field NMR centers are now part of INSTRUCT ERIC (pan-European research infrastructure in structural biology) following a major expansion of the Spanish center to enhance international access and foster collaborative research across multidisciplinary platforms.

2.14. SRF in São Paulo, Brazil (José Fabián Schneider)

In Brazil, the NMR-Lab of São Carlos Institute of Physics, University of São Paulo hosts a multiuser solid-state NMR facility, which was established in 2021 with funding from the São Paulo State Foundation for Research Support (FAPESP). A Bruker Avance Neo 400 MHz wide-bore spectrometer is among the heavily used instruments for external users. The facility is overseen by administrative and users committees. This spectrometer is equipped with multiple probes (1.3 mm HX, 2.5 mm HXY, 3.2 mm XY low gamma, 4 mm H/FXY, 5 mm HX, and 7 mm HX) and capable of spinning speeds up to 60 kHz. Additional resources include a Bruker 600 MHz, a Varian 300 MHz, and an Agilent 250 MHz system. Instruments operate year-round on a 24/7 basis, with ~5% time dedicated to maintenance. Several modes of access and services are provided: reduced fees for researchers supported by federal or state grants, free of charge on a collaborative basis with NMR-Lab staff, or by paid packages for industry. Technical and academic staff assist users, many of them non-NMR specialists, in selecting experiments and ensuring high-quality data. In the past four years, over 50 users accessed the facility. Research areas include materials science (40%), physical chemistry (30%), structural chemistry (10%), and environmental chemistry (20%). Research projects and samples include ceramics, glasses, biomaterials, nanoparticles, concretes, polymers, small molecules, and hybrid materials [157–162].

2.15. SRF in Córdoba, Argentina (Gustavo Monti, Rodolfo Acosta, Horacio Pastawski)

In Argentina, the National Laboratory for Research and Services in Solid-State Nuclear Magnetic Resonance (LANAIS-RMS) was established at the National University of Córdoba in 1990 as part of Argentina's network of open national facilities for research and services. The laboratory is centered on a Bruker AVANCE NEO 300 WB spectrometer equipped with various probes, including 4 mm CP/MAS, wide-line proton and X-nuclei, as well as diffusion and micro-imaging options. Research activities encompass quantum phenomena, pharmaceuticals, polymers, porous media for petrophysics, battery electrodes, and pulse sequence development [163–171]. Complementary resources include low-field solid-state NMR at 20 MHz (Bruker Minispec mq20) and 60 MHz (custom-built with a Magritek KEA console), single-sided NMR (Magritek profile MOUSE PM25/PM5), and a Magritek Spinsolve80

Ultra Multi X for solution-state NMR. LANAIS-RMS provides extensive services in materials characterization, with emphasis on catalysis and food polymers. More than 120 researchers from Argentina, Brazil, the U. S. A., and Europe have collaborated at the facility, which also serves 13 private companies (including pharmaceutical and oil industries) and 41 academic laboratories. Access is offered through collaborative projects or paid services, supported by the National University of Córdoba and national research grants. Services include training, experiment design, data acquisition, analysis, and simulations.

2.16. Multiple SRFs in Russia (Yury G. Kolyagin)

Multiple sites in Russia host NMR-based SRFs including Lomonosov Moscow State University, St. Petersburg State University, Kazan Federal University, Novosibirsk State University, and the Russian Academy of Sciences institutes. These SRFs provide open access to solid-state NMR facilities to researchers through collaborative arrangements. These facilities are equipped with multinuclear solid-state probes, MAS capabilities, and advanced pulse sequence libraries, enabling studies of both crystalline and amorphous solids. Applications of solid-state NMR in Russia are diverse, ranging from fundamental investigations of mineral structures, catalysts, and porous materials to studies of polymers, pharmaceuticals, and biomolecular assemblies. Particular emphasis has been placed on porous materials, as well as heterogeneous catalysts and clays of geological interest. Shared access models allow researchers from smaller institutions to benefit from advanced NMR infrastructure. Recent studies include NMR studies of metal-organic polyhedral frameworks [172–174], NMR crystallography of salts and zeolite materials [175–177], and methodological advances in solid-state NMR of quadrupolar nuclei such as ^{51}V , ^{93}Nb , and ^{95}Mo [177,178]. Notably, research groups are developing *in-situ* and *operando* solid-state NMR approaches, often combined with MAS under batch or flow conditions, to investigate heterogeneous catalysts [179–182] and/or hydrothermal synthesis [182]. Beyond enabling access to SRFs, these facilities act as training hubs, offering workshops and technical support that broaden expertise in solid-state NMR across the community. Annually, an estimated 30–50 users access these facilities, contributing to the publication of over 200 papers between 2015 and 2024. By fostering collaboration and efficient use of resources, Russia's shared NMR facilities play a vital role in multidisciplinary innovation at the national level, and knowledge exchange.

2.17. SRFs in Abu Dhabi, U. A. E., and Jeddah, S. A. (Brijith Thomas)

In the Gulf, Middle-East and North Africa regions (broader MENA), the NMR infrastructure is booming and facilitating access to users from different research areas. Regionally, a high-field solid-state NMR facility in Saudi Arabia supports access to 14 instruments with magnetic fields up to 21 T (^1H = 900 MHz), including variable-temperature capabilities and a 400 MHz DNP-NMR system. At the King Abdullah University of Science and Technology (KAUST), researchers access instrumentation at core facilities via a Request for Service (RFS) platform and receive independent training, facilitating research in wider chemistry areas such as porous materials and membranes, for example [183,184]. The facility supports approximately 150 users, of whom about 10% are external, including industrial partners such as SABIC and ARAMCO. The KAUST maintains active collaborations with universities across Saudi Arabia and frequently hosts external researchers and industrial partners for training activities and instrument access. In addition, the shared core NMR platform at New York University Abu Dhabi (NYU-AD) houses four moderate field NMR spectrometers (11.7 T–14.1 T or 500–600 MHz) with solution and solid-state 600 MHz and a 600 MHz DNP NMR instrument, supporting about 10 internal users and external collaborations with academic institutions and industry partners. Operated by the Core Technology Platform (CTP) at NYU-AD, the primary research focus includes solid-state NMR studies of supramolecular assemblies, organic

crystals, MOFs and COFs, catalysts, membranes, polymers related materials, dynamic and self-healing crystals, NMR crystallography, and methodology development [185–194]. The overall facility usage rate is ~ 60%, with the 600 MHz wide-bore solid-state NMR being the most heavily utilized.

2.18. Multiple user facilities in South Africa

The NMR facilities at Stellenbosch University (Central Analytical Facilities, CAF), the University of the Witwatersrand, and Nelson Mandela University, are among the characterization hubs with solid-state NMR capabilities in South Africa. These labs support studies on clays, minerals, cements, catalysts, polymers, and biomaterials, addressing structural problems in energy, mining, and environmental sustainability [195–197]. Applications also include probing local structure in aluminosilicates, tracking cement hydration, studying catalytic active sites, and investigating drug–excipient interactions in pharmaceuticals.

2.19. Multiple SRFs in India (Vipin Agarwal)

India's progress in solid-state NMR research has been driven by growing technical expertise and improved access to both commercial solution and solid-state NMR instrumentation. Several SRFs are already in place at national research institutes and centers: Indian Institute of Science (IISc, Bangalore) and technologies (IITs), Indian Institute of Science and Research (IISER's), Tata Institute of Fundamental Research (Hyderabad), Centre of Biomedical Research (CBMR, Lucknow), Indian Institute of Chemical Technology (IICT, Hyderabad), and the National Magnetic Resonance Facility (NMRF, CSIR-NCL Pune). These SRFs have been proving access to support a broad range of experiments including static and MAS, multidimensional correlation spectroscopy including studies of spin-half and quadrupolar nuclei [23,198], and actively involved in designing novel solid-state NMR methods from first-principles [198–203]. The facilities are geographically distributed to enhance accessibility across the country. The IISc and the TIFR SRFs were established as early as the 1980s and these facilities today host eight spectrometers ranging from 400 to 800 MHz, with solution- and solid-state capabilities, while NCL and IICT SRFs host 9 and 10 spectrometers, respectively, aligned with the mission of supporting NMR access to researchers both in academia and industry nationwide. In addition, the Industrial Research and Consultancy Centre at IIT Powai hosts a 750 MHz solid-state NMR spectrometer, accessible to both academic and industrial users. Most of the other centers operate solid-state NMR instruments in the 300–700 MHz range and are equipped with 1.3 mm MAS probes capable of spinning up to 65 kHz. Two facilities also house fast MAS probes operating at speeds up to 111 kHz, enabling high-resolution proton detection in solids. Many of these SRFs actively collaborate and support non-NMR researchers working in the area of pharmaceuticals, energy storage, catalysis, polymers, COFs, MOFs, biomolecules, and materials characterization [204–207]. Researchers affiliated with these SRFs maintain close collaboration through the National Magnetic Resonance Society (NMRS-India), established in 1995, which brings together more than 1,000 lifetime members from India and around the world including scientists from NMR, EPR, NQR, quantum computing, and related fields.

The SRFs in India also offer access to researchers, often at minimal cost, in order to promote wider scientific participation and community building, making an impact across disciplines. To promote awareness of basic principles, recent advancements, and applications in NMR, the SRFs frequently organize national-level training and workshops for users. TIFR biannually organizes the advanced workshop to train and attract new talent in the field, and NMRS organizes annual national symposia hosted at different centers across the country and also supports a variety of local events focused on specific areas within these disciplines, and to date featured over 30 annual symposia. Strengthening collaboration among academia, industry, and instrument

manufacturers, along with upgrading SRFs, are projected to be an important aspect optimizing resource sharing and promoting interdisciplinary innovation. Recently, academia-industry allies have significantly enhanced structural insight and quality assurance in applied sectors such as pharma, oil, renewable energy, and materials companies. To continue advancing the field, strategic investments are envisaged to expand access to dynamic nuclear polarization (DNP), ultrafast MAS, and high-field magnets.

2.20. Multiple SRFs in China (Guangjin Hou, Feng Deng)

Multiple user facilities for solid-state NMR have been established across China, supporting cutting-edge research in chemistry, material science, and structural biology. Major shared NMR facilities include the Beijing NMR Center, the Center for Physicochemical Analysis and Measurement (Beijing), the State Key Laboratory of Magnetic Resonance Spectroscopy and Imaging (Wuhan), the Shanghai Key Laboratory of Magnetic Resonance (Shanghai), the NMR center in Dalian Institute of Chemical Physics (DICP, Dalian), and the Lanzhou NMR center (Lanzhou), University of Science and Technology of China (USTC, Hefei), among others. These NMR centers host multiple standard solid-state NMR spectrometers operating in 300–600 MHz, with some facilities offering magnetic field up to 18.8 T (800 MHz), such as those in Beijing, Wuhan, and Dalian. These platforms are fully equipped for conventional solid-state NMR analysis as well as advanced experiments such as fast MAS, MAS-CAT, Laser heating VT experiments. Notably, three DNP MAS NMR instruments have been installed in Wuhan (400 MHz), Beijing (600 MHz) and Shanghai (400 MHz), with several additional systems currently on order by other NMR centers. Research groups within these centers, particularly in Dalian, Wuhan and Shanghai, are actively engaged in the development of advanced solid-state NMR methodologies [208–211]. Custom-built NMR equipment is also in operation, including hyperpolarized ^{129}Xe NMR (Wuhan, Dalian), parahydrogen-induced polarization NMR (Wuhan, Shanghai), high-temperature/high-pressure *in situ* MAS NMR (Dalian), *in situ* electrochemistry NMR probes (Dalian, Shanghai), among others. All of these solid-state NMR SRFs are accessible to domestic and international users from both academic and industrial sectors, either free of charge or at moderate fees according to the policies of each center.

Most SRFs maintain active collaborations with multidisciplinary researchers in the field of chemistry, materials science and the life sciences, and have supported numerous publications in areas such as catalysis, energy storage materials, functional materials, polymers, and biomolecules [212–216]. While China has now built a robust solid-state NMR infrastructure comprising over 100 spectrometers dedicated to solid-state NMR studies (most of which are within SRF platforms), the community continues to work toward a more efficient, equitable and user-friendly SRF network. Ongoing efforts aim to enhance resource and expertise sharing, and to promote cross-disciplinary research and innovation both nationally and globally.

In addition to the shared NMR platforms (primarily based on commercial spectrometers), China is also advancing the research and development of high-field NMR spectrometers. A prominent example is the high-field NMR facility located in Beijing, which is dedicated to developing an advanced hybrid superconducting magnet that integrates both low- and high-temperature superconducting technologies to achieve magnetic fields ranging from 25 to 32 T (some currently under construction), enabling NMR experiments at ^1H frequencies exceeding 1 GHz. Complementing this effort, the High Magnetic Field Laboratory (CHMFL) in Hefei operates steady-state magnets, including resistive, superconducting, and hybrid systems, with current capabilities reaching up to 45.22 T. The CHMFL hosts solid-state NMR systems operating between 18.8 and 23.2 T, and supports high-end applications in chemistry, materials, and medical imaging through the integration of hybrid magnets for NMR/MRI and surface-sensitive microscopy techniques for molecular-level investigations [35,211,213,217–229].

2.21. SRF in Singapore (Kai Xue)

The use of solid-state NMR spectroscopy is rapidly growing in Singapore's shared facilities and centers. For example, the high field NMR Centre at Nanyang Technological University (NTU), is emerging as a national hub for advanced NMR capabilities. The center hosts moderate to high-field ranging from 14.1 T to 18.8 T (600 MHz and 800 MHz) solid-state NMR spectrometers. For the solid-state NMR measurements, the facility is equipped with a comprehensive set of MAS probes (7 mm, 4 mm, 3.2 mm, 1.9 mm, and 0.7 mm) with varying MAS capabilities, a high-resolution (HR)-MAS probe, a cryogenic probe for solution-state NMR detection, and a microimaging system. During 2022–2024, the center supported a wide range of research projects across disciplines such as materials science, civil engineering, chemistry, physics, and structural biology. Since 2019, the center has contributed to over 50 publications in biological studies [230,231], catalysis [232], and materials [233–235]. In addition to serving NTU researchers, the center provides access to users from other national institutions, including the National University Singapore (NUS), the Agency for Science, Technology and Research (A*STAR), and maintains active collaborations with international partners from the UK, Germany, Belgium, Saudi Arabia, Malaysia, Thailand, and China [236–241]. In response to growing demand, the center is exploring future expansion, with a focus on advanced high-field NMR applications including 2D correlation experiments with indirect detection via proton and the study of low- γ nuclei and insensitive quadrupolar nuclei.

2.22. Multiple facilities in Australia and New Zealand

Examples of solid-state NMR facilities in Australia include University of New South Wales (UNSW), Monash University (MU), Deakin University, University of Queensland, and the Australian National University (ANU). These hubs offer multinuclear NMR instrumentation with MAS, variable-temperature control, and a suite of advanced pulse sequences, enabling users to characterize local structures and their impact in driving bulk properties in molecular solids, polymers, inorganic materials, and biological systems [242,243]. Facilities at UNSW and ANU involve in characterization of catalysts, battery and semiconductor materials, glasses, polymers, and porous frameworks [244,245].

In New Zealand, the distributed NMR platforms at the University of Auckland, Victoria University of Wellington, the University of Otago, and the University of Canterbury feature solid-state NMR instrumentation and expert staff who provide training. Applications span energy materials (batteries, catalysts, perovskites), organic electronics and polymers, geological and cementitious solids, pharmaceuticals, and agro/food systems (notably dairy and plant-based matrices) [246], often in partnership with Research Institutes such as GNS Science, Scion, and AgResearch.

2.23. SRF in Yokohama, Japan (Takanori Kigawa)

In Japan, the RIKEN Yokohama NMR Research Infrastructure (YNMR) offers access to thirteen high-performance NMR systems for researchers within and outside RIKEN, including those from industry, supporting diverse scientific fields such as chemistry, life sciences, materials science, environmental science, drug discovery, and healthcare [247–253]. As one of Japan's leading research infrastructures, YNMR houses three 900 MHz NMR systems, including a solid-state NMR system equipped with multiple probe heads with access to examine spin-half and quadrupolar nuclei including ^{17}O , ^{27}Al , ^{45}Sc , ^{93}Nb , and ^{139}La , to name a few. In addition, YNMR serves as the lead institution of the “NMR Platform,” a national network comprising eight NMR research infrastructures and four partner companies, supported by Japan's Ministry of Education, Culture, Sports, Science and Technology (MEXT). In the past five years (2020–2024), YNMR infrastructure has enabled access to 128 projects with 7253 access days, contributing to 108

publications. Among them, 24 projects and publications involve solid-state NMR research across a wide range of areas. In addition, efforts have also been expended to construct high field magnets for NMR applications [254].

3. Funding, organization, governance, and management

The national or regional SRFs are typically governed through multi-tiered structures of management, which strategically align with national research priorities. Facility directors and scientific advisory committees provide strategic and technical leadership, while oversight boards including representatives from academia, industry, and funding bodies, ensure alignment with broader scientific, economic, and societal goals. Engagement with industry can further strengthen the impact of SRFs by supporting commercially relevant research and development. Financial support for research, equipment, and infrastructure is usually provided by national or regional governments, which reflects the value of SRFs as public investments in scientific capacity, training, and competitiveness. In many cases, this funding covers the acquisition and installation of large-scale instrumentation, as well as ongoing needs such as maintenance, consumables, upgrades, user training, outreach, administrative costs, and sustainability initiatives.

Operationally, SRFs rely on strong coordination between administrative and scientific teams. Management oversees compliance, budgeting, safety, data governance, and strategic planning, while scientific staff maintain instrumentation, schedule magnet time, develop methods, and support user access, training, and data analysis. Instrument time (magnet days) is often organized thematically or methodologically to provide consistent access across disciplines and user groups.

Access to SRFs is typically regulated through open calls and peer-reviewed proposal systems. Researchers submit proposals detailing their scientific objectives, resource needs and specifications, and required spectrometer time. Proposals are evaluated on merit, feasibility, and alignment with SRF capabilities, while also considering user diversity (e.g., students vs. early-career scientists vs. senior PIs; industrial collaborations vs. fundamental research in strategic areas).

To sustain excellence in research dissemination and deliverables, SRFs prioritize robust feedback and careful monitoring of scientific output (i.e., publications, conference presentations, Ph.D. theses, etc.). In this regard, the SRFs provide much more than access to instrumentation and data collection. Training programs, such as workshops, tutorials, hands-on experimentation, and direct participation in outreach activities are of great value as they promote knowledge sharing, build operational proficiency, and support effective dissemination of research outcomes and facility capabilities. During such training programs, technical and scientific staff offer expert guidance tailored to user end-goals, help design experiments, and support data analysis using custom-built pipelines and software platforms. Staff play a central role in guiding experiment design, troubleshooting methodologies, and supporting data processing, which is especially important for researchers from institutions with limited in-house expertise. Remote-operation systems further expand access and collaborative capacity, a need highlighted and accelerated during the COVID-19 pandemic. Initiatives such as the HORIZON-EU REMOTE-NMR (R-NMR, a project by a consortium of 26 European NMR centers) aim to establish standardized remote access platforms across European partners.

In addition to supporting individual research groups and projects, SRFs are playing an increasingly important role in advancing open-science practices. Many facilities are implementing standardized data formats, rich metadata, and digital archiving frameworks aligned with FAIR (Findability, Accessibility, Interoperability and Reuse) principles. Since 2021, the MagLab encouraged users to adapt to the principles of FAIR data management, which strive to make high-field NMR data more accessible, interpretable, and reusable. Researchers are also encouraged to use the Open Science Framework (OSF), with support from partners including NSF, FSU and UF. In addition, the Network for Advanced NMR

(NAN, funded by the NSF) has taken a step forward with the launch of the NAN Data Archive and NAN Data Transport System (NDTS). These platforms ensure that NMR data are automatically saved, securely archived, organized, and linked to sample information, making them readily accessible and easier to track.

4. Impact of SRFs on multidisciplinary research

The broader impact of large-scale solid-state NMR SRFs stems from key advancements: (i) enhancements in sensitivity and resolution for investigating mass- or volume-limited samples, including experiments involving insensitive, unresponsive nuclides [22,26,55,255–265] and (ii) the integration of solid-state NMR with other complementary analytical techniques and computational modeling [42,266–275]. These advances, along with new real-time, *in situ* and *in operando* experiments, enable the investigation of molecular-level structure, dynamics, and reaction kinetics and their relation to physicochemical properties and emergent functions of materials—this extends from small organic molecules to complex hybrid materials to the largest polymeric and/or biological macromolecules [18,32,34,276,277]. In addition, solid-state NMR studies are increasingly investigating systems in their functional states such as biomacromolecules at the cellular levels, thin films and membranes of importance in energy storage and harvesting devices including batteries and solar cells, and the natural biomass and biominerals such as wood and bone [14,15,17,22,278–284].

A survey of the literature from 2015 to 2024 demonstrates a steady increase in the use of solid-state NMR across multiple domains (Fig. 2), including materials science, chemistry, pharmaceutical science, energy storage, biomolecular sciences, and catalysis. Certain research fields require substantially more NMR measurement time, staff expertise, and iterative experiments to generate publishable data or resolve complex structures, making direct comparisons between publication output and instrument-time allocation challenging. Productivity per instrument hour varies significantly across disciplines. Although it is difficult to

precisely quantify the contributions of SRFs to the outcomes shown in Fig. 2, they play a central role in enabling broad access to advanced instrumentation, technical knowledge, and specialized methodologies that are often difficult for individual groups to realize and implement. For example, collaborative efforts across European SRFs have supported major scientific achievements within the PANACEA network, demonstrating the benefits of coordinated SRF support in driving community-wide advances during its first three years of activity [148], and discussion featured above highlight similar contributions from individual SRFs. In this way, SRFs complement the topic-specific research programs led by individual principal investigators.

A search using Clarivate® Web of Science with keywords “solid state NMR” and “solid-state NMR” displayed approximately 29k and 26k publications, respectively. These 29k publications span a broad range of chemistry topics (including physical, analytical, applied, organic, and inorganic chemistry) accounting for ~45 % of the total. Other areas include materials science (~11 %), biological/biomedical/pharmaceutical sciences (~7 %), multidisciplinary physics (~10 %), nanoscience (~5 %), and polymers (~4 %). While solid-state NMR is also employed in engineering, environmental science, and condensed-matter physics, its activity is limited in these domains. Overall, this distribution highlights the multidisciplinary breadth of solid-state NMR and its increasingly important role across diverse areas of contemporary research.

Fig. 3 shows publication trends over the past decade reflecting the use of solid-state NMR spectroscopy in different research areas. In chemistry and materials science, solid-state NMR has become indispensable for elucidating local structure and intermolecular interactions, with results consistently reported in 600–800 publications annually. The rise in materials characterization is particularly noteworthy, increasing from ~400 publications in 2015 to over 600 in 2024. Other fields including nanoscience/technology, catalysis and pharmaceuticals have seen growth, while outcomes of biomacromolecules and polymers have remained steady.

In material solids, properties such as crystallinity, defect structures,




Fig. 2. Distribution of solid-state NMR spectroscopy usage by research area (2015–2024), based on ~29k records from Clarivate® Web of Science.




Fig. 3. Trends in publication outcome across selected research areas (2015–2024).

grain boundaries, and phase heterogeneity are among the determinants of their performance. High-field solid-state NMR and DNP-enhanced NMR, in combination with complementary analytical techniques, make the connections between structures, properties, and function that are necessary for understanding complex systems, including semiconductor films, nano- and quantum materials, functional polymers, and organic-inorganic hybrid materials [15,285–293]. Solid-state NMR can also be used to monitor phase transitions and the effects of electronic, optical, and mechanical inputs and stressors, which are important for applications in electronics, photonics, and spintronics. *In situ* solid-state NMR studies under extreme conditions (temperature, pressure, illumination, etc.) can provide real-time insights into the effects of atomic and molecular rearrangements on phase transitions, chemical transformations, and/or degradation. Integration with other techniques and computational modeling (e.g., Monte-Carlo methods, molecular dynamics, first-principles calculations, ShiftML, AI/ML crystal growth calculations) serves to accelerate materials discovery in the aforementioned areas [42,266–274]. In nanotechnology, solid-state NMR has advanced understanding of hybrid nanomaterials, nanoparticle surfaces, and host–guest architectures, especially for energy storage and delivery [294–297]. Studies of glasses and disordered systems has advanced with multinuclear, and multidimensional experiments acquired with conventional or DNP NMR spectrometers, enabling unprecedented resolution of atomic-scale structure and dynamics [298–303]. These applications extend to the energy storage systems [304–306]. Combined with computational modeling, these approaches reveal local environments, medium-range order, and dynamics, providing insights into structure–property relationships in such amorphous materials. In addition, research on emerging molecular semiconductor such as metal halide perovskites [17,131,286,288,307–313], and organic semiconductors [262,314–321] benefit from solid-state NMR spectroscopy measurements and analysis for gaining insights into synthesis–structure–property relationships.

Protein and biomolecular applications have expanded from microcrystalline and fibrillar systems to complex assemblies such as membrane proteins, amyloids, biomaterials and large supramolecular complexes, providing atomic-level insights into structure and dynamics in native-like environments [258,284,322–326]. The SRFs equipped with solid-state NMR instrumentation have significantly contributed to structural biology, particularly in the study of amyloid fibrils and aggregates associated with neurodegenerative diseases such as Alzheimer's and Parkinson's. Solid-state NMR has uniquely enabled

atomic-resolution insights into complex and dynamic membrane proteins, information often inaccessible by means of other analytical techniques. Additionally, research on naturally occurring biopolymers, such as plant cellulose, nucleosides, and biomass-derived polymers, equally benefits from solid-state NMR or DNP NMR measurements and analysis [327–330].

In polymers, solid-state NMR spectroscopy enables quantitative assessment of crystallinity, chain packing, segmental mobility, and phase separation, directly linking microstructure to mechanical, thermal, and electronic properties [331–335]. In this way, soft matter and polymers, with their dynamic and multiscale structures, benefit from the ability of solid-state NMR to resolve conformational difference/changes, inter- and intramolecular interactions, and heterogeneous domains [293,336–342]. For thin films and coatings, it can probe buried interfaces, dopants, additives, detect subtle chemical modifications, and reveal molecular orientation, all of which are important for applications in electronics, packaging, and protective layers [262,318,321]. Its ability to study interfaces without selective labeling is particularly valuable for multi-component and heterogeneous systems. Another booming research area is thin film polymeric organic semiconductors for wider optoelectronics and energy paradigms [282,314,320,343,344]. Analysis of interfaces between multi-layer thin-films particularly benefit from solid-state NMR and DNP NMR [280], though challenges remain. The inherently low sensitivity of NMR can be limiting for the nanometer-to-micrometer thicknesses typical of thin films and coatings, where only microgram sample quantities are available. Achieving high resolution in complex, disordered, or multiphase materials often requires advanced isotopic labeling or fast sample spinning, which may be costly or impractical for routine use. Interfaces present additional difficulties due to signal overlap from bulk phases and low interfacial volume fractions.

Combined solid-state and DNP NMR investigations of heterogeneous catalysts in conjunction with modelling techniques have enabled atomic-level insights into these materials, guiding the design of efficient and robust catalysts for chemical conversions [345–350]. Complementary surface-sensitive techniques, such as X-ray photoelectron spectroscopy (XPS) and scanning tunneling microscopy (STM), work well with solid-state NMR to reveal transient states and interfacial interactions critical to catalytic function. In addition, porous materials such as MOFs and COFs have greatly benefited from the conventional solid-state and DNP NMR analysis, and modelling techniques, lending insight into not only the complex molecular frameworks, but also into

host-guest interactions and mobile components within MOFs that may have value in future applications in sensors or data storage devices [351–355].

In pharmaceutical research, solid-state NMR is entering a golden age, enabling the differentiation of polymorphs, identification of unique solvates, hydrates, and salts, and increasing our understanding of complex drug formulations [129,356–368]. Active pharmaceutical ingredients (APIs) and nutraceuticals can be investigated in both their “bulk” forms as well as in dosage formulations [252,357,367,369–371], especially using multinuclear solid-state NMR methods. Structure elucidation of both can be accomplished using 2D NMR analysis and/or NMR crystallographic approaches [372,373]. The DNP NMR has recently been shown to be useful for understanding the crystallization processes of APIs, as well as for the detection of APIs in low concentrations. An exciting frontier is the use of solid-state NMR to investigate protein-drug interactions and perform pharmacokinetics analyses—these may be game changers for both manufacturers and consumers of a variety of drug and nutraceutical products.

Industrially, solid-state NMR spectroscopy plays an increasingly vital role in the characterization of pharmaceutical polymorphs, drug formulations, catalysts, composite materials, and energy storage materials. Its non-destructive nature, sensitivity to the local structure of each atomic environment, and chemical specificity, make solid-state NMR an indispensable tool for quality control, failure analysis, regulatory compliance, and driving innovation across a plethora of commercial sectors. Noteworthy techniques include HR-MAS based diffusion NMR spectroscopy [374,375], dynamic magnetic resonance imaging (MRI) in chemical process and reaction engineering [34,276,277], enabling the *in situ* reaction monitoring and chemical separation. Importantly, several SRFs have enabled this research in the context of industrial research and developments by closely coordinating with pharma, chemical, energy, and advanced materials companies [80,280,376–379].

The spectrometer vendors such as Bruker and JEOL, related hardware vendors like Doty Scientific, Phoenix NMR, BlueSky NMR, ePROBE, and NMR SERVICE, have all played significant roles in shaping the trajectory of solid-state NMR research by driving both hardware and software innovations. These advances include static and MAS probes, higher magnetic fields, and low-temperature technologies have dramatically improved spectral resolution and sensitivity, and *in situ* and *operando* capabilities that enable time-resolved NMR data acquisition, permitting to carry out experiments that were previously inaccessible [21,25,51,54,380–387]. Beyond instrumentation, these vendors have fostered strong partnerships with the SRFs through collaborative projects, beta tests, workshops, sponsorships and training programs at conferences, thereby accelerating the adoption of cutting-edge methods. The SRFs also help producing spin-offs such as BlackFox (U.S. probe construction), QuadruFY (Canada, NQR), and Bridge12 (DNP). Their sustained contributions have not only advanced methodology, but also expanded the reach of solid-state NMR into emerging areas of chemistry, life sciences and materials paradigm.

In a broader context, multidisciplinary research areas involving chemistry, biochemistry, and advanced materials play a role in addressing Sustainable Development Goals (UNSDG) [388,389], where solid-state NMR is expected to find opportunities and challenges (Fig. 4). Among the list of 17 goals as of the time of writing this article, the aforementioned research publications and outcomes are linked to good health and well-being (~60 %), affordable and clean energy (~11 %), and climate action (~6 %). Research on biology, biomedical and pharmaceuticals particularly contribute towards these goals, while understanding structure-property relationships helps better design molecular/materials solids for clean energy and climate action. Additionally, industry innovation and infrastructure (UNSDG-09) are also expected to benefit from research on materials solids.




Fig. 4. Usage of solid-state NMR by research area (2015–2024) linked to the Sustainable Development Goals. Approximately 29k hits are partitioned based on Clarivate® Web of Science analysis.

5. Summary and looking forward

Looking ahead, current and future SRFs are expected to evolve as innovation centers that not only provide access to advanced instrumentation but enhance progress in magnetic resonance spectroscopy and imaging, enabling high-throughput characterization across disciplines. Sustainability will be central to this evolution—particularly for SRFs operating GHz-class magnets, cryogenic probes, and high-field DNP platforms—requiring continued efforts to minimize helium consumption, improve cryogenic efficiency, and modernize infrastructure in line with environmental and financial responsibility. Probe design will remain an important aspect in solid-state and DNP NMR applications by directly influencing sensitivity, resolution, and experimental capabilities at high magnetic fields and extreme conditions (*i.e.*, MAS frequencies >100 kHz, cryogenic temperatures, and high microwave-power DNP setups). The rotor size, bearing design, and RF coil configuration must be precisely optimized to ensure mechanical stability, high RF efficiency, and high-sensitivity detection on multiple channels, as well as accommodating microwave irradiation from high-power gyrotrons in the case of DNP MAS NMR probes. Moreover, customized probes and sample tubes (rotors) expand the types of experiments that can be performed, including multi-resonance detection (*e.g.*, HCN, HFX), *in situ* monitoring, or sample environments that mimic harsh conditions (*e.g.*, temperature, gas flow, new generation of cryogenic MAS probes, or electrochemical control). As materials, biological systems, and microstructured films become more complex—and sample quantities decrease—such tailored probe solutions will be crucial. Meeting these challenges will also require closer integration with complementary characterization techniques and computational modeling to link NMR observables with functional performance across materials, pharmaceutical, biological, catalytic, and energy-related fields.

Digital transformation including, real-time collaboration, and AI/ML-assisted experiment design, data collection, reducing access barriers and analysis is expected to gain popularity in NMR research [390–392]. While many of these capabilities are still emerging and implemented to varying degrees, they hold great promise for the future of the field. Emerging applications in autonomous NMR control, predictive experiment planning, automated analysis pipelines, and NMR-guided structure determination draw inspiration from parallel advances in materials informatics and biomedical imaging. Interlinked SRFs with shared expertise and complementary resources will play an increasingly important role in enabling these capabilities.

At the same time, SRFs may face mounting pressures from rising hardware costs, rapid technological turnover, and uneven user demand can strain resources and accelerate instrument usage. High-field NMR systems and DNP platforms demand frequent upgrades and place significant burdens on cryogen supply chains, energy use, and maintenance

budgets. Addressing these issues requires sustained investment in helium recovery, efficient cooling technology, predictive maintenance, and, importantly, specialized scientific and technical staff. Without dedicated personnel, even state-of-the-art SRFs risk reduced uptime, limited training capacity, and under-realized scientific impact. Achieving a balance between sustainability, operational reliability, and financial constraints will be essential to ensure the long-term viability of SRFs with current (and future) funding models.

In summary, solid-state NMR will remain indispensable for molecular-level characterization across scientific disciplines. To stay competitive and maximize impact, SRFs must continue to evolve strategically, strengthen collaborations, and embrace sustainable, technologically forward models of operation. If successful, they will provide significant benefits to fields spanning materials science, sustainable energy, health, catalysis, and quantum technologies, while contributing to a more collaborative, open, and resilient global research ecosystem.

Notes

The manuscript was written through contributions from all authors. The views and opinions expressed in this article are those of the authors and do not necessarily reflect those of the publisher.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

Research featuring the NHMFL NMR Facilities is supported by the National Science Foundation Cooperative Agreements (DMR-1644779, DMR-2128556) and the State of Florida. The NMR instruments housed in the OSU CCIC NMR Facility and the National Gateway Ultrahigh Field NMR Center were purchased using funds provided by the mid-scale research infrastructure award from the National Science Foundation (NSF DBI-1935913, Mid-Scale RI-1: 1.2 GHz NMR Spectrometer for National Gateway Ultrahigh Field NMR Center, PI: R.P. Bruschweiler; Co-PIs: M.P. Foster, P.J. Grandinetti, C.P. Jaroniec, B.S. Tolbert), the National Institutes of Health, the State of Ohio and The Ohio State University. NMRFAM at University of Wisconsin-Madison is supported by NIH P41GM136463 and R24GM141526 and the 1.1 GHz instrument was purchased with funds from the NSF Mid-Scale Research Infrastructure-II Award 1946970 (PI: J. Hoch; Co-PIs: A. Edison, K. Henzler-Wildman, C. M. Rienstra). Canadian SRFs acknowledge support from the Canada Foundation for Innovation, the Natural Sciences and Engineering Research Council of Canada, the National Research Council Canada, Laboratories Canada, the University of Ottawa, Bruker Biospin, and several other partners. Research featuring the UK High-Field Solid-State NMR Facility was funded by EPSRC and BBSRC (EP/T015063/1), as well as the University of Warwick including via part funding through Birmingham Science City Advanced Materials Projects 1 and 2 supported by Advantage West Midlands (AWM) and the European Regional Development Fund (ERDF) enabling access to 850 MHz, as well as, for the 1 GHz instrument, (EP/R029946/1). The French national research infrastructure Infranalytics acknowledges the support from CNRS Chemistry Department, and facilities features therein benefit from the regional (Fonds européen de développement régional, FEDER), national and European funding. SwedNMR is funded by the Swedish Research Council (contract 2021-00167) as a national research infrastructure and involves nine participating NMR partners: University of Gothenburg, Chalmers University of Technology, Lund University, Linköping University, Royal Institute of Technology (KTH), Stockholm University, Swedish University of Agricultural Sciences (SLU), Umeå University, Uppsala University. The Italian Research Facility CERM

acknowledge The project “Potentiating the Italian Capacity for Structural Biology Services in Instruct-ERIC, Acronym “ITACA.SB” (Project no. IR0000009) within the call MUR 3264/2021 PNRR M4/C2/L3.1.1, funded by the European Union – NextGenerationEU; MUR Dipartimenti di Eccellenza 2023-2027 (DICUS 2.0) to the Department of Chemistry “Ugo Schiff” of the University of Florence, and financial support by project INFRAIA-02-2020 PANACEA (H2020, contract number 101008500); Fondazione CR Firenze (<https://fondazionecrfirenze.it/en/>) is also acknowledged. This work benefited of the support and the use of resources of Instruct-ERIC, and specifically of the CERM/CIRMP Italy Centre. Portuguese NMR Centre benefits from funding from CICECO (Centre for Research in Ceramics and Composite Materials)–Aveiro Institute of Materials, and European Union ERC (Grant agreement ID: 865974) and PANACEA grants. Research benefited from PANACEA project acknowledges the funding from the EU Horizon 2020 INFRAIA-02-2020 call (*Integrating Activities for Starting Communities*). For the SRFs at Poland, the authors are grateful for the financial support from the EU Regional Operational Program of the Lodz Voivodeship, RPLD.01.01.00-10-0008/18, which enabled the purchase and upgrading of NMR equipment. The 1 GHz instrument and major upgrades to other high field instruments at University of Barcelona have been financed by grants ICT2021-006875 and ICT2022-7829 from the Ministry of Science, Innovation and Universities with funds from the European Union Next Generation EU/PRTR. The solid-state instruments in Madrid have been funded by the Spanish AEI and Fundación Ramon Areces. The solid-state NMR research in the UAE is supported by the Core Technology Platforms of New York University Abu Dhabi and Tamkeen. The solid-state NMR research at KAUST (King Abdullah University of Science and Technology) is supported by Core-lab facilities. Research involving solid-state NMR facilities is primarily supported by the Government of India through intramural funding from various research organizations, as well as through central science funding agencies such as the Department of Atomic Energy (DAE), Science and Engineering Research Board (SERB), Department of Science and Technology (DST), Department of Biotechnology (DBT), Council of Scientific and Industrial Research (CSIR), University Grants Commission (UGC), and the Indian Council of Medical Research (ICMR), among others. Research carried out on SRFs in China acknowledge the support from the National Natural Science Foundation of China (NSFC), the Ministry of Science and Technology of China (MOST), the Chinese Academy of Sciences (CAS), and the Ministry of Education of China (MOE), among others. Research facilities at Japan are supported by Japan’s Ministry of Education, Culture, Sports, Science and Technology (MEXT). In Argentina, funds for NMR facilities are provided by the National University of Cordoba and grants from national research agencies.

Data availability

Data will be made available on request.

References

- [1] E.R. Andrew, E. Szczesniak, A historical account of NMR in the solid state, *Prog. Nucl. Magn. Reson. Spectrosc.* 28 (1995) 11–36.
- [2] W.K. Rhim, A. Pines, J.S. Waugh, Violation of the spin-temperature hypothesis, *Phys. Rev. Lett.* 25 (1970) 218–220.
- [3] A. Pines, M.G. Gibby, J.S. Waugh, Proton-enhanced NMR of dilute spins in solids, *J. Chem. Phys.* 59 (1973) 569–590.
- [4] F.D. Doty, Probe Design and Construction, 2007 eMagRes.
- [5] R.W. Martin, J.E. Kelly, J.I. Kelz, Advances in instrumentation and methodology for solid-state NMR of biological assemblies, *J. Struct. Biol.* 206 (2019) 73–89.
- [6] D.L. Bryce, G.M. Bernard, M. Gee, M.D. Lumsden, K. Eichele, R.E. Wasylshen, Practical aspects of modern routine solid-state multinuclear magnetic resonance spectroscopy: one-dimensional experiments, *Can. J. Anal. Sci. Spect.* 46 (2001) 46–81.
- [7] D.D. Laws, H.-M.L. Bitter, A. Jerschow, Solid-state NMR spectroscopic methods in chemistry, *Angew. Chem. Int. Ed.* 41 (2002) 3096–3129.
- [8] B. Reif, S.E. Ashbrook, L. Emsley, M. Hong, Solid-state NMR spectroscopy, *Nat. Rev. Methods Primers* 1 (2021) 2.

- [9] D.C. Apperley, in: David C. Apperley, Robin K. Harris, Paul Hodgkinson (Eds.), *Solid-State NMR : Basic Principles & Practice*, first ed., Momentum Press, LLC, New York, 2012.
- [10] M.J. Duer, *Solid-state NMR spectroscopy principles and applications. Solid-State NMR Spectroscopy Principles and Applications*, 2001, pp. 563–567.
- [11] A. Medek, J.S. Harwood, L. Frydman, Multiple-quantum magic-angle spinning NMR: a new method for the study of quadrupolar nuclei in solids, *J. Am. Chem. Soc.* 117 (1995) 12779–12787.
- [12] A.P.M. Kentgens, A practical guide to solid-state NMR of half-integer quadrupolar nuclei with some applications to disordered systems, *Geoderma* 80 (1997) 271–306.
- [13] T.J. Simmons, J.C. Mortimer, O.D. Bernardinelli, A.-C. Pöppler, S.P. Brown, E. R. deAzevedo, R. Dupree, P. Dupree, Folding of xylan onto cellulose fibrils in plant cell walls revealed by solid-state NMR, *Nat. Commun.* 7 (2016) 13902.
- [14] X. Kang, A. Kirui, M.C. Dickwella Widanage, F. Mentink-Vigier, D.J. Cosgrove, T. Wang, Lignin-polysaccharide interactions in plant secondary cell walls revealed by solid-state NMR, *Nat. Commun.* 10 (2019) 347.
- [15] M. Seifrid, G.N.M. Reddy, B.F. Chmelka, G.C. Bazan, Insight into the structures and dynamics of organic semiconductors through solid-state NMR spectroscopy, *Nat. Rev. Mater.* 5 (2020) 910–930.
- [16] E.W. Zhao, T. Liu, E. Jónsson, J. Lee, I. Temprano, R.B. Jethwa, A. Wang, H. Smith, J. Carretero-González, Q. Song, C.P. Grey, In situ NMR metrology reveals reaction mechanisms in redox flow batteries, *Nature* 579 (2020) 224–228.
- [17] D.J. Kubicki, S.D. Stranks, C.P. Grey, L. Emsley, NMR spectroscopy probes microstructure, dynamics and doping of metal halide perovskites, *Nat. Rev. Chem.* 5 (2021) 624–645.
- [18] D. Lyu, Y. Jin, P.C.M.M. Magusin, S. Sturniolo, E.W. Zhao, S. Yamamoto, S. T. Keene, G.G. Malliaras, C.P. Grey, Operando NMR electrochemical gating studies of ion dynamics in PEDOT:PSS, *Nat. Mater.* 22 (2023) 746–753.
- [19] P. Wikus, W. Frantz, R. Kümmerle, P. Vonlanthen, Commercial gigahertz-class NMR magnets, *Supercond. Sci. Technol.* 35 (2022) 033001.
- [20] M.H. Levitt, Symmetry-Based Pulse Sequences in Magic-Angle Spinning Solid-State NMR, 2007 eMagRes.
- [21] A. Samoson, H.-MAS, J. Magn. Reson. 306 (2019) 167–172.
- [22] Z. Sun, C. Ollier, A. Rancz, B. Thienpont, K. Grohe, L. Becker, A. Pura, F. Engelke, S. Wegner, J. Sturgis, T. Polenova, T. Le Marchand, G. Pintacuda, Pushing the boundaries of resolution in solid-state nuclear magnetic resonance of biomolecules with 160 kHz magic-angle spinning, *J. Am. Chem. Soc.* 147 (2025) 19433–19437.
- [23] Y. Nishiyama, G. Hou, V. Agarwal, Y. Su, A. Ramamoorthy, Ultrafast magic angle spinning solid-state NMR spectroscopy: advances in methodology and applications, *Chem. Rev.* 123 (2023) 918–988.
- [24] T. Gopinath, K. Shin, Y. Tian, W. Im, J. Struppe, B. Perrone, A. Hassan, F. M. Marassi, Solid-state NMR MAS CryoProbe enables structural studies of human blood protein vitronectin bound to hydroxyapatite, *J. Struct. Biol.* 216 (2024) 108061.
- [25] B. Perrone, M. Gunnarsson, D. Bernin, E. Sparr, D. Topgaard, Cryogenic probe technology enables multidimensional solid-state NMR of the stratum corneum without isotope labeling, *Solid State Nucl. Magn. Reson.* 134 (2024) 101972.
- [26] S. Andrés-Campos, G.A. Titaux-Delgado, F.C. Escobedo-González, M. Mompeán, High-quality 13C-detected structural analysis of mass-limited amyloid samples using a CPMAS CryoProbe and moderate magnetic fields, *Solid State Nucl. Magn. Reson.* 139 (2025) 102028.
- [27] A.B. Barnes, G. De Paëpe, P.C.A. van der Wel, K.N. Hu, C.G. Joo, V.S. Bajaj, M. L. Mak-Jurkaskas, J.R. Sirigiri, J. Herzfeld, R.J. Temkin, R.G. Griffin, High-field dynamic nuclear polarization for solid and solution biological NMR, *Appl. Magn. Reson.* 34 (2008) 237–263.
- [28] A.J. Rossini, A. Zagdoun, M. Lelli, A. Lesage, C. Copéret, L. Emsley, Dynamic nuclear polarization surface enhanced NMR spectroscopy, *Acc. Chem. Res.* 46 (2013) 1942–1951.
- [29] A.G.M. Rankin, J. Trébosc, F. Pourpoint, J.-P. Amoureux, O. Lafon, Recent developments in MAS DNP-NMR of materials, *Solid State Nucl. Magn. Reson.* 101 (2019) 116–143.
- [30] C. Bessada, A.-L. Rollet, A. Rakhmatullin, I. Nuta, P. Florian, D. Massiot, In situ NMR approach of the local structure of molten materials at high temperature, *C. R.Chim.* 9 (2006) 374–380.
- [31] R. Tycko, NMR at low and ultralow temperatures, *Acc. Chem. Res.* 46 (2013) 1923–1932.
- [32] O. Pecher, J. Carretero-González, K.J. Griffith, C.P. Grey, Materials' methods: NMR in battery research, *Chem. Mater.* 29 (2017) 213–242.
- [33] L. Qi, R. Alamillo, W.A. Elliott, A. Andersen, D.W. Hoyt, E.D. Walter, K.S. Han, N. M. Washton, R.M. Rioux, J.A. Dumesic, S.L. Scott, Operando solid-state NMR observation of solvent-mediated adsorption-reaction of carbohydrates in zeolites, *ACS Catal.* 7 (2017) 3489–3500.
- [34] Q. Zheng, J.H. Williams, S.V. Elgersma, M.D. Mantle, A.J. Sederman, G. L. Bezemer, C.M. Guédon, L.F. Gladden, Operando characterisation of the products of fischer-tropsch synthesis in a fixed-bed reactor studied by magnetic resonance, *Catal. Today* 428 (2024) 114416.
- [35] D. Wu, X. Yao, L. Liang, Q. Deng, L. Han, G. Hou, W. Dai, Operando two-dimensional solid-state NMR unveils dual-template collaboration and competition in SAPO-34 zeolite crystallization, *J. Am. Chem. Soc.* 147 (2025) 18864–18875.
- [36] D.L. Bryce, *Modern NMR Crystallography: Concepts and Applications*, Royal Society of Chemistry, 2025.
- [37] S.E. Ashbrook, Concluding remarks: faraday discussion on NMR crystallography (and references therein), *Faraday Discuss* 255 (2025) 583–601.
- [38] L.J. Mueller, M.F. Dunn, NMR crystallography of enzyme active sites: probing chemically detailed, three-dimensional structure in tryptophan synthase, *Acc. Chem. Res.* 46 (2013) 2008–2017.
- [39] F.M. Paruzzo, A. Hofstetter, F. Musil, S. De, M. Ceriotti, L. Emsley, Chemical shifts in molecular solids by machine learning, *Nat. Commun.* 9 (2018) 4501.
- [40] A.R. Oganov, C.J. Pickard, Q. Zhu, R.J. Needs, Structure prediction drives materials discovery, *Nat. Rev. Mater.* 4 (2019) 331–348.
- [41] M. Cordova, P. Moutzouri, B. Simões de Almeida, D. Torodii, L. Emsley, Pure isotropic proton NMR spectra in solids using deep learning, *Angew. Chem. Int. Ed.* 62 (2023) e202216607.
- [42] C. Bonhomme, C. Gervais, F. Babonneau, C. Coelho, F. Pourpoint, T. Azaïs, S. E. Ashbrook, J.M. Griffin, J.R. Yates, F. Mauri, C.J. Pickard, First-principles calculation of NMR parameters using the gauge including projector augmented wave method: a chemist's point of view, *Chem. Rev.* 112 (2012) 5733–5779.
- [43] P. Moutzouri, M. Cordova, B. Simões de Almeida, D. Torodii, L. Emsley, Two-dimensional pure isotropic proton solid state NMR, *Angew. Chem. Int. Ed.* 62 (2023) e202301963.
- [44] M. Cordova, E.A. Engel, A. Stefaniuk, F. Paruzzo, A. Hofstetter, M. Ceriotti, L. Emsley, A machine learning model of chemical shifts for chemically and structurally diverse molecular solids, *J. Phys. Chem. C* 126 (2022) 16710–16720.
- [45] M. Kellner, J.B. Holmes, R. Rodriguez-Madrid, F. Viscosi, Y. Zhang, L. Emsley, M. Ceriotti, A deep learning model for chemical shieldings in molecular organic solids including anisotropy, *J. Phys. Chem. Lett.* 16 (2025) 8714–8722.
- [46] J.R. Miller, M.D. Bird, S. Bole, A. Bonito-Oliva, Y. Eyssa, W.J. Kenney, T. A. Painter, H.J. Schneider-Muntau, L.T. Summers, S.W.v. Sciver, S. Welton, R. J. Wood, J.E.C. Williams, S. Bobrov, Y. Iwasa, M. Leupold, V. Stejskal, R. Weggel, An overview of the 45-T hybrid magnet system for the new national high magnetic field laboratory, *IEEE Trans. Magn.* 30 (1994) 1563–1571.
- [47] Z. Gan, I. Hung, X. Wang, J. Paulino, G. Wu, I.M. Litvak, P.L. Gor'kov, W.W. Brey, P. Lendi, J.L. Schiano, M.D. Bird, I.R. Dixon, J. Toth, G.S. Boebinger, T.A. Cross, NMR spectroscopy up to 35.2T using a series-connected hybrid magnet, *J. Magn. Reson.* 284 (2017) 125–136.
- [48] R. Fu, W.W. Brey, K. Shetty, P. Gor'kov, S. Saha, J.R. Long, S.C. Grant, E. Y. Chekmenev, J. Hu, Z. Gan, M. Sharma, F. Zhang, T.M. Logan, R. Brüschweiler, A. Edison, A. Blue, I.R. Dixon, W.D. Markiewicz, T.A. Cross, Ultra-wide bore 900MHz high-resolution NMR at the national high magnetic field laboratory, *J. Magn. Reson.* 177 (2005) 1–8.
- [49] T. Dubroca, A.N. Smith, K.J. Pike, S. Froud, R. Wylde, B. Trociewicz, J. McKay, F. Mentink-Vigier, J. van Tol, S. Wi, W. Brey, J.R. Long, L. Frydman, S. Hill, A quasi-optical and corrugated waveguide microwave transmission system for simultaneous dynamic nuclear polarization NMR on two separate 14.1 T spectrometers, *J. Magn. Reson.* 289 (2018) 35–44.
- [50] P.L. Gor'kov, E.Y. Chekmenev, C. Li, M. Cotten, J.J. Buffry, N.J. Traaseth, G. Veglia, W.W. Brey, Using low-E resonators to reduce RF heating in biological samples for static solid-state NMR up to 900MHz, *J. Magn. Reson.* 185 (2007) 77–93.
- [51] S. Wang, T. Ravula, J.A. Stringer, P.L. Gor'kov, O.A. Warmuth, C.G. Williams, A. F. Thome, L.J. Mueller, C.M. Rienstra, Ultrahigh-resolution solid-state NMR for high-molecular weight proteins on GHz-scale spectrometers, *Sci. Adv.* 11 (2025) ead6016.
- [52] B.D. Harding, A.M. Barclay, D.W. Piehl, A. Hielt, O.A. Warmuth, R. Han, K. Henzler-Wildman, C.M. Rienstra, Cross polarization stability in multidimensional NMR spectroscopy of biological solids, *J. Magn. Reson.* 365 (2024) 107724.
- [53] C.G. Borcik, B. DeZonia, T. Ravula, B.D. Harding, R. Garg, C.M. Rienstra, OPTO: automated optimization for solid-state NMR spectroscopy, *J. Am. Chem. Soc.* 147 (2025) 3293–3303.
- [54] J. Schönzart, R. Han, T. Gennett, C.M. Rienstra, J.A. Stringer, Magnetic susceptibility modeling of magic-angle spinning modules for part per billion scale field homogeneity, *J. Magn. Reson.* 364 (2024) 107704.
- [55] R. Han, C.G. Borcik, S. Wang, O.A. Warmuth, K. Geohring, C. Mullen, M. Incitti, J. A. Stringer, C.M. Rienstra, Solid-state NMR ¹³C sensitivity at high magnetic field, *J. Magn. Reson.* 365 (2024) 107709.
- [56] A. Barde, R. Han, M.A. Olson, M. Tonelli, C.M. Rienstra, K.A. Henzler-Wildman, T. Ravula, In situ light-driven pH modulation for NMR studies, *Angew. Chem. Int. Ed.* 64 (2025) e202501440.
- [57] D.W. Conroy, Y. Xu, H. Shi, N. Gonzalez Salguero, R.N. Purusottam, M. D. Shannon, H.M. Al-Hashimi, C.P. Jaronec, Probing watson-crick and hoogsteen base pairing in duplex DNA using dynamic nuclear polarization solid-state NMR spectroscopy, *Proc. Natl. Acad. Sci. U. S. A.* 119 (2022) e2200681119.
- [58] S.T. Smrt, N. Gonzalez Salguero, J.K. Thomas, M. Zandian, M.G. Poirier, C. P. Jaronec, Histone H3 core domain in chromatin with different DNA linker lengths studied by ¹H-Detected solid-state NMR spectroscopy, *Front. Mol. Biosci.* 9 (2023), 2022.
- [59] J. Cheng, S. Venkatesh, K. Ke, M.J. Barratt, J.I. Gordon, A human gut *Faecalibacterium prausnitzii* fatty acid amide hydrolase, *Science* 386 (2024) eado6828.
- [60] P. Gupta, C.E. Moore, C. Turro, Photo- and electrocatalytic hydrogen evolution by heteroleptic Dirhodium(II,II) complexes: role of the bridging and diimine ligands, *J. Am. Chem. Soc.* 146 (2024) 27161–27172.
- [61] S.C. Yuan, Z. Álvarez, S.R. Lee, R.Z. Pavlović, C. Yuan, E. Singer, S.J. Weigand, L. C. Palmer, S.I. Stupp, Supramolecular motion enables chondrogenic bioactivity of a cyclic peptide mimetic of transforming growth Factor- β 1, *J. Am. Chem. Soc.* 146 (2024) 21555–21567.

- [62] R. Muzquiz, C. Jamshidi, D.W. Conroy, C.P. Jaroniec, M.P. Foster, Insights into ligand-mediated activation of an oligomeric ring-shaped gene-regulatory protein from solution- and solid-state NMR, *J. Mol. Biol.* 436 (2024) 168792.
- [63] V. Sridharan, T. George, D.W. Conroy, Z. Shaffer, W.K. Surewicz, C.P. Jaroniec, Copper binding alters the core structure of amyloid fibrils formed by Y145Stop human prion protein, *Phys. Chem. Chem. Phys.* 26 (2024) 26489–26496.
- [64] N. Kumar, T.J. Finnegan, S. Taneja, D.C. Rostam, A.L. Hansen, C.E. Ward, S. Uçar, J.R. Parquette, J.D. Badjić, Dendritic molecular baskets for selective binding of toxic methotrexate, *Angew. Chem. Int. Ed.* 64 (2025) e202420574.
- [65] Z. Du, H. Wang, S. Luo, Z. Yun, C. Wu, W. Yang, M. Buck, W. Zheng, A.L. Hansen, H.-Y. Kao, S. Yang, The sequence–structure–function relationship of intrinsic ERα disorder, *Nature* 638 (2025) 1130–1138.
- [66] C.P. Romao, F.A. Perras, U. Werner-Zwanziger, J.A. Lussier, K.J. Miller, C. M. Calahoo, J.W. Zwanziger, M. Bieringer, B.A. Marinkovic, D.L. Bryce, M. A. White, Zero thermal expansion in $\text{ZrMgMo}_3\text{O}_{12}$: NMR crystallography reveals origins of thermoelastic properties, *Chem. Mater.* 27 (2015) 2633–2646.
- [67] K. Zhu, C.A. O'Keefe, V.N. Vukotic, R.W. Schurko, S.J. Loeb, A molecular shuttle that operates inside a metal–organic framework, *Nat. Chem.* 7 (2015) 514–519.
- [68] H. Fenniri, G.A. Tikhomirov, D.H. Brouwer, S. Bouatra, M. El Bakkari, Z. Yan, J.-Y. Cho, T. Yamazaki, High field solid-state NMR spectroscopy investigation of ^{15}N -Labeled Rosette Nanotubes: Hydrogen Bond Network and Channel-Bound Water, *J. Am. Chem. Soc.* 138 (2016) 6115–6118.
- [69] J. Lu, I. Hung, A. Brinkmann, Z. Gan, X. Kong, G. Wu, Solid-state ^{17}O NMR Reveals Hydrogen-Bonding Energetics: Not All Low-Barrier Hydrogen Bonds Are Strong, *Angew. Chem. Int. Ed.* 56 (2017) 6166–6170.
- [70] B.E.G. Lucier, S. Chen, Y. Huang, Characterization of metal–organic frameworks: unlocking the potential of solid-state NMR, *Acc. Chem. Res.* 51 (2018) 319–330.
- [71] S. Chen, S. Mukherjee, B.E.G. Lucier, Y. Guo, Y.T.A. Wong, V.V. Terskikh, M. J. Zaworotko, Y. Huang, Cleaving carboxyls: understanding thermally triggered hierarchical pores in the metal–organic framework MIL-121, *J. Am. Chem. Soc.* 141 (2019) 14257–14271.
- [72] A. Bahadoor, A. Brinkmann, J.E. Melanson, ^{13}C -Satellite decoupling strategies for improving accuracy in quantitative nuclear magnetic resonance, *Anal. Chem.* 93 (2021) 851–858.
- [73] A. Karmakar, A. Bhattacharya, D. Sarkar, G.M. Bernard, A. Mar, V.K. Michaelis, Influence of hidden halogen mobility on local structure of $\text{CsSn}(\text{Cl}_{1-x}\text{Br}_x)_3$ mixed-halide perovskites by solid-state NMR, *Chem. Sci.* 12 (2021) 3253.
- [74] D. Sarkar, A. Bhattacharya, J. Meyer, A.M. Kirchberger, V. Mishra, T. Nilges, V. K. Michaelis, Unraveling sodium-ion dynamics in honeycomb-layered $\text{Na}_2\text{Mg}_x\text{Zn}_{2-x}\text{TeO}_6$ solid electrolytes with solid-state NMR, *J. Am. Chem. Soc.* 145 (2023) 19727–19745.
- [75] T. Nag, V.V. Terskikh, D.L. Bryce, Experimental evidence for non-Fermi-contact J coupling across chalcogen bonds in ionic salt cocrystal polymorphs, *Angew. Chem. Int. Ed.* 63 (2024) e202402441.
- [76] S.A. Southern, V. Kumar, V. Terskikh, D.L. Bryce, A. Brinkmann, NMR crystallographic investigation coupled with molecular dynamics simulations reveals the nature of disorder in chlorpromazine hydrochloride solvatomorphs, *Mol. Pharm.* 22 (2025) 4693–4707.
- [77] H.A. Davies, J. Madine, D.A. Middleton, Comparisons with amyloid-beta; reveal an aspartate residue that stabilizes fibrils of the aortic amyloid peptide medin, *J. Biol. Chem.* 290 (2015) 7791–7803.
- [78] J.M. Lamley, C. Öster, R.A. Stevens, J.R. Lewandowski, Intermolecular interactions and protein dynamics by solid-state NMR spectroscopy, *Angew. Chem. Int. Ed.* 54 (2015) 15374–15378.
- [79] A. Pugliese, M. Toresco, D. McNamara, D. Iuga, A. Abraham, M. Tobyn, L. E. Hawarden, F. Blanc, Drug–polymer interactions in acetaminophen/hydroxypropylmethylcellulose acetyl succinate amorphous solid dispersions revealed by multidimensional multinuclear solid-state NMR spectroscopy, *Mol. Pharm.* 18 (2021) 3519–3531.
- [80] P.M.J. Szell, Z. Rehman, B.P. Tatman, L.P. Hughes, H. Blade, S.P. Brown, Exploring the potential of multinuclear solid-state ^1H , ^{13}C , and ^{35}Cl Magnetic Resonance To Characterize Static and Dynamic Disorders in Pharmaceutical Hydrochlorides, *ChemPhysChem* 24 (2023) e202200558.
- [81] G.N.M. Reddy, G.M. Peters, B.P. Tatman, T.S. Rajan, S.M. Kock, J. Zhang, B. G. Frenguelli, J.T. Davis, A. Marsh, S.P. Brown, Magic-angle spinning NMR spectroscopy provides insight into the impact of small molecule uptake by G-quartet hydrogels, *Mater. Adv.* 1 (2020) 2236–2247.
- [82] L. Yu, Y. Yoshimi, R. Cresswell, R. Wightman, J.J. Lyczakowski, L.F.L. Wilson, K. Ishida, K. Stott, X. Yu, S. Charalambous, J. Wurman-Rodrich, O.M. Terrett, S. P. Brown, R. Dupree, H. Temple, K.B.R.M. Krogh, P. Dupree, Eudicot primary cell wall glucomannan is related in synthesis, structure, and function to xyloglucan, *Plant Cell* 34 (2022) 4600–4622.
- [83] M. Bourdon, J.J. Lyczakowski, R. Cresswell, S. Amsbury, F. Vilaplana, M.-J. Le Guen, N. Follain, R. Wightman, C. Su, F. Alatorre-Cobos, M. Ritter, A. Liszka, O. M. Terrett, S.R. Yadav, A. Vatén, K. Nieminen, G. Eswaran, J. Alonso-Serra, K. H. Müller, D. Iuga, P.C. Miskolczi, L. Kalmbach, S. Otero, A.P. Mähönen, R. Bhalerao, V. Bulone, S.D. Mansfield, S. Hill, I. Burgert, J. Beaupierre, Y. Benitez-Alfonso, R. Dupree, P. Dupree, Y. Helariutta, Ectopic callose deposition into woody biomass modulates the nano-architecture of macrofibrils, *Nat. Plant.* 9 (2023) 1530–1546.
- [84] C.L. Jones, C.E. Hughes, H.H.M. Yeung, A. Paul, K.D.M. Harris, T.L. Easun, Exploiting in situ NMR to monitor the formation of a metal–organic framework, *Chem. Sci.* 12 (2021) 1486–1494.
- [85] Z. Wang, A.M. Sheveleva, D. Lee, Y. Chen, D. Iuga, W.T. Franks, Y. Ma, J. Li, L. Li, Y. Cheng, L.L. Daemen, S.J. Days, A.J. Ramirez-Cuesta, B. Han, A.S. Eggeman, E. J.L. McInnes, F. Tuna, S. Yang, M. Schröder, Modulation of uptake and reactivity of nitrogen dioxide in metal–organic framework materials, *Angew. Chem. Int. Ed.* 62 (2023) e202302602.
- [86] L. Corti, D. Iuga, J.B. Claridge, M.J. Rosseinsky, F. Blanc, Disorder and oxide ion diffusion mechanism in $\text{La}_{1.54}\text{Sr}_{0.46}\text{Ga}_3\text{O}_{7.27}$ melilite from nuclear magnetic resonance, *J. Am. Chem. Soc.* 145 (2023) 21817–21831.
- [87] B. Darminto, G.J. Rees, J. Cattermull, K. Hashi, M. Diaz-Lopez, N. Kuwata, S. J. Turrell, E. Milan, Y. Chart, C. Di Mino, H. Jeong Lee, A.L. Goodwin, M. Pasta, On the origin of the non-arrhenius Na-ion conductivity in Na_3OBr , *Angew. Chem. Int. Ed.* 62 (2023) e202314444.
- [88] C.M. Rice, O.J. Dovernor, R.E. Morris, S.E. Ashbrook, Investigations of the ADOR process using solid-state NMR spectroscopy, *Cryst. Growth Des.* 23 (2023) 8991–9000.
- [89] S. Zhou, B.M. Gallant, J. Zhang, Y. Shi, J. Smith, J.N. Drysdale, P. Therdkatanyuphong, M. Taddei, D.P. McCarthy, S. Barlow, R.C. Kilbride, A. Dasgupta, A.R. Marshall, J. Wang, D.J. Kubicki, D.S. Ginger, S.R. Marder, H. J. Snaith, Reactive passivation of wide-bandgap organic–inorganic perovskites with benzylamine, *J. Am. Chem. Soc.* 146 (2024) 27405–27416.
- [90] Z. Wang, H.R. Felstead, R.I. Troup, B. Linclau, P.T.F. Williamson, Lipophilicity modulations by fluorination correlate with membrane partitioning, *Angew. Chem. Int. Ed.* 62 (2023) e202301077.
- [91] B.P. Tatman, W.T. Franks, S.P. Brown, J.R. Lewandowski, Nuclear spin diffusion under fast magic-angle spinning in solid-state NMR, *J. Chem. Phys.* 158 (2023) 184201.
- [92] C. Legein, B.J. Morgan, F. Fayon, T. Koketsu, J. Ma, M. Body, V. Sarou-Kanian, X.-K. Wei, M. Heggen, O.J. Borkiewicz, P. Strasser, D. Dambournet, Atomic insights into aluminium-ion insertion in defective anatase for batteries, *Angew. Chem. Int. Ed.* 59 (2020) 19247–19253.
- [93] C. Doyen, E. Larquet, P.-D. Coeureux, O. Frances, F. Herman, S. Sablé, J.-P. Burnouf, C. Sizun, E. Lescop, Nuclear magnetic resonance spectroscopy: a multifaceted toolbox to probe structure, dynamics, interactions, and real-time in situ release kinetics in peptide-liposome formulations, *Mol. Pharm.* 18 (2021) 2521–2539.
- [94] G.G. Moreira, F.-X. Cantrelle, A. Quezada, F.S. Carvalho, J.S. Cristóvão, U. Sengupta, N. Puangmalai, A.P. Carapeto, M.S. Rodrigues, I. Cardoso, G. Fritz, F. Herrera, R. Kaye, I. Landrieu, C.M. Gomes, Dynamic interactions and Ca^{2+} -binding modulate the holdase-type chaperone activity of S100B preventing tau aggregation and seeding, *Nat. Commun.* 12 (2021) 6292.
- [95] A. Lends, N. Birlirakis, X. Cai, A. Daskalov, J. Shenoy, M.B. Abdul-Shukoor, M. Berbon, F. Ferrage, Y. Liu, A. Loquet, K.O. Tan, Efficient 18.8 T MAS-DNP NMR reveals hidden side chains in amyloid fibrils, *J. Biomol. NMR* 77 (2023) 121–130.
- [96] L.M. Becker, M. Berbon, A. Vallet, A. Grelard, E. Morvan, B. Bardiaux, R. Lichtenacker, M. Ernst, A. Loquet, P. Schanda, The rigid core and flexible surface of amyloid fibrils probed by magic-angle-spinning NMR spectroscopy of aromatic residues, *Angew. Chem. Int. Ed.* 62 (2023) e202219314.
- [97] P. Raval, M.A. Akhavan Kazemi, J. Ruellou, J. Trébosc, O. Lafon, L. Delevoye, F. Sauvage, G.N.M. Reddy, Examining a year-long chemical degradation process and reaction kinetics in pristine and defect-passivated lead halide perovskites, *Chem. Mater.* 35 (2023) 2904–2917.
- [98] A. Maity, J. Wulfel, I. Ayala, A. Favier, Y. Adam, D. Bourgeois, B. Brutscher, Structural heterogeneity in a phototransformable fluorescent protein impacts its photochemical properties, *Adv. Sci.* 11 (2024) 2306272.
- [99] L. Niccoli, G. Casano, G. Menzildjian, M. Yulikov, T. Robinson, S.-E. Akrial, Z. Wang, C. Reiter, A. Pura, D. Siri, A. Venkatesh, L. Emson, D. Gajan, M. Lelli, O. Ouari, A. Lesage, Efficient DNP at high fields and fast MAS with antenna-sensitized dinitroxides, *Chem. Sci.* 15 (2024) 16582–16593.
- [100] T.J. Matemb M Ntep, M. Wahiduzzaman, E. Laurenz, I. Cornu, G. Mouchaham, I. Dovgaliuk, S. Nandi, K. Knop, C. Jansen, F. Nouar, P. Florian, G. Földner, G. Maurin, C. Janiak, C. Serre, When polymorphism in metal–organic frameworks enables water sorption profile tunability for enhancing heat allocation and water harvesting performance, *Adv. Mater.* 36 (2024) 2211302.
- [101] K. Weinhäupl, L. Meuret, S. Desrat, F. Roussi, N. Morellet, S. Beaupierre, C. Guillou, C. van Heijenoort, O. Abian, S. Vega, I. Wolf, T. Akopian, O. Krandor, E. Rubin, A. Velazquez-Campoy, D. Gauto, H. Fraga, Identification of new ClpC1-NTD binders for Mycobacterium tuberculosis drug development, *Sci. Rep.* 15 (2025) 4146.
- [102] A. Peach, N. Fabregue, C. Erre, T.-X. Métro, D. Gajan, F. Mentink-Vigier, F. Scott, J. Trébosc, F. Voron, N. Patris, C. Gervais, D. Laurencin, Capturing and labeling CO_2 in a jar: mechanochemical ^{17}O -enrichment and ssNMR study of sodium and potassium (bi)carbonate salts, *Chem. Sci.* 16 (2025) 10731–10741.
- [103] A.U. Mushtaq, J. Áden, L.A. Clifton, H. Wacklin-Knecht, M. Campana, A.P. G. Dingeldein, C. Persson, T. Sparrman, G. Gröbner, Neutron reflectometry and NMR spectroscopy of full-length Bcl-2 protein reveal its membrane localization and conformation, *Commun. Biol.* 4 (2021) 507.
- [104] H. Serk, M.B. Nilsson, J. Figueira, J.P. Krüger, J. Leifeld, C. Alewell, J. Schleucher, Organochemical characterization of peat reveals decomposition of specific hemicellulose structures as the main cause of organic matter loss in the acrotelm, *Environ. Sci. Technol.* 56 (2022) 17410–17419.
- [105] D.H. De Oliveira, V. Gowda, T. Sparrman, L. Gustafsson, R. Sanches Pires, C. Riekel, A. Barth, C. Lendel, M. Hedhammar, Structural conversion of the spidroin C-terminal domain during assembly of spider silk fibers, *Nat. Commun.* 15 (2024) 4670.
- [106] K. Nam, A.R.A. Thodika, S. Tschlik, C. Phoeurk, T.M. Nagy, L. Schierholz, J. Áden, P. Rogne, M. Drescher, A.E. Sauer-Eriksson, M. Wolf-Watz, Magnesium induced structural reorganization in the active site of adenylate kinase, *Sci. Adv.* 10 (2024) eado5504.

- [107] F. Guerroudj, L. Fliri, J. Bengtsson, L.C. Gomes, T. Gazzola, M. Hummel, D. Bernin, Applications of NMR based methodologies investigating the behavior of lignin and cellulose towards bio-based carbon fibers production, *Solid State Nucl. Magn. Reson.* 134 (2024) 101977.
- [108] P. Petermeier, J.P. Bittner, T. Jonsson, P. Domínguez de María, E. Byström, S. Kara, Integrated preservation of water activity as key to intensified chemoenzymatic synthesis of bio-based styrene derivatives, *Commun. Chem.* 7 (2024) 57.
- [109] V.M. Dinh, S.G. Khokarale, P.O. May, T. Sparman, K. Irgum, J.-P. Mikkola, Ionic liquid strategy for chitosan production from chitin and molecular insights, *RSC Sustain.* 2 (2024) 1154–1164.
- [110] N. Altincekic, N. Jores, F. Löhr, C. Richter, C. Ehrhardt, M.J.J. Blommers, H. Berg, S. Öztürk, S.L. Gande, V. Linhard, J. Orts, M.J. Abi Saad, M. Bütkofer, J. Kaderli, B.G. Karlsson, U. Brath, M. Hedenström, G. Gröbner, U.H. Sauer, A. Perrakis, J. Langer, L. Banci, F. Cantini, M. Fragai, D. Grifagni, T. Barthel, J. Wollenhaupt, M.S. Weiss, A. Robertson, A. Bax, S. Sreeramulu, H. Schwalbe, Targeting the main protease (mpro, nsp5) by growth of fragment scaffolds exploiting structure-based methodologies, *ACS Chem. Biol.* 19 (2024) 563–574.
- [111] A. Thirunavukkarasu, M. Hedenström, T. Sparman, M.B. Nilsson, J. Schleucher, M. Öquist, Unraveling the dynamics of lignin chemistry on decomposition to understand its contribution to soil organic matter accumulation, *Plant Soil* 511 (2025) 1485–1502.
- [112] S.-M. Argyri, L. Svenningsson, F. Guerroudj, D. Bernin, L. Evenäs, R. Bordes, Contact-free magnetic resonance imaging and spectroscopy with acoustic levitation, *Nat. Commun.* 16 (2025) 3917.
- [113] T. Kobayashi, M. Kanmi, P. Piotr, N.-K. Agnieszka, S. Justyna, N. Yusuke, G. D T, M.J. P, P. Marek, Study of intermolecular interactions in the corrole matrix by solid-state NMR under 100 kHz MAS and theoretical calculations, *Angew. Chem. Int. Ed.* 52 (2013) 14108–14111.
- [114] P. Paluch, T. Pawlak, J.-P. Amoureux, M.J. Potrzebowski, Simple and accurate determination of X-H distances under ultra-fast MAS NMR, *J. Magn. Reson.* 233 (2013) 56–63.
- [115] T. Pawlak, M.J. Potrzebowski, Fine refinement of solid-state molecular structures of Leu- and met-enkephalins by NMR crystallography, *J. Phys. Chem. B* 118 (2014) 3298–3309.
- [116] E. Skorupska, P. Paluch, A. Jeziorna, M.J. Potrzebowski, NMR study of BA/FBA cocrystal confined within mesoporous silica nanoparticles employing thermal solid phase transformation, *J. Phys. Chem. C* 119 (2015) 8652–8661.
- [117] P. Paluch, T. Pawlak, M. Oszejka, W. Lasocha, M.J. Potrzebowski, Fine refinement of solid state structure of racemic form of phospho-tyrosine employing NMR crystallography approach, *Solid State Nucl. Magn. Reson.* 65 (2015) 2–11.
- [118] P. Paluch, T. Pawlak, A. Jeziorna, J. Trébosc, G. Hou, A.J. Vega, J.-P. Amoureux, M. Dracinsky, T. Polenova, M.J. Potrzebowski, Analysis of local molecular motions of aromatic sidechains in proteins by 2D and 3D fast MAS NMR spectroscopy and quantum mechanical calculations, *Phys. Chem. Chem. Phys.* 17 (2015) 28789–28801.
- [119] K. Trzeciak, S. Kaźmierski, E. Wielgus, M.J. Potrzebowski, DiSUpLo - new extremely easy and efficient method for loading of active pharmaceutical ingredients into the pores of MCM-41 mesoporous silica particles, *Microporous Mesoporous Mater.* 308 (2020) 110506.
- [120] T. Pawlak, I. Sudgen, G. Bujacz, D. Iuga, S.P. Brown, M.J. Potrzebowski, Synergy of solid-state NMR, single-crystal X-ray diffraction, and crystal structure prediction methods: a case study of teriflunomide (TFM), *Cryst. Growth Des.* 21 (2021) 3328–3343.
- [121] T. Pawlak, M. Oszejka, M. Szczesio, M.J. Potrzebowski, Solid-state study of the structure, dynamics, and thermal processes of safinamide mesylate—a new generation drug for the treatment of neurodegenerative diseases, *Mol. Pharm.* 19 (2022) 287–302.
- [122] M.K. Dudek, K. Trzeciak, L. Tajber, J. Zajac, S. Kaźmierski, E. Pindelska, T. Makowski, M. Svyntkivska, M.J. Potrzebowski, A new look at the mechanism of cocrystal formation and cofomers exchange in processes forced by mechanical and/or thermal stimuli – ex situ and in situ studies of low-melting eutectic mixtures, *Chem. Eur. J.* 30 (2024) e202302138.
- [123] T. Kobayashi, K. Mao, P. Paluch, A. Nowak-Król, J. Sniechowska, Y. Nishiyama, D. T. Gryko, M.J. Potrzebowski, M. Pruski, Study of intermolecular interactions in the corrole matrix by solid-state NMR under 100 kHz MAS and theoretical calculations, *Angew. Chem. Int. Ed.* 52 (2013) 14108–14111.
- [124] B.A. Atterberry, P. Paluch, A.R. Lamkins, W. Huang, A.J. Rossini, Rapid acquisition of ^{103}Rh solid-state NMR spectra by ^{31}P detection and sideband selective methods, *J. Am. Chem. Soc.* 147 (2025) 14411–14421.
- [125] I. Bertini, A. Bhaumik, G. De Paëpe, R.G. Griffin, M. Lelli, J.R. Lewandowski, C. Luchinat, High-resolution solid-state NMR structure of a 17.6 kDa protein, *J. Am. Chem. Soc.* 132 (2010) 1032–1040.
- [126] I. Bertini, L. Gonnelli, C. Luchinat, J. Mao, A. Nesi, A new structural model of A β 40 fibrils, *J. Am. Chem. Soc.* 133 (2011) 16013–16022.
- [127] L. Cerofolini, E. Ravera, S. Bologna, T. Wiglenda, A. Böddrich, B. Purfürst, I. Benilova, M. Korsak, G. Gallo, D. Rizzo, L. Gonnelli, M. Fragai, B. De Strooper, E.E. Wanker, C. Luchinat, Mixing A β (1–40) and A β (1–42) peptides generates unique amyloid fibrils, *Chem. Commun.* 56 (2020) 8830–8833.
- [128] R. Shukla, F. Lavore, S. Maity, M.G.N. Derks, C.R. Jones, B.J.A. Vermeulen, A. Melcrová, M.A. Morris, L.M. Becker, X. Wang, R. Kumar, J. Medeiros-Silva, R. A.M. van Beekveld, A.M.J.J. Bonvin, J.H. Lorent, M. Lelli, J.S. Nowick, H. D. MacGillivray, A.J. Peoples, A.L. Spoering, L.L. Ling, D.E. Hughes, W.H. Roos, E. Breukink, K. Lewis, M. Weingarh, Teixobactin kills bacteria by a two-pronged attack on the cell envelope, *Nature* 608 (2022) 390–396.
- [129] R. Shukla, A.J. Peoples, K.C. Ludwig, S. Maity, M.G.N. Derks, S. De Benedetti, A. M. Krueger, B.J.A. Vermeulen, T. Harbig, F. Lavore, R. Kumar, R.V. Honorato, F. Grein, K. Nieselt, Y. Liu, A.M.J.J. Bonvin, M. Baldus, U. Kubitschek, E. Breukink, C. Achorn, A. Nitti, C.J. Schwalen, A.L. Spoering, L.L. Ling, D. Hughes, M. Lelli, W.H. Roos, K. Lewis, T. Schneider, M. Weingarh, An antibiotic from an uncultured bacterium binds to an immutable target, *Cell* 186 (2023) 4059–4073.e4027.
- [130] F. Bruno, L. Gigli, G. Ferraro, A. Cavallo, V.K. Michaelis, G. Goobes, E. Fratini, E. Ravera, Lysozyme is sterically trapped within the silica cage in bioinspired silica-lysozyme composites: a multi-technique understanding of elusive protein-material interactions, *Langmuir* 38 (2022) 8030–8037.
- [131] A. Karmakar, M.S. Dodd, S. Agnihotri, E. Ravera, V.K. Michaelis, Cu(II)-Doped Cs $_2$ SbAgCl $_6$ double perovskite: a lead-free, low-bandgap material, *Chem. Mater.* 30 (2018) 8280–8290.
- [132] G. Provinciali, N.A. Consoli, R. Caliendo, V. Mangini, L. Barba, C. Giannini, G. Tuci, G. Giambastiani, M. Lelli, A. Rossin, Ammonia borane and hydrazine bis (borane) confined within zirconium bithiazole and bipyridyl metal-organic frameworks as chemical hydrogen storage materials, *J. Phys. Chem. C* 129 (2025) 6094–6108.
- [133] A. Lund, G. Casano, G. Menzildjian, M. Kaushik, G. Stevanato, M. Yulikov, R. Jabbour, D. Wisser, M. Renom-Carrasco, C. Thieuleux, F. Bernada, H. Karoui, D. Siri, M. Rosay, I.V. Sergeyev, D. Gajan, M. Lelli, L. Emsley, O. Ouari, A. Lesage, TinyPols: a family of water-soluble binitroxides tailored for dynamic nuclear polarization enhanced NMR spectroscopy at 18.8 and 21.1 T, *Chem. Sci.* 11 (2020) 2810–2818.
- [134] G. Liu, M. Levien, N. Karschin, G. Parigi, C. Luchinat, M. Bennati, One-thousand-fold enhancement of high field liquid nuclear magnetic resonance signals at room temperature, *Nat. Chem.* 9 (2017) 676–680.
- [135] A. Zagdoun, G. Casano, O. Ouari, M. Schwarzwälder, A.J. Rossini, F. Aussenac, M. Yulikov, G. Jeschke, C. Copéret, A. Lesage, P. Tordo, L. Emsley, Large molecular weight nitroxide biradicals providing efficient dynamic nuclear polarization at temperatures up to 200 K, *J. Am. Chem. Soc.* 135 (2013) 12790–12797.
- [136] D. Pereira, M. Sardo, I. Marín-Montesinos, L. Mafra, One-shot resin 3D-Printed stators for low-cost fabrication of magic-angle spinning NMR probeheads, *Anal. Chem.* 95 (2023) 10384–10389.
- [137] D. Pereira, M. Sardo, R. Vieira, I. Marín-Montesinos, L. Mafra, Enhancing CO $_2$ capture via fast microwave-assisted synthesis of the CALF-20 metal-organic framework, *Inorg. Chem.* 64 (2025) 3302–3310.
- [138] A. Santamaria, D. Pereira, N. Pawar, B.T. Kelly, J. Carrascosa-Tejedor, M. Sardo, L. Mafra, G. Fragneto, D.J. Owen, I. Marín-Montesinos, E. Guzmán, N.R. Zaccari, A. Maestro, Structural molecular details of the endocytic adaptor protein CALM upon binding with phosphatidylinositol 4,5-bisphosphate-containing model membranes, *Commun. Chem.* 8 (2025) 219.
- [139] P. Zelenovskii, M. Soares, C. Bornes, I. Marín-Montesinos, M. Sardo, S. Kopyl, A. Kholkin, L. Mafra, F. Figueiredo, Detection of helical water flows in sub-nanometer channels, *Nat. Commun.* 15 (2024) 5516.
- [140] M. Sardo, T. Morais, M. Soares, R. Vieira, M. Ilkaeva, M.A.O. Lourenço, I. Marín-Montesinos, L. Mafra, Unravelling the structure of CO $_2$ in silica adsorbents: an NMR and computational perspective, *Chem. Commun.* 60 (2024) 4015–4035.
- [141] M.M.A. Sacramento, M.B. Oliveira, J.R.B. Gomes, J. Borges, B.R. Freedman, D. J. Mooney, J.M.M. Rodrigues, J.F. Mano, Natural polymer-polyphenol bioadhesive coacervate with stable wet adhesion, antibacterial activity, and On-Demand detachment, *Adv. Healthcare Mater.* 13 (2024) 2304587.
- [142] S. Magina, S. Marques, F. Gírio, A. Lourenço, A. Barros-Timmons, D.V. Evtuguin, Chemical composition and structural features of cellolignin from steam explosion followed by enzymatic hydrolysis of Eucalyptus globulus bark, *Ind. Crop. Prod.* 211 (2024) 118217.
- [143] R.P. Monteiro, I.B. Calhau, A.C. Gomes, A.D. Lopes, J.P. Da Silva, I.S. Gonçalves, M. Pillinger, β -Cyclodextrin and cucurbit[7]uril as protective encapsulation agents of the CO-releasing molecule [CpMo(CO) $_3$ Me], *Dalton Trans* 54 (2025) 166–180.
- [144] M.M. Antunes, A.C. Gomes, I.S. Gonçalves, M. Pillinger, A.A. Valente, Catalytic sulfoxidation and epoxidation using oxomolybdenum compounds generated from tricarboxyl(1-methylimidazole)molybdenum(0), *Appl. Catal., A* 704 (2025) 120372.
- [145] R. Bastos, I. Marín-Montesinos, S.S. Ferreira, F. Mentink-Vigier, M. Sardo, L. Mafra, M.A. Coimbra, E. Coelho, Covalent connectivity of glycogen in brewer's spent yeast cell walls revealed by enzymatic approaches and dynamic nuclear polarization NMR, *Carbohydr. Polym.* 324 (2024) 121475.
- [146] D. Pereira, R. Fonseca, I. Marín-Montesinos, M. Sardo, L. Mafra, Understanding CO $_2$ adsorption mechanisms in porous adsorbents: a solid-state NMR survey, *Curr. Opin. Colloid Interface Sci.* 64 (2023) 101690.
- [147] R. Fonseca, R. Vieira, M. Sardo, I. Marín-Montesinos, L. Mafra, Exploring molecular dynamics of adsorbed CO $_2$ species in amine-modified porous silica by solid-state NMR relaxation, *J. Phys. Chem. C* 126 (2022) 12582–12591.
- [148] R. Coulon, D. Gajan, W. Papawassiliou, A.J. Pell, J. Schlagnitweit, F. Fayon, P. Florian, D. Massiot, A. Afrough, D.W. Juhl, T. Vosegaard, L. Cerofolini, M. Lelli, M. Lucci, C. Luchinat, R.L.E.G. Aspers, J.S. Gómez, A.P.M. Kentgens, S.F. H. Lambregts, Y.T. Angel Wong, L. Mafra, I. Marín-Montesinos, J. Rocha, M. Sardo, U. Brath, G. Karlsson, A.C. Pinon, S. Schantz, M. Šoltésová, S. Bachmann, S.P. Brown, D. Iuga, W. Trent Franks, A. Menakath, L. Frydman, F. Mentink-Vigier, R.W. Schurko, K. Grohe, F. Engelke, J. Kempf, A. Purea, C. Reiter, S. Wegner, V. Castro, C. Cobas, D. Jeannerat, F. Seoane, E. Vaz, D. Jardon Alvarez, M. Leskes, F. De Biasi, G. Pintacuda, A. Lesage, Transforming

- solid-state nuclear magnetic resonance towards a chemistry-ready technique, *Solid State Nucl. Magn. Reson.* 140 (2025) 102048.
- [149] I. Serrano-Chacón, B. Mir, L. Cupellini, F. Colizzi, M. Orozco, N. Escaja, C. González, pH-Dependent capping interactions induce large-scale structural transitions in i-Motifs, *J. Am. Chem. Soc.* 145 (2023) 3696–3705.
- [150] J. Carrasco, R. Antón, A. Valbuena, D. Pantoja-Uceda, M. Mukhi, R. Hervás, D. V. Laurents, M. Gasset, J. Oroz, Metamorphism in TDP-43 prion-like domain determines chaperone recognition, *Nat. Commun.* 14 (2023) 466.
- [151] M.P. Lenza, L. Egia-Mendikute, A. Antoñana-Vildosola, C.O. Soares, H. Coelho, F. Corzana, A. Bosch, P. Manisha, J.I. Quintana, I. Oyenarte, L. Unione, M. J. Moure, M. Azkargorta, U. Atxabal, K. Sobczak, F. Elortza, J.D. Sutherland, R. Barrio, F. Marcelo, J. Jiménez-Barbero, A. Palazon, J. Ereño-Orbea, Structural insights into Siglec-15 reveal glycosylation dependency for its interaction with T cells through integrin CD11b, *Nat. Commun.* 14 (2023) 3496.
- [152] M.P. Lenza, I. Oyenarte, T. Diercks, J.I. Quintana, A. Gimeno, H. Coelho, A. Diniz, F. Peccati, S. Delgado, A. Bosch, M. Valle, O. Millet, N.G.A. Abrescia, A. Palazon, F. Marcelo, G. Jiménez-Osés, J. Jiménez-Barbero, A. Ardá, J. Ereño-Orbea, Structural characterization of N-Linked glycans in the receptor binding domain of the SARS-CoV-2 spike protein and their interactions with human lectins, *Angew. Chem. Int. Ed.* 59 (2020) 23763–23771.
- [153] R. Gil-Redondo, R. Conde, C. Bruzzzone, M.L. Seco, M. Bizkarguenaga, B. González-Valle, A. de Diego, A. Laín, H. Habisch, C. Haudum, N. Verheyen, B. Obermayer-Pietsch, S. Margarita, S. Pelusi, I. Verde, N. Oliveira, A. Sousa, A. Zabala-Letona, A. Santos-Martin, A. Loizaga-Iriarte, M. Unda-Urzaiz, J. Kazenwadel, G. Berezchnoy, T. Geisler, M. Gawaz, C. Cannet, H. Schäfer, T. Diercks, C. Trautwein, A. Carracedo, T. Madl, L. Valenti, M. Spraul, S.C. Lu, N. Embade, J.M. Mato, O. Millet, MetSCORE: a molecular metric to evaluate the risk of metabolic syndrome based on serum NMR metabolomics, *Cardiovasc. Diabetol.* 23 (2024) 272.
- [154] M. Arbesú, M. Maffei, T.N. Cordeiro, J.M.C. Teixeira, Y. Pérez, P. Bernadó, S. Roche, M. Pons, The unique domain forms a fuzzy intramolecular complex in src family kinases, *Structure* 25 (2017) 630–640.e634.
- [155] A. Fernández, M. Gairí, M.T. González, M. Pons, A fast method to monitor tyrosine kinase inhibitor mechanisms, *J. Med. Chem.* 67 (2024) 20571–20579.
- [156] B. Mir, M. Gairí, M.T. González, A. Vila-Planas, M. Mayzel, M. Pons, M. Poggio, M. Terrazas, NMR characterization of the diastereomeric composition of a model therapeutic oligonucleotide, *J. Pharm. Sci.* 114 (2025).
- [157] L.C. Maia, L.C. Soares, M.M.C.E. Carvalho, G.R. dos Santos, E.R. de Azevedo, J.d. S. Soares, L.V.A. Gurgel, Synthesis and application of an unprecedented bioadsorbent for removing arsenic from aqueous systems, *Sep. Purif. Technol.* 324 (2023) 124495.
- [158] G. Dognani, A.S. Gomes, G. de Oliveira Setti, C.S. Martin, R.J.G. Rubira, C.J. L. Constantino, Functionalized cellulose microfibers as reductant agent to gold nanoparticles and its application on SERRS for sensitive detection of phthalocyanine dye, *J. Environ. Chem. Eng.* 11 (2023) 111266.
- [159] R. Borges, T. Zambanini, A.M. Pelosine, G.Z. Justo, A.C.S. Souza, J. Machado, J. F. Schneider, D.R. de Araujo, J. Marchi, A colloidal hydrogel-based drug delivery system overcomes the limitation of combining bisphosphonates with bioactive glasses: in vitro evidence of a potential selective bone cancer treatment allied with bone regeneration, *Biomater. Adv.* 151 (2023) 213441.
- [160] M.J. Ferreira, J.F. Schneider, Solid-state vitrification of LiPO₃ through mechanical milling, *Mater. Chem. Phys.* 328 (2024) 129958.
- [161] C.C.d.M. Capetti, V.O.A. Pellegrini, M.C. Espirito Santo, A.A. Cortez, M. Falvo, A. A.d.S. Curvelo, E. Campos, J.G. Filgueiras, F.E.G. Guimaraes, E.R. de Azevedo, I. Polikarpov, Enzymatic production of xylooligosaccharides from corn cobs: assessment of two different pretreatment strategies, *Carbohydr. Polym.* 299 (2023) 120174.
- [162] A. Gaddam, G. Tricot, P. Gołębiewski, H.R. Fernandes, R. Buczyński, J.M. F. Ferreira, H. Eckert, Structural organization of phase-separated bioactive glasses and the clustering of Si, P, B, Na and F atoms investigated by solid-state NMR and monte carlo simulations, *Acta Mater.* 259 (2023) 119203.
- [163] R. Frenzel, D. Morales, G. Romanelli, G. Sathicq, M. Blanco, L. Pizzio, Synthesis, characterization and catalytic evaluation of H₃PW₁₂O₄₀ included in acrylic acid/acrylamide polymer for the selective oxidation of sulfides, *J. Mol. Catal. Chem.* 420 (2016) 124–133.
- [164] V. Luca, J. Veliscek-Carolan, New insights into the radiolytic stability of metal(IV) phosphonate hybrid adsorbent materials, *Phys. Chem. Chem. Phys.* 22 (2020) 17027–17032.
- [165] C. Leal Marchena, G.P. Ortenzi, L.B. Pierella, L.R. Pizzio, Synthesis of novel photocatalysts based on polyoxovanadates supported onto zeolites employed for azo dye degradation, *ChemistrySelect* 9 (2024) e202404545.
- [166] R.N. Costa, A.L. Reviglio, S. Siedler, S.G. Cardoso, Y.G. Linck, G.A. Monti, A.M. G. Carvalho, J.A.L.C. Resende, M.H.C. Chaves, H.V.A. Rocha, D. Choquesillo-Lazarte, L. Infantes, S.L. Cuffini, New multicomponent forms of the antiretroviral nevirapine with improved dissolution performance, *cryst. Growth Des.* 20 (2020) 688–698.
- [167] C.M. Sánchez, A.K. Chattah, K.X. Wei, L. Buljubasich, P. Cappellaro, H. M. Pastawski, Perturbation independent decay of the loschmidt echo in a many-body system, *Phys. Rev. Lett.* 124 (2020) 030601.
- [168] A. Ruderman, S. Smrekar, M.V. Bracamonte, E.N. Primo, G.L. Luque, J. Thomas, E. Leiva, G.A. Monti, D.E. Barraco, F. Vaca Chávez, Unveiling the stability of Sn/Si/graphite composites for Li-ion storage by physical, electrochemical and computational tools, *Phys. Chem. Chem. Phys.* 23 (2021) 3281–3289.
- [169] E. Fuentes-Quezada, S.A. Maldonado Ochoa, R.H. Acosta, M. Bruno, E. de la Llave, M.P. Longinotti, F. Vaca Chávez, H.R. Corti, Effect of the carbon mesoporous structure on the transport properties of confined lithium chloride aqueous solutions, *Microporous Mesoporous Mater.* 323 (2021) 111255.
- [170] C.M. Sánchez, A.K. Chattah, H.M. Pastawski, Emergent decoherence induced by quantum chaos in a many-body system: a Loschmidt echo observation through NMR, *Phys. Rev. A* 105 (2022) 052232.
- [171] Y. Garro Linck, G. Martínez Delfa, J.A. Donadelli, E.V. Silletta, M.I. Velasco, M. B. Franzoni, G.A. Monti, C. Smal, R.H. Acosta, Total organic carbon determination by ¹³C CP/MAS Solid-State Solid-sN Unconventional Oil/Gas Source ROSS: A Case Study from the Neuquén and Cuyo Basins in Argentina, *Energy Fuel.* 38 (2024) 5098–5105.
- [172] Y. Yan, D.I. Kolokolov, I. da Silva, A.G. Stepanov, A.J. Blake, A. Dailly, P. Manuel, C.C. Tang, S. Yang, M. Schröder, Porous metal-organic polyhedral frameworks with optimal molecular dynamics and pore geometry for methane storage, *J. Am. Chem. Soc.* 139 (2017) 13349–13360.
- [173] A. Knebel, B. Geppert, K. Volkmann, D.I. Kolokolov, A.G. Stepanov, J. Twiefel, P. Heitjans, D. Volkmer, J. Caro, Defibrillation of soft porous metal-organic frameworks with electric fields, *Science* 358 (2017) 347–351.
- [174] D.I. Kolokolov, A.G. Stepanov, H. Jobic, Mobility of the 2-Methylimidazole linkers in ZIF-8 probed by ²H NMR: Saloon doors for the guests, *J. Phys. Chem. C* 119 (2015) 27512–27520.
- [175] Y.G. Kolyagin, A.V. Yakimov, S. Tolborg, P.N.R. Vennestrom, I.I. Ivanova, Direct observation of tin in different T-Sites of Sn-BEA by One- and two-dimensional ¹¹⁹Sn MAS NMR Spectroscopy, *J. Phys. Chem. Lett.* 9 (2018) 3738–3743.
- [176] Y.G. Kolyagin, A.V. Yakimov, D.S. Zasukhin, I.I. Ivanova, Direct identification of zeolite beta polymorphs by solid-state ²⁹Si Nuclear Magnetic Resonance Spectroscopy, *J. Phys. Chem. Lett.* 13 (2022) 10793–10798.
- [177] E. Papulovskiy, S.D. Kirik, D.F. Khabibulin, A.A. Shubin, V.M. Bondareva, A. S. Samoilov, O.B. Lapina, Condensation of ammonium niobium oxalate studied by NMR crystallography and X-ray powder diffraction, *Catal. Today* 354 (2020) 26–35.
- [178] O.B. Lapina, Modern ssNMR for heterogeneous catalysis, *Catal. Today* 285 (2017) 179–193.
- [179] A.A. Gabrienko, S.S. Arzumano, A.V. Toktarev, I.G. Danilova, I.P. Prosvirni, V. V. Krivontsov, V.I. Zaikovskii, D. Freude, A.G. Stepanov, Different efficiency of Zn²⁺ and ZnO species for methane activation on Zn-Modified zeolite, *ACS Catal.* 7 (2017) 1818–1830.
- [180] Z.N. Lashchinskaya, A.A. Gabrienko, A.G. Stepanov, Mechanisms of light alkene oligomerization on metal-modified zeolites, *ACS Catal.* 14 (2024) 4984–4998.
- [181] I.I. Ivanova, Y.G. Kolyagin, I.A. Kasyanov, A.V. Yakimov, T.O. Bok, D.N. Zarubin, Time-resolved in situ MAS NMR monitoring of the nucleation and growth of zeolite BEA catalysts under hydrothermal conditions, *Angew. Chem. Int. Ed.* 56 (2017) 15344–15347.
- [182] I.I. Ivanova, Y.G. Kolyagin, Application of multinuclear MAS NMR for the in situ monitoring of hydrothermal synthesis of zeolites, *Chem. Eur. J.* 27 (2021) 14143–14167.
- [183] S. Zhou, O. Shekhah, A. Ramírez, P. Lyu, E. Abou-Hamad, J. Jia, J. Li, P.M. Bhatt, Z. Huang, H. Jiang, T. Jin, G. Maurin, J. Gascon, M. Eddaoudi, Asymmetric pore windows in MOF membranes for natural gas valorization, *Nature* 606 (2022) 706–712.
- [184] I. Yarullina, K. De Wispelaere, S. Bailleul, J. Goetze, M. Radersma, E. Abou-Hamad, I. Vollmer, M. Goesten, B. Mezari, E.J.M. Hensen, J.S. Martínez-Espín, M. Morten, S. Mitchell, J. Perez-Ramirez, U. Olsbye, B.M. Weckhuysen, V. Van Speybroeck, F. Kapteijn, J. Gascon, Structure–performance descriptors and the role of Lewis acidity in the methanol-to-propylene process, *Nat. Chem.* 10 (2018) 804–812.
- [185] R. Mathew, J. Gupta, M.D. Devadarsan, S. Mavi, M. Jasil, J.A. Fereiro, B. Joseph, K.N. Parida, B. Thomas, Dynamic nuclear polarization of TEMPO radical cross conjugated with a thioanisole scaffold, *Solid State Nucl. Magn. Reson.* 137 (2025) 102005.
- [186] A. Jrad, G. Das, N. Alkhatib, T. Prakasam, F. Benyettou, S. Varghese, F. Gándara, M. Olson, S. Kirmizialtin, A. Trabolsi, Cationic covalent organic framework for the fluorescent sensing and cooperative adsorption of perfluorooctanoic acid, *Nat. Commun.* 15 (2024) 10490.
- [187] R. Mathew, B. Thomas, Deciphering the topology of sitagliptin using an integrated approach, *ACS Omega* 10 (2025) 2289–2295.
- [188] K. Vinod, R. Mathew, C. Jandl, B. Thomas, M. Hariharan, Electron diffraction and solid-state NMR reveal the structure and exciton coupling in a eumelanin precursor, *Chem. Sci.* 15 (2024) 16015–16024.
- [189] R. Mathew, A. Mazumder, P. Kumar, J. Matula, S. Mohamed, P. Brazda, M. Hariharan, B. Thomas, Unveiling the topology of partially disordered micro-crystalline nitro-perylene diimide with X-aggregate stacking: an integrated approach, *Chem. Sci.* 15 (2024) 490–499.
- [190] R. Jabbour, S. Raran-Kurussi, V. Agarwal, A. Equbal, Tailoring solid-state DNP methods to the study of α -synuclein LLPS, *Biophys. Chem.* 313 (2024) 107303.
- [191] G. Das, S. Singha Roy, F. Abou Ibrahim, A. Merhi, H.N. Dirawi, F. Benyettou, A. Kumar Das, T. Prakasam, S. Varghese, S. Kumar Sharma, S. Kirmizialtin, R. Jagannathan, F. Gándara, S. Aouad, M.A. Olson, S. Kundu, B.R. Kaafarani, A. Trabolsi, Electrocatalytic water splitting in isoindigo-based covalent organic frameworks, *Angew. Chem. Int. Ed.* 64 (2025) e202419836.
- [192] P. Commins, M.B. Al-Handawi, D.P. Karothu, G. Raj, P. Naumov, Efficiently self-healing boronic ester crystals, *Chem. Sci.* 11 (2020) 2606–2613.
- [193] S. Schramm, D.P. Karothu, N.M. Lui, P. Commins, E. Ahmed, L. Catalano, L. Li, J. Weston, T. Moriwaki, K.M. Soltsev, P. Naumov, Thermochemiluminescent peroxide crystals, *Nat. Commun.* 10 (2019) 997.
- [194] P. Commins, A. Natarajan, C.-K. Tsai, S.I. Khan, N.K. Nath, P. Naumov, M. A. Garcia-Garibay, Structure–reactivity correlations and mechanistic

- understanding of the photorearrangement and photosalt effect of α -Santonin and its derivatives in solutions, crystals, and nanocrystalline suspensions, *Cryst. Growth Des.* 15 (2015) 1983–1990.
- [195] K. Kutzner, M. Meinken, Retrospective determination of exposure temperature of standing trees during wildfires with solid-state NMR, *Sci. Rep.* 14 (2024) 12821.
- [196] C. Reichhardt, L.-M. Joubert, K.V. Clemons, D.A. Stevens, L. Cegelski, Integration of electron microscopy and solid-state NMR analysis for new views and compositional parameters of *Aspergillus fumigatus* biofilms, *Med. Mycol.* 57 (2019) S239–S244.
- [197] M.E. Matthews, H. Duveskog, A.J. van Reenen, An investigation into the applicability of wide-line NMR for the determination of oil in waxes, *Macromol. Mater. Eng.* 309 (2024) 2300372.
- [198] S. Ahlawat, K.R. Mote, N.-A. Lakomek, V. Agarwal, Solid-state NMR: methods for biological solids, *Chem. Rev.* 122 (2022) 9643–9737.
- [199] V. Agarwal, The origin of negative cross-peaks in proton-spin diffusion spectrum of fully protonated solids at fast MAS: coherent or incoherent effect? *J. Magn. Reson.* 311 (2020) 106661.
- [200] M.G. Jain, K.R. Mote, J. Hellwagner, G. Rajalakshmi, M. Ernst, P.K. Madhu, V. Agarwal, Measuring strong one-bond dipolar couplings using REDOR in magic-angle spinning solid-state NMR, *J. Chem. Phys.* 150 (2019) 134201.
- [201] S. Ray, V.S. Redrouth, A. Equbal, S.K. Jain, Optimal control-based nuclear spin cross-polarization in the presence of complicating anisotropic interactions, *Phys. Chem. Chem. Phys.* 27 (2025) 7016–7027.
- [202] M.K. Pandey, Y. Nishiyama, A one-dimensional solid-state NMR approach for $^{15}\text{NH}/^{14}\text{NH}$ overtone correlation through $^1\text{H}/^1\text{H}$ mixing under fast MAS, *Phys. Chem. Chem. Phys.* 20 (2018) 25849–25853.
- [203] N.T. Duong, S. Raran-Kurussi, Y. Nishiyama, V. Agarwal, Quantitative ^1H - ^1H Distances in Protonated Solids by Frequency-selective Recoupling at Fast Magic-Angle Spinning NMR, *J. Phys. Chem. Lett.* 9 (2018) 5948–5954.
- [204] B. Patra, S. Narayanan, S. Halder, M. Sharma, D. Sachdeva, N. Ravishanker, S. K. Pati, S.K. Jain, P. Senguttuvan, NASICON- $\text{NaV}_{0.25}\text{Al}_{0.25}\text{Nb}_{1.5}(\text{PO}_4)_3/\text{C}$: a high-rate and robust anode for fast charging and long-life sodium-ion batteries, *Adv. Mater.* 37 (2025) 2419417.
- [205] S. Karak, K. Koner, A. Karmakar, S. Mohata, Y. Nishiyama, N.T. Duong, N. Thomas, T.G. Ajithkumar, M.S. Hossain, S. Bandyopadhyay, S. Kundu, R. Banerjee, Morphology tuning via linker modulation: metal-free covalent organic nanostructures with exceptional chemical stability for electrocatalytic water splitting, *Adv. Mater.* 36 (2024) 2209919.
- [206] Y.-H. Lin, N. Sakai, P. Da, J. Wu, H.C. Sansom, A.J. Ramadan, S. Mahesh, J. Liu, R. D.J. Oliver, J. Lim, L. Aspitarte, K. Sharma, P.K. Madhu, A.B. Morales-Vilches, P. K. Nayak, S. Bai, F. Gao, C.R.M. Grovenor, M.B. Johnston, J.G. Labram, J. R. Durrant, J.M. Ball, B. Wenger, B. Stannowski, H.J. Snaith, A piperidinium salt stabilizes efficient metal-halide perovskite solar cells, *Science* 369 (2020) 96–102.
- [207] C. Hareendran, P.S. Shaligram, R. Gonnade, T.G. Ajithkumar, A solid-state NMR method for characterization of pharmaceutical eutectics, *Phys. Chem. Chem. Phys.* 26 (2024) 3800–3803.
- [208] L. Liang, K. Chen, G. Hou, Highly efficient heteronuclear polarization transfer using dipolar-echo edited R-symmetry sequences in solid-state NMR, *Chem. Sci.* 16 (2025) 2251–2257.
- [209] L. Liang, Y. Ji, Z. Zhao, C.M. Quinn, X. Han, X. Bao, T. Polenova, G. Hou, Accurate heteronuclear distance measurements at all magic-angle spinning frequencies in solid-state NMR spectroscopy, *Chem. Sci.* 12 (2021) 11554–11564.
- [210] Q. Wang, J. Trébosc, Y. Li, O. Lafon, S. Xin, J. Xu, B. Hu, N. Feng, J.-P. Amoureux, F. Deng, Uniform signal enhancement in MAS NMR of half-integer quadrupolar nuclei using quadruple-frequency sweeps, *J. Magn. Reson.* 293 (2018) 92–103.
- [211] H. Xiao, J. Wang, H. Tan, Y. Gan, W. Liu, Y. Zhang, Z. Zhang, J. Yang, Robust heteronuclear correlations for sub-milligram protein in ultrafast magic-angle spinning solid-state NMR, *J. Am. Chem. Soc.* 147 (2025) 6384–6389.
- [212] G. Li, C. Foo, R. Fan, M. Zheng, Q. Wang, Y. Chu, J. Li, S. Day, P. Steadman, C. Tang, T.W.B. Lo, F. Deng, S.C.E. Tsang, Atomic Locations and Adsorbate Interactions of Al Single and Pair Sites in H-ZSM-5 Zeolite, vol. 387, 2025, pp. 388–393.
- [213] M.-Q. Zhang, Y. Zhou, R. Cao, S. Tian, Y. Jiao, Z. Guo, M. Wang, H. Peng, B. Sun, B. Xu, M. Wang, S. Xu, D. Ma, In-line NMR guided orthogonal transformation of real-life plastics, *Nature* 643 (2025) 395–403.
- [214] X. Qian, L. Chen, L. Yin, Z. Liu, S. Pei, F. Li, G. Hou, S. Chen, L. Song, K.H. Thebo, H.-M. Cheng, W. Ren, CdPS_3 nanosheets-based membrane with high proton conductivity enabled by Cd vacancies 370 (2020) 596–600.
- [215] Y. Ji, P. Gao, Z. Zhao, D. Xiao, Q. Han, H. Chen, K. Gong, K. Chen, X. Han, X. Bao, G. Hou, Oxygenate-based routes regulate syngas conversion over oxide-zeolite bifunctional catalysts, *Nat. Catal.* 5 (2022) 594–604.
- [216] F. Pan, K. Ni, T. Xu, H. Chen, Y. Wang, K. Gong, C. Liu, X. Li, M.-L. Lin, S. Li, X. Wang, W. Yan, W. Yin, P.-H. Tan, L. Sun, D. Yu, R.S. Ruoff, Y. Zhu, Long-range ordered porous carbons produced from C_{60} , *Nature* 614 (2023) 95–101.
- [217] W. Cai, C. Wang, Y. Chu, M. Hu, Q. Wang, J. Xu, F. Deng, Unveiling the Brønsted acid mechanism for Meerwein-Ponndorf-Verley reduction in methanol conversion over ZSM-5, *Nat. Commun.* 15 (2024) 8736.
- [218] D. Chen, L. Ma, Y. Chen, X. Zhou, S. Xing, Y. Deng, Y. Hao, C. Pu, X. Kong, Y. Jin, Electrochemically stable ligands of ZnO electron-transporting layers for quantum-dot light-emitting diodes, *Nano Lett.* 23 (2023) 1061–1067.
- [219] L. Gao, J. Chen, Q. Chen, X. Kong, The chemical evolution of solid electrolyte interface in sodium metal batteries, *Sci. Adv.* 8 (2022) eabm4606.
- [220] B. Niu, H. Wu, J. Yin, B. Wang, G. Wu, X. Kong, B. Yan, J. Yao, C.-Z. Li, H. Chen, Mitigating the lead leakage of high-performance perovskite solar cells via in situ polymerized networks, *ACS Energy Lett.* 6 (2021) 3443–3449.
- [221] Y. Sun, L. Liang, M. Yang, Y. Ji, G. Hou, K. Chen, Identification and quantification of Al pairs and their impact on dealumination in zeolites, *J. Am. Chem. Soc.* 147 (2025) 10160–10171.
- [222] H. Tan, W. Zhao, M. Duan, Y. Zhao, Y. Zhang, H. Xie, Q. Tong, J. Yang, Native cellular membranes facilitate channel activity of MscL by enhancing slow collective motions of its transmembrane helices, *J. Am. Chem. Soc.* 146 (2024) 31472–31485.
- [223] C. Wang, M. Zheng, M. Hu, W. Cai, Y. Chu, Q. Wang, J. Xu, F. Deng, Unraveling spatially dependent hydrophilicity and reactivity of confined carbocation intermediates during methanol conversion over ZSM-5 zeolite, *J. Am. Chem. Soc.* 146 (2024) 8688–8696.
- [224] Y. Xu, P. Peng, H. Wang, H. Xiong, Z. Xu, X. Chen, Y. Li, A. Zheng, Y. Wei, Z. Yan, S. Xu, Z. Liu, Revealing the crucial roles of pore interconnectivity between zeolitic and nonzeolitic components in enhancing diffusion and catalytic efficiency of industrial zeolite-based catalysts, *J. Am. Chem. Soc.* 147 (2025) 18550–18562.
- [225] M. Zheng, Q. Wang, Y. Chu, W. Wang, Y. Xi, Y. Wang, G. Qi, J. Xu, S. B. Hong, F. Deng, Revealing the Brønsted acidic nature of penta-coordinated aluminum species in dealuminated zeolite Y with solid-state NMR spectroscopy, *J. Am. Chem. Soc.* 146 (2024) 29417–29428.
- [226] X. Cui, X. Hu, C. Cui, Y. Zhang, Q. Wang, Research on the fiber intensity modulation model of superconducting magnetic levitation rotor, *Rev. Sci. Instrum.* 95 (2024) 055113.
- [227] X. Wu, X. Lou, C. Hu, F. Geng, J. Li, C. Li, B. Hu, Coupling of oxygen dimer and trapped O_2 with strong magnetic frustration in layered Li-rich cathodes, *ACS Energy Lett.* 10 (2025) 459–468.
- [228] F. Geng, B. Hu, C. Li, C. Zhao, O. Lafon, J. Trébosc, J.-P. Amoureux, M. Shen, B. Hu, Anionic redox reactions and structural degradation in a cation-disordered rock-salt $\text{Li}_{1.2}\text{Ti}_{0.4}\text{Mn}_{0.4}\text{O}_2$ cathode material revealed by solid-state NMR and EPR, *J. Mater. Chem. A* 8 (2020) 16515–16526.
- [229] G. Li, C. Foo, R. Fan, M. Zheng, Q. Wang, Y. Chu, J. Li, S. Day, P. Steadman, C. Tang, T.W.B. Lo, F. Deng, S.C.E. Tsang, Atomic locations and adsorbate interactions of Al single and pair sites in H-ZSM-5 zeolite, *Science* 387 (2025) 388–393.
- [230] M. Shoaib, Q. Chen, X. Shi, N. Nair, C. Prasanna, R. Yang, D. Walter, K. S. Frederiksen, H. Einarsson, J.P. Svensson, C.F. Liu, K. Ekwall, M. Lerdurp, L. Nordenskiöld, C.S. Sørensen, Histone H4 lysine 20 mono-methylation directly facilitates chromatin openness and promotes transcription of housekeeping genes, *Nat. Commun.* 12 (2021) 4800.
- [231] X. Shi, C. Prasanna, T. Nagashima, T. Yamazaki, K. Pervushin, L. Nordenskiöld, Structure and dynamics in the nucleosome revealed by solid-state NMR, *Angew. Chem. Int. Ed.* 57 (2018) 9734–9738.
- [232] C. Lv, L. Zhong, H. Liu, Z. Fang, C. Yan, M. Chen, Y. Kong, C. Lee, D. Liu, S. Li, J. Liu, L. Song, G. Chen, Q. Yan, G. Yu, Selective electrocatalytic synthesis of urea with nitrate and carbon dioxide, *Nat. Sustain.* 4 (2021) 868–876.
- [233] T.R. Rana, M. Abbas, E. Schwartz, F. Jiang, M.Y. Yaman, Z. Xu, D.S. Ginger, D. MacKenzie, Scalable passivation strategies to improve efficiency of slot die-coated perovskite solar cells, *ACS Energy Lett.* 9 (2024) 1888–1894.
- [234] L.Y. Ee, S.Y.R. Chia, K. Xue, S.Y. Chin, C.A.H. Cho, X.Y. Tan, S.F.Y. Li, Hyperbranched nanocellulose enabling rapid boron removal from aqueous environment, *Chem. Eng. J.* 454 (2023) 140218.
- [235] K. Ye, S.Y. Chin, N.L. Xi, B. Sharma, Y. Lu, K. Xue, Characterizing the behavior of water interacting with a nano-pore material: a structural investigation in native environment using magnetic resonance approaches, *ChemPhysChem* 25 (2024) e202400053.
- [236] S.Y. Chin, L. Zhao, Y. Chen, Z. Zhai, X. Shi, K. Xue, Nanosecond molecular motion in pH1 α liquid-liquid phase separation captured by solid-state NMR, *J. Phys. Chem. Lett.* 16 (2025) 1150–1156.
- [237] S.Y. Chin, Y. Lu, W. Di, K. Ye, Z. Li, C. He, Y. Cao, C. Tang, K. Xue, Regulating polystyrene glass transition temperature by varying the hydration levels of aromatic ring/ Li^+ interaction, *Phys. Chem. Chem. Phys.* 25 (2023) 30223–30227.
- [238] W. Di, K. Xue, J. Cai, Z. Zhu, Z. Li, H. Fu, H. Lei, W. Hu, C. Tang, W. Wang, Y. Cao, Single-molecule force spectroscopy reveals Cation- π interactions in aqueous media are highly affected by cation dehydration, *Phys. Rev. Lett.* 130 (2023) 118101.
- [239] T.J.N. Hooper, B. Febriansyah, T. Krishnamoorthy, W.P.D. Wong, K. Xue, J. W. Ager, N. Mathews, Multinuclear solid-state NMR investigation of structurally diverse low-dimensional hybrid metal halide perovskites, *J. Mater. Chem. A* 12 (2024) 23461–23474.
- [240] S.Y. Chin, Y. Chen, L. Zhao, C.-P. Chng, J. Kuang, Z. Zhai, Y. Lu, C. Huang, C. Shi, X. Shi, K. Xue, Role of trifluoroacetic acid anions in pH1 α liquid-liquid phase separations, *J. Phys. Chem. B* 129 (2025) 7964–7971.
- [241] S. Mahalingam, A. Manapp, K.S. Lau, R. Rabeya, C.H. Chia, K. Xue, P. Chelvanathan, Recombination suppression in TiO_2 /boron-doped reduced graphene oxide-based dye-sensitized solar cells, *Renew. Sustain. Energy Rev.* 209 (2025) 115088.
- [242] X. Wang, C. Zhang, M. Sawczyk, J. Sun, Q. Yuan, F. Chen, T.C. Mendes, P. C. Howlett, C. Fu, Y. Wang, X. Tan, D.J. Searles, P. Král, C.J. Hawker, A. K. Whittaker, M. Forsyth, Ultra-stable all-solid-state sodium metal batteries enabled by perfluoropolyether-based electrolytes, *Nat. Mater.* 21 (2022) 1057–1065.
- [243] Y. García, L.A. O'Dell, Understanding the interfacial region in organic ionic plastic crystal composite electrolyte materials by solid-state NMR, *Curr. Opin. Colloid Interface Sci.* 61 (2022) 101632.
- [244] Y. Shang, A. Rawal, D. Kundu, High-voltage and highly reversible redox chemistry of in situ cross-linked polydiphenylamine as an aqueous zinc battery cathode, *Mater. Today Energy* 46 (2024) 101702.

- [245] Y. Chen, X. Yang, T. Wang, X. Tang, D. Li, S. Wang, Y. Lei, Y. Han, S. Chen, M. Armand, D. Aurbach, G. Wang, Fluoroether design enables high-voltage all-solid-state lithium metal batteries, *Adv. Mater.* (2025) 2506020.
- [246] Z. Zujovic, P.A. Kilmartin, J. Travas-Sejdic, The applications of solid-state NMR to conducting polymers. The special case on polyaniline, *Molecules* 25 (2020) 444.
- [247] X. Wu, Z. Liu, H. Guo, Y.-I. Hong, B. Xu, K. Zhang, Y. Nishiyama, W. Jiang, S. Horike, S. Kitagawa, G. Zhang, Host-guest assembly of H-Bonding networks in covalent organic frameworks for ultrafast and anhydrous proton transfer, *ACS Appl. Mater. Interfaces* 13 (2021) 37172–37178.
- [248] S. Suzuki, S. Kumagai, T. Nagashima, T. Yamazaki, T. Okitsu, A. Wada, A. Naito, K. Katayama, K. Inoue, H. Kandori, I. Kawamura, Characterization of retinal chromophore and protonated schiff base in Thermoplasmatales archaeon heliorhodopsin using solid-state NMR spectroscopy, *Biophys. Chem.* 296 (2023) 106991.
- [249] Y. Suzuki, T. Higashi, T. Yamamoto, H. Okamura, T.K. Sato, T. Asakura, Presence of β -Turn structure in recombinant spider silk dissolved in formic acid revealed with NMR, *Molecules* 27 (2022) 511.
- [250] K. Koner, S. Das, S. Mohata, N.T. Duong, Y. Nishiyama, S. Kandambeth, S. Karak, S.M. Reddy, R. Banerjee, Viscoelastic covalent organic nanotube fabric via macroscopic entanglement, *J. Am. Chem. Soc.* 144 (2022) 16052–16059.
- [251] S. Miyata, D. Aoki, Y. Matsushita, M. Takeuchi, K. Fukushima, Evaluation of guaiacyl lignin aromatic structures using $^{13}\text{C}_2$ administered Ginkgo biloba L. xylem by quantitative solid- and liquid-state ^{13}C NMR, *Holzforschung* 77 (2023) 230–239.
- [252] Y.-I. Hong, G.N.M. Reddy, Y. Nishiyama, Selective detection of active pharmaceutical ingredients in tablet formulations using solid-state NMR spectroscopy, *Solid State Nucl. Magn. Reson.* 106 (2020) 101651.
- [253] N. Hiraishi, T. Gondo, Y. Shimada, R. Hill, F. Hayashi, Crystallographic and physicochemical analysis of bovine and human teeth using X-ray diffraction and solid-state nuclear magnetic resonance, *J. Funct. Biomater.* 13 (2022) 254.
- [254] Y. Yanagisawa, M. Hamada, K. Hashi, H. Maeda, Review of recent developments in ultra-high field (UHF) NMR magnets in the Asia region, *Supercond. Sci. Technol.* 35 (2022) 044006.
- [255] M.J. Jaroszewicz, A.R. Altenhof, R.W. Schurko, L. Frydman, Sensitivity enhancement by progressive saturation of the proton reservoir: a solid-state NMR analogue of chemical exchange saturation transfer, *J. Am. Chem. Soc.* 143 (2021) 19778–19784.
- [256] A.R. Altenhof, S. Wi, R.W. Schurko, Broadband adiabatic inversion cross-polarization to integer-spin nuclei with application to deuterium NMR, *Magn. Reson. Chem.* 59 (2021) 1009–1023.
- [257] N.T. Duong, Z. Gan, Y. Nishiyama, Selective ^1H - ^{14}N Distadce Measurements by ^{14}N Over-tone Solid-otate NMRsolid-soscopy atsFast MAS, *Front. Mol. Biosci.* 8 (2021), 2021.
- [258] I. Hung, Z. Gan, G. Wu, Two- and three-dimensional ^{13}C - ^{17}O Heteronuclear Correlatioc NMR Spectroscopy for Studying Osganic ano BiologicalbSolids, *J. sPhys. Chem. Lett.* 12 (2021) 8897–8902.
- [259] I. Hung, Z. Gan, On the use of single-frequency versus double-frequency satellite-transition pulses for MQMAS, *J. Magn. Reson.* 328 (2021) 106994.
- [260] R. Bayzou, J. Trébosc, I. Hung, Z. Gan, O. Lafon, J.-P. Amoureux, Indirect NMR detection via proton of nuclei subject to large anisotropic interactions, such as ^{14}N , ^{195}Pt , and ^{35}Cl , using the T-HMQC sequence, *J. Chem. Phys.* 156 (2022).
- [261] R.W. Dorn, A.L. Paterson, I. Hung, P.L. Gor'kov, A.J. Thompson, A.D. Sadow, Z. Gan, A.J. Rossini, Dipolar heteronuclear correlation solid-state NMR experiments between half-integer quadrupolar nuclei: the case of ^{11}B - ^{17}O , *J. Phys. Chem. C* 126 (2022) 11652–11666.
- [262] S. Yoon, N. Schopp, D.G. Choi, H. Wakidi, K. Ding, H. Ade, H. Vezin, G.N. M. Reddy, T.-Q. Nguyen, Influences of metal electrodes on stability of non-fullerene acceptor-based organic photovoltaics, *Adv. Funct. Mater.* (2023) 2308618.
- [263] S.T. Holmes, J. Schönzart, A.B. Philips, J.J. Kimball, S. Termos, A.R. Altenhof, Y. Xu, C.A. O'Keefe, J. Autschbach, R.W. Schurko, Structure and bonding in rhodium coordination compounds: a ^{103}Rh solid-state NMR and relativistic DFT study, *Chem. Sci.* 15 (2024) 2181–2196.
- [264] S.T. Holmes, Y. Xu, A.B. Philips, S. Termos, J.J. Kimball, J. Autschbach, R. W. Schurko, ^{99}Ru solid-state nuclear magnetic resonance spectroscopy of organometallic compounds: linking nuclear magnetic resonance parameters with metal-ligand bonding, *J. Am. Chem. Soc.* 147 (2025) 20865–20877.
- [265] I. Hung, Z. Gan, High-resolution $^1\text{H}/^{17}\text{O}$ NMR correlation under fast magic-angle spinning: Revisiting cross-polarization for quadrupolar nuclei, *J. Phys. Chem. Lett.* 16 (2025) 7036–7041.
- [266] C.J. Pickard, F. Mauri, All-electron magnetic response with pseudopotentials: NMR chemical shifts, *Phys. Rev. B* 63 (2001) 245101.
- [267] E. Salager, G.M. Day, R.S. Stein, C.J. Pickard, B. Elena, L. Emsley, Powder crystallography by combined crystal structure prediction and high-resolution ^1H solid-state NMR spectroscopy, *J. Am. Chem. Soc.* 132 (2010) 2564–2566.
- [268] S.T. Holmes, C.S. Vojvodin, R.W. Schurko, Dispersion-corrected DFT methods for applications in nuclear magnetic resonance crystallography, *J. Phys. Chem. A* 124 (2020) 10312–10323.
- [269] A.H. Mazurek, Ł. Szeleszczuk, D.M. Pisklak, Periodic DFT calculations—review of applications in the pharmaceutical sciences, *Pharmaceutics* 12 (2020) 415.
- [270] D. Schäfer, K. Hankins, M. Allion, U. Krewer, F. Karcher, L. Derr, R. Schuster, J. Maibach, S. Mück, D. Kramer, R. Mönig, F. Jeschull, S. Daboss, T. Philipp, G. Neusser, J. Romer, K. Palanisamy, C. Kranz, F. Buchner, R.J. Behm, A. Ahmadian, C. Kübel, I. Mohammad, A. Samoson, R. Witter, B. Smarsly, M. Rohnke, Multiscale investigation of sodium-ion battery anodes: analytical techniques and applications, *Adv. Energy Mater.* 14 (2024) 2302830.
- [271] A.A. Peach, C.H. Fleischer, K. Levin, S.T. Holmes, J.E. Sanchez, R.W. Schurko, Quadrupolar NMR crystallography guided crystal structure prediction (QNMRX-CSP), *CrystEngComm* 26 (2024) 4782–4803.
- [272] J. Autschbach, Analysis of NMR spectroscopic parameters, in: G.A. Aucar (Ed.), *NMR Spectroscopic Parameters: Theories and Models, Computational Codes and Calculations*, Royal Society of Chemistry, 2025.
- [273] M. Cordova, M. Balodis, A. Hofstetter, F. Paruzzo, S.O. Nilsson Lill, E.S. E. Eriksson, P. Berruyer, B. Simões de Almeida, M.J. Quayle, S.T. Norberg, A. Svensk Ankarberg, S. Schantz, L. Emsley, Structure determination of an amorphous drug through large-scale NMR predictions, *Nat. Commun.* 12 (2021) 2964.
- [274] H. Park, A. Onwuli, A. Walsh, Exploration of crystal chemical space using text-guided generative artificial intelligence, *Nat. Commun.* 16 (2025) 4379.
- [275] J. Koppe, A.V. Yakimov, D. Gioffrè, M.-E. Usteri, T. Vosegaard, G. Pintacuda, A. Lesage, A.J. Pell, S. Mitchell, J. Pérez-Ramírez, C. Copéret, Coordination environments of Pt single-atom catalysts from NMR signatures, *Nature* 642 (2025) 613–619.
- [276] M.D. Mantle, A.J. Sederman, Dynamic MRI in chemical process and reaction engineering, *Prog. Nucl. Magn. Reson. Spectrosc.* 43 (2003) 3–60.
- [277] J.D. Cooper, N.P. Ramskill, A.J. Sederman, L.F. Gladden, A. Tsolakis, E.H. Stitt, A. P.E. York, Insights into automotive particulate filters using magnetic resonance imaging, *Johnson Matthey Technology Review* 64 (2020) 165–179.
- [278] D. Beriashvili, J. Zhou, Y. Liu, G.E. Folkers, M. Baldus, Cellular applications of DNP solid-state NMR – state of the art and a look to the future, *Chem. Eur J.* 30 (2024) e202400323.
- [279] B.J. Albert, C. Gao, E.L. Sesti, E.P. Saliba, N. Alaniva, F.J. Scott, S.T. Sigurdsson, A.B. Barnes, Dynamic nuclear polarization nuclear magnetic resonance in human cells using fluorescent polarizing agents, *Biochemistry* 57 (2018) 4741–4746.
- [280] G.N.M. Reddy, P. Selter, Y. Makita, S. Arai, M. Miyagawa, H. Nakano, H. Wang, J. A. Gerbec, F. Shimizu, B.F. Chmelka, Nanoscale surface compositions and structures of plasma-modified poly(ethylene terephthalate) thin films, *J. Phys. Chem. C* 125 (2021) 20658–20669.
- [281] A. Bertarello, P. Berruyer, M. Artelsmair, C.S. Elmore, S. Heydarkhan-Hagvall, M. Schade, E. Chiarparin, S. Schantz, L. Emsley, In-Cell quantification of drugs by magic-angle spinning dynamic nuclear polarization NMR, *J. Am. Chem. Soc.* 144 (2022) 6734–6741.
- [282] Z. Du, H.M. Luong, S. Sabury, A.L. Jones, Z. Zhu, P. Panoy, S. Chae, A. Yi, H. J. Kim, S. Xiao, V.V. Brus, G.N.M. Reddy, J.R. Reynolds, T.-Q. Nguyen, High-performance wearable organic photodetectors by molecular design and green solvent processing for pulse oximetry and photoplethysmography, *Adv. Mater.* 36 (2024) 2310478.
- [283] K.H. Mroue, Y. Nishiyama, M. Kumar Pandey, B. Gong, E. McNerny, D.H. Kohn, M.D. Morris, A. Ramamoorthy, Proton-detected solid-state NMR spectroscopy of bone with ultrafast magic angle spinning, *Sci. Rep.* 5 (2015) 11991.
- [284] R. Yadav, B. Patra, R. Rai, N. Sinha, C. Singh, Solid-state NMR spectroscopy for unraveling structure and dynamics in biomaterials, *Solid State Nucl. Magn. Reson.* 140 (2025) 102045.
- [285] R. Ji, Z. Zhang, M. Deconinck, Y.J. Hofstetter, J. Shi, F. Paulus, P. Raval, G.N. M. Reddy, Y. Vaynzof, Spontaneous Formation of 1D/3D Perovskite Heterojunctions for Efficient Inverted Perovskite Solar Cells, *Adv. Energy Mater.* 14 (2024) 2304126.
- [286] C.J. Dahlman, D.J. Kubicki, G.N.M. Reddy, Interfaces in metal halide perovskites probed by solid-state NMR spectroscopy, *J. Mater. Chem. A* 9 (2021) 19206–19244.
- [287] C. Greve, P. Ramming, M. Griesbach, N. Leupold, R. Moos, A. Köhler, E.M. Herzig, F. Panzer, H. Grüninger, To stop or to shuttle halides? The role of an ionic liquid in thermal halide mixing of hybrid perovskites, *ACS Energy Lett.* 8 (2023) 5041–5049.
- [288] S. Sidhik, I. Metcalf, W. Li, T. Kodalle, C.J. Dolan, M. Khalili, J. Hou, F. Mandani, A. Torma, H. Zhang, R. Garai, J. Persaud, A. Marciel, I.A. Muro Puente, G.N. M. Reddy, A. Balvanz, M.A. Alam, C. Katan, E. Tsai, D. Ginger, D.P. Fenning, M. G. Kanatzidis, C.M. Sutter-Fella, J. Even, A.D. Mohite, Two-dimensional perovskite templates for durable, efficient formamidinium perovskite solar cells, *Science* 384 (2024) 1227–1235.
- [289] A. Marchetti, J. Chen, Z. Pang, S. Li, D. Ling, F. Deng, X. Kong, Understanding surface and interfacial chemistry in functional nanomaterials via solid-state NMR, *Adv. Mater.* 29 (2017) 1605895.
- [290] A. Gabbani, E. Della Latta, A. Mohan, A. Scarperi, X. Li, M. Ruggeri, F. Martini, F. Biccari, M. Kociak, M. Geppi, S. Borsacchi, F. Pineider, Direct determination of carrier parameters in indium tin oxide nanocrystals, *ACS Nano* 18 (2024) 15139–15153.
- [291] L. Pan, Z. Liu, C. Welton, V.V. Klepov, J.A. Peters, M.C. De Siena, A. Benadia, I. Pandey, A. Miceli, D.Y. Chung, G.N.M. Reddy, B.W. Wessels, M.G. Kanatzidis, Ultrahigh-flux X-ray detection by a solution-grown perovskite CsPbBr_3 single-crystal semiconductor detector, *Adv. Mater.* 35 (2023) 2211840.
- [292] A. Azmy, S. Li, G.K. Angeli, C. Welton, P. Raval, M. Li, N. Zibouche, L. Wojtas, G. N.M. Reddy, P. Guo, P.N. Trikalitis, I. Spanopoulos, Porous and water stable 2D hybrid metal halide with broad light emission and selective H_2O vapor sorption, *Angew. Chem. Int. Ed.* 62 (2023) e202218429.
- [293] H.J. Kantrow, C. Welton, N. Thomas, V. Sarou-Kanian, T. Pawlak, N. Stingelin, G. N.M. Reddy, A close look at the local structure of functional polymers: the example of Poly(Vinylidene fluoride), *Adv. Funct. Mater.* 35 (2025) 2422354.
- [294] H.J. Jakobsen, H. Bildsøe, M. Bondegaard, B.B. Iversen, M. Brorson, F.H. Larsen, Z. Gan, I. Hung, Exciting opportunities for solid-state ^{95}Mo NMR studies of MoS_2 nanostructures in materials research from a low to an ultrahigh magnetic field (35.2 T), *J. Phys. Chem. C* 125 (2021) 7824–7838.

- [295] E. Bellan, F. Maleki, M. Jakoobi, P. Fau, K. Fajerwerger, D. Lagarde, A. Balocchi, P. Lecante, J. Trébosc, Y. Xu, Z. Gan, L. Pautrot-d'Alençon, T. Le Mercier, H. Nagashima, G. Pacchioni, O. Lafon, Y. Coppel, M.L. Kahn, Ultra-high-field ^{67}Zn and ^{35}P NMR studies coupled with DFT calculations reveal the structure of ZnS nanoplatelets prepared by an organometallic approach, *J. Phys. Chem. C* 127 (2023) 17809–17819.
- [296] Z. Xiong, H.P. Andaraarachchi, J.T. Held, R.W. Dorn, Y.-J. Jeong, A. Rossini, U. R. Kortshagen, Inductively coupled nonthermal plasma synthesis of size-controlled $\gamma\text{-Al}_2\text{O}_3$ nanocrystals, *Nanomaterials* 13 (2023) 1627.
- [297] R.E. Ortega, R.B. Smith, J.E. Kuszynski, A. Bayles, S.A. McGill, N.J. Halas, R. W. Schurko, G.F. Strouse, Observing metallic carriers in highly faceted plasmonic Cd_2SnO_4 inverse spinel nanocrystals, *Adv. Opt. Mater.* 12 (2024) 2400388.
- [298] S. Sen, I. Hung, J.M. Lovi, Z. Gan, Application of high-resolution ^{95}Mo solid state NMR spectroscopy in structural studies of complex alkali molybdate crystals and glasses, *J. Phys. Chem. C* 129 (2025) 6869–6881.
- [299] S. Sen, R.F. Lancelotti, I. Hung, Z. Gan, Characterization of the Pb coordination environment and its connectivity in lead silicate glasses: results from 2D ^{207}Pb NMR Spectroscopy, *J. Phys. Chem. B* 128 (2024) 2811–2820.
- [300] R.K. Swanson, J.F. Stebbins, T.M. Yeo, Y. Xu, I. Hung, Z. Gan, S.J. McCormack, S. Sen, Structure and fragility of normal and invert lanthanum borate glasses: results from ^{11}B and ^{17}O NMR spectroscopy and calorimetry, *J. Non-Cryst. Solids* 603 (2023) 122119.
- [301] S. Sen, J.F. Stebbins, S. Kroeker, I. Hung, Z. Gan, Evidence for mixed Mg coordination environments in silicate glasses: results from ^{25}Mg NMR Spectroscopy at 35.2 T, *J. Phys. Chem. B* 127 (2023) 10659–10666.
- [302] S. Sen, J.F. Stebbins, Y. Xu, I. Hung, Z. Gan, Germanate anomaly and its temperature dependence: an ultra-high field ^{17}O NMR spectroscopic study of sodium germanate glasses, *J. Non-Cryst. Solids* 18 (2023) 100175. X.
- [303] A. Singh, D. Dayton, D.M. Ladd, G. Reuveni, P. Paluch, L. Montagne, J. Mars, O. Yaffe, M. Toney, G.N.M. Reddy, D.B. Mitzi, Local structure in crystalline, glass and melt states of a hybrid metal halide perovskite, *J. Am. Chem. Soc.* 146 (2024) 25656–25668.
- [304] M.A. Hope, B.L.D. Rinkel, A.B. Gunnarsdóttir, K. Märker, S. Menkin, S. Paul, I. V. Sergeyev, C.P. Grey, Selective NMR observation of the SEI-Metal interface by dynamic nuclear polarisation from lithium metal, *Nat. Commun.* 11 (2020) 2224.
- [305] A. Harchol, G. Reuveni, V. Ri, B. Thomas, R. Carmieli, R.H. Herber, C. Kim, M. Leskes, Endogenous dynamic nuclear polarization for sensitivity enhancement in solid-state NMR of electrode materials, *J. Phys. Chem. C* 124 (2020) 7082–7090.
- [306] Y. Jin, N.-J.H. Kneusels, L.E. Marbella, E. Castillo-Martínez, P.C.M.M. Magusin, R. S. Weatherup, E. Jönsson, T. Liu, S. Paul, C.P. Grey, Understanding Fluoroethylene Carbonate and Vinylene Carbonate Based Electrolytes for Si Anodes in Lithium Ion Batteries with NMR Spectroscopy, *J. Am. Chem. Soc.* 140 (2018) 9854–9867.
- [307] W.M.J. Fransen, A.P.M. Kentgens, Solid-state NMR of hybrid halide perovskites, *Solid State Nucl. Magn. Reson.* 100 (2019) 36–44.
- [308] H. Grüniger, M. Bokdam, N. Leupold, P. Tinnemans, R. Moos, G.A. De Wijs, F. Panzer, A.P.M. Kentgens, Microscopic (Dis)Order and dynamics of cations in mixed FA/MA lead halide perovskites, *J. Phys. Chem. C* 125 (2021) 1742.
- [309] L. Piveteau, V. Morad, M.V. Kovalenko, Solid-state NMR and NQR spectroscopy of lead-halide perovskite materials, *J. Am. Chem. Soc.* 142 (2020) 19413–19437.
- [310] D. Sarkar, R.W. Hooper, A. Karmakar, A. Bhattacharya, A. Pominov, V. Terskikh, V.K. Michaelis, Metal halide perovskite and perovskite-like materials through the lens of ultra-wideline $^{35/37}\text{Cl}$ NMR Spectroscopy, *ACS Mater. Lett.* 4 (2022) 1255.
- [311] R.W. Hooper, D. Sarkar, V.K. Michaelis, Bulk and nanoscale semiconducting materials: structural advances using solid-state NMR spectroscopy, *Curr. Opin. Colloid Interface Sci.* 62 (2022) 101631.
- [312] P. Fu, M.A. Quintero, E.S. Vasileiadou, P. Raval, C. Welton, M. Kepenekian, G. Volonakis, J. Even, Y. Liu, C. Malliakas, Y. Yang, C. Laing, V.P. Dravid, G.N. M. Reddy, C. Li, E.H. Sargent, M.G. Kanatzidis, Chemical behavior and local structure of the Ruddlesden-Popper and Dion-Jacobson alloyed Pb/Sn bromide 2D perovskites, *J. Am. Chem. Soc.* 145 (2023) 15997–16014.
- [313] P. Fu, M.A. Quintero, C. Welton, X. Li, B. Cucco, M.C. De Siena, J. Even, G. Volonakis, M. Kepenekian, R. Liu, C.C. Laing, V. Klepov, Y. Liu, V.P. Dravid, G.N.M. Reddy, C. Li, M.G. Kanatzidis, Short aromatic diammonium ions modulate distortions in 2D lead bromide perovskites for tunable white-light emission, *Chem. Mater.* 34 (2022) 9685–9698.
- [314] S.R. Chaudhari, J.M. Griffin, K. Broch, A. Lesage, V. Lemaure, D. Dudenko, Y. Olivier, H. Sirringhaus, L. Emsley, C.P. Grey, Donor-acceptor stacking arrangements in bulk and thin-film high-mobility conjugated polymers characterized using molecular modelling and MAS and surface-enhanced solid-state NMR spectroscopy, *Chem. Sci.* 8 (2017) 3126–3136.
- [315] A.L. Dixon, H. Vezin, T.-Q. Nguyen, G.N.M. Reddy, Structural insights into lewis acid- and F4TCNQ-doped conjugated polymers by solid-state magnetic resonance spectroscopy, *Mater. Horiz.* 9 (2022) 981–990.
- [316] B.R. Luginbuhl, S.-J. Ko, N.A. Ran, H. Hu, S.M. Becwar, A. Karki, M. Seifrid, T. Okubo, M. Wang, H.W. Ade, B.F. Chmelka, G.C. Bazan, G.N.M. Reddy, T.-Q. Nguyen, Low voltage-loss organic solar cells light the way for efficient semitransparent photovoltaics, *Sol. RRL* 6 (2022) 2200135.
- [317] Z. Du, H.M. Luong, S. Sabury, P. Therdkatanyuphong, S. Chae, C. Welton, A. L. Jones, J. Zhang, Z. Peng, Z. Zhu, S. Nanayakkara, V. Coropceanu, D.G. Choi, S. Xiao, A. Yi, H.J. Kim, J.-L. Bredas, H. Ade, G.N.M. Reddy, S.R. Marder, J. R. Reynolds, T.-Q. Nguyen, Additive-free molecular acceptor organic solar cells processed from a biorenewable solvent approaching 15% efficiency, *Mater. Horiz.* 10 (2023) 5564–5576.
- [318] H.M. Luong, S. Chae, A. Yi, K. Ding, J. Huang, B.M. Kim, C. Welton, J. Chen, H. Wakidi, Z. Du, H.J. Kim, H. Ade, G.N.M. Reddy, T.-Q. Nguyen, Impact of thermal stress on device physics and morphology in organic photodetectors, *ACS Energy Lett.* (2023) 2130–2140.
- [319] S. Yoon, B. Reyes-Suárez, S.T. Pham, H. Vezin, Y.A. Tobon, M. Lee, S. Mugiraneza, B.M. Kim, M.Y.T. Oide, S. Yoo, S. Lee, S.H. Wang, S.M. Collins, C.M. Bates, Y. Park, B. Kim, G.N.M. Reddy, T.-Q. Nguyen, Molecular cross-linking enhances stability of non-fullerene acceptor organic photovoltaics, *ACS Energy Lett.* 10 (2025) 541–551.
- [320] B. Reyes-Suárez, N.T.Z. Du, Z.-z. Qu, T. Pawlak, R. Vasant, D. Vu, S. Yoon, S. Mugiraneza, Y. Lai, J. Fishburn, S. Chae, A. Yi, P. Raval, S. Mukherjee, Y. Qin, S. Xiao, H.J. Kim, H. Ade, T.-Q. Nguyen, G.N.M. Reddy, Solvent-induced polymorphism influences efficiency in organic solar cells, *ACS Energy Lett.* 10 (2025) 4872–4881.
- [321] P. Raval, M. Dhennin, H. Vezin, T. Pawlak, P. Roussel, T.-Q. Nguyen, G.N. M. Reddy, Understanding the p-doping of spiroOMeTAD by tris (pentafluorophenyl)borane, *Electrochim. Acta* 424 (2022) 140602.
- [322] Y. Shin, R. Prasad, N. Das, J.A. Taylor, H. Qin, W. Hu, Y.-Y. Hu, R. Fu, R. Zhang, H.-X. Zhou, T.A. Cross, Mycobacterium tuberculosis CrgA forms a dimeric structure with its transmembrane domain sandwiched between cytoplasmic and periplasmic β -Sheets, enabling multiple interactions with other diverse proteins, *J. Am. Chem. Soc.* 147 (2025) 11117–11131.
- [323] C.-H. Byeon, T. Kinney, H. Saricayir, K. Holst Hansen, F.J. Scott, S. Srinivasa, M. K. Wells, F. Mentink-Vigier, W. Kim, Ü. Akbey, Ultrasensitive characterization of native bacterial biofilms via dynamic nuclear polarization-enhanced solid-state NMR, *Angew. Chem. Int. Ed.* 64 (2025) e202418146.
- [324] N. Dwivedi, B. Patra, F. Mentink-Vigier, S. Wi, N. Sinha, Unveiling charge-pair salt-bridge interaction between GAGs and collagen protein in cartilage: atomic evidence from DNP-enhanced ssNMR at natural isotopic abundance, *J. Am. Chem. Soc.* 146 (2024) 23663–23668.
- [325] W. Qiang, M.K. Kengwerere, W. Zhao, F.J. Scott, X. Wutou-Hughes, T. Wang, F. Mentink-Vigier, Heterotypic interactions between the 40- and 42-Residue isoforms of β -Amyloid peptides on lipid bilayer surfaces, *ACS Chem. Neurosci.* 14 (2023) 4153–4162.
- [326] F. Liu, A.I. Greenwood, Y. Xiong, R.T. Miceli, R. Fu, K.W. Anderson, S. A. McCallum, M. Mihailescu, R. Gross, M.L. Cotten, Host defense peptide piscidin and yeast-derived glycolipid exhibit synergistic antimicrobial action through concerted interactions with membranes, *JACS Au* 3 (2023) 3345–3365.
- [327] O.M. Terrett, J.J. Lyczakowski, L. Yu, D. Iuga, W.T. Franks, S.P. Brown, R. Dupree, P. Dupree, Molecular architecture of softwood revealed by solid-state NMR, *Nat. Commun.* 10 (2019) 4978.
- [328] W. Zhao, E.C. Thomas, D. Debnath, F.J. Scott, F. Mentink-Vigier, J.R. White, R. L. Cook, T. Wang, Enriched molecular-level view of saline wetland soil carbon by sensitivity-enhanced solid-state NMR, *J. Am. Chem. Soc.* 147 (2025) 519–531.
- [329] N. Thomas, S. Moussaoui, B. Reyes-Suárez, O. Lafon, G.N.M. Reddy, Dual cross-linked cellulose based hydrogel films, *Mater. Adv.* 5 (2024) 9210–9219.
- [330] A. Javed, R. Jabbour, W. Zia, A. Eubal, Direct polarization transfer to remote nuclei: expanding the reach of cross-effect dynamic nuclear polarization, *Solid State Nucl. Magn. Reson.* 140 (2025) 102049.
- [331] M.R. Hansen, R. Graf, H.W. Spiess, Interplay of structure and dynamics in functional macromolecular and supramolecular systems as revealed by magnetic resonance spectroscopy, *Chem. Rev.* 116 (2016) 1272–1308.
- [332] J.C. Martínez-López, M. Santos Rodríguez, V. Oliver Cuenca, G. Silva Testa, E. van Eck, E.W. Zhao, Á.E. Lozano, C. Álvarez, J. Carretero-González, Dibenzodioxin-based polymers of intrinsic microporosity with enhanced transport properties for lithium ions in aqueous media, *Macromolecules* 57 (2024) 9442–9456.
- [333] A. Poulhazan, A. Baer, G. Daliaho, F. Mentink-Vigier, A.A. Arnold, D.C. Browne, L. Hering, S. Archer-Hartmann, L.E. Pepi, P. Azadi, S. Schmidt, G. Mayer, I. Marcotte, M.J. Harrington, Peculiar phosphonate modifications of velvet worm slime revealed by advanced nuclear magnetic resonance and mass spectrometry, *J. Am. Chem. Soc.* 145 (2023) 20749–20754.
- [334] T. Olla, R. Jabbour, A. Labiod, O. Boyron, S. Méry, B. Heinrich, T. Heiser, D. Jacquemin, P. Lévêque, A. Lesage, N. Leclerc, How halogenation impacts the polymer backbone conformation: learning from combination of solid-state MAS NMR and X-Ray scattering, *Adv. Funct. Mater.* 32 (2022) 2204929.
- [335] M. Seifrid, A. Karki, H. Wakidi, H. Vezin, C. Welton, G.C. Bazan, B.F. Chmelka, T.-Q. Nguyen, G.N.M. Reddy, Importance of short-range order in governing thin film morphology and electronic properties of polymeric organic semiconductors, *Chem. Mater.* 36 (2024) 1214–1227.
- [336] Z. Miao, S.A. Gonsales, C. Ehm, F. Mentink-Vigier, C.R. Bowers, B.S. Sumerlin, A. S. Veige, Cyclic polyacetylene, *Nat. Chem.* 13 (2021) 792–799.
- [337] Z. Miao, D. Konar, B.S. Sumerlin, A.S. Veige, Soluble polymer precursors via ring-expansion metathesis polymerization for the synthesis of cyclic polyacetylene, *Macromolecules* 54 (2021) 7840–7848.
- [338] P. Rassolov, A. Scigliani, H. Mohammadigoushki, Kinetics of shear banding flow formation in linear and branched wormlike micelles, *Soft Matter* 18 (2022) 6079–6093.
- [339] M. Wagner, R. Pigliapochi, V. Di Tullio, J. Catalano, N. Zumbulyadis, S. A. Centeno, X. Wang, K. Chen, I. Hung, Z. Gan, M.R. Dworzak, G.P.A. Yap, C. Dybowski, Multi-technique structural analysis of zinc carboxylates (soaps), *Dalton Trans.* 52 (2023) 6152–6165.
- [340] X. Zhang, S.F. Marxsen, P. Ortmann, S. Mecking, R.G. Alamo, Crystallization of long-spaced precision polyacetalates II: effect of polymorphism on isothermal crystallization kinetics, *Macromolecules* 53 (2020) 7899–7913.

- [341] P. Raval, N. Thomas, L. Hamdouna, L. Delevoye, O. Lafon, G.N.M. Reddy, Boron adsorption kinetics of microcrystalline cellulose and polymer resin, *Langmuir* 39 (2023) 5384–5395.
- [342] A.L. Jones, C.H.Y. Ho, S.A. Schneider, J. Zhang, Y. Pei, J. Wang, X. Zhan, S. R. Marder, M.F. Toney, F. So, G.N.M. Reddy, J.R. Reynolds, Insights into the local bulk-heterojunction packing interactions and donor–acceptor energy level offsets in scalable photovoltaic polymers, *Chem. Mater.* 34 (2022) 6853–6867.
- [343] B.R. Luginbuhl, P. Raval, T. Pawlak, Z. Du, T. Wang, G. Kugan, N. Schopp, S. Chae, S. Yoon, A. Yi, H. Jung Kim, V. Coropceanu, J.-L. Brédas, T.-Q. Nguyen, G.N.M. Reddy, Resolving atomic-scale interactions in nonfullerene acceptor organic solar cells with solid-state NMR spectroscopy, crystallographic modelling, and molecular dynamics simulations, *Adv. Mater.* 34 (2022) 2105943.
- [344] P. Mondelli, P. Kaienburg, F. Silvestri, R. Scatena, C. Welton, M. Grandjean, V. Lemaire, E. Solano, M. Nyman, P.N. Horton, S.J. Coles, E. Barrena, M. Riede, P. Radaelli, D. Beljonne, G.N.M. Reddy, G. Morse, Understanding the role of non-fullerene acceptor crystallinity in the charge transport properties and performance of organic solar cells, *J. Mater. Chem. A* 11 (2023) 16263–16278.
- [345] J. Wang, X. You, C. Xiao, X. Zhang, S. Cai, W. Jiang, S. Guo, S. Cao, Z. Chen, Small-sized Pt nanoparticles supported on hybrid structures of MoS₂ nanoflowers/graphene nanosheets: highly active composite catalyst toward efficient ethanol oxidation reaction studied by in situ electrochemical NMR spectroscopy, *Appl. Catal. B Environ.* 259 (2019) 118060.
- [346] W. Gao, Q. Qi, Q. Wang, W. Wang, S. Li, I. Hung, Z. Gan, J. Xu, F. Deng, Dual active sites on Molybdenum/ZSM-5 catalyst for methane dehydroaromatization: insights from solid-state NMR spectroscopy, *Angew. Chem. Int. Ed.* 60 (2021) 10709–10715.
- [347] R.W. Dorn, M.C. Cendejas, K. Chen, I. Hung, N.R. Altvater, W.P. McDermott, Z. Gan, I. Hermans, A.J. Rossini, Structure determination of boron-based oxidative dehydrogenation heterogeneous catalysts with ultrahigh field 35.2 T ¹¹B Solid-State NMR spectroscopy, *ACS Catal.* 10 (2020) 13852–13866.
- [348] R. Wang, Z. Zhao, P. Gao, K. Chen, Z. Gan, Q. Fu, G. Hou, Impact of adsorption configurations on alcohol dehydration over alumina catalysts, *J. Phys. Chem. C* 126 (2022) 10073–10080.
- [349] M. Zheng, Y. Chu, Q. Wang, Y. Wang, J. Xu, F. Deng, Advanced solid-state NMR spectroscopy and its applications in zeolite chemistry, *Prog. Nucl. Magn. Reson. Spectrosc.* 140–141 (2024) 1–41.
- [350] T. Kobayashi, F.A. Perras, I.I. Slowing, A.D. Sadow, M. Pruski, Dynamic nuclear polarization solid-state NMR in heterogeneous catalysis research, *ACS Catal.* 5 (2015) 7055–7062.
- [351] J.L. Belmonte-Vázquez, D.P. Karothu, C.H. Fleischer, D. Galicia-Badillo, M. Maldonado-Domínguez, R.W. Schurko, L. Li, P. Naumov, B. Rodríguez-Molina, Low-temperature dynamics at Nano- and macroscales: organic crystal that exhibits low-temperature molecular motion and the thermosalt effect, *Chem. Mater.* 37 (2025) 3373–3383.
- [352] B.H. Wilson, L.M. Abdulla, R.W. Schurko, S.J. Loeb, Translational dynamics of a non-degenerate molecular shuttle imbedded in a zirconium metal–organic framework, *Chem. Sci.* 12 (2021) 3944–3951.
- [353] A. Gonzalez-Nelson, S. Mula, M. Siménas, S. Balčiūnas, A.R. Altenhof, C. S. Vojvodin, S. Canossa, J.R. Banys, R.W. Schurko, F.-X. Coudert, M.A. van der Veen, Emergence of coupled rotor dynamics in metal–organic frameworks via tuned steric interactions, *J. Am. Chem. Soc.* 143 (2021) 12053–12062.
- [354] R. Jabbour, C.W. Ashling, T.C. Robinson, A.H. Khan, D. Wisser, P. Berruyer, A. C. Ghosh, A. Ranscht, D.A. Keen, E. Brunner, J. Canivet, T.D. Bennett, C. Mellot-Draznieks, A. Lesage, F.M. Wisser, Unravelling the molecular structure and confining environment of an organometallic catalyst heterogenized within amorphous porous polymers, *Angew. Chem. Int. Ed.* 62 (2023) e202310878.
- [355] J. Perego, A. Daolio, C.X. Bezuidehout, S. Piva, G. Prando, B. Costarella, P. Carretta, L. Marchiò, D. Kubicki, P. Sozzani, S. Bracco, A. Comotti, Solid state machinery of multiple dynamic elements in a metal–organic framework, *Angew. Chem. Int. Ed.* 63 (2024) e202317094.
- [356] S. Jekhmene, M.G.N. Derks, S. Maity, C.J. Slingerland, K.H.M.E. Tehrani, J. Medeiros-Silva, V. Charitou, D. Ammerlaan, C. Fetz, N.A. Consoli, R.V. K. Cochrane, E.J. Matheson, M. van der Weide, B.O.W. Elenbaas, F. Lavore, R. Cox, J.H. Lorent, M. Baldus, M. Künzler, M. Lelli, S.A. Cochrane, N.I. Martin, W.H. Roos, E. Breukink, M. Weingarth, Host defence peptide plectasin targets bacterial cell wall precursor lipid II by a calcium-sensitive supramolecular mechanism, *Nat. Microbiol.* 9 (2024) 1778–1791.
- [357] S.T. Holmes, J.M. Hook, R.W. Schurko, Nutraceuticals in bulk and dosage forms: analysis by ³⁵Cl and ¹⁴N Solid-State NMR and solid-sculations, *Mol. Pharm.* 19 (2022) 440–455.
- [358] P. Hodgkinson, NMR crystallography of molecular organics, *Prog. Nucl. Magn. Reson. Spectrosc.* 118–119 (2020) 10–53.
- [359] S.T. Holmes, C.S. Vojvodin, N. Veinberg, E.M. Iacobelli, D.A. Hirsh, R.W. Schurko, Hydrates of active pharmaceutical ingredients: a ³⁵Cl and ²H solid-state NMR and DFT study, *Solid State Nucl. Magn. Reson.* 122 (2022) 101837.
- [360] L.M. Abdulla, A.A. Peach, S.T. Holmes, Z.T. Dowdell, L.K. Watanabe, E. M. Iacobelli, D.A. Hirsh, J.M. Rawson, R.W. Schurko, Synthesis and characterization of xylazine hydrochloride polymorphs, hydrates, and cocrystals: a ³⁵Cl Solid-State NMR and DFT Study, *Cryst. Growth Des.* 23 (2023) 3412–3426.
- [361] I. Hung, Z. Gan, Pushing the limit of MQMAS for low-γ quadrupolar nuclei in pharmaceutical hydrochlorides, *J. Magn. Reson.* 350 (2023) 107423.
- [362] A.A. Peach, S.T. Holmes, L.R. MacGillivray, R.W. Schurko, The formation and stability of fluoxetine HCl cocrystals investigated by multicomponent milling, *CrystEngComm* 25 (2023) 213–224.
- [363] A.J. Stirk, S.T. Holmes, F.E.S. Souza, I. Hung, Z. Gan, J.F. Britten, A.W. Rey, R. W. Schurko, An unusual ionic cocrystal of ponatinib hydrochloride: characterization by single-crystal X-ray diffraction and ultra-high field NMR spectroscopy, *CrystEngComm* 26 (2024) 1219–1233.
- [364] G. Wu, Y. Dai, I. Hung, Z. Gan, V. Tersikh, ¹H/¹⁷O chemical shift waves in carboxyl-bridged hydrogen bond networks in organic solids, *J. Phys. Chem. A* 128 (2024) 4288–4296.
- [365] Z.T. Dowdell, A.A. Peach, J.P. Purdie, S.T. Holmes, L.K. Watanabe, J.M. Rawson, R.W. Schurko, Mechanochemical synthesis and ³⁵Cl NMR crystallography of ionic cocrystals of phenothiazine drugs, *CrystEngComm* 27 (2025) 3752–3768.
- [366] C.H. Fleischer, S.T. Holmes, K. Levin, S.L. Veinberg, R.W. Schurko, Characterization of ephedrine HCl and pseudoephedrine HCl using quadrupolar NMR crystallography guided crystal structure prediction, *Faraday Discuss* 255 (2025) 88–118.
- [367] N. Thomas, C. Welton, T. Pawlak, P. Raval, J. Trébosc, S.K. Jain, G.N.M. Reddy, Deuteron-proton isotope correlation spectroscopy at high magnetic fields, *Solid State Nucl. Magn. Reson.* 136 (2025) 101993.
- [368] S.T. Holmes, R.A. Wiscons, K. Parks, S. Nickel, H.S. Ankeny, A.M. Viggiano, D. Bedillion, D. Shoup, R.J. Iulucci, Q. Wang, R.W. Schurko, R. Quiñones, A novel solid form of erlotinib: synthesis by heterogeneous complexation and characterization by NMR crystallography, *Cryst. Growth Des.* 25 (2025) 3219–3228.
- [369] D.A. Hirsh, A.V. Wijesekara, S.L. Carnahan, I. Hung, J.W. Lubach, K. Nagapudi, A. J. Rossini, Rapid characterization of formulated pharmaceuticals using fast MAS ¹H Solid-State NMR Spectroscopy, *Mol. Pharm.* 16 (2019) 3121–3132.
- [370] M. Li, W. Xu, Y. Su, Solid-state NMR spectroscopy in pharmaceutical sciences, *TrAC, Trends Anal. Chem.* 135 (2021) 116152.
- [371] C. Welton, P. Raval, J. Trébosc, G.N.M. Reddy, Chemical exchange of labile protons by deuterium enables selective detection of pharmaceuticals in solid formulations, *Chem. Commun.* 58 (2022) 11551–11554.
- [372] D. Torodii, M. Cordova, J.B. Holmes, P. Moutzouri, T. Casalini, S.O. Nilsson Lill, A.C. Pinon, C.S. Knee, A. Svensk Ankarberg, O.D. Putra, S. Schantz, L. Emsley, Three-Dimensional atomic-level structure of an amorphous glucagon-like Peptide-1 receptor agonist, *J. Am. Chem. Soc.* 147 (2025) 17077–17087.
- [373] C.H. Fleischer, S.T. Holmes, X. Lin, R.W. Schurko, De novo crystal structure determination of L-alaninamide HCl by quadrupolar NMR crystallography guided crystal structure prediction (QNMR-CSP), *Solid State Nucl. Magn. Reson.* 140 (2025) 102034.
- [374] S. Viel, Z. Fabio, C. Stefano, Enhanced diffusion-edited NMR spectroscopy of mixtures using chromatographic stationary phases, *Proc. Natl. Acad. Sci. USA* 100 (2003) 9696–9698.
- [375] G.N.M. Reddy, R. Ballesteros-Garrido, J. Lacour, S. Caldarelli, Determination of labile chiral supramolecular ion pairs by chromatographic NMR spectroscopy, *Angew. Chem. Int. Ed.* 52 (2013) 3255–3258.
- [376] M.B. Schmithorst, S. Prasad, A. Moini, B.F. Chmelka, Direct detection of paired aluminum heteroatoms in chabazite zeolite catalysts and their significance for methanol dehydration reactivity, *J. Am. Chem. Soc.* 145 (2023) 18215–18220.
- [377] C. Gioiosa, K. Sharma, T.C. Robinson, D. Gajan, S. Jannin, S. Laage, A. Lesage, Elucidating the substitution pattern of a hydroxyethyl cellulose thickening agent by DNP-enhanced solid-state NMR, *Anal. Chem.* 97 (2025) 16161–16167.
- [378] J. Viger-Gravel, F.M. Paruzzo, C. Cazaux, R. Jabbour, A. Leleu, F. Canini, P. Florian, F. Ronzon, D. Gajan, A. Lesage, Atomic-Scale description of interfaces between antigen and aluminum-based adjuvants used in vaccines by dynamic nuclear polarization (DNP) enhanced NMR spectroscopy, *Chem. Eur. J.* 26 (2020) 8976–8982.
- [379] G.N.M. Reddy, J.A. Gerbec, F. Shimizu, B.F. Chmelka, Nanoscale surface compositions and structures influence boron adsorption properties of anion exchange resins, *Langmuir* 35 (2019) 15661–15673.
- [380] E. Keijji, JEOL, NMR and ESR: a 65 year evolution, *J. Magn. Reson.* 306 (2019) 17–26.
- [381] R.G. Griffin, T.M. Swager, R.J. Temkin, High frequency dynamic nuclear polarization: new directions for the 21st century, *J. Magn. Reson.* 306 (2019) 128–133.
- [382] H. Maeda, Y. Yanagisawa, Future prospects for NMR magnets: a perspective, *J. Magn. Reson.* 306 (2019) 80–85.
- [383] E. Kryukov, A.J. Perez Linde, S. Raghunathan, S. Burgess, P. Jonsen, J. Good, On the magnetic field stability of cryogen-free magnets for magnetic resonance applications, *Solid State Nucl. Magn. Reson.* 105 (2020) 101639.
- [384] K. Xu, O. Pecher, M. Braun, J. Schmedt auf der Günne, Stable magic angle spinning with low-cost 3D-Printed parts, *J. Magn. Reson.* 333 (2021) 107096.
- [385] E. Sić, D. Fredericks, O. Pecher, S. Wegner, H. Breitzke, V. Singh, G. Buntkowsky, T. Gutmann, Towards routine ⁷Li In Situ solid-state NMR Studies of Electrochemical Processes in Li/LiPF₆/LFP Cells, *Appl. Magn. Reson.* 55 (2024) 575–583.
- [386] T.J.N. Hooper, R. de Oliveira-Silva, D. Sakellariou, High-resolution in situ photo-irradiation MAS NMR: application to the UV-polymerization of n-butyl acrylate, *J. Mater. Chem. A* 13 (2025) 933–939.

- [387] N. Thomas, J. Dhainaut, A. Moissette, O. Lafon, G.N.M. Reddy, Resolving transformative reactions in zeolites under catalytic hydrothermal conditions, *J. Phys. Chem. C* 128 (2024) 10428–10439.
- [388] S. Axon, D. James, The UN sustainable development goals: how can sustainable chemistry contribute? A view from the chemical industry, *Curr. Opin. Green Sustainable Chem.* 13 (2018) 140–145.
- [389] A.M. Noce, How chemistry can help meet the UN's sustainable development goals, *C&EN Global Enterprise* 96 (2018), 43–43.
- [390] V.K. Shukla, G.T. Heller, D.F. Hansen, Biomolecular NMR spectroscopy in the era of artificial intelligence, *Structure* 31 (2023) 1360–1374.
- [391] S. Kern, S. Liehr, L. Wander, M. Bornemann-Pfeiffer, S. Müller, M. Maiwald, S. Kowarik, Artificial neural networks for quantitative online NMR spectroscopy, *Anal. Bioanal. Chem.* 412 (2020) 4447–4459.
- [392] D. Guo, S. Li, J. Liu, Z. Tu, T. Qiu, J. Xu, L. Feng, D. Lin, Q. Hong, M. Lin, Y. Lin, X. Qu, CloudBrain-NMR: an intelligent cloud-computing platform for NMR spectroscopy processing, reconstruction, and analysis, *IEEE Trans. Instrum. Meas.* 73 (2024) 1–11.