



Investigation of pesticides on honey bee carbonic anhydrase inhibition

Ercan Soydan, Ahmet Can Olcay, Gürkan Bilir, Ömer Taş, Murat Şentürk, Deniz Ekinci & Claudiu T. Supuran

To cite this article: Ercan Soydan, Ahmet Can Olcay, Gürkan Bilir, Ömer Taş, Murat Şentürk, Deniz Ekinci & Claudiu T. Supuran (2020) Investigation of pesticides on honey bee carbonic anhydrase inhibition, Journal of Enzyme Inhibition and Medicinal Chemistry, 35:1, 1923-1927, DOI: [10.1080/14756366.2020.1835885](https://doi.org/10.1080/14756366.2020.1835885)

To link to this article: <https://doi.org/10.1080/14756366.2020.1835885>



© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 20 Oct 2020.



Submit your article to this journal [↗](#)



Article views: 1388



View related articles [↗](#)



View Crossmark data [↗](#)




Citing articles: 9 View citing articles [↗](#)

RESEARCH PAPER



Investigation of pesticides on honey bee carbonic anhydrase inhibition

Ercan Soydan^a, Ahmet Can Olcay^a, Gürkan Bilir^a, Ömer Taş^a, Murat Şentürk^b, Deniz Ekinci^a and Claudiu T. Supuran^c 

^aFaculty of Agriculture, Department of Agricultural Biotechnology, Ondokuz Mayıs University, Samsun, Turkey; ^bPharmacy Faculty, Department of Biochemistry, Agri Ibrahim Cecen University, Agri, Turkey; ^cNeurofarba Department, University of Florence, Firenze, Italy

ABSTRACT

Carbonic anhydrase (CA, EC 4.2.1.1) plays crucial physiological roles in many different organisms, such as in pH regulation, ion transport, and metabolic processes. CA was isolated from the European bee *Apis mellifera* (AmCA) spermatheca and inhibitory effects of pesticides belonging to various classes, such as carbamates, thiophosphates, and pyrethroids, were investigated herein. The inhibitory effects of methomyl, oxamyl, deltamethrin, cypermethrin, dichlorodiphenyltrichloroethane (DDT) and diazinon on AmCA were analysed. These pesticides showed effective *in vitro* inhibition of the enzyme, at sub-micromolar levels. The IC₅₀ values for these pesticides ranged between of 0.0023 and 0.0385 μM. The CA inhibition mechanism with these compounds is unknown at the moment, but most of them contain ester functionalities which may be hydrolysed by the enzyme with the formation of intermediates that can either react with amino acid residues or bid to the zinc ion with the active site.

ARTICLE HISTORY

Received 15 September 2020
Revised 28 September 2020
Accepted 5 October 2020

KEYWORDS

Carbonic anhydrase; bee; pesticides; inhibition; pyrethroids

1. Introduction

Pesticides are chemical compounds that are used against various pests as a biological control agent. Some pesticides are persistent organic contaminants in the soil and environment. They also constitute one of the most significant causes of pollution worldwide¹. The rapid growth of agriculture and animal processing has caused bees to be exposed to such contaminants with which they had never previously come into contact. Growing food demand has forced farmers to use more mineral fertilisers and pesticides for producing higher yields². In recent decades the increasing concern about the effect of pesticides on pollinators has been expressed in the scientific literature³. For this reason, some new data were collected from laboratory and semi-field studies on the toxic effects of pesticides on bees, especially bumble bees^{3,4}. A variety of articles have highlighted the significance of bees as natural pollinators not only for our crops but also for wildflowers and woodland plants, in temperate and tropical habitats^{5–7}. For example, it has been reported that about 60 crop plant species may not bear growing fruit without bees as impollinators⁸; with devastating economic implications.

Carbonic anhydrases (Cas, EC 4.2.1.1) are found in almost all living organisms and control pH and CO₂/bicarbonate levels⁹. Many different CA isoenzymes have been identified in higher vertebrates, although these enzymes are less investigated in other species, such as the arthropods, including insects^{10–12}. The physiological role of CA isozymes is to promote CO₂ to HCO₃⁻ interconversion, thus, playing vital functions in various biochemical/physiological processes, including physiological pH regulation, gas balancing, calcification, photosynthesis, metabolism, etc^{9–12}. Additionally, in vertebrates, CAs play a significant role in the eye, kidneys, central nervous system (CNS), inner ear, and many other

organs, in terms of ion transfer, pH regulation, and metabolism^{13–15}.

Pesticides and fungicides interfering with rainwater, irrigation water, or groundwater, plants, can inhibit particular enzymes^{13–15}. In our group's previous research, we investigated the effects of several widely used pesticides, such as tebuconazole, propoxur, carbaryl, carbofuran, simazine, and atrazine on the recently discovered bee CA, termed AmCA¹⁵. However, little is known about the effects of other chemical agents on this enzyme. Therefore, in this study, we purified honey bee spermatheca AmCA enzyme and analysed its interactions with pesticides and fungicides belonging to other classes.

2. Materials and methods

2.1. Chemicals

All chemicals for the affinity system were provided by Sigma-Aldrich (St. Louis, MO). Other reagents were obtained from Merck (Darmstadt, Germany).

2.2. Homogenate

Bee spermatheca samples were washed three times with 50 mM Tris/Sulphate (pH 7.8). Spermatheca samples taken from 200 bees were combined and homogenised with liquid nitrogen, then placed in the same buffer and centrifuged at 4 °C, 15,000g for 30 min. The precipitate portion was discarded and the supernatant portion was separated and used in subsequent studies.

CONTACT Ercan Soydan  esoydan@omu.edu.tr  Department of Agricultural Biotechnology, Ondokuz Mayıs University, Samsun 55139, Turkey; Claudiu T. Supuran  claudiu.supuran@unifi.it  Neurofarba Department, University of Florence, Firenze 50121, Italy

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Table 1. AmCA inhibition data from our previous study¹⁵.

Inhibitor	IC ₅₀ (μM)
Tebuconazole	0.0030
Carbaryl	0.0031
Carbonfuran	0.0087
Atrazine	0.0165
Simazin	0.0273
Propoxur	0.0321

2.3. Purification of the enzyme

The enzyme was purified using Sepharose-4B-sulfanilamide affinity gel prepared by our group¹⁵. Aniline was used as a spacer arm in the chromatography column. The column was then equilibrated with 25 mM Tris-HCl/0.1M Na₂SO₄ (pH 8.7) and the gel was washed with 25 mM Tris-HCl/22 mM Na₂SO₄ (pH 8.7). Finally, elution was performed with 1 M NaCl/25 mM Na₂HPO₄ (pH 6.3). The temperature was kept at 4 °C during all experiments.

2.4. In vitro inhibition experiments for AmCA enzyme

The effects of methomyl, oxamyl, deltamethrin, dichlorodiphenyltrichloroethane (DTT), cypermethrin and diazinon on AmCA activities were assayed colorimetrically using the CO₂ hydrase assay¹⁶, in triplicate at each concentration of inhibitor (ranging from 1 nM to 10 mM). Control enzyme activity was taken as 100%. An activity % versus inhibitor graph was drawn for all pesticides (Microsoft Office 2000, Excel). Pesticide concentrations that caused 50% inhibition (IC₅₀) of the enzyme activity were thus obtained graphically using a regression software. The enzyme concentration in the assay system was 8.9 nM.

3. Results and discussion

The collapse of honeybee (*Apis mellifera*) colonies poses an important problem in many developed countries^{17–20}. In fact, bee colonies are economically important for honey and wax products. The collapse of hive clones is generally related to two phenomena: (i) parasites, such as virus²¹, nosema infections²², mites²³, and hive insects²⁴, which attack the colony; and (ii) pesticides use, which negatively impacts the insects due to their toxicity²⁵. Furthermore, such low pesticide levels can also make bees an easier target to biological infections^{20–28}.

The majority of the problems in beehives faced by beekeepers are due to biological factors²⁷. However, these factors are improbable to be the most important reason for the recent decrease of bumblebees in North America and Europe or the disappearing of several wild beespecies^{22,28}. Agrochemicals, including pesticides, are probably the most significant factors that provoke honeybee and wild bee colonies extinction. Much research has been conducted ultimately in order to understand this problem in North America²⁹, France³⁰, Spain³¹ and India³². Such research focussed on determining the amount and prevalence of pesticide residues in honey, pollen, wax, and other different beehive matrices (e.g. combs)^{29–31}. A dataset was thus established for explaining the effects of pesticide residue both on honeybees and, potentially wild bees^{33,34}.

Recently, we isolated and characterised the α-CA enzyme from *Apis mellifera* spermatheca, AmCA¹⁵. In that study, we were able to purify the enzyme in a single step using Sepharose 4B tyrosine-sulfanilamide affinity chromatography¹⁵. Furthermore, pesticides such as tebuconazole, carbaryl, carbonfuran, atrazine, simazin, and propoxur were tested on the amCA enzyme and IC₅₀ values were

determined as 0.0030, 0.0031, 0.0087, 0.0165, 0.0273, and 0.0321 μM, respectively (Table 1)¹⁵. The efficiency order of pesticides AmCA inhibition was: tebuconazole > carbaryl > carbofuran > atrazine > simazine > propoxur.

Here we report the inhibitory effects of 6 pesticides belonging to various classes (Figure 1) on AmCA. Indeed, carbamates, thiophosphates, and pyrethroids were considered as potential CA inhibitors in the present study (Figure 1).

Besides the enzyme purification already reported in the previous work¹⁵, the inhibitory effects of six different pesticides on honeybee spermatheca CA and their IC₅₀ parameters were investigated here. The activity of AmCA was inhibited by pesticides shown in Figure 1 at low micromolar concentrations. Indeed, the IC₅₀ values were determined to be: 0.0023 ± 0.0001, 0.0025 ± 0.0001, 0.0028 ± 0.0001, 0.0034 ± 0.0001, 0.0078 ± 0.0003, and 0.0385 ± 0.0012 μM for methomyl, oxamyl, deltamethrin, dichlorodiphenyltrichloroethane (DTT), cypermethrin, and diazinon, respectively (Table 2).

It should be mentioned that a detailed study on pesticide residues in bees was conducted by Sanchez-Bayo and Goka²⁰. In this study, topical LD₅₀ and oral LD₅₀ values of pesticides against honey bees and bumblebees were reported²⁰. The topical lethal dose (LD₅₀) rates found in this study²⁰ were as follows: deltamethrin 0.02 μg/bee, cypermethrin 0.03 μg/bee, methomyl 0.49 μg/bee, and diazinon 0.38 μg/bee. In the same study, the oral dose rates were as follows: cypermethrin 0.06 μg/bee, methomyl 3.38 μg/bee, diazinon 0.21 μg/bee, and DDT 5.08 μg/bee²⁰. Compared to this literature results²⁰, the results obtained in the present study show a low level of IC₅₀ for the inhibition of AmCA with pesticides presented in Figure 1 and widely used in agriculture (Table 2). Thus, our results may prove that pesticides as those presented in Figure 1 may induce a strong inhibition on honey bee spermatheca CA enzyme. It is not known at the moment whether this inhibition may lead to physiological consequences but studies are ongoing in the field in our laboratories.

There has been a great interest in recent studies focussed on the effects of several classes of chemicals on CA enzymes^{35–37}. CA is a classical metalloenzyme, whose isozymes have significant roles in many tissues, that has been characterized and purified from several organisms, including animals^{35–37}. A huge number of pollutants, such as metals, acids, bases, and other toxic compounds³⁸ are being mixed in water sources and also in the atmosphere, which increasingly damages our environment. The potent AmCA inhibitory effects of compounds shown in Figure 1 may represent a potential explanation of why bees (domestic and wild ones) are under pressure worldwide with an increasing level of extinction of many diverse such insect species.

This also brings us to the possible mechanism of action of these pesticides investigated here. The carbamates, thiophosphates, and pyrethroids investigated here (Figure 1) possess ester bonds which can be hydrolysed by the esterase activity of the α-CAs. In fact, it has been thoroughly documented that these enzymes are esterases/thioesterases/selenoesterases with carboxylic, phosphoric, thiocarboxylic, and even seleno esters^{39–42}. Only DDT does not have this functionality, but this compound was reported by Bitman et al. to act as a CAI in the 70s⁴³.

The two carbamates from Figure 1, methomyl, and oxamyl can be substrates of CAs which may hydrolyse their ester/thioester bonds with the formation of small molecules which can bind to the metal centre (acetate, methyl-thiol, or Me₂N-COOH in the case of the second carbamate). The pyrethroids may also be hydrolysed at their ester functionality, with the generation of carboxylic acids and alcohols which were shown to act as CAIs^{12,13,36}.




Figure 1. Chemical structures of the pesticides tested as inhibitors of AmCA in this study.

Table 2. AmCA inhibition data for new pesticides shown in Figure 1.

Inhibitor	IC ₅₀ (μM)
Methomyl	0.0023 ± 0.0001
Oxamyl	0.0025 ± 0.0001
Deltamethrin	0.0028 ± 0.0001
DDT	0.0034 ± 0.0001
Cypermethrin	0.0078 ± 0.0003
Diazinon	0.0385 ± 0.0012

Diazinon has thiophosphate functionalities which were shown to be hydrolysed by the esterase activity of CAs in previous work from this laboratory, leading to suicide inhibitors of the enzyme³⁹. However, these hypotheses should be verified by X-ray crystallography, one of the most powerful techniques useful to assess CA inhibition mechanisms, especially the innovative ones. This technique also has its weak points, especially when used in an inattentive manner. The best example is the report by Liljas' group that cyanide and cyanate do not coordinate the metal ion from CA active site⁴⁴. Subsequent work from other laboratories showed those data to be false, as both cyanate and cyanide were observed coordinated to the metal ion, as most other anion inhibitors investigated to date^{45,46}. Furthermore, cyanate was also shown to be a suicide substrate that can be hydrolysed by the CA activity with the formation of carbamate⁴⁵. However, bitter and dubious comments from the above-mentioned crystallography group continued even 30 years later⁴⁷.

4. Conclusions

AmCA was purified from *Apis mellifera* spermatheca by affinity chromatography. Pesticides such as methomyl, oxamyl, deltamethrin, and DDT showed inhibition effects, comparable to those of cypermethrin and diazinon. The IC₅₀ values were determined as 0.0023 ± 0.0001, 0.0025 ± 0.0001, 0.0028 ± 0.0001, 0.0034 ± 0.0001, 0.0078 ± 0.0003 and 0.0385 ± 0.0012 μM, respectively (Table 2). Our results showed that pesticides inhibit AmCA activity with the following order: methomyl > oxamyl > deltamethrin > DDT > cypermethrin > diazinon. Our findings indicate these pesticides to act as potent inhibitors of AmCA, which might cause undesirable biological effects in bees, by disrupting their acid-base regulation as well as salt transport. Further studies are warranted in order to understand whether these inhibition data are relevant for the significant diminution of domestic and wild bee species worldwide. Several studies on bees as well as other organisms showed that sperm life depends on different parameters, among which the pH. It was observed that sperm loses its vitality if the pH level increases. This supports the relevance of the present findings. However, the data presented here need a careful *in vivo* (or in the hive) validation.

Disclosure statement

No potential conflict of interest was reported by the author(s).

ORCIDClaudiu T. Supuran  <http://orcid.org/0000-0003-4262-0323>**References**

- Ekinci D, Senturk M, Interactions of fungicides and pesticides with specific enzymes. In: Carisse O. ed. Fungicides. London, UK: In-Tech Open Publisher, 2010: 38–404.
- Sattari SZ, Bouwman AF, Martinez Rodríguez R, et al. Negative global phosphorus budgets challenge sustainable intensification of grasslands. *Nat Commun* 2016;7:10696.
- Osborne JL. Ecology: bumblebees and pesticides. *Nature* 2012;491:43–5.
- Marletto F, Patetta A, Manino A. Laboratory assessment of pesticide toxicity to bumble bees. *Bull Insectology* 2003;56: 155–8.
- Klein A-M, Vaissière BE, Cane JH, et al. Importance of pollinators in changing landscapes for world crops. *P Roy Soc B-Biol Sci* 2007;274:303–13.
- Allen-Wardell G, Bernhardt P, Bitner R, et al. The potential consequences of pollinator declines on the conservation of biodiversity and stability of food crop yields. *Conserv Biol* 1998;12:8–17.
- Potts SG, Biesmeijer JC, Kremen C, et al. Global pollinator declines: trends, impacts and drivers. *Trends Ecol Evol* 2010; 25:345–53.
- Heard TA. The role of stingless bees in crop pollination. *Annu Rev Entomol* 1999;44:183–206.
- (a) Supuran CT. Carbonic anhydrases: novel therapeutic applications for inhibitors and activators. *Nature Rev Drug Discov* 2008;7:168. (b) Supuran CT. Exploring the multiple binding modes of inhibitors to carbonic anhydrases for novel drug discovery. *Expert Opin Drug Discov* 2020;15: 671–86.
- (a) Demirdag R, Comakli V, Senturk M, et al. Characterization of carbonic anhydrase from sheep kidney and effects of sulfonamides on enzyme activity. *Bioorg Med Chem* 2013;21: 1522–5. (b) Supuran CT. Structure and function of carbonic anhydrases. *Biochem J* 2016;473:2023–32. (c) Alterio V, Di Fiore A, D'Ambrosio K, et al. Multiple binding modes of inhibitors to carbonic anhydrases: how to design specific drugs targeting 15 different isoforms? *Chem Rev* 2012;112: 4421–68.
- (a) Ekinci D, Kurbanoglu NI, Salamci E, et al. Carbonic anhydrase inhibitors: inhibition of human and bovine isoenzymes by benzenesulphonamides, cyclitols and phenolic compounds. *J Enzyme Inhib Med Chem* 2012;27:845–8. (b) Neri D, Supuran CT. Interfering with pH regulation in tumours as a therapeutic strategy. *Nat Rev Drug Discov* 2011;10:767–77.
- (a) Özdemir ZÖ, Şentürk M, Ekinci D. Inhibition of mammalian carbonic anhydrase isoforms I, II and VI with thiamine and thiamine-like molecules. *J Enzyme Inhib Med Chem* 2013;28:316–9. (b) Supuran CT. Advances in structure-based drug discovery of carbonic anhydrase inhibitors. *Expert Opin Drug Discov* 2017;12:61–88. (c) Nocentini A, Supuran CT. Advances in the structural annotation of human carbonic anhydrases and impact on future drug discovery. *Expert Opin Drug Discov* 2019;14:1175–97. (d) Supuran CT. How many carbonic anhydrase inhibition mechanisms exist? *J Enzyme Inhib Med Chem* 2016;31:345–60.
- (a) Ceyhun SB, Şentürk M, Yerlikaya E, et al. Purification and characterization of carbonic anhydrase from the teleost fish *Dicentrarchus labrax* (European seabass) liver and toxicological effects of metals on enzyme activity. *Environ Toxicol Pharmacol* 2011;32:69–74. (b) Supuran CT. Carbonic anhydrases and metabolism. *Metabolites* 2018;8:25. (c) Supuran CT. Carbonic anhydrase inhibitors and their potential in a range of therapeutic areas. *Expert Opin Ther Pat* 2018;28: 709–12. (d) Supuran CT. Applications of carbonic anhydrases inhibitors in renal and central nervous system diseases. *Expert Opin Ther Pat* 2018;28:713–21. (e) Supuran CT, Altamimi ASA, Carta F. Carbonic anhydrase inhibition and the management of glaucoma: a literature and patent review 2013–2019. *Expert Opin Ther Pat* 2019;29:781–92.
- Ceyhun SB, Şentürk M, Ekinci D, et al. Deltamethrin attenuates antioxidant defense system and induces the expression of heat shock protein 70 in rainbow trout. *Comp Biochem Phys C* 2010;152:215–23.
- Soydan E, Guler A, Biyik S, et al. Carbonic anhydrase from *Apis mellifera*: purification and inhibition by pesticides. *J Enzyme Inhib Med Chem* 2017;32:47–50.
- Wilbur KM, Anderson NG. Electrometric and colorimetric determination of carbonic anhydrase. *J Biol Chem* 1948;176: 147–54.
- Wang D, Hu J, Bobulescu IA, et al. A sperm-specific Na⁺/H⁺ exchanger (sNHE) is critical for expression and in vivo bicarbonate regulation of the soluble adenylyl cyclase (sAC). *Proc Natl Acad Sci USA* 2007;104:9325–30.
- Pettis JS, Rice N, Joselow K, et al. Colony failure linked to low sperm viability in honey bee (*Apis mellifera*) queens and an exploration of potential causative factors. *PLoS One* 2016;11:e0155833.
- Paynter E, Harvey M, Welch M, et al. Insights into the molecular basis of long-term storage and survival of sperm in the honeybee (*Apis mellifera*). *Sci Rep* 2017;7:40236.
- Sanchez-Bayo F, Goka K. Pesticide residues and bees – a risk assessment. *PLoS One* 2014;9:e94482.
- Cox-Foster DL, Conlan S, Holmes EC, et al. A metagenomic survey of microbes in honey bee colony collapse disorder. *Science* 2007;318:283–7.
- Cameron SA, Lozier JD, Strange JP, et al. Patterns of widespread decline in North American bumble bees. *Proc Natl Acad Sci USA* 2011;108:662–7.
- Thompson H, Ball R, Brown M, et al. *Varroa destructor* resistance to pyrethroid treatments in the United Kingdom. *Bull Insect* 2003;56:175–81.
- Underwood RM, vanEngelsdorp D. Colony Collapse Disorder: have we seen this before? *Bee Culture* 2007;135:13–15.
- Maini S, Medrzycki P, Porrini C. The puzzle of honey bee losses: a brief review. *Bull Insect* 2010;63:153–60.
- Vidau C, Diogon M, Aufauvre J, et al. Exposure to sublethal doses of fipronil and thiacloprid highly increases mortality of honeybees previously infected by *Nosema ceranae*. *PLoS One* 2011;6:e21550.
- Williams GR, Tarpy DR, vanEngelsdorp D, et al. Colony collapse disorder in context. *BioEssays* 2010;32:845–6.
- Goulson D, Lye GC, Darvill B. Decline and conservation of bumble bees. *Annual Rev Entomol* 2008;53:191–208.
- Stoner KA, Eitzer BD. Using a hazard quotient to evaluate pesticide residues detected in pollen trapped from honey bees (*Apis mellifera*) in Connecticut. *PLoS One* 2013;8: e77550.
- Chauzat MP, Carpentier P, Martel AC, et al. Influence of pesticide residues on honey bee (Hymenoptera: Apidae) colony health in France. *Environ Entomol* 2009;38:514–23.

31. Bernal J, Garrido-Bailon E, Nozal MJd, et al. Overview of pesticide residues in stored pollen and their potential effect on bee colony (*Apis mellifera*) losses in Spain. *J Econ Entomol* 2010;103:1964–71.
32. Choudhary A, Sharma DC. Dynamics of pesticide residues in nectar and pollen of mustard (*Brassica juncea* (L.) Czern.) grown in Himachal Pradesh (India). *Environ Monit Assess* 2008;144:143–50.
33. Thompson HM. Assessing the exposure and toxicity of pesticides to bumblebees (*Bombus* sp.). *Apidologie* 2001;32:305–21.
34. Greig-Smith PW, Thompson HM, Hardy AR, et al. Incidents of poisoning of honeybees (*Apis mellifera*) by agricultural pesticides in Great Britain 1981–1991. *Crop Prot* 1994;13:567–81.
35. (a) Ekinci D, Şentürk M, Küfrevioğlu Öİ. Salicylic acid derivatives: synthesis, features and usage as therapeutic tools. *Expert Opin Ther Pat* 2011;21:1831–41. (b) Innocenti A, Vullo D, Scozzafava A, Supuran CT. Carbonic anhydrase inhibitors. Interactions of phenols with the 12 catalytically active mammalian isoforms (CA I – XIV). *Bioorg Med Chem Lett* 2008;18:1583–7. (c) Innocenti A, Vullo D, Scozzafava A, Supuran CT. Carbonic anhydrase inhibitors. Inhibition of mammalian isoforms I – XIV with a series of substituted phenols including paracetamol and salicylic acid. *Bioorg Med Chem* 2008;16:7424–8.
36. (a) Ekinci D, Ceyhun SB, Şentürk M, et al. Characterization and anions inhibition studies of an α -carbonic anhydrase from the teleost fish *Dicentrarchus labrax*. *Bioorg Med Chem* 2011;19:744–8. (b) De Simone G, Supuran CT. (In)organic anions as carbonic anhydrase inhibitors. *J Inorg Biochem* 2012;111:117–29. (c) Supuran CT. Carbon- versus sulphur-based zinc binding groups for carbonic anhydrase inhibitors? *J Enzyme Inhib Med Chem* 2018;33:485–95.
37. (a) Dizdaroglu Y, Albay C, Arslan T, et al. Design, synthesis and molecular modelling studies of some pyrazole derivatives as carbonic anhydrase inhibitors. *J Enzyme Inhib Med Chem* 2020;35:289–97. (b) Nocentini A, Supuran CT, Winum JY. Benzoxaborole compounds for therapeutic uses: a patent review (2010-2018). *Expert Opin Ther Pat* 2018;28:493–504. (c) Supuran CT. Carbonic anhydrase activators. *Future Med Chem* 2018;10:561–73.
38. Bielmyer GK, Brix KV, Grosell A. Is Cl⁻ protection against silver toxicity due to chemical speciation? *Aquat Toxicol* 2008;87:81–7.
39. (a) Innocenti A, Scozzafava A, Parkkila S, et al. Investigations of the esterase, phosphatase, and sulfatase activities of the cytosolic mammalian carbonic anhydrase isoforms I, II, and XIII with 4-nitrophenyl esters as substrates. *Bioorg Med Chem Lett* 2008;18:2267–71. (b) Innocenti A, Supuran CT. Paraoxon, 4-nitrophenyl phosphate and acetate are substrates of α - but not of β -, γ - and ζ -carbonic anhydrases. *Bioorg Med Chem Lett* 2010;20:6208–12.
40. (a) Lopez M, Vu H, Wang CK, et al. Promiscuity of carbonic anhydrase II. Unexpected ester hydrolysis of carbohydrate-based sulfamate inhibitors. *J Am Chem Soc* 2011;133:18452–62. (b) Truppo E, Supuran CT, Sandomenico A, et al. Carbonic anhydrase VII is S-glutathionylated without loss of catalytic activity and affinity for sulfonamide inhibitors. *Bioorg Med Chem Lett* 2012;22:1560–4.
41. Tanc M, Carta F, Scozzafava A, Supuran CT. α -Carbonic anhydrases possess thioesterase activity. *ACS Med Chem Lett* 2015;6:292–5.
42. Angeli A, Carta F, Donnini S, et al. Selenolesterase enzyme activity of carbonic anhydrases. *Chem Commun* 2020;56:4444–7.
43. Bitman J, Cecil HC, Fries GF. DDT-induced inhibition of avian shell gland carbonic anhydrase: a mechanism for thin eggshells. *Science* 1970;168:594–6.
44. Lindahl M, Svensson LA, Liljas A. Metal poison inhibition of carbonic anhydrase. *Proteins* 1993;15:177–82.
45. Supuran CT, Conroy CW, Maren TH. Is cyanate a carbonic anhydrase substrate? *Proteins* 1997;27:272–8.
46. West D, Pinaud MA, Tu C, et al. Human carbonic anhydrase II-cyanate inhibitor complex: putting the debate to rest. *Acta Crystallogr F Struct Biol Commun* 2014;70:1324–7.
47. Jonsson BH, Liljas A. Perspectives on the classical enzyme carbonic anhydrase and the search for inhibitors. *Biophys J* 2020;119:1275–80.