



FLORE Repository istituzionale dell'Università degli Studi di Firenze

Interior a priori estimates for supersolutions of fully nonlinear subelliptic equations under geometric conditions

Questa è la Versione finale referata (Post print/Accepted manuscript) della seguente pubblicazione:

Original Citation:

Interior a priori estimates for supersolutions of fully nonlinear subelliptic equations under geometric conditions / Goffi, Alessandro. - In: BULLETIN OF THE LONDON MATHEMATICAL SOCIETY. - ISSN 0024-6093. - STAMPA. - 56:(2024), pp. 1385-1398. [10.1112/blms.13001]

Availability:

This version is available at: 2158/1383949 since: 2024-11-28T16:42:44Z

Published version: DOI: 10.1112/blms.13001

Terms of use: Open Access

La pubblicazione è resa disponibile sotto le norme e i termini della licenza di deposito, secondo quanto stabilito dalla Policy per l'accesso aperto dell'Università degli Studi di Firenze (https://www.sba.unifi.it/upload/policy-oa-2016-1.pdf)

Publisher copyright claim:

(Article begins on next page)

INTERIOR A PRIORI ESTIMATES FOR SUPERSOLUTIONS OF FULLY NONLINEAR SUBELLIPTIC EQUATIONS UNDER GEOMETRIC CONDITIONS

ALESSANDRO GOFFI

ABSTRACT. In this note, we prove interior a priori first- and second-order estimates for solutions of fully nonlinear degenerate elliptic inequalities structured over the vector fields of Carnot groups, under the main assumption that u is semiconvex along the fields. These estimates for supersolutions are new even for linear subelliptic inequalities in nondivergence form, whereas in the nonlinear setting they do not require neither convexity nor concavity on the second derivatives. We complement the analysis exhibiting an explicit example showing that horizontal $W^{2,q}$ regularity of Calderón-Zygmund type for fully nonlinear subelliptic equations posed on the Heisenberg group cannot be in general expected in the range q < Q, Q being the homogeneous dimension of the group.

1. INTRODUCTION

Recently, E. Braga and D. Moreira obtained an optimal $C^{1,\alpha}$ regularity result for semiconvex supersolutions to fully nonlinear uniformly elliptic equations with an unbounded source term $f \in L^q$, q > n, n being the dimension of the ambient space, see Theorem 3.6 in [9]. This extended a previous a priori estimate on the modulus of continuity of the gradient for semiconvex supersolutions to linear uniformly elliptic equations proved by L. Caffarelli, R. Kohn, L. Nirenberg and J. Spruck [12], cf. Corollary of Lemma 2.2. An improvement of such a result up to q = n, along with the optimal interior regularity of the convex envelope of L^n -viscosity supersolutions to Pucci's extremal equations with unbounded coefficients, were investigated by E. Braga, A. Figalli and D. Moreira in [8].

Given a family $\mathcal{X} = \{X_1, ..., X_m\}$ of C^2 vector fields of Carnot-type, the purpose of this note, inspired by the aforementioned works, is to establish a priori high-order estimates in second order Sobolev spaces (and in first-order Hölder spaces via Morrey-type embeddings as a byproduct) for solutions to degenerate fully nonlinear subelliptic inequalities of the form

(1)
$$G(x, (D_{\mathcal{X}}^2 u)^*) \le f(x) \text{ in } \Omega,$$

where $(D^2_{\mathcal{X}}u)^*$ stands for the symmetrized horizontal Hessian along the vector fields $(D^2_{\mathcal{X}}u)^* = \frac{X_i X_j u + X_j X_i u}{2}$. Here, $G: \Omega \times \mathcal{S}_m \to \mathbb{R}, \Omega \subseteq \mathbb{R}^n, n \ge m, \mathcal{S}_m$ being the space of $m \times m$ symmetric matrices, is continuous and uniformly subelliptic in the sense that

(2)
$$\mathcal{M}^{-}_{\lambda,\Lambda}(X-Y) \leq G(x,X) - G(x,Y) \leq \mathcal{M}^{+}_{\lambda,\Lambda}(X-Y), X, Y \in \mathcal{S}_m,$$

Date: January 14, 2024.

²⁰²⁰ Mathematics Subject Classification. 35D40, 35B65, 35H20.

The author is member of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). The author was partially supported by the INdAM-GNAMPA Project 2022 "Proprietà quantitative e qualitative per EDP non lineari con termini di gradiente" and by the King Abdullah University of Science and Technology (KAUST) project CRG2021-4674 "Mean-Field Games: models, theory and computational aspects".

 $\mathcal{M}_{\lambda,\Lambda}^{-}, \mathcal{M}_{\lambda,\Lambda}^{+}$ being the Pucci's extremal operators, see (4)-(5) below. A longstanding open problem in the regularity theory of such equations or, more generally, subelliptic equations in nondivergence form with bounded measurable coefficients, is the validity of an analogue of the Krylov-Safonov Harnack inequality. This is mainly due to the lack of Aleksandrov-Bakel'man-Pucci (briefly ABP) maximum principles, see the introduction of [17] for a thorough discussion. In this framework, it was only shown that ABP type estimates cannot be produced, at least in Heisenberg-type groups, when the right-hand side of the equation lies in L^q , q < Q, Q being the homogeneous dimension [17], in accordance with the classical results of A. D. Aleksandrov and C. Pucci [3, 30]. Nonetheless, some invariant Harnack inequalities have been obtained under smallness conditions on the ratio among the ellipticity constants (this can be referred to Landis/Cordes-type conditions) for nondivergence structure elliptic problems posed on Carnot groups of Heisenberg-type, see [23, 31, 1], and also for parabolic Kolmogorov and kinetic operators with diffusion in nondivergence form, cf. [2]. For weighted ABP estimates and Harnack inequalities on Grushin geometries see [29]. We also mention that a global Krylov-Safonov Harnack inequality was proved in [10] for equations in nondivergence form on Riemannian manifolds under curvature lower-bounds. Note that the results in [31, 1] immediately lead to an extension to the fully nonlinear setting, and also provide low-regularity in Hölder spaces using classical arguments, under suitable restrictions on the ellipticity constants. Still, a subelliptic version of the $C^{1,\alpha}$ estimate of Cordes-Nirenberg type seems not available, and to our knowledge no results appeared in the context of degenerate fully nonlinear subelliptic equations of second order in horizontal $C_{\mathcal{X}}^{2,\alpha}$ and $W_{\mathcal{X}}^{2,q}$ spaces, even for smooth solutions and smooth functionals. Indeed, the structure of the problem over horizontal Hessians prevents from the use of classical linearization arguments, based for example on the Bernstein method, see e.g. [14, 12, 32, 19]. Typically, when \mathcal{X} are the Euclidean vector fields, see [19], if u smooth solves the model problem $F(D^2u) = 0$, with $F \in C^1$ and uniformly elliptic in the sense of [14], then $v = \partial_e u, e \in \mathbb{R}^n$ with |e| = 1, solves a linear nondivergence structure equation of the form

$$a_{ij}(x)\partial_{ij}v = 0$$
, $a_{ij}(x) = F_{ij}(D^2u).$

Then, the Krylov-Safonov Hölder regularity applies to v (since no regularity properties on a_{ij} are required) leading thus to $C^{1,\alpha}$ estimates under no assumptions on the nonlinearity other than the uniform ellipticity.

However, the non-commutative structure of sub-Riemannian geometries and the absence of the Krylov-Safonov theory do not allow to reproduce the previous approach. Similarly, the Bernstein method, based on a linearization argument, cannot be performed: this is easily seen by the recent Bochner-type formulas obtained in [21] for step-2 Carnot groups, which give rise to additional commutator terms.

In the classical regularity theory for fully nonlinear equations, though $C^{1,\alpha}$, $\alpha < 1$, estimates are obtained under essentially no assumptions on $F = F(D^2u)$, $C^{1,1}$ estimates (and higherorder $C^{2,\alpha}$ estimates through the Evans-Krylov theorem) usually require concavity/convexity type assumptions on F. However, $C^{1,1}$ estimates were proved in [24] for strictly elliptic inequalities under the geometric assumption that the unknown function is convex: this was obtained through the properties of the convex envelope found in [4], without requiring neither convexity nor concavity on the fully nonlinear operator. The aforementioned works [9, 8] lead to a further development, proving indeed a generalization of the results in [12, 24] for more general semiconvex semisolutions with unbounded coefficients in Lebesgue spaces.

In the subelliptic context, a generalized notion of convexity (and, thus, semiconvexity) along

the vector fields, called \mathcal{X} -convexity, was discussed in [5], see also [16] for Carnot groups. The results in [5] say that $u \in \text{USC}$ is convex along the fields of \mathcal{X} if and only if $(D^2_{\mathcal{X}}\varphi(x))^* \geq 0$ for all smooth φ and $x \in \operatorname{argmax}(u - \varphi)$.

Therefore, having such a notion at our disposal and following [9], we prove the following model result for horizontal Hessian inequalities structured over the fields of a Carnot group: if u is \mathcal{X} -semiconvex according to [5] and a smooth solution to

$$G((D^2_{\mathcal{X}}u)^*) \le f \text{ in } B_1$$

with $f \in L^{\infty}(B_1)$, G satisfying (2) with G(0) = 0 (with no other assumptions on G = G(X)), where B_r is a metric ball of radius r, then the following interior a priori estimate holds:

$$\|(D_{\mathcal{X}}^2 u)^*\|_{L^q(B_{\frac{1}{n}})} \lesssim C(\|u\|_{L^q(B_1)}, \|f\|_{L^q(B_1)}), \ q \in (1, \infty).$$

When q is larger than a certain threshold involving the homogeneous dimension associated to the fields we get $C^{1,\alpha}$ estimates with respect to the corresponding sub-Riemannian distance. As stated in the previous results, throughout the paper we work with classical solutions, though all the estimates are independent of the smoothness and depend only on the structural constants and the integrability of the data. We briefly discuss some one-side second order estimates, that can be thus regarded as regularity estimates, in the viscosity sense, see Remark 3.2.

The prototype example to which our results apply is the (degenerate) Isaacs equation, i.e. a PDE of the form

$$G((D_{\mathcal{X}}^2 u)^*) = \sup_{\beta} \inf_{\alpha} \operatorname{Tr}(A_{\alpha\beta}(x) \ (D_{\mathcal{X}}^2 u)^*) = \sup_{\beta} \inf_{\alpha} \operatorname{Tr}(A_{\alpha\beta}(x)\sigma^T(x)D^2 u\sigma(x)),$$

where $\sigma \in \mathbb{R}^{n \times m}$ is a (degenerate) matrix having the fields X_i as columns, α, β belong to some control sets A, B respectively, while $D^2 u \in S_n$ is the standard Hessian of the unknown function u. It is well-known that every nonlinear operator $G = G((D^2_{\mathcal{X}}u)^*)$ satisfying (2) can be written in Isaacs form as follows, cf. [11]: by (2) one has for $X, Y \in S_m$ and $\mathcal{A}_{\lambda,\Lambda} = \{A \in S_m, \lambda I_m \leq A \leq \Lambda I_m\}$,

$$G(X) - G(Y) \le \mathcal{M}^+_{\lambda,\Lambda}(X - Y) = \sup_{A \in \mathcal{A}_{\lambda,\Lambda}} \operatorname{Tr}(A(X - Y)).$$

Since the equality is attained when X = Y, one obtains

$$G(X) = \min_{Y \in \mathcal{S}_m} \max_{A \in \mathcal{A}_{\lambda,\Lambda}} \{ \operatorname{Tr}(AX) + G(Y) - \operatorname{Tr}(AY) \}$$

By taking then $X = (D^2_{\mathcal{X}} u)^*$ we have

$$G((D^2_{\mathcal{X}}u)^*) = \min_{Y \in \mathcal{S}_m} \max_{A \in \mathcal{A}_{\lambda,\Lambda}} \{ \operatorname{Tr}(A(D^2_{\mathcal{X}}u)^*) + G(Y) - \operatorname{Tr}(AY) \}.$$

Note that the previous operator is degenerate elliptic if expressed in Euclidean coordinates. However, the previous announced results turn out to be new even for general linear subelliptic equations of the form $G((D_{\mathcal{X}}^2 u)^*) = \operatorname{Tr}(A(x)D_{\mathcal{X}}^2 u)^*)$, where A satisfies $\lambda I_m \leq A \leq \Lambda I_m$ and has bounded measurable entries, and, finally, apply to any Carnot group.

Some comments on the results are now in order. Though the (geometric) a priori requirement on the solution could appear strong (and thus the consequent estimate conditional to the geometric bound), the procedure we are going to implement allows to conclude the estimate for semisolutions, and not only for solutions. Indeed, $W^{2,q}$ estimates are typically obtained

for solutions to uniformly elliptic PDEs (which indicates a two-side analytic control on the problem, since, roughly speaking, G = f with $|f| \leq C$ means $-C \leq G \leq C$). Indeed, classical results for the Poisson equation show that if u solves $\Delta u = f \in L^{\infty}$, i.e. $-C_f \leq \Delta u \leq C_f$ for some $C_f > 0$, then $u \in C^{1,\alpha} \cap W^{2,q}$ and the following estimates hold

$$\begin{aligned} \|u\|_{W^{2,q}(B_{\frac{1}{2}})} &\leq C(n)(\|u\|_{L^{\infty}(B_{1})} + C_{f}), \\ \|u\|_{C^{1,\alpha}(B_{\frac{1}{2}})} &\leq C(n)(\|u\|_{L^{\infty}(B_{1})} + C_{f}). \end{aligned}$$

Motivated by these results, one can ask whether it is possible to expect some (high) level of regularity of u when one has only an analytic control from above, i.e. $\Delta u \leq C$, based on the regularity properties of the right-hand side and a geometric control from below involving second derivatives, such as a convexity-type assumption.

Besides, the analysis of the regularity properties of semisolutions to fully nonlinear degenerate equations satisfying geometric conditions like convexity is motivated by the understanding of a regularity theory for such degenerate PDEs. Indeed, a crucial step in the proof of the ABP maximum principle for equations over Euclidean vector fields in [13] is to show that the convex envelope of supersolutions to $\mathcal{M}_{\lambda,\Lambda}^{-}(D^2u) \leq f$ is $C_{\text{loc}}^{1,1}$ when $f \in L^{\infty}$, cf. Chapter 3 of [14]. Thus, a tightly related question is the study of the regularity of supersolutions satisfying convexity constraints and linear or nonlinear partial differential inequalities in nondivergence form. In this direction, a study of the horizontal convex envelope in the Heisenberg group, along with its application to the study of horizontal convexity properties of solutions to fully nonlinear equations was performed in [28]. This analysis has its roots in the earlier work by O. Alvarez, J.-M. Lasry and P.-L. Lions [4]. Based on this study, in the course of Section 3.2 we provide some sufficient conditions to obtain convexity estimates on solutions, providing an instance of the full result unconditional to the geometric bound. Another paper dealing with such convexity preserving properties for nondivergence structure uniformly parabolic PDEs is [26].

Other than the previous motivations, this seems the first (a priori) regularity result concerning fully nonlinear subelliptic equations in higher-order spaces. Few results are available in the literature of nondivergence subelliptic equations: the works [31, 1] allow to deduce a priori estimates at the level of C^{α} spaces for subelliptic Hessian equations with bounded right-hand side. Indeed, in the smooth setting one can write

$$G((D_{\mathcal{X}}^2 u)^*) = \int_0^1 G_{ij}(t(D_{\mathcal{X}}^2 u)^*)) dt \ (D_{\mathcal{X}}^2 u)_{ij}^* + G(0) = \operatorname{Tr}(A(x)(D_{\mathcal{X}}^2 u)^*) + G(0),$$

 \mathbf{SO}

$$G((D_{\mathcal{X}}^2 u)^*) = 0 \iff \operatorname{Tr}(A(x)(D_{\mathcal{X}}^2 u)^*) = -G(0),$$

and apply e.g. the results from [31, 1] valid when the ellipticity constants are sufficiently close to each other.

We also prove, as a further step towards a possible development of a regularity theory for fully nonlinear subelliptic equations, that $W^{2,q}_{\mathcal{X}}$ estimates for the uniformly subelliptic equation

$$G((D^2_{\mathcal{X}}u)^*) = f \in L^q$$

fail when q < Q, at least in the Heisenberg group, Q being its homogeneous dimension. This shows that the best possible integrability order to get such estimates is q = Q. This is done adapting a counterexample proposed by L. Caffarelli in [13]. We emphasize that in this sub-Riemannian context there is a dimensional discrepancy between the dimension of the horizontal layer, say m, and the homogeneous dimension Q (and even the topological dimension n), so it is unclear the right integrability order to deduce such an estimate. This shares some similarities with a result obtained in [17] for linear subelliptic equations in nondivergence form with bounded measurable coefficients in H-type groups, concerning the validity of the uniqueness of solutions in the Sobolev space $W_{\mathcal{X}}^{2,Q}$. The example in [17] also implies that ABP-type estimates cannot hold when the right-hand of the equation does not belong to L^q , $q \geq Q$.

The paper is organized as follows. Section 2 gives some preliminaries on sub-Riemannian geometries and convexity along vector fields. Section 3 is devoted to the proof of the a priori estimate for elliptic inequalities. Section 4 ends the paper with the failure of second order estimates in L^q spaces when q < Q.

2. Preliminaries on subelliptic structures

We denote by $\mathcal{X} = \{X_1(x), ..., X_m(x)\}$ a family of C^2 -vector fields in \mathbb{R}^n and $\Omega \subset \mathbb{R}^n$ an open and connected set. We recall that the Carnot-Carathéodory (briefly CC), or sub-Riemannian distance d on \mathbb{R}^n associated to the vector fields of the family \mathcal{X} , is the length of the shortest horizontal curves connecting two-points, and is denoted by d. The pair (\mathbb{R}^n, d) is a sub-Riemannian geometry if $d(x, y) < +\infty$ for all $x, y \in \mathbb{R}^n$ and the vector fields \mathcal{X} are the generators. Important examples of such spaces are Carnot groups.

A stratified group (or Carnot group) [7, Definition 2.2.3] is a connected and simply connected Lie group whose Lie algebra \mathcal{G} admits a stratification $\mathcal{G} = \bigoplus_{i=1}^{r} V_i$, where the layers V_i satisfy the relations $[V_i, V_{i-1}] = V_i$ for $2 \leq i \leq r$ and $[V_1, V_r] = 0$, with $[V, W] := \operatorname{span}\{[v, w] : v \in V, w \in W\}$. Here, r is called the step of the group, and we set $m = n_1 = \dim(V_1)$, which stands for the dimension of the horizontal layer, $n_i = \dim(V_i)$, $2 \leq i \leq r$. A stratified group can be identified with a homogeneous Carnot group up to an isomorphism by means of [7, Section 2.2.3]. A homogeneous Carnot group \mathbb{G} (see e.g. [7, Definition 1.4.1]) can be identified with $\mathbb{R}^n = \mathbb{R}^{n_1} \times \ldots \times \mathbb{R}^{n_r}$ $(n = \sum_{i=1}^r n_i)$ endowed with a group law \star if for any $\lambda > 0$ the dilation $\delta_{\lambda} : \mathbb{R}^n \to \mathbb{R}^n$ of the form $\delta_{\lambda}(x) = (\lambda x^{(1)}, \ldots, \lambda^r x^{(r)})$ is an automorphism of the group, where $x = (x^{(1)}, \ldots, x^{(r)})$, $x^{(i)} \in \mathbb{R}^{n_i}$. Then the vector fields in \mathbb{R}^n of the family $\mathcal{X} = \{X_1, \ldots, X_m\}$ generate the homogeneous Carnot group $(\mathbb{R}^n, \star, \delta_{\lambda})$ if they are left-invariant on \mathbb{G} and such that $X_j(0) = \partial_{x_j}|_0$ for $j = 1, \ldots, n_1$ span \mathbb{R}^n at every point $x \in \mathbb{R}^n$. In this case we say that \mathbb{G} has step r and $m = n_1$ generators. We also denote by $\Delta_{\mathcal{X}} = \sum_{i=1}^m X_i^2$ the operator sum of squares of vector fields, usually known as sub-Laplacian on the Carnot group \mathbb{G} . We also denote with

$$Q := \sum_{i=1}^{r} i n_i = \sum_{i=1}^{r} i \dim(V_i)$$

the homogeneous dimension of the group, with

$$D_{\mathcal{X}}u := (X_1u, \dots, X_mu)$$

the horizontal gradient along the vector fields, and with $D_{\mathcal{X}}^2 u = X_i X_j u$, i, j = 1, ..., m, the horizontal Hessian built over the frame \mathcal{X} . Finally, $(D_{\mathcal{X}}^2 u)^* = \frac{X_i X_j u + X_j X_i u}{2} \in \mathcal{S}_m$ is the symmetrized horizontal Hessian.

We point out that it was proved in [7] that the fields of a Carnot group can be written as

$$X_j = \partial_{x_j} + \sum_{i=m+1}^n b_{ij}(x)\partial_{x_i}, j = 1, ..., m.$$

where $b_{ij}(x) = \sigma_{ij}(x_1, ..., x_{i-1})$ are homogeneous polynomials of degree less than or equal to n - m. A model example that we will analyze in Section 4 is the Heisenberg group in $\mathbb{H}^d \simeq \mathbb{R}^{2d+1}$, n = 2d + 1, m = 2d, Q = 2d + 2, where the fields of the horizontal layer are

$$X_i = \partial_{x_i} + 2x_{i+d}\partial_{2d+1}, X_{i+d} = \partial_{x_i} - 2x_i\partial_{2d+1}, i = 1, ..., d$$

We denote by $B_r(x) = \{y \in \mathbb{G} : d(x, y) < r\}$ the metric ball of radius r.

We recall that for u smooth, the convexity of u along the fields of the family \mathcal{X} is equivalent to the positive semidefiniteness of the symmetrized horizontal Hessian $(D^2_{\mathcal{X}}u)^*$. This has been extended to the viscosity setting in [5] to which we refer for further details.

3. Interior a priori estimates for elliptic inequalities under \mathcal{X} -semiconvexity conditions

3.1. High-order local estimates under geometric conditions. Let \mathcal{X} be a family of vector fields of Carnot-type, and let $X_1, ..., X_m$ be the generators of the group, where m is the dimension of the horizontal layer. We assume that $G : \Omega \times \mathbb{R} \times \mathbb{R}^m \times \mathcal{S}_m \to \mathbb{R}, \Omega \subseteq \mathbb{R}^n, n \geq m$, is continuous and uniformly subelliptic, i.e.

(3)
$$\mathcal{M}^{-}_{\lambda,\Lambda}(X-Y) \le G(x,r,p,X) - G(x,r,p,Y) \le \mathcal{M}^{+}_{\lambda,\Lambda}(X-Y)$$

for $X, Y \in \mathcal{S}_m$ and $(x, r, p) \in \Omega \times \mathbb{R} \times \mathbb{R}^m$, where

(4)
$$\mathcal{M}^+_{\lambda,\Lambda}(M) = \Lambda \sum_{e_k > 0} e_k + \lambda \sum_{e_k < 0} e_k = \sup\{\operatorname{Tr}(AM), \lambda I_m \le A \le AI_m\}$$

(5)
$$\mathcal{M}^{-}_{\lambda,\Lambda}(M) = \lambda \sum_{e_k > 0} e_k + \Lambda \sum_{e_k < 0} e_k = \inf\{\operatorname{Tr}(AM), \lambda I_m \le A \le A I_m\}$$

and e_k denotes the k-th eigenvalue of $M \in \mathcal{S}_m$.

We prove the following interior a priori estimate in the horizontal spaces $W^{2,q}_{\mathcal{X}}(\Omega) = \{u \in L^q(\Omega) : X_I u \in L^q(\Omega) \text{ for any } |I| \leq 2\}$ and $C^{1,\alpha}_{\mathcal{X}}$, the classical first-order Hölder space with respect to the sub-Riemannian distance. Note that even if $u \in C^2$ and f is bounded, bounds will depend only on the summability of the data, so this can be regarded as an a priori estimate under the one-side geometric condition that u is semiconvex along the fields of the family \mathcal{X} .

Theorem 3.1. Let $u \in C^2(B_1) \cap L^q(B_1)$, $q \in (1, \infty)$, be a classical and \mathcal{X} -semiconvex solution (with constant 4C) to the inequality

(6)
$$G(x, u, D_{\mathcal{X}}u, (D_{\mathcal{X}}^2u)^*) \le f \text{ in } B_1$$

with $f \in L^{\infty}(B_1)$, G satisfying (3) with $G(x, u, D_{\mathcal{X}}u, 0)$ bounded (independently of u), i.e.

$$|G(x, u, D_{\mathcal{X}}u, 0)| \le M,$$

with M independent of u. Then, there exist a constant \tilde{C} depending on $C, m, \lambda, \Lambda, M, q$ and a universal constant $K_1 = K_1(m, q, \lambda, \Lambda)$ such that

$$\|u\|_{W^{2,q}_{\mathcal{X}}(B_{\frac{1}{2}})} \le K_1(\tilde{C} + \|u\|_{L^q(B_1)} + \|f\|_{L^q(B_1)}).$$

If, in addition, q > Q and $u \in L^{\infty}(B_1)$, we have the estimate

$$\|u\|_{C^{1,1-\frac{Q}{q}}_{\mathcal{X}}(B_{\frac{1}{2}})} \le K_2(\tilde{C} + \|u\|_{L^{\infty}(B_1)} + \|f\|_{L^q(B_1)}).$$

Here, K_1 and K_2 are universal constants depending on m, q, λ, Λ , while K_2 depends also on the constant of the corresponding Sobolev embedding.

Proof. We follow [9, Remark 1.2]. Since u is smooth and \mathcal{X} -semiconvex with constant 4C, then $v = u + 2C \sum_{i=1}^{m} x_i^2 = u + z$ is \mathcal{X} -convex. Therefore, recalling that, cf. [14],

(7)
$$\mathcal{M}^{-}_{\lambda,\Lambda}(X+Y) \ge \mathcal{M}^{-}_{\lambda,\Lambda}(X) + \mathcal{M}^{-}_{\lambda,\Lambda}(Y)$$

and

(8)
$$\mathcal{M}^{-}_{\lambda,\Lambda}(-X) = -\mathcal{M}^{+}_{\lambda,\Lambda}(X)$$

we have, using first (3) with r = v - z, $p = D_{\mathcal{X}}(v - z)$, $X = (D_{\mathcal{X}}^2(v - z))^*$ and Y = 0, then (7)-(8) and also that v is \mathcal{X} -convex

$$\begin{split} f \geq G(x, v-z, D_{\mathcal{X}}(v-z), (D_{\mathcal{X}}^{2}(v-z))^{*}) \geq \mathcal{M}_{\lambda,\Lambda}^{-}((D_{\mathcal{X}}^{2}(v-z))^{*}) + G(x, v-z, D_{\mathcal{X}}(v-z), 0) \\ \geq \mathcal{M}_{\lambda,\Lambda}^{-}((D_{\mathcal{X}}^{2}v)^{*}) + \mathcal{M}_{\lambda,\Lambda}^{-}((D_{\mathcal{X}}^{2}(-z))^{*}) + G(x, u, D_{\mathcal{X}}u, 0) \\ = \mathcal{M}_{\lambda,\Lambda}^{-}((D_{\mathcal{X}}^{2}v)^{*}) - \mathcal{M}_{\lambda,\Lambda}^{+}((D_{\mathcal{X}}^{2}z)^{*}) + G(x, u, D_{\mathcal{X}}u, 0) \\ \geq \lambda \mathrm{Tr}((D_{\mathcal{X}}^{2}v)^{*}) - 4mC\Lambda + G(x, u, D_{\mathcal{X}}u, 0) \\ = \lambda \left(\sum_{i=1}^{m} X_{i}^{2}u + 4mC\right) - 4mC\Lambda + G(x, u, D_{\mathcal{X}}u, 0) \\ \geq \lambda \sum_{i=1}^{m} X_{i}^{2}u - 4mC(\Lambda - \lambda) - M. \end{split}$$

Therefore, the last estimate together with the \mathcal{X} -semiconvexity give the pointwise bound

(9)
$$-4Cm \le \sum_{i=1}^{m} X_i^2 u(x) \le \lambda^{-1} (f(x) + 4mC(\Lambda - \lambda) + M), x \in B_1.$$

Therefore, using that u is $\mathcal X\text{-semiconvex},$ we get for a constant K=K(m)>0 and i,j=1,...,m

(10)
$$|X_i X_j u| \le \sum_{i,j=1}^m |X_i X_j u| \le K(\Delta_{\mathcal{X}} u + 4Cm).$$

Indeed, denoting by $||X||_{\text{spec}} = \max\{e_i(X)\}\$ the spectral norm of $X \in S_m$ and by $||X||_1 = \sum_{i,j=1}^m |x_{ij}|$, since $X \in S_m$, we know that for a constant L(m) > 0 it holds

 $||X||_1 \le L(m) ||X||_{\text{spec}}.$

This implies, since v is \mathcal{X} -convex, the following inequality

$$|X_i X_j v| \le L(m) || (D_{\mathcal{X}}^2 v)^* ||_{\text{spec}} \le L(m) \text{Tr}(X_i X_j v) = L(m) \Delta_{\mathcal{X}} v = L(m) (\Delta_{\mathcal{X}} u + 4Cm)$$

We conclude by noting that

$$|X_i X_j v| = |X_i X_j u + 4C\delta_{ij}| \ge |X_i X_j u|.$$

By interpolation estimates, see e.g. the proof of Proposition 2.9 in [6], for every $\delta > 0$ there exists a constant $c_q > 0$ such that

(11)
$$\|D_{\mathcal{X}}u\|_{L^{q}(B_{\frac{1}{2}})} \leq \delta \|\Delta_{\mathcal{X}}u\|_{L^{q}(B_{1})} + \frac{c_{q}}{\delta} \|u\|_{L^{q}(B_{1})}.$$

Recalling the definition of the $W^{2,q}_{\mathcal{X}}$ norm and combining (9)-(10)-(11), we obtain for a universal constant $K_1 = K_1(m,q,\lambda,\Lambda)$ and a constant $\tilde{C} = \tilde{C}(\lambda,\Lambda,m,C,q)$, both independent of u,

$$\|u\|_{W^{2,q}_{\mathcal{X}}(B_{\frac{1}{2}})} \le K_1(\tilde{C} + \|u\|_{L^q(B_1)} + \|f\|_{L^q(B_1)} + M).$$

If q > Q and u is bounded, by the horizontal Morrey embedding, see e.g. [20], we have

$$\|u\|_{C^{1,1-\frac{Q}{q}}_{\mathcal{X}}(B_{\frac{1}{2}})} \le K_2(\tilde{C} + \|u\|_{L^{\infty}(B_1)} + \|f\|_{L^q(B_1)})$$

for a different universal constant K_2 depending on K_1 and C_s , the latter being the constant of the embedding $W^{2,q}_{\mathcal{X}} \hookrightarrow C^{1,\alpha}_{\mathcal{X}}$.

Some remarks are in order:

Remark 3.2. The estimate in the right-hand side of (9) can be made rigorous in the weak setting of viscosity solutions by means of the transitivity of the viscosity inequalities. More precisely, adapting Lemma 2.12 in [14] in the degenerate setting of Hörmander vector fields, one can prove that if $u \in \text{LSC}(B_1)$ is a viscosity solution to (6) with $f \in C(B_1)$, then there exists a constant K depending on $\sup_{B_1} f$, $\lambda, \Lambda, |G(x, u, D_X u, 0)|, m, C$ such that

$$\sum_{i=1}^{m} X_i X_j \varphi(x) \le K$$

provided $\varphi \in C^2$ and $u - \varphi$ has a local minimum at x.

Remark 3.3. The estimates in Theorem 3.1 can be obtained for the smaller class of \mathcal{X} -convex supersolutions, but under the weaker condition of strict (sub)ellipticity

$$\lambda \operatorname{Tr}(X - Y) + G(x, r, p, Y) \le G(x, r, p, X)$$

or the one-side condition

$$G(x, r, p, X) \ge G(x, r, p, 0) + \mathcal{M}^{-}_{\lambda, \Lambda}(X)$$

For instance, the linear operator G(x, X) = Tr(A(x)X) satisfies the first condition when $A \ge \lambda I_m$. In this case one can get $C_{\mathcal{X}}^{1,1}$ estimates along vector fields, as it is done in Theorems 3 and 4 of [24]. It is enough to prove that u is a solution to the inequality

$$\lambda \Delta_{\mathcal{X}} u + G(x, u, D_{\mathcal{X}} u, 0) \le 0 \text{ in } \mathbb{R}^n.$$

To see the validity of $C^{1,1}$ estimates when u is smooth, it is sufficient to exploit the strict ellipticity, together with the fact that $(D^2_{\mathcal{X}}u)^* \geq 0$ holds in the classical sense and that, for $A \geq 0, A \in \mathcal{S}_m$, we have the inequality $A \leq C(m) \operatorname{Tr}(A) I_m$. The following estimate then follows for \mathcal{X} -Lipschitz solutions by exploiting the equation

$$\|(D_{\mathcal{X}}^2 u)^*\| \le \frac{C(m)}{\lambda} |G(x, u, D_{\mathcal{X}} u, 0)|.$$

A similar estimate holds when u is \mathcal{X} -semiconvex.

Remark 3.4. It is worth noting that one cannot expect the validity of the above a priori estimates when q < Q. An example is the following, cf. Example 3.4 in [8]: consider in the Heisenberg group $\mathbb{H}^d \simeq \mathbb{R}^{2d+1}$ the function $u(x) = \rho(x) - 1$ in B_1 , $\rho(x) = (|x_H|^4 + x_V^2)^{\frac{1}{4}}$, $x_H = (x_1, \dots, x_{2d})$ and $x_V = x_{2d+1}$. Then u is \mathcal{X} -convex for $|x_H| \neq 0$ by Theorem 6.6 of [16]

Heisenberg group $\mathbb{H}^d \simeq \mathbb{R}^{2d+1}$ the function $u(x) = \rho(x) - 1$ in B_1 , $\rho(x) = (|x_H|^4 + x_V^2)^{\frac{1}{4}}$, $x_H = (x_1, ..., x_{2d})$ and $x_V = x_{2d+1}$. Then, u is \mathcal{X} -convex for $|x_H| \neq 0$ by Theorem 6.6 of [16] and solves $\Delta_{\mathcal{X}} u = \frac{Q-1}{\rho} |D_{\mathcal{X}}\rho|^2 =: f(x)$. Then, $f \in L^q(B_1)$ if and only if q < Q, but u is not differentiable at points where $(x_H, x_V) = (0, 0)$ since for i = 1, ..., d we have

$$X_i \rho = x_i \frac{|D_{\mathcal{X}}\rho|^2}{\rho} + \frac{x_{i+d}x_{2d+1}}{\rho^3}, \quad X_{i+d}\rho = x_{i+d} \frac{|D_{\mathcal{X}}\rho|^2}{\rho} - \frac{x_i x_{2d+1}}{\rho^3}.$$

3.2. Convexity and semiconvexity estimates for fully nonlinear equations. In the theory of elliptic and parabolic equations, geometric one-side second order estimates play a crucial role. To our knowledge there are three methods allowing to prove convexity and semiconvexity properties of solutions for elliptic and parabolic equations in the Euclidean setting. The first one dates back to [25], where it was developed the so-called concavity maximum principle, later adapted to the weak setting of viscosity solutions [22]. Another method, based again on the comparison principle, was introduced in [4] and it is based on the fact that the convex envelope of a viscosity supersolution remains a supersolution to the starting equation. The more recent one, recently introduced in [18] and based on integral methods, allows to prove certain concavity-type preserving estimates for nonlinear viscous problems under appropriate assumptions on the nonlinearity. This also applies to fully nonlinear diffusions under appropriate regularity assumptions on the nonlinear terms. The extensions of these properties to a degenerate setting such as those of Carnot groups is by no means immediate. The work [27] shows by tripling the number of variables à la Ishii-Lions a right-invariant convexity property for evolution equations in nondivergence form. The more recent paper [28] studies the (left) horizontal convexity preserving property and proved the

a right-invariant convexity property for evolution equations in nondivergence form. The more recent paper [28] studies the (left) horizontal convexity preserving property and proved the following result for fully nonlinear subelliptic equations posed on the first Heisenberg group in \mathbb{R}^3 : if G(x, r, p, X) is proper, concave in all the variables and symmetric with respect to (x, p)(see assumptions (A3) and (A4) in [28]), continuous solutions to fully nonlinear subelliptic equations on the Heisenberg group are horizontally convex provided that the comparison principle holds for coercive solutions, cf. [28, Theorem 5.7]. By Lemma 4.1 in [5] we would conclude that u is \mathcal{X} -convex along the fields of the Heisenberg group. Consequently, we have the following result:

Theorem 3.5. Suppose that G is uniformly subelliptic, proper, concave in all the variables and symmetric with respect to x, p. Then, coercive solutions to $G(x, u, D_{\mathcal{X}}u, (D_{\mathcal{X}}^2u)^*) = 0$ posed on the first Heisenberg group \mathbb{H} satisfy the a priori estimates of Remark 3.3.

We remark that convexity and semiconvexity type properties hold for the more restrictive class of solutions and typically require additional conditions on the nonlinearity (such as concavity), as it happens when the equation is driven by the sole (sub-)Laplacian in the Euclidean setting.

4. Impossibility of $W^{2,q}_{\mathcal{X}}$ estimates for solutions fully nonlinear subelliptic equations when q < Q

In this section we turn to the smaller class of solutions to elliptic equations, without imposing geometric conditions on them. We prove the following

Theorem 4.1. There exists a fully nonlinear subelliptic operator modeled on the horizontal Hessian of the Heisenberg vector fields, i.e. $G(X) : S_m \to \mathbb{R}$, such that $W^{2,q}_{\mathcal{X}}$ a priori estimates for the equation $G((D^2_{\mathcal{X}}u)^*) = f(x)$ fail when q < Q.

To prove the result we premise the following algebraic lemma, referring for its proof to [15].

Lemma 4.2. Let \mathcal{X} be the Heisenberg vector fields in \mathbb{R}^{2d+1} and ρ be the homogeneous norm defined as

(12)
$$\rho(x) = \left(\left(\sum_{i=1}^{2d} x_i^2 \right)^2 + x_{2d+1}^2 \right)^{\frac{1}{4}}.$$

Then, for $|x_H| \neq 0$, being $x_H = (x_1, ..., x_{2d})$,

$$D_{\mathcal{X}}\rho = \frac{\eta}{\rho^3}, \qquad |D_{\mathcal{X}}\rho|^2 = \frac{|x_H|^2}{\rho^2} \le 1,$$

where $\eta \in \mathbb{R}^{2d}$ is defined by

$$\eta_i := x_i |x_H|^2 + x_{i+d} x_{2d+1} , \qquad \eta_{i+d} := x_{i+d} |x_H|^2 - x_i x_{2d+1}.$$

for i = 1, ..., d. In addition

$$D_{\mathcal{X}}^2 \rho)^* = -\frac{3}{\rho} D_{\mathcal{X}} \rho \otimes D_{\mathcal{X}} \rho + \frac{1}{\rho} |D_{\mathcal{X}} \rho|^2 I_{2d} + \frac{2}{\rho^3} \begin{pmatrix} B & C \\ -C & B \end{pmatrix} ,$$

where the matrices $B = (b_{ij}) \in \mathbb{R}^{d \times d}$ and $C = (c_{ij}) \in \mathbb{R}^{d \times d}$ are defined as follows

$$b_{ij} := x_i x_j + x_{d+i} x_{d+j}$$
, $c_{ij} := x_i x_{d+j} - x_j x_{d+i}$

for i, j = 1, ..., d and satisfy $B = B^T$ and $C^T = -C$. Finally, for a radial function $\psi = \psi(\rho)$ we have

$$(D_{\mathcal{X}}^{2}\psi)^{*}(\rho) = \frac{\psi'(\rho)|D_{\mathcal{X}}\rho|^{2}}{\rho}I_{2d} + 2\frac{\psi'(\rho)}{\rho^{3}}\begin{pmatrix} B & C\\ -C & B \end{pmatrix} + \left(\psi''(\rho) - 3\frac{\psi'(\rho)}{\rho}\right)D_{\mathcal{X}}\rho \otimes D_{\mathcal{X}}\rho ,$$

and its eigenvalues are $\psi''(\rho)|D_{\mathcal{X}}\rho|^2$, $3\psi'(\rho)\frac{|D_{\mathcal{X}}\rho|^2}{\rho}$, which are simple, and $\psi'(\rho)\frac{|D_{\mathcal{X}}\rho|^2}{\rho}$ with multiplicity 2d-2.

Proof of Theorem 4.1. This is inspired from the work by L. Caffarelli [13]. Consider, for $q < Q, \varepsilon > 0, 0 < \alpha < 1$, the family of functions $u_{\varepsilon,\alpha}(x) = \psi(\rho(x))$ defined as

$$u_{\varepsilon,\alpha}(x) = \begin{cases} 1 - (\rho(x))^{\alpha} & \text{for } \rho > \varepsilon\\ 1 - \alpha \varepsilon^{\alpha - 2} (\rho(x))^2 - (1 - \alpha) \varepsilon^{\alpha} & \text{for } \rho \le \varepsilon, \end{cases}$$

 ρ being the homogeneous norm of the Heisenberg group defined in (12). Then, we have

$$\psi_{\rho\rho} = \begin{cases} \alpha(1-\alpha)\rho^{\alpha-2} & \text{for } \rho > \varepsilon \\ -2\alpha\varepsilon^{\alpha-2} & \text{for } \rho < \varepsilon. \end{cases}$$

and

$$\frac{1}{\rho}\psi_{\rho} = \begin{cases} -\alpha\rho^{\alpha-2} & \text{for } \rho > \varepsilon\\ -2\alpha\varepsilon^{\alpha-2} & \text{for } \rho < \varepsilon. \end{cases}$$

Then, define, for $\alpha < 1$ fixed, the constants $\Lambda = \frac{1}{1-\alpha}$ and $\lambda = \frac{1}{Q-1}$ (note that $\Lambda > \lambda$) and consider the fully nonlinear operator $\mathcal{M}^+_{\lambda,\Lambda}(X)$. Note also that it is a convex degenerate

operator. We apply it to the symmetrized horizontal Hessian of the family $u_{\varepsilon,\alpha}$, i.e. $X = (D_{\mathcal{X}}^2 u_{\varepsilon,\alpha})^*$. Then, by Lemma 4.2, the eigenvalues of $(D_{\mathcal{X}}^2 u_{\varepsilon,\alpha})^*$ (where $u_{\varepsilon,\alpha}$ is seen as a radial function of ρ) are $|D_{\mathcal{X}}\rho|^2 \psi_{\rho\rho}$, $3|D_{\mathcal{X}}\rho|^2 \frac{1}{\rho}\psi_{\rho}$ which are simple, and $|D_{\mathcal{X}}\rho|^2 \frac{1}{\rho}\psi_{\rho}$ with multiplicity 2d-2. Therefore, being B_{ε} the homogeneous metric ball of radius ε , we have for $\rho < \varepsilon$ and $|x_H| \neq 0$

$$\mathcal{M}^+_{\lambda,\Lambda}((D^2_{\mathcal{X}}u_{\varepsilon,\alpha})^*) = \lambda |D_{\mathcal{X}}\rho|^2 (-2\alpha\varepsilon^{\alpha-2} - 2\alpha\varepsilon^{\alpha-2}(Q-1)) = -C_1(\alpha,Q)|D_{\mathcal{X}}\rho|^2\varepsilon^{\alpha-2}\chi_{B_{\varepsilon}} =: f_{\varepsilon,\alpha}(x),$$

where $C_1(\alpha, Q) = \frac{2\alpha Q}{Q-1}$ and $\chi_{B_{\varepsilon}}$ is the indicator function of B_{ε} . Instead, when $\rho > \varepsilon$ we conclude

$$\mathcal{M}^+_{\lambda,\Lambda}((D^2_{\mathcal{X}}u_{\varepsilon,\alpha})^*) = \Lambda |D_{\mathcal{X}}\rho|^2 \alpha (1-\alpha)\rho^{\alpha-2} - \lambda(Q-1)\alpha |D_{\mathcal{X}}\rho|^2 \rho^{\alpha-2} = 0.$$

When $|x_H| = 0$ we have $\mathcal{M}^+_{\lambda,\Lambda}((D^2_{\mathcal{X}}u_{\varepsilon,\alpha})^*) = 0$. Therefore, using that $|D_{\mathcal{X}}\rho| \leq 1$ and applying Proposition 5.4.4 of [7] (denoting by $\omega_q = |B_1|$), it follows that for $\alpha < 1$ fixed

$$\begin{split} \|f_{\varepsilon,\alpha}\|_{L^q}^q &= \int_{B_{\varepsilon}} C_1^q(\alpha, Q) |D_{\mathcal{X}}\rho|^{2q} \varepsilon^{(\alpha-2)q} \, dx \\ &\leq C_1^q(\alpha, Q) \varepsilon^{(\alpha-2)q} \int_{B_{\varepsilon}} dx = C_1^q(\alpha, Q) \varepsilon^{(\alpha-2)q} Q \omega_q \int_0^{\varepsilon} s^{Q-1} \, ds \\ &= C_2(\alpha, q, Q) \varepsilon^{(\alpha-2)q+Q} \to 0 \text{ as } \varepsilon \to 0 \end{split}$$

for any

$$q < \frac{Q}{2-\alpha}.$$

Therefore, for $|x_H| \neq 0$, $||f_{\varepsilon,\alpha}||_q \leq \overline{K}$ remains bounded for all $\varepsilon \in (0,1]$, whereas for such value of q the norm $||u_{\varepsilon,\alpha}||_{W^{2,q}_{\mathcal{X}}(\mathbb{R}^n)}$ blows-up because of the definition of $\psi_{\rho\rho}$ for $\rho > \varepsilon$, showing that second order estimates in L^q cannot be achieved in the range q < Q.

Remark 4.3. The same counterexample can be built on H-type groups using the computations of $(D_{\chi}^2 \rho)^*$ carried out in [31].

References

- F. Abedin, C. E. Gutiérrez, and G. Tralli. Harnack's inequality for a class of non-divergent equations in the Heisenberg group. Comm. Partial Differential Equations, 42(10):1644–1658, 2017.
- [2] F. Abedin and G. Tralli. Harnack inequality for a class of Kolmogorov-Fokker-Planck equations in nondivergence form. Arch. Ration. Mech. Anal., 233(2):867–900, 2019.
- [3] A. D. Aleksandrov. Impossibility of general estimates of solutions and of uniqueness conditions for linear equations with norms weaker than those of L_n . Vestnik Leningrad. Univ., 21(13):5–10, 1966.
- [4] O. Alvarez, J.-M. Lasry, and P.-L. Lions. Convex viscosity solutions and state constraints. J. Math. Pures Appl. (9), 76(3):265–288, 1997.
- [5] M. Bardi and F. Dragoni. Convexity and semiconvexity along vector fields. Calc. Var. Partial Differential Equations, 42(3-4):405-427, 2011.
- [6] S. Biagi, A. Bonfiglioli, and M. Bramanti. Global estimates in Sobolev spaces for homogeneous Hörmander sums of squares. J. Math. Anal. Appl., 498(1):Paper No. 124935, 19, 2021.
- [7] A. Bonfiglioli, E. Lanconelli, and F. Uguzzoni. Stratified Lie groups and potential theory for their sub-Laplacians. Springer Monographs in Mathematics. Springer, Berlin, 2007.
- [8] J. E. M. Braga, A. Figalli, and D. Moreira. Optimal regularity for the convex envelope and semiconvex functions related to supersolutions of fully nonlinear elliptic equations. *Comm. Math. Phys.*, 367(1):1–32, 2019.
- [9] J. E. M. Braga and D. Moreira. Inhomogeneous Hopf-Oleĭnik lemma and regularity of semiconvex supersolutions via new barriers for the Pucci extremal operators. Adv. Math., 334:184–242, 2018.

- [10] X. Cabré. Nondivergent elliptic equations on manifolds with nonnegative curvature. Comm. Pure Appl. Math., 50(7):623–665, 1997.
- [11] X. Cabré and L. A. Caffarelli. Interior $C^{2,\alpha}$ regularity theory for a class of nonconvex fully nonlinear elliptic equations. J. Math. Pures Appl. (9), 82(5):573-612, 2003.
- [12] L. Caffarelli, J. J. Kohn, L. Nirenberg, and J. Spruck. The Dirichlet problem for nonlinear second-order elliptic equations. II. Complex Monge-Ampère, and uniformly elliptic, equations. *Comm. Pure Appl. Math.*, 38(2):209–252, 1985.
- [13] L. A. Caffarelli. Interior a priori estimates for solutions of fully nonlinear equations. Ann. of Math. (2), 130(1):189–213, 1989.
- [14] L. A. Caffarelli and X. Cabré. Fully nonlinear elliptic equations, volume 43 of American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, RI, 1995.
- [15] A. Cutrì and N. Tchou. Barrier functions for Pucci-Heisenberg operators and applications. Int. J. Dyn. Syst. Differ. Equ., 1(2):117–131, 2007.
- [16] D. Danielli, N. Garofalo, and D.-M. Nhieu. Notions of convexity in Carnot groups. Comm. Anal. Geom., 11(2):263–341, 2003.
- [17] D. Danielli, N. Garofalo, and D.-M. Nhieu. On the best possible character of the L^Q norm in some a priori estimates for non-divergence form equations in Carnot groups. Proc. Amer. Math. Soc., 131(11):3487–3498, 2003.
- [18] L. C. Evans. Adjoint and compensated compactness methods for Hamilton-Jacobi PDE. Arch. Ration. Mech. Anal., 197(3):1053–1088, 2010.
- [19] X. Fernández-Real and X. Ros-Oton. Regularity theory for elliptic PDE. Zurich Lectures Notes in Advanced Mathematics. European Mathematical Society, 2022.
- [20] G. B. Folland. Subelliptic estimates and function spaces on nilpotent Lie groups. Ark. Mat., 13(2):161–207, 1975.
- [21] N. Garofalo. A note on monotonicity and Bochner formulas in Carnot groups. arXiv:2204.13073, to appear in Proc. Roy. Soc. Edinburgh Sect. A, 2022.
- [22] Y. Giga, S. Goto, H. Ishii, and M.-H. Sato. Comparison principle and convexity preserving properties for singular degenerate parabolic equations on unbounded domains. *Indiana Univ. Math. J.*, 40(2):443–470, 1991.
- [23] C. E. Gutiérrez and F. Tournier. Harnack inequality for a degenerate elliptic equation. Comm. Partial Differential Equations, 36(12):2103–2116, 2011.
- [24] C. Imbert. Convexity of solutions and C^{1,1} estimates for fully nonlinear elliptic equations. J. Math. Pures Appl. (9), 85(6):791–807, 2006.
- [25] N. J. Korevaar. Convex solutions to nonlinear elliptic and parabolic boundary value problems. Indiana Univ. Math. J., 32:603–614, 1983.
- [26] P.-L. Lions and M. Musiela. Convexity of solutions of parabolic equations. C. R. Math. Acad. Sci. Paris, 342(12):915–921, 2006.
- [27] Q. Liu, J. J. Manfredi, and X. Zhou. Lipschitz continuity and convexity preserving for solutions of semilinear evolution equations in the Heisenberg group. *Calc. Var. Partial Differential Equations*, 55(4):Art. 80, 25, 2016.
- [28] Q. Liu and X. Zhou. Horizontal convex envelope in the Heisenberg group and applications to sub-elliptic equations. Ann. Sc. Norm. Super. Pisa, Cl. Sci. (5), 22(4):2039–2076, 2021.
- [29] A. Montanari. Harnack inequality for a subelliptic PDE in nondivergence form. Nonlinear Anal., 109:285– 300, 2014.
- [30] C. Pucci. Operatori ellittici estremanti. Ann. Mat. Pura Appl. (4), 72:141-170, 1966.
- [31] G. Tralli. A certain critical density property for invariant Harnack inequalities in H-type groups. J. Differential Equations, 256(2):461–474, 2014.
- [32] N. S. Trudinger. Fully nonlinear, uniformly elliptic equations under natural structure conditions. Trans. Amer. Math. Soc., 278(2):751–769, 1983.

DIPARTIMENTO DI MATEMATICA "TULLIO LEVI-CIVITA", UNIVERSITÀ DEGLI STUDI DI PADOVA, VIA TRI-ESTE 63, 35121 PADOVA (ITALY)

Email address: alessandro.goffi@unipd.it