

Key Points:

- Garnet U–Pb, mica Rb–Sr dating constrain exhumation of Rotondo granite from amphibolite facies at 34–20 Ma to greenschist facies at 15–14 Ma
- Fast exhumation (~3 mm/yr) accommodated by weak ductile shear zones localized on pre-collisional brittle deformation structures
- The European crust was relatively weak during collision, especially where rheology was controlled by localized shear zones

Supporting Information:

Supporting Information may be found in the online version of this article.

Correspondence to:

A. Ceccato,
aceccato@erdw.ethz.ch

Citation:

Ceccato, A., Behr, W. M., Zappone, A. S., Tavazzani, L., & Giuliani, A. (2024). Structural evolution, exhumation rates, and rheology of the European crust during Alpine collision: Constraints from the Rotondo granite—Gotthard nappe. *Tectonics*, 43, e2023TC008219. <https://doi.org/10.1029/2023TC008219>

Received 14 DEC 2023

Accepted 6 JUN 2024

Author Contributions:

Conceptualization: A. Ceccato, W. M. Behr
Data curation: A. Ceccato, L. Tavazzani, A. Giuliani
Formal analysis: A. Ceccato, L. Tavazzani, A. Giuliani
Funding acquisition: W. M. Behr
Investigation: A. Ceccato, W. M. Behr, A. S. Zappone
Project administration: W. M. Behr, A. S. Zappone
Supervision: W. M. Behr
Validation: A. Ceccato, W. M. Behr, A. S. Zappone, L. Tavazzani, A. Giuliani

© 2024 The Authors.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial License](#), which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

Structural Evolution, Exhumation Rates, and Rheology of the European Crust During Alpine Collision: Constraints From the Rotondo Granite—Gotthard Nappe

A. Ceccato¹ , W. M. Behr¹ , A. S. Zappone² , L. Tavazzani³ , and A. Giuliani³ 

¹Department of Earth Sciences, Structural Geology and Tectonics Group, Geological Institute, ETH Zurich, Zurich, Switzerland, ²Swiss Seismological Service, ETHZ, Zurich, Switzerland, ³Department of Earth Sciences, Institute of Geochemistry and Petrology, ETH Zurich, Zurich, Switzerland

Abstract The rheology of crystalline units controls the large-scale deformation geometry and dynamics of collisional orogens. Defining a time-constrained rheological evolution of such units may help unravel the details of collisional dynamics. Here, we integrate field analysis, pseudosection calculations and in situ garnet U–Pb and mica Rb–Sr geochronology to define the structural and rheological evolution of the Rotondo granite (Gotthard nappe, Central Alps). We identify a sequence of four (D₁–D₄) deformation stages. Pre-collisional D₁ brittle faults developed before Alpine peak metamorphism, which occurred at 34–20 Ma (U–Pb garnet ages) at 590 ± 25°C and 0.9 ± 0.1 GPa. The reactivation of D₁ structures controlled the rheological evolution, from D₂ reverse mylonitic shearing at amphibolite facies (520 ± 40°C and 0.8 ± 0.1 GPa) at 18–20 Ma (white mica Rb–Sr ages), to strike-slip, brittle-ductile shearing at greenschist-facies D₃ (395 ± 25°C and 0.4 ± 0.1 GPa) at 14–15 Ma (white mica and biotite Rb–Sr ages), and then to D₄ strike-slip faulting at shallow conditions. Although highly misoriented for the Alpine collisional stress orientation, D₁ brittle structures controlled the localization of D₂ ductile mylonites accommodating fast (~3 mm/yr) exhumation rates due to their weak shear strength (<10 MPa). This structural and rheological evolution is common across External Crystalline Massifs (e.g., Aar, Mont Blanc), suggesting that the European upper crust was extremely weak during Alpine collision, its strength controlled by weak ductile shear zones localized on pre-collisional deformation structures, that in turn controlled localized exhumation at the scale of the orogen.

1. Introduction

During mountain-building events, rheological contrasts between lithospheric plates are first-order controls on the geometry of collision (Candioti et al., 2021; Faccenda et al., 2008; Vogt et al., 2018), the development of topography (Cook & Royden, 2008; Wolf et al., 2022), and the styles and rates of regional deformation and metamorphism (Piccolo et al., 2018; Willingshofer et al., 2005). Rheological contrasts may result from different crustal compositions, ages, geological histories, and/or thermal regimes of the lithospheric plates involved in collision (Audet & Bürgmann, 2011; Mouthereau et al., 2013). For example, depending on the composition and the fluid content, lithospheric plates may present different mechanical behavior (brittle vs. viscous deformation) and strength at the same depth and temperature conditions during collision (Behr & Platt, 2014; Bürgmann & Dresen, 2008; Jamtveit et al., 2019; Menegon et al., 2011). Furthermore, the occurrence of anisotropic structural fabrics (foliations and fractures), strictly related to the geological history of crustal sections, may promote or hinder deformation depending on their suitability to be reactivated, and/or their ability to promote or hinder fluid infiltration (Ceccato, Menegon, et al., 2020; Zertani et al., 2023). Pressure, temperature, fluid, and structural fabrics evolve with the tectono-metamorphic evolution of a collisional orogen, and so do their effects on the rheological contrast between colliding plates (Behr & Platt, 2013; Bellanger et al., 2014; Ceccato, Menegon, et al., 2020; Groome et al., 2008).

Most of the deformation and shortening in the core of collisional belts is accommodated through deformation of crystalline basement units (Lacombe & Mouthereau, 2002; Pfiffner, 2016; Rosenberg & Kissling, 2013). Pre-collisional events such as lithospheric rifting, prograde burial, and subduction, lead to the development of additional deformation structures (e.g., rift-related normal fault zones) and tectonic fabrics (e.g., prograde foliations), which may introduce rheological heterogeneity that later influences collisional dynamics (Mohn et al., 2014). In the European Alps, both inherited compositional variations, as well as pre-collisional tectonics, are known to have strongly influenced later syn-orogenic development. For example, rheological contrast

Visualization: A. Ceccato, L. Tavazzani, A. Giuliani

Writing – original draft: A. Ceccato

Writing – review & editing: A. Ceccato, W. M. Behr, A. S. Zappone, L. Tavazzani, A. Giuliani

between the upper (Adriatic) and lower (European) crust varies along the strike of the orogen, and resulted in different patterns of strain partitioning, amounts of shortening and exhumation, and collisional styles between the Western, Central and Eastern Alps (Bellahsen et al., 2014; Rosenberg & Kissling, 2013). In the Central Alps, in particular, the Adriatic upper plate, represented by the almost undeformed Ivrea-Verbano complex, indents into the weaker European crust (Rosenberg & Kissling, 2013). The European crust is composed of stacked slices of crystalline basement derived from the thinned Mesozoic European margin, now exposed in the Aar massif, Gotthard nappe, and Lepontine dome (Figure 1). However, constraints on the factors controlling the “weakness” of the European crust are sparse, including whether the crust was weak since the beginning of burial and subduction, or if it was initially strong and then progressively weakened during collision. Both tectonic inheritance related to Mesozoic rifting (Bellahsen et al., 2014), as well as Variscan orogeny (Schwartz et al., 2024), and syn-collisional Barrovian metamorphism (Rosenberg & Kissling, 2013) might have contributed to the weakening of the European continental crust in this part of the Alps.

Previous investigations have revealed a recurrent brittle-to-ductile structural evolution (i.e., ductile shear zone related to collisional processes overprinting pre-existent brittle faults and fractures) of crystalline basement units in the Western and Central Alps (e.g., Mont Blanc: Guermani & Pennacchioni, 1998; Gran Paradiso: Menegon & Pennacchioni, 2010; Aar-Gotthard: Oliot et al., 2014; Rolland et al., 2009; Wehrens et al., 2016; Lepontine Dome: Goncalves et al., 2016). Several hypotheses were proposed to explain such brittle-to-ductile evolution, including the occurrence of prograde brittle deformation during burial (Guermani & Pennacchioni, 1998), and mid-crustal seismicity at peak metamorphic conditions (Wehrens et al., 2016). Previous authors have also speculated on the occurrence of pervasive extensional faulting related to the Mesozoic rifting of the European margin, providing field evidence for limited reactivation of structures inherited from rifting (Ballèvre et al., 2018; Dall’Asta et al., 2022; Herwegh et al., 2017, 2020; Musso Piantelli et al., 2022; Nibourel et al., 2021).

Here we present an integrated field and petrochronological study of the deformation features of the Rotondo granite (RG) in the Gotthard nappe (Figure 1). The Gotthard nappe represents a sliver of European crust now exposed in the Central Swiss Alps. The RG is a Post-Variscan pluton (i.e., not affected by Variscan tectono-metamorphic events), intruded into the European polymetamorphic crust. Differently from its host polymetamorphic host rock, the lack of Variscan pervasive fabrics (foliations) and the homogeneous texture of the granite allow us to define a sequence of localized deformation structures probably related to Alpine deformation. We use structural and petrochronological data to:

1. Define the pressure-temperature-time-deformation (*P-T-t-d*) path of the RG;
2. Examine the time-constrained structural and rheological evolution of the thickened crust of the lower plate during Alpine continental collision;
3. Investigate the geological factors that affect the rheological evolution of the crystalline unit.

2. Geological Setting

The European Alps (Figure 1) are a double-verging orogen resulting from the continental collision between Europe and Adria, following the closure and subduction of the Mesozoic Tethys ocean (Dal Piaz et al., 2003). European and Adriatic polymetamorphic crustal sections were each strongly modified by the Variscan orogeny during the formation of the Pangean supercontinent. The Permo-Mesozoic breakup of Pangea led to the development of the Tethys Ocean, including the Liguro-Piemontese ocean and Valais trough. The development of the Liguro-Piemontese ocean divided Europe from Adria (210–140 Ma) by 200–400 km (Ballèvre et al., 2018; Beltrando et al., 2014). A second, more short-lived rifting phase took place on the European margin to the north of the Liguro-Piemontese ocean and led to the development of the Valais trough during Late-Jurassic to Early Cretaceous (140–120 Ma), separating the European distal margin from the southern Briançonnais microcontinent (Beltrando et al., 2012; Célini et al., 2023; Handy et al., 2010). This former paleogeography is now preserved in the Internal (Penninic) domains of the Central and Western Alps, exposing the remnants of the Valaisan, Briançonnais and Liguro-Piemontese units (Figure 1). The proximal European passive margin is now exposed in the External Crystalline Massifs (ECMs), including: Aar, Mont Blanc, Aiguille Rouges, Belledonne, Pelvoux-Oisan massifs as well as in the Gotthard nappe (Figure 1a; Lemoine et al., 1986).

At the regional scale, the European crust was affected by the development of Permo-Mesozoic transtensional basins, resulting eventually in the formation of the Valais trough in the Jurassic-Cretaceous period (Ballèvre et al., 2018; Célini et al., 2023; Handy et al., 2010). The units now included in the Gotthard nappe were part of the




Figure 1. Geological setting of the study area. (a) Tectonic sketch of the Central-Western Alps (redrawn from Ballèvre et al., 2018, Schmid et al., 2004; Schwartz et al., 2024). AA: Aar; AG: Argentera; AR: Aiguilles rouges; BD: Belledonne; DM: Dora Maira; GP: Gran Paradiso; GT: Gotthard; IVZ: Ivrea-Verbano Zone; LD: Leptontine Dome; MB: Mont Blanc; MR: Monte Rosa; PE: Pelvoux. (b) Tectonic sketch of the Central-Western Gotthard massif showing the spatial distribution of the meta-granitoid intrusion (Rotondo, Fiabba, Gamsboden). (c) Geological section across the Central Swiss Alps.

distal European passive margin located north of the Valais trough (Schmid et al., 2004). From the Late Cretaceous onwards, convergence between Europe and Adria led to the subduction of the Liguro-Piemontese ocean and to progressive development of the Penninic accretionary wedge facing the advancing Adriatic upper plate (Dal Piaz et al., 2003). Progressive convergence led to burial and underthrusting of the European passive margin, eventually leading to continental collision.

The study area is located in the Gotthard nappe (Swiss Alps, Figure 1a). The Gotthard nappe includes a series of polymetamorphic Ordovician-Silurian crystalline units intruded by late-Variscan granitoids (Berger et al., 2017).

The crystalline units include high-grade gneisses of the Val Nalps, Paradis and Streifegneis complexes (Figure 2a; Berger et al., 2017). The Val Nalps and Paradis Complexes preserve evidence of an Early to Mid-Ordovician (~470 Ma) high grade metamorphism, later affected by Silurian (~440 Ma) magmatism (Berger et al., 2017). Between 340 and 300 Ma, these complexes were affected by Variscan amphibolite facies metamorphism and transpressional shearing (Bühler et al., 2022; Simonetti et al., 2020; Vanardois et al., 2022). Post-Variscan magmatism led to the intrusion of several granitic plutons into the polymetamorphic basement, including the Cristallina granodiorite, the Fibbia and Gamsboden granite-gneisses and the RG (Figure 1b; Berger et al., 2017). The Permo-Mesozoic tectonic evolution of the Gotthard nappe is still poorly constrained. The Gotthard nappe was buried beneath the advancing Penninic accretionary wedge around 35 Ma (Handy et al., 2010), reaching greenschist-facies conditions between 35 and 22 Ma (Herwegh et al., 2020; Janots et al., 2009). Subsequently, continental collision between Europe and Adria led to the rapid exhumation of the crystalline units of the Gotthard-Aar massifs at around 22–17 Ma, through the activation of greenschist facies sub-vertical ductile shear zones with reverse kinematics ($T = 450\text{--}500^\circ\text{C}$ and $P = 0.7\text{--}0.8$ GPa; Challandes et al., 2008; Goncalves et al., 2012; Herwegh et al., 2017; Oliot et al., 2010; Rolland et al., 2008, 2009). From 14 Ma onward, the Gotthard nappe was then affected by regional strike-slip tectonics related to the activity of the Simplon-Rhone transtensional fault system (Campani et al., 2010; Herwegh et al., 2017; Rolland et al., 2009). Shallow brittle faulting has affected the Gotthard nappe since the Late Miocene, leading to the activation of brittle gouge-bearing faults up to recent times (Kralik et al., 1992; Pleuger et al., 2012).

2.1. The Rotondo Granite

The RG is an Early Permian (295 Ma, U–Pb on zircon, Rast et al., 2022) peraluminous granite, crosscut by mafic dykes (290–285 Ma, U–Pb on zircon, Bussien et al., 2008). It includes two main magmatic facies (equigranular RG_1 and porphyritic RG_2) both composed of $\text{Qz} + \text{Kfs} + \text{Pl} + \text{Bt} \pm \text{Wm} \pm \text{Grt} \pm \text{Ep} \pm \text{Chl} \pm \text{Zr} \pm \text{Spn} \pm \text{Cal} \pm \text{Py}$ (Rast et al., 2022, mineral abbreviations from Whitney & Evans, 2010; Wm: white mica). RG_1 and RG_2 facies only differ by mineral proportions and the occurrence of a Bt-Kfs foliation in RG_2 (Rast et al., 2022). This meso-scale bulk foliation has been attributed to an Alpine greenschist facies overprint, based on field and microstructural observations (Gapais et al., 1987; Steck, 1976; Steck & Burri, 1971). Another evidence of Alpine greenschist facies metamorphism is the occurrence of atoll-like garnets in the RG (Steck, 1976; Steck & Burri, 1971). The peculiar atoll-like shape, and their Ca-rich composition, have been interpreted by Steck and Burri (1971) to reflect two metamorphic growth stages at different temperature and/or fluid activity conditions. However, the textural relationship between the atoll-garnets and the bulk foliation was not addressed in detail. A set of steep, NW-dipping ductile shear zones, with top-to-SE dip-slip reverse kinematics developed during the same Alpine retrograde event (Lützenkirchen & Loew, 2011). The ductile shear zones have been classified in two main groups (Rast et al., 2022): (a) granitic shear zones, composed of fine-grained mylonite with feldspar augens in a biotite-bearing foliation; and (b) quartz-biotite-rich shear zones, characterized by the occurrence of sigmoidal quartz veins with rigid cm-sized calcite clasts. Ductile shear zones were exploited as nucleation sites for late brittle faulting at upper crustal levels, as inferred from the stability of syn- to post-kinematic zeolite minerals, and the formation of clay-rich gouges (Lützenkirchen & Loew, 2011). Despite the general understanding of the regional and local scale tectonic evolution, a detailed and holistic description of the structural and tectonometamorphic features, and absolute timing of deformation events in this area are still missing.

3. Material and Methods

3.1. Field Structural Analysis

This work further extends the previous work of Lützenkirchen and Loew (2011) and Rast et al. (2022), improving the detail of structural description, and adding absolute age constraints on the deformation structures, with implications on regional tectonic and rheological evolution. It provides a detailed description of the structural evolution and inventory of the deformation structures affecting the rock massif hosting the Bedretto Underground Laboratory for Geosciences and Geoenergies (BULGG; Ma et al., 2022). Field survey was focused on the analyses of deformation features and the collection of structural data at 205 structural stations (“Waypoints”—WP in Figure 2a), resulting in a georeferenced data set of 473 structural measurements, each of which includes a structural description, orientation of shear plane (Dip/Dip direction) and lineation (Trend/Plunge), kinematics, mineralogy, deformation fabric (brittle vs. ductile), thickness, length, and throw. These structures were then subdivided into sets based on kinematic compatibility, mineralogy, and texture. Oriented samples were collected




Figure 2.

for further microstructural and petrochronological analysis. The geographic coordinates of relevant waypoints are reported in Table S1 in Supporting Information S2. The Structural data set is available in Data set S1.

3.2. Optical/Scanning Electron Microscopy and Electron Probe Micro Analyses

Thin sections were cut parallel to the lineation direction (X kinematic direction) and perpendicular to the foliation plane (XY kinematic plane). Backscattered electron images and Energy Dispersion Spectrometry mapping were performed at ScopeM (ETH) with a Hitachi SU5000 Scanning Electron Microscope (SEM). Quantitative compositional analyses were performed at the Institute for Geochemistry and Petrology (ETH) with a JEOL JXA-8230 Electron Probe Microanalyzer equipped with five Wavelength Dispersion Spectrometers. Further details on analytical conditions are reported in Text S1 in Supporting Information S1. Mineral compositions are reported in Table S2 in Supporting Information S2.

3.3. P-T Pseudosection Calculation

The bulk rock compositions adopted for pseudosection calculation were obtained by X-Ray Fluorescence spectroscopy at the Institute for Geochemistry and Petrology (ETH) with a WD-XRF PANalytical AXIOS equipped with five diffraction crystals (bulk compositions are reported in Table S3 in Supporting Information S2). Pressure-temperature pseudosection calculations were performed with Perple_X 6.9.1 (Connolly, 2005) adopting the thermodynamic database of pure end-members from Holland and Powell (2011; hp62ver.dat). Adopted solid solution models and computational details are reported in Text S1 in Supporting Information S1. The chemical system used for the calculation is $\text{MnO}-\text{Na}_2\text{O}-\text{CaO}-\text{K}_2\text{O}-\text{FeO}-\text{MgO}-\text{Al}_2\text{O}_3-\text{SiO}_2-\text{H}_2\text{O}-\text{TiO}_2-\text{Fe}_2\text{O}_3$ (MNCKFMASHTO). In the text, the term “observed” refers to the paragenesis observed in thin section and to the phase chemistry obtained from EPMA analyses; the term “computed” refers to the chemistry and mineral paragenesis calculated by pseudosection computations. Results of pseudosections and related files are available in Data set S2.

3.4. In Situ LA-ICP-MS U–Pb and Trace Element Analyses

In situ Garnet U–Pb dating, and trace element analyses were performed on polished thin sections by laser ablation-inductively coupled plasma-mass spectrometry (LA-ICP-MS) at the ERDW department of ETH Zurich using an ASI RESolution S-155 excimer (ArF, 193 nm) laser ablation system coupled to a Thermo Scientific Element XR sector-field ICP-MS (Guillong et al., 2014). Instrumentation and data acquisition parameters for U–Pb dating are summarized in Data set S3 reporting standards of Horstwood et al. (2016). All data from the session, including details on the data reduction strategies and results of validation reference materials can be found in Data set S3.

3.5. In Situ LA-ICP-MS Rb–Sr Analyses

In situ Rb–Sr isotope analyses of mica in thin section were undertaken using an ASI RESolution 193 nm excimer laser probe interfaced to an Agilent 8800 ICP-MS/MS at ETH Zurich following the procedure outlined in Giuliani et al. (2024). This method employs an isochronous in-house mica reference material from the Wimbledon lamproite (Sarkar et al., 2023) to calibrate the Rb/Sr fractionation in mica unknowns following initial calibration of $^{87}\text{Sr}/^{86}\text{Sr}$ and $^{87}\text{Rb}/^{86}\text{Sr}$ using the silicate glass reference material NIST 610. This method is validated by analyses of micas from the Bultfontein kimberlite with ages independently constrained by isotope-dilution Rb–Sr dating (Fitzpayne et al., 2020). All the details pertaining analytical conditions, reference materials and data processing can be found in Text S1 in Supporting Information S1. The Rb–Sr age data are summarized in Table 1 and all the Rb–Sr analyses can be found in Data set S4.

Figure 2. Structural map and data of the surveyed area in the Rotondo Granite (RG). (a) Structural field map of the southern rim of the RG summarizing the field observations and showing the location of investigated areas (modified after Berger et al., 2017). (b)–(i) Equal area, lower-hemisphere stereographic projections of the structural data for each set of deformation structures. Great circles: slip planes (S); Dots and contour: lineations (L). Blue and red planes and dots represent dextral and sinistral kinematics, respectively. Contours are calculated as Area percentage, minimum contour is 5 area%—computed with Stereonet 11 (<https://www.rickallmendinger.net/stereonet>). (b) D_1 shear fractures, cataclasites and breccias; (c) D_1 plane-parallel thick quartz veins; (d) D_2 ductile shear zones, dip-slip, top-to-SE reverse kinematics; (e) low-angle quartz veins kinematically related to D_2 shear zones; (f) D_{3A} ductile shear zones (black great circles) showing strike-slip reactivation and associated extensional wing cracks and quartz-veins developed in dilational jogs (orange great circles); (g) D_{3B} conjugate, brittle-ductile shear zones (solid great circles) and extensional veins (dashed great circles); (h) D_{3C} normal faults; (i) D_4 Zeolite- and gouge-bearing brittle fault zones.

Table 1
Summary of the Results From U–Pb in Garnet and Rb–Sr in Mica Analyses

Sample	Description	Age (Ma)	Uncert. Int. ($\pm 2\sigma$, Ma)	Uncert. with ssys ($\pm 2\sigma$, Ma) ^a	Relative uncertainty ($\pm 2\sigma$, %)	Number of analyses	MSWD	p (χ^2)
ACB18	U–Pb Grt	23.23	3.58	3.60	15.48%	45/55	3.40	0.0
ACB35	U–Pb Grt, Zr > 25 ppm	127.97	9.08	9.28	7.25%	26/36	2.90	0.0
	U–Pb Grt, Zr < 25 ppm	34.04	4.37	4.40	12.93%	25/35	2.80	0.0
ACB_Sp3b	U–Pb Grt	26.92	1.25	1.31	4.88%	54/59	1.90	0.0
ACB_Sp7b	U–Pb Grt	30.51	6.79	6.81	22.31%	57/58	0.47	1.0
ACB_Sp8c	U–Pb Grt	24.04	2.07	2.10	8.74%	68/71	0.92	0.7
ACB_Sp8d	U–Pb Grt	20.09	1.00	1.04	5.20%	33/33	0.86	0.7
B19-1417	Bt, Undeformed RG1	15.15	–	0.71	4.69%	27/29	0.34	1.0
ACB3b	Bt, D2 in RG1	15.25	–	0.88	5.77%	28/30	0.65	0.9
ACB12b	Wm, Bulk RG2 fol	18.59	–	1.87	10.06%	35/37	0.76	0.8
ACB14c	Bt, Bulk RG2 fol	15.01	–	1.17	7.79%	30/30	0.42	1.0
ACB27a	Wm, D2	18.14	–	0.8	4.41%	38/39	1.50	0.0
ACB37b	Wm, D3	14.68	–	1.55	10.56%	35/36	1.30	0.1
ACB_Sp6	Bt, Bulk RG2 fol	14.77	–	0.74	5.01%	30/30	0.65	0.9

^aSystematic, long-term excess variance used for propagation is 1.5%. *Note.* Samples in bold are discussed in the main text. Ages reported for U–Pb in garnet are lower intercept ages on the concordia curve.

4. Results

4.1. Field Observations—Sequence of Localized Deformation Structures

In the following, we describe the sequence of subsolidus deformation structures, numbered following their relative chronology (from the oldest D_1 , to the youngest D_4), as inferred from field analyses of crosscutting and overprinting relationships, mineralogy, texture, and kinematics. Structural data are summarized in Figures 2b–2i. Field images of the described structure sets are reported in Figures 3–5. In Text S2 in Supporting Information S1, additional data about the magmatic structures (aplitic and mafic dykes), and the tectonometamorphic evolution of the RG-host rocks are presented.

4.1.1. D_1 Brittle Shear Fractures, Cataclasites and Breccias

The D_1 set consists of brittle shear fractures (Figure 3a), cataclasites (Figure 3b) and breccias (Figures 3c and 3d) containing a dark, fine-grained matrix that surrounds angular clasts of undeformed granite (Figures 3b–3e). Milky quartz veins are common along these structures, ranging in thickness from a few mm to >1 m, and showing mutual overprinting relationship with the dark fine-grained matrix (Figure 3f). In some cases, large breccia bodies are observed, characterized by a transitional texture from crackle-breccias to fine grained cataclasites. A peculiar feature of D_1 structures is the occurrence of mm-size garnets overgrowing the dark matrix (Figure 3e). D_1 structures are steeply dipping, SE-verging, and ENE–WSW-striking (Figures 2b and 2c). A subset of D_1 cataclasites (23 planar measurements) presents an orientation at high angle to the main set (Figure 2b). The kinematics of set D_1 structures is rather difficult to constrain, given that they are overprinted by the following stage of ductile deformation. A dip-slip lineation L_1 is observed on the exposed surface of the matrix (Figure 3g), and incipient breccias and shear fractures commonly show either strike-slip dextral or normal-sense displacement of crosscut markers in the present orientation (Figure 3h). Garnet is only observed to overgrow the breccia matrix in D_1 structures not heavily overprinted by the ductile deformation related to D_2 shear zones (Figures 3b–3e), as described below.

4.1.2. D_2 Dip-Slip, Reverse Ductile Shear Zones

The D_2 set consists of mylonitic ductile shear zones. D_2 shear zones exploit as nucleation site the pre-existing structural and/or compositional heterogeneities in the host RG, such as aplitic and mafic dykes, veins, and D_1




Figure 3.

structures (Figure 4). Deformed aplitic and mafic dykes develop an oblique homogeneous foliation abruptly terminating at the dyke selvage against the undeformed host RG. D_2 shear zones exploiting D_1 brittle structures preserve the geometric and textural complexity of the precursor, developing an heterogeneous Bt-Wm-bearing foliation wrapping around low-strain granite clasts and lithons (Figure 4b). D_2 structures strike ENE–WSW, showing a dip-slip, L_2 Bt-Wm-bearing lineation (Figure 2d). The dominant kinematics is reverse, top-to-SE, even though dip-slip normal kinematics are observed on rare SE-steeply-dipping shear planes. Overall, the D_2 ductile shear zones form a large-scale network defined by subparallel zones of high strain surrounding high-aspect-ratio lozenges of undeformed granite (Gapais et al., 1987). Sheared quartz + calcite veins are common in high strain D_2 ductile shear zones (Figure 4a; see also Figure 11 of Rast et al., 2022). Locally, calcite-rich shear zones are observed, showing thick homogeneous calcite-rich layers close to layers clearly resembling sheared calcite-bearing breccias (Figure 4d). A set of shallowly NW-dipping quartz + feldspar veins and non-mineralized joints also occur (Figures 2e and 4b), and commonly abut major mylonitic shear zones, with vein tips dragged into the main mylonitic foliation (Figure 4b).

4.1.3. D_3 Strike-Slip, Brittle-Ductile Shear Zones

D_3 structures include: (a) D_{3A} , strike-slip mylonitic shear zones reactivating pre-existent D_1 – D_2 structures; (b) D_{3B} , a set of conjugate, brittle-ductile strike-slip faults; (c) D_{3C} , a set of conjugate, normal dip-slip faults. D_1 – D_2 structures are reactivated under brittle-ductile conditions with a dominant strike-slip kinematics (Figures 4e–4h), as inferred from the development of a sub-horizontal L_{3A} lineation overprinting the pre-existing dip-slip L_2 (Figure 4e). The orientation of D_{3A} structures reflects the pre-existent ENE–WSW mylonitic shear zones (Figure 2f). A heterogeneous S-C fabric is developed in major mylonitic zones, showing dominant dextral kinematics (Figures 4e and 4f, same outcrop of the shear zone a few meters west from Figure 4a). Tensile wing cracks (Figures 2f and 4g), brecciated dilational jogs (Figure 4h), and quartz + feldspar veins develop at high angle to D_1 – D_2 shear fractures and mylonites during re-shearing under strike-slip orientations. These veins and dilational breccias can be distinguished from previous D_1 quartz-veins and breccias based on their orientation and their content of coarse quartz clear crystals.

The D_{3B} structures consist of brittle-ductile discrete faults and mylonites, arranged in conjugate sets (Figure 5a). The N–S–striking set of sinistral, strike-slip fault planes is conjugated to a WNW–ESE-striking set of dextral strike-slip fault planes (Figure 2g). In both cases, the L_{3B} lineation is oblique (Figure 2g), shallowly plunging toward ENE or WNW, respectively. The conjugate fault sets crop out in low-strain domains bounded by major D_{3A} shear zones. A very localized mylonitic foliation is observed along the discrete fault planes. The conjugate set of brittle-ductile faults is associated with subvertical, NW–SE-striking tensional $Qz + Chl + Wm + Py + Hem$ veins (Figure 5a). Such mineralization and veins are observed also in extensional jogs between overlapping en-echelon fault segments (Figure 5a). Episyenites (i.e., quartz-depleted, vuggy altered granites) are observed close to mineralized veins and shearing planes (Pennacchioni et al., 2016; Rossi & Rolland, 2014).

D_{3C} structures consist of NW–SE striking faults with a dip-slip, $Qz + Wm$ -bearing L_{3C} lineation with normal kinematics (Figures 2h and 5b). They occur as discrete shear planes, with a near-constant spacing on the m-scale (Figure 5b). They crop out mainly in the southern part of the RG, and they are less developed elsewhere. The dominant set of shear planes dips NE, with a pure dip-slip L_{3C} lineation. Conjugate, SW-dipping shear planes with normal kinematics also occur. In some cases, the shear plane is also characterized by a weak ductile foliation in the host rock. D_{3C} shows mutual crosscutting relationships with D_{3B} structures.

4.1.4. D_4 Zeolite- and Gouge-Bearing Brittle Faults

The latest set of deformation structures D_4 includes zeolite- and gouge-bearing brittle faults (Figures 5c and 5d). They form gullies and valleys in the topography of the RG, and thus their exposure is very limited and, when

Figure 3. D_1 structures. Note that D_1 brittle structures contain clast and lithons only of undeformed granite. White arrow points to garnet in the matrix. (a) Set of quartz-biotite-bearing shear fractures showing en-echelon spatial arrangement (Wp004). (b) A cataclaste, showing limited ductile reactivation and preserving angular clasts of undeformed granite (Wp078). (c)–(d) Example of breccias (outcrop of sample ACB35; Wp149). (e)–(e.iii) Magnification of selected areas in b,c,d showing the garnets overgrowing the breccia matrix. (f) Sheared breccia showing moderate ductile reactivation and sheared clasts (resulting from the reworking of a thick quartz vein (Wp149)). (g) Example of lineated surfaces in the breccia matrix (Wp149). (h) Cataclasites displacing with dextral (and normal) strike-slip kinematics a subvertical pegmatitic-aplitic dyke (Wp199). Handlens (3 cm) for scale.




Figure 4.

present, badly preserved. When observed, the L_4 lineation is subhorizontal, consistent with a dominant dextral kinematics inferred from the few planes showing offset markers (Figure 2i). Zeolite-bearing, fine-grained breccias are observed on anastomosing planes exploiting pre-existent D_1 shear fractures. Gouge-bearing fault zones are observed to develop at the contact between major quartzo-feldspathic mylonites and the undeformed granite (Figures 5c and 5d). D_4 structures exploit pre-existent structural discontinuities, and the structural data in Figure 2i show that D_4 structures actually reactivate and exploit the entire set of pre-existent structures.

4.2. Microstructures and Petrochronology

We report here the results of microstructural (Figure 6), chemical and petrochronological characterization of representative samples from three of the four classes of deformation structures, including: (a) ACB35, D_1 brittle breccia (pseudosections presented in Figure 7; U–Pb dating on garnet results presented in Figure 8); (b) ACB27a, major dip-slip D_2 ductile shear zone with tip-to-SE kinematics (pseudosection and Rb–Sr dating presented in Figure 9); (c) ACB37b, D_{3A} brittle-ductile shear zones with dextral strike-slip reactivation (pseudosection and Rb–Sr dating presented in Figure 9). Microstructure and petrography are presented together with the mineral composition obtained from EPMA. Representative mineral compositions, diagrams, and bulk rock chemical compositions are reported in the Tables S2–S3 in Supporting Information S2, and in Figures S2–S5 in Supporting Information S1. Si content in white mica is reported as Wm(Si) [apfu]; Mg and Ti content in biotite are abbreviated as Bt(Mg#) and Bt(Ti) [apfu], respectively.

4.2.1. Microstructures, Mineral Paragenesis and Pseudosection Calculations

4.2.1.1. D_1 Qz-Bt-Bearing Breccia

D_1 breccias are composed of cm-size angular granite clasts, weakly flattened and sheared, embedded in a fine-grained matrix showing homogeneous grain size and a weak pervasive foliation defined by Bt (Figure 6b). The matrix mineral paragenesis in ACB35 includes $Qz + Bt + Kfs + Grt + Ep/Aln + Ab \pm Pl \pm Wm \pm Chl + Ap + Zrc + Nb - Y - REE$ oxides and silicates (Gadolinite group). Mg and Ti content in biotite are Mg# = 0.40–0.45 and Ti = 0.02 apfu. Rare Mnz grains are observed scattered in the recrystallized matrix. Grt crystals range from 100 μ m to 5–10 mm, showing a wide range of crystal morphologies, from euhedral grains containing angular inclusions (Figure 6a), to grains showing anhedral shapes, resorbed rims, and poikiloblastic/honeycomb textures (Figure 6b). In most samples of pristine D_1 breccia, euhedral Grt includes a random pattern of $Qz + Pl + Kfs$ angular inclusions with no shape preferred orientation (Figures 6a and 6b). These inclusions are only preserved within euhedral garnets and are here interpreted to reflect the fine-grained cataclastic matrix of the D_1 breccia on which Grt grew. There is no evidence of Grt growth and shearing coeval with D_1 brecciation. Grt grains containing angular inclusions are typically enveloped by a weak mylonitic foliation overprinting the breccia matrix (Figures 6a and 7a). In some cases, Grt is weakly pleochroic, suggesting a likely non-cubic crystal symmetry (Cesare et al., 2019). The Grt in sample ACB35 shows an elongated shape, parallel to the foliation, with an honeycomb texture, characterized by a heterogeneous distribution of oriented inclusions (Figure 7a). The inclusions are mainly euhedral Ep and Aln with a peculiar texture (Figures 6c and 6d). Aln is observed at the core, surrounded by Ep forming the euhedral rim (Figure 6d). Ep in the recrystallized breccia matrix shows a concentric, rhythmic zoning with brighter rims, without Aln cores (Figures 6e and 6f). Ep aggregates in the recrystallized matrix contain spongy Zrc crystals, and partially destabilized Thorite-Xenotime at their core (Figure 6f).

Grt compositional variability ranges from $Alm_{36}Sps_{30}Grs_{32}Prp_2$ (Grt_A in Figures 6c and 7a) to $Alm_{36}Sps_{26}Grs_{36}Prp_2$ (Grt_B in Figures 6c and 7a). Garnets with similar compositions are also observed in sheared

Figure 4. D_2 – D_3 structures. (a) Dip-slip D_2 shear zone localized on a mafic dyke, parallel to a set of D_1 shear fractures (left hand side of the image) and characterized by the pervasive occurrence of sheared quartz + calcite veins. (Wp170). (b) Heterogeneous D_2 ductile shear zone (foliation marked by white arrows) with reverse kinematics and the associated low-angle Qz-vein (black arrows) (Wp087). (c) Sheared D_2 quartz + feldspar vein showing homogeneous internal foliation, suggesting top-to-SE reverse kinematics (Wp020). (d) D_2 calcite-bearing mylonite showing the boundary between a homogeneous calcite-mylonite (bottom) and a sheared breccia (top) with granitoid clasts (Wp182). (e) D_{3A} mylonitic shear zone showing S-C fabric related to strike-slip reactivation of a former dip-slip D_2 ductile shear zone (Wp171). (f) Detail of the D_{3A} shear zone reported in panel (e) showing the occurrence of the L_{3A} , strike-slip lineation and the L_2 -dip-slip lineation on adjacent shear planes; L_{3A} lineation occurs on S-C planes of the mylonitic shear zone wrapping around lensoid domains where L_2 is still preserved. (g) D_{3A} shear fractures, showing the development of quartz + feldspar wing cracks suggesting dextral strike-slip reactivation (Wp178). (h) D_{3A} brecciated dilational jog between two reactivated D_1 shear fractures (Wp137).




Figure 5.

D₁ shear fractures and veins (samples ACB_Sp3d and ACB18, Alm_{34–41}Sp_{26–18}Gr_{38–39}Prp₁; Figure S2a in Supporting Information S1). Grt in ACB35 shows resorbed rims and embayment at the contact with the sheared granular matrix and phyllosilicates (Figures 6c and 6d).

The bulk composition adopted for the calculation of the pseudosection for sample ACB35 (Figure 7b) reflects the recrystallized matrix of the breccia (Table S2 in Supporting Information S2; Figure 7a). Its composition is enriched in SiO₂ compared to the undeformed granite (Table S3 in Supporting Information S2). The observed paragenesis Qz + Bt + Kfs + Grt + Ep + Ab is computed to be stable over a wide range of *P*–*T* conditions (*T* < ~550°C, *P* < 0.9 GPa). The variation of computed Bt composition is limited and not useful to further constrain *P*–*T* conditions of apparent equilibrium. The observed Grt_A and Grt_B compositions are computed to be stable at *T* = ~590–600°C, and *P* = 0.9–1.0 GPa (Figures 7c–7e). These conditions are close to the stability field of Qz + Bt + Kfs + Grt + Ep + Pl + Ab + H₂O. Similar *P*–*T* estimates are retrieved from pseudosection calculations using samples ACB18 (590 ± 25°C, 1.0 ± 0.1 GPa), as well as ACB25 (575 ± 15°C, 0.9 ± 0.1 GPa), the latter representing the sheared contact between granite and the host rock (Texts S2–S3 in Supporting Information S1).

4.2.1.2. D₂ Dip-Slip, Reverse Shear Zones

The sample ACB27a represents a high-strain domain of a D₂ ductile shear zone. D₂ ductile shear zones are characterized by a pervasive and homogeneous mylonitic foliation including Qz + Kfs + Wm + Ep/Aln + Bt + Ab ± Pl enveloping mm-sized Wm + Bt porphyroblasts (Figure 6g). Si content in white mica is 6.6–6.8 apfu. Mg and Ti content in biotite are Mg# = 0.55 and Ti = 0.04 apfu. The observed paragenesis Qz + Bt + Wm + Kfs + Ep + Ab is computed to be stable over a wide range of *P*–*T* conditions, at H₂O-saturated conditions (Figure 9a). Computed Wm(Si) is comparable to the observed composition. Considering the observed Wm(Si), Bt(Ti) = 0.03 and Bt(Mg#) > 0.5, the stability field of the observed paragenesis is constrained to *T* = 520 ± 40°C and *P* = 0.8 ± 0.1 GPa (Figure 9a).

4.2.1.3. D₃ Strike-Slip, Dextral Shear Zones

The ACB37b mineral paragenesis includes Qz + Kfs + Wm + Bt + Pl + Ab + Ep/Aln + Ap ± Chl (Figure 6h). Mg and Ti content in biotite are Mg# = 0.58 and Ti = 0.05 apfu. The main foliation is defined by anastomosing S–C shear planes of fine-grained Wm, incorporating Bt and Ep/Aln inclusions, wrapping around recrystallized Qz + Pl + Ab + Kfs lenses. Wm(Si) is variable, forming two main compositional groups which have no microstructural correspondence: Wm(Si)₁ = 6.40 apfu and Wm(Si)₂ = 6.65–6.85 apfu (Figure S5a and S5b in Supporting Information S1). The observed paragenesis is stable over a wide range of *P*–*T* conditions (Figure 9b). Computed Wm(Si)₁, and Bt(Ti) isopleths define a field centered at *T* = 395 ± 25°C and *P* = 0.4 ± 0.1 GPa (Figure 9b). At these conditions the computed Bt(Mg#) ~ 0.55, similar to the observed Mg# for biotite. Spn is predicted in very small amounts (<1 vol%) but not observed.

4.2.2. In Situ U–Pb on Garnet

We report here the results of U–Pb and trace element analyses from a set of samples representing D₁ breccias (ACB35; ACB_Sp8; ACB_Sp7), and sheared Qz-veins and fractures (ACB18, ACB_Sp3) (Figure 6; Figure S2 in Supporting Information S1).

The analyses of Grt in ACB35 show two distinct U–Pb populations, defining two separate trends in a Tera–Wasserburg concordia diagram (Figure 8a). A first population of U–Pb data forms a linear array defining a lower intercept age of 128.0 ± 9.3 Ma (*n* = 26, MSWD = 2.9). A second population can be fitted by a regression line with a lower intercept corresponding to an age of 34.0 ± 4.4 Ma (*n* = 25, MSWD = 2.8). The combination of U–Pb ratios and trace element concentrations reveals that the first population contains elevated Zr amounts (up to

Figure 5. D₃–D₄ structures. (a) D_{3B} shear plane of a sinistral brittle-ductile fault, showing the oblique L_{3B} lineation and the occurrence of mineralized jogs (quartz vein) (Wp092). (b) D_{3C} brittle-ductile normal fault characterized by cm-scale heterogeneous foliation along the shear plane. (c–d) Examples of field outcrops of episyenites, showing the high porosity related to quartz dissolution (e.g., Pennacchioni et al., 2016). (e) Outcrop close to Poncione di Cassina Baggio, (d) outcrop close to Passo di Manio. (e) D₄ zeolite-bearing cataclasite (delimited by dashed white curves) localized at the contact between a major quartzo-feldspathic D₂ mylonite (left-hand side of the picture) and the undeformed granite (right-hand side of the picture, Wp047). (f) D₄ gouge-bearing fault localized on D₂ mylonitic foliation (Wp158).




Figure 6.

7,000 ppm), which are indicative of contamination of the analysis by ablation of zircon inclusions in the garnet. As shown in the trace element maps of Figure 8c, some of the ablation spots fall adjacent to or on top of areas where high amounts of Zr are detected. Therefore, to avoid contamination artifacts, we have excluded from the final age calculation all the U–Pb data with Zr content >20–30 ppm (depending on the sample). This approach filters out the anomalously old, spurious U–Pb intercept age (~128 Ma) defined above.

The pooled lower intercept Tera–Wasserburg age obtained from analyses of (inclusion-free) Grt grains for this (ACB35) and the other samples are reported in Table 1. Other samples range between 26.9 ± 1.3 Ma ($n = 54$, MSWD = 1.9) for ACB_Sp3b (Figure 8d), and 20.1 ± 1.0 Ma ($n = 33$, MSWD = 0.86) for ACB_Sp8d (Figure 8e). Additional details of U–Pb analyses, including Tera–Wasserburg plots, are reported in Text S2 and Figure S2 in Supporting Information S1.

The REE patterns of analyzed garnets (Figure 8b, insets in Figures 8d and 8e) are characterized by a high variability in REE contents, ranging between two end-member compositions characterized by distinct LREE contents (Grt-Type1 and Grt-Type2 in Figure 8b inset). Similar variations are also observed in samples ACB18, ACB_Sp3b, ACB_Sp8c/d, ACB_Sp7d (Figures insets in Figures 8d and 8e and Figure S3 in Supporting Information S1).

4.2.3. In Situ Rb–Sr on White Mica and Biotite

We report here the results of in situ Rb–Sr analyses of white mica defining the main foliation in the samples ACB27a, ACB37b, already described before, as well as in an additional sample ACB12b. Then, Rb–Sr data for biotite in the undeformed granite B19-1417 (sample from Rast et al., 2022) are presented.

4.2.3.1. D_2 Dip-Slip, Reverse Shear Zone

In ACB27a the mica grains show a large spread in $^{87}\text{Rb}/^{86}\text{Sr}$ from 119 to 3,125 with corresponding variations in $^{87}\text{Sr}/^{86}\text{Sr}$ between 0.82 and 1.42 ($n = 39$). A regression through these data generates an isochron with a slope corresponding to an age of 18.1 ± 0.9 Ma (2se, $n = 38$, MSWD = 1.5) and a $^{87}\text{Sr}/^{86}\text{Sr}$ intercept of 0.799 ± 0.010 (Figure 9e).

4.2.3.2. D_3 Strike-Slip, Dextral Shear Zone

The analyses of mica in sample ACB37b provide different age results. Most of the analyzed grains cluster at $^{87}\text{Rb}/^{86}\text{Sr}$ between 700 and 1,000 with four measurements extending to higher $^{87}\text{Rb}/^{86}\text{Sr}$ (up to 3,030) and one to lower $^{87}\text{Rb}/^{86}\text{Sr}$ (4.1). $^{87}\text{Sr}/^{86}\text{Sr}$ values are similarly clustered between 0.86 and 0.99 with three higher (up to 1.44) and one lower value (0.77). The corresponding Rb–Sr isochron provides an age of 14.7 ± 1.5 Ma and an initial $^{87}\text{Sr}/^{86}\text{Sr}$ of 0.767 ± 0.018 ($n = 35$, MSWD = 1.3; Figure 9f). These values change marginally if the high Sr analysis, for which contribution by a Sr-bearing phase is likely, is removed (e.g., 13.1 ± 2.1 Ma).

4.2.3.3. D_2 Bulk Foliation in RG_2 Granite

Sample ACB12b represents the foliated RG_2 granite, where the Wm-bearing bulk foliation wraps around the porphyric Kfs and the partially recrystallized quartz domains. The recrystallized matrix along the foliation is mainly composed of $\text{Qz} + \text{Wm} + \text{Ab} + \text{Ep}$. White mica shows limited $^{87}\text{Rb}/^{86}\text{Sr}$ spread (≤ 352) including two relatively low values of 8.0 due to ablation of Sr-rich impurities. $^{87}\text{Sr}/^{86}\text{Sr}$ is also limited compared to the previous samples (≤ 0.83) with $^{87}\text{Sr}/^{86}\text{Sr}$ of 0.73 for the two low-Rb/Sr samples. The age of the Rb–Sr isochron (18.6 ± 1.9 Ma; $n = 35$; MSWD = 0.76; $^{87}\text{Sr}/^{86}\text{Sr}_i = 0.729 \pm 0.005$) is largely constrained by the two low Rb/Sr

Figure 6. Microstructures of the analyzed deformation zones. (a) Optical plane-polarized light micrograph of D_1 breccias. Dashed curves delimit mm-to-cm clasts of the host granite. (b) Backscattered electron (BSE) image of the garnet in panel (a) showing the detail of the inclusion pattern. Note the angular shape of the inclusions and the occurrence of a weak preferential shape orientation perpendicular to the foliation surrounding the garnet (dashed white lines). (c) BSE image of Grt in sample ACB35 showing the alignment of Ep/Aln inclusions. See text for explanations. (d) BSE image of the Ep/Aln inclusions in the Grt of sample ACB35, showing the Aln, inclusion-rich cores with resorption textures and euhedral Ep rims. (e) BSE image of Ep crystals included in Bt in the fine-grained matrix of sample ACB35, showing rhythmic zoning between Ep-rich and Aln-rich layers. (f) BSE image of Ep aggregates along the mylonitic foliation in sample ACB35, including Zircon (Zrc), Nb–Y-oxides (NbYox), and Xtm/Thr aggregates likely resulting from the destabilization of Monazite. (g) BSE image of sample ACB27a showing the Wm porphyroblast and mylonitic foliation. Note the craters due to laser ablation analyses. (h) BSE image of the mylonitic foliation of sample ACB37b showing the occurrence of fine-grained aggregates of Wm + Bt. Note the craters due to laser ablation analyses.




Figure 7. Microstructure and results of pseudosection calculation for sample ACB35—D₁ breccias. (a) Optical plane-polarized light micrograph of the recrystallized matrix of a D₁ breccia (sample ACB35), showing the analyzed honeycomb Grt. (b) Computed pseudosections for sample ACB35. (c–e) Simplified sketch of the pseudosection in panel (b) showing the isopleths for each end-member of the garnet solid solution: (c) Grt(Alm)—almandine, (d) Grt(Sps)—spessartine, (e) Grt(Grs)—grossular. Exagons marked by A and B represent the *P*-*T* conditions of stability of Grt_A and Grt_B compositions reported in the text.

analyses. Exclusion of these two analyses returns a similar although much less precise of 16.3 ± 3.7 Ma. For all these samples (ACB27a, ACB37b, and ACB12b) we prefer to consider the more precise isochron ages that include the high-Sr analyses because the high Sr impurities included in these analyses appear to be sourced from minerals (epidote) cogenetic with mica.




Figure 8.

4.2.3.4. Undeformed RG₁ Granite

Sample B19-1417 represents the undeformed RG₁ granite. Biotite in this undeformed granite provides similar ages to those of white mica in sheared sample ACB37b. This biotite shows very high and highly variable ⁸⁷Rb/⁸⁶Sr (2350–45720) corresponding to elevated ⁸⁷Sr/⁸⁶Sr between 1.22 and 10.5. The large spread in Rb–Sr data point results in a relatively precise isochron corresponding to an age of 15.2 ± 0.7 Ma ($n = 28/30$; MSWD = 0.34) but a poorly defined ⁸⁷Sr/⁸⁶Sr intercept of 0.68 ± 0.11 (Figure 8f).

Biotite in RG₁ (sample B19-1417), foliated RG₂ (sample ACB14c, Text S4, Figure S5 in Supporting Information S1) and D₂ localized shear zone in RG₂ (sample ACB3b, Text S4, Figure S5 in Supporting Information S1) yielded indistinguishable ages (Table 1) which are all within uncertainty of the Rb–Sr age of white mica from the D_{3A} localized shear zone (Figure 8d).

5. Discussion

Here, we discuss and interpret the field observation and petrochronological data to (a) define the time-constrained *P-T-d* path and (b) to characterize the rheological evolution of the RG, including the factors controlling it, during collisional tectonics.

5.1. P-T-t-d Path and Tectonic Evolution

The *P-T-t-d* path summarizing the structural, petrochronological, and rheological evolution of the granite is presented in Figure 10.

5.1.1. Brittle-To-Ductile Evolution and Alpine Peak Metamorphic Conditions

The oldest structures observed in the granite consist of D₁ shear fractures, cataclasites and breccias. Similar brittle structures pre-dating ductile shear zones are reported from several other crystalline units of the Alps (External and Internal Crystalline Massifs: Bertini et al., 1985; Ceccato et al., 2022; Goncalves et al., 2012; Guermani & Pennacchioni, 1998; Menegon & Pennacchioni, 2010; Oliot et al., 2014; Rolland et al., 2009; Wehrens et al., 2016; Tauern Window: Leydier et al., 2019; Mancktelow & Pennacchioni, 2020; Suretta nappe: Goncalves et al., 2016). In many cases these brittle structures were interpreted to have formed in the biotite stability field, at relatively high *T* (>350°C) and mid-to-lower crustal conditions (Goncalves et al., 2016; Wehrens et al., 2016). Accordingly, they have been interpreted to represent either (a) a prograde phase of brittle Alpine deformation (Guermani & Pennacchioni, 1998), or (b) brittle (seismogenic) deformation at mid-crustal depths at the Alpine peak metamorphic conditions (Leydier et al., 2019; Mancktelow & Pennacchioni, 2020; Wehrens et al., 2016). It is interesting to note that such brittle-to-ductile evolution at peak metamorphic conditions has been reported from different crystalline units across the Alps spanning the whole range of “peak metamorphic conditions” recorded for the different case studies, from sub-greenschist to high-pressure amphibolite facies (see Ceccato et al., 2022; Figure 10a). Conversely, recent studies proposed an inherited origin for similar brittle structures occurring in the ECMs, suggesting their development during the Permo-Mesozoic rifting or previous Paleozoic collision (Ballèvre et al., 2018; Dall’Asta et al., 2022; Herwegh et al., 2020; Schwartz et al., 2024).

Although we do not have any quantitative constraints on the timing or exact *P-T* conditions of formation of D₁ structures, their micro and macro-structural relationships provide several clues about their relative timing with respect to Alpine collision, and their deformation conditions. In terms of relative timing, first, a key observation is that D₁ structures are overprinted by D₂ mylonitic shear zones without mutual cross-cutting relationship. Second, in the present orientation, D₁ structures are oriented at high angles (70–80°) with respect to the long-term NNW–SSE shortening direction and maximum principal stress σ_1 during Alpine convergence and D₂ reverse shearing.

Figure 8. Petrochronological data for sample of D₁ structures. (a) Tera-Wasserburg (TW) diagram of U–Pb garnet and trace element data for samples ACB35, showing the two populations of data, with analyses plotted according to their Zr content. (b) Chondrite-normalized (McDonough & Sun, 1995) REE-patterns of the analyzed Grt in samples ACB35. The inset shows the compositions of the two identified end-members (Grt-Type1, Grt-Type2) characterized by either flat LREE or steep LREE profile. (c) Laser ablation maps for selected isotopic masses, including ⁵⁵Mn, ³¹P, ⁹⁰Zr, ¹³⁹La, ¹⁷²Yb, ²³⁸U, on ACB35 Grt (see area delimited in Figure S4a in Supporting Information S1). (d–f) TW diagram for Grt in ACB_Sp3b and ACB_Sp8d, respectively, also displaying the Chondrite-normalized REE pattern in the inset. Two profiles representative of the Grt-Type1 and Grt-Type2 are reported as black curves.




Figure 9. *P-T-t* data for sample ACB27a–ACB37b. (a–b) Computed pseudosections for sample ACB27a and ACB37b, respectively. (c) ACB27a, (d) ACB37b, (e) ACB12b, and (f) B19-1417 mica Rb–Sr isochrones including their corresponding ages (slope) and initial $^{87}\text{Sr}/^{86}\text{Sr}$ composition (intercept). The size of the ellipses indicates internal 2SE (standard error). Isochronous regressions are plotted as black lines with their 95% confidence level as gray envelopes. All plots were generated using IsoplotR (Vermeesch, 2018).




Figure 10.

This high-angle orientation (much larger than the $\sim 30^\circ$ expected for Andersonian thrust faults) makes it difficult to explain the origin of D_1 structures as reverse brittle faults during Alpine convergence. As discussed by Herwegh et al. (2020) for similar brittle structures occurring in the Aar massif, exaggerated rotation ($>60^\circ$) of the entire massif would be necessary to re-orient low-angle thrust planes into the observed D_1 orientation. Indeed, such high-angle orientation would be more consistent with the development of faults and fractures under a strike-slip or extensional tectonic regime (Sibson, 2003). The steep orientation of D_1 structures is a common feature of many of the shear zones presenting a brittle-to-ductile evolution in the ECMs (Bertini et al., 1985; Guermani & Pennacchioni, 1998; Herwegh et al., 2017, 2020; Oliot et al., 2014; Rolland et al., 2009; Wehrens et al., 2016). If we extrapolate our observations from the Rotondo to the other massifs, the common D_1 steep orientation suggests that little or no reorientation occurred regionally, and that this orientation might represent an original feature of the brittle deformation structures at the regional scale. These observations suggest the D_1 structures pre-date Alpine convergence, rather than being synkinematic with cyclical brittle-ductile deformation at peak metamorphic conditions, as instead proposed for other case studies (Herwegh et al., 2017; Mancktelow & Pennacchioni, 2020; Wehrens et al., 2016).

The geochemical and age relationships observed in Grt porphyroblasts that overprint D_1 structures provide further constraints on the earliest stages of RG deformation and the transition from D_1 to D_2 structures. First, Grt postdates breccia formation, statically overprinting the pre-existing texture (Figures 6a and 6b). Later deformation during peak to D_2 retrograde shearing led to foliation development in the matrix surrounding the Grt. The foliation-parallel, elongated, and honeycomb-like crystal shapes (Figure 7a) may suggest that Grt partially re-equilibrated or crystallized synkinematically to early D_2 shearing at conditions close to peak metamorphism. However, most of the D_2 shear zones do not contain Grt, probably indicating its metastability during the main phase of retrograde D_2 shearing. Thus, we define three main stages of early Rotondo evolution: (a) D_1 brecciation and cataclasis, (b) Grt growth at a post-kinematic stage relative to D_1 ; (c) shearing and likely Grt destabilization during D_2 .

The REE and U–Pb systematics suggest that Grt growth occurred over variable and evolving conditions. The variability in trace element (REE) composition of garnet (Grt-Type1, Grt-Type2) can be explained by Grt growth in equilibrium with different LREE-bearing accessory phases, such as Aln, Ep, Ap, and Mnz, which are stable at different P–T metamorphic conditions (Hentschel et al., 2020; Janots et al., 2008, 2009; Spear, 2010). For example, the prograde-to-peak destabilization of LREE-rich Aln (observed as inclusions in Grt) into LREE-poor Mnz, as well as the opposite reaction during retrograde metamorphism, could explain the change in LREE availability and the crystallization of Grt with variable LREE content (Hentschel et al., 2020; Janots et al., 2008).

U–Pb dating of Grt results in scattered ages ranging from ~ 34 to ~ 20 Ma, which are broadly consistent with the ages for the regional peak metamorphism obtained from other case studies in the Gotthard nappe (22–19 Ma, Boston et al., 2017; Janots & Rubatto, 2014; Janots et al., 2009) and nearby Lepontine Dome (32–22 Ma, Rubatto et al., 2009; Tagliaferri et al., 2023). The peak metamorphic conditions are constrained by pseudosections

Figure 10. *P–T–t* path for the Rotondo Granite (RG) and sketch of its tectonic and rheological evolution. (a) Diagram summarizing the *P–T* conditions of deformation retrieved from thermodynamic modeling. Light blue boxes report the *P–T* conditions of peak/retrograde shear zone from other External Crystalline Massifs (AA: Aar/Grimsel, Goncalves et al., 2012; AR: Aiguille Rouges, Egli et al., 2017; MB: Mont Blanc: Rolland et al., 2008; GT: Gotthard-Fibbia, Oliot et al., 2014; Zeo: zeolite-faults from Lützenkirchen & Loew, 2011). Exhumation and cooling rates are reported. (b–c) Sketch (not to scale) of the tectonic evolution of the RG and Gotthard nappe through the D_2 – D_4 deformation stages. GT: Gotthard nappe; Penninic a.w.: Penninic accretionary wedge. (d–f) Mohr plots describing the rheological evolution of the RG during D_2 (d), D_3 (e), and D_4 (f) deformation events. σ_V : vertical principal stress; σ_H : major horizontal stress; σ_1 : minor horizontal stress. Mohr plots computed with MohrPlotter (<https://www.rickallmendinger.net/mohrplotter>). Gray-shaded Mohr circles represent the stress conditions at depth, light-blue circles represent the effective stress conditions. Mohr-Coulomb failure envelope (black curve) computed considering a cohesion $C = 30$ MPa and a frictional coefficient $\mu = 0.75$. The dashed black line represents the cohesionless re-shearing criterion in panels (e) and (f) with frictional coefficient $\mu > 0.6$ (Byerlee values, e) or $\mu < 0.6$ (f). The differential stress and shear stress values reported in the main text are retrieved from the intersection, or tangent points between the state of stress and the failure criteria (1) Mohr-Coulomb composite envelope for intact granite; (2) cohesionless re-shear criterion for frictional reactivation of cohesionless planes. The misoriented planes are reported along the Mohr circle as orange dots. Points A–B in panel (d) represent the shear stress conditions for reactivation during ductile shearing retrieved from field constraints (see text for explanation). Point A represents the shear stress conditions on D_1 planes oriented at 80° to σ_1 considering a differential stress of 60 MPa, which represents the upper strength limit for the intact granite due to the formation of extensional veins. Point B represents the shear stress conditions on D_1 planes oriented at 80° to σ_1 considering a differential stress of 10 MPa, which represents the differential stress conditions inferred for shearing along ductile shear zones (see text for explanation). (e) The white star represents the minimum effective stress conditions for the formation of the conjugate set of D_{3B} strike-slip faults in the intact granite. The misoriented D_1 – D_2 – D_{3A} planes (represented by the orange dot) are far away from the cohesionless re-shearing criterion (dashed line). (f) The blue circles represent the variable effective stress states during reactivation of the misoriented D_1 – D_2 – D_{3A} (intersection between orange points and the cohesionless re-shearing criterion).

calculated for different samples (ACB35, ACB25, ACB18) at $590 \pm 15^\circ\text{C}$ and 0.9 ± 0.1 GPa. The obtained P - T conditions are consistent with recent estimates of Alpine peak metamorphic conditions from the southern Aar massif and Gotthard nappe (Berger et al., 2020; Janots et al., 2008, 2009; Nibourel et al., 2021; Wiederkehr et al., 2011), as well as the Penninic units of the Northern Lepontine dome (Boston et al., 2017; Galli et al., 2007). The 34–20 Ma age spread would describe a prolonged thermal peak in the Gotthard nappe lasting for ~ 10 Myrs. This conclusion is consistent with geochronological data supporting the occurrence of a prolonged thermal peak starting at ~ 32 –34 Ma in several other ECMs and Penninic units, lasting until 22–17 Ma when the main phase of exhumation occurred (Boston et al., 2017; Cenki-Tok et al., 2014; Egli et al., 2016; Girault et al., 2020; Janots et al., 2008, 2009; Rolland & Rossi, 2016; Rubatto et al., 2009; Sanchez et al., 2011; Tagliaferri et al., 2023).

5.1.2. Exhumation— D_2 Shear Zones

The reverse kinematics of D_2 shear zones and the occurrence of associated shallowly dipping tensional veins constrain a subhorizontal σ_1 , parallel to a NW–SE trending maximum shortening axis ϵ_1 , and perpendicular to a subvertical σ_3 (Figure 10b). Based on the results of pseudosection calculation, these shear zones were already active at $520 \pm 40^\circ\text{C}$ and 0.8 ± 0.1 GPa. The conditions of re-equilibration of the analyzed samples (ACB27a $T > 500^\circ\text{C}$) are similar to the closure temperature for Rb–Sr in white mica inferred for similar case studies of granitoid shear zones ($T \leq 500$ – 550°C ; e.g., Egli et al., 2016; Ribeiro et al., 2023). Therefore, it is very likely that the obtained Rb–Sr date of 18.1 ± 0.8 Ma (Figure 9) reflects the (re-) crystallization of white mica in D_2 ductile shear zones. D_2 ductile shear zones accommodate the main phase of tectonic exhumation of the Gotthard nappe through reverse shearing on NW-steeply dipping planes. Rb–Sr dating of white mica indicates that the bulk foliation of RG_2 developed during the same amphibolite-facies deformation event at 18.6 ± 1.9 Ma (sample ACB12b, Figure 9e). However, biotite in D_2 shear zones (15.3 ± 0.9 Ma, sample ACB3b, Text S4, Figure S5 in Supporting Information S1) provides a younger Rb–Sr age which reflects the lower closure temperature of the Rb–Sr isotope system in biotite (≤ 350 – 400°C ; e.g., Jenkin et al., 2001) compared to white mica ($\leq 500^\circ\text{C}$). The exhumation of the ECMs in the Central and Western Alps have been accommodated by similar steeply dipping, reverse shear zones developed during retrograde greenschist facies conditions between 22 and 17 Ma (Cenki-Tok et al., 2014; Goncalves et al., 2012; Herwegh et al., 2020; Rolland et al., 2008).

5.1.3. Strike-Slip Tectonics— D_3 – D_4

The kinematics and geometrical relationships of D_{3A} and D_{3B} structures constrain their development under a transpressional strain field developed during NW–SE convergence (D_{3A} – D_{3B}), associated with NE–SW-directed extension (D_{3B} tensional veins and D_{3C} normal faults; Figure 10c). The kinematics of D_3 shear zones and tensional veins constrain a subhorizontal σ_1 and maximum shortening axis ϵ_1 oriented $\sim$ NW–SE and a subhorizontal, NE–SW-oriented σ_3 (Figure 10c). Strike-slip D_{3A} shearing is constrained to develop at $395 \pm 25^\circ\text{C}$ and 0.4 ± 0.1 GPa. At these temperature conditions, the Rb–Sr chronometer applied to white mica constrains the age of mica (re-)crystallization because the closure temperature of Sr diffusion in white mica is considerably higher (see previous section). Hence, the 14.7 ± 1.6 Ma age of white mica in D_{3A} sample ACB37b (Figure 9) probably constrains the age of this deformation event. The occurrence of similar Rb–Sr ages for white mica (ACB37b, D_{3A}) and biotite (B19-1417, undeformed granite: 15.5 ± 0.7 Ma; ACB3b and ACB14c, foliated granite: 15.3 ± 0.9 Ma and 15.0 ± 1.2 Ma, respectively; Table 1, Figure 9, and Figure S5 in Supporting Information S1), regardless of the intensity of sample deformation, suggests that the D_{3A} deformation event occurred at conditions broadly corresponding to the closure temperature of the Rb–Sr chronometer in biotite, that is $\sim \leq 350^\circ\text{C}$ (e.g., Jenkin et al., 2001). If the temperatures were substantially higher (e.g., 400 – 500°C), white mica would have recorded an older age due to its higher closure temperature for the Rb–Sr isotope system. In other words, the ~ 15 Ma age of biotite in the undeformed RG represents a cooling age. The overlap of white mica and biotite Rb–Sr ages implies that ductile reactivation of D_2 under strike-slip conditions leading to D_{3A} shear zones occurred during a very short time period at ~ 15 Ma. Further deformation during D_3 strike-slip tectonics was accommodated by brittle-ductile transpressional and extensional faults. The NW–SE-striking extensional veins associated to this brittle-ductile deformation event in the Lepontine dome and Aar-Gotthard area are consistently dated to < 14 Ma (Bergemann et al., 2020), as well as to < 9 Ma in Mont Blanc massif (Rossi & Rolland, 2014). The contemporaneous (or cyclic) development of D_{3B} – D_{3C} extensional and transpressional structures is consistent with the regional tectonic setting during the activity of the Rhone-Simplon fault system (Campani et al., 2010), accommodating NE–SW extensional tectonics under a constant dominant NW–SE transpression.

Late D₄ zeolite- and gouge-bearing brittle faults reactivated the pre-existing, steeply dipping structural discontinuities under strike-slip conditions. Their activity is constrained to have occurred between 12 and 3 Ma based on K–Ar illite dating (Kralik et al., 1992; Pleuger et al., 2012) at upper crustal levels ($T < 200^{\circ}\text{C}$, depth < 7 km; Lützenkirchen & Loew, 2011). These chronological constraints are consistent with the prolonged Neogene activity of the Periadriatic-Simplon-Rhone fault system (Ricchi et al., 2019).

In conclusion, the brittle-ductile-brittle evolution inferred from the sequence of deformation structures in the RG is the result of pre-Alpine tectonics overprinted by the peak-and-retrograde collisional Alpine tectonics. Pre-collisional structures apparently controlled the localization and accommodation of collisional strain in the crystalline unit.

5.1.4. Exhumation and Cooling Rates During Alpine Collision

Rates of tectonic exhumation and cooling can be calculated considering the constraints on P , T , and age of deformation provided above (Figure 10a). Rates are computed considering a geothermal gradient of $25^{\circ}\text{C}/\text{km}$ and a lithostatic pressure gradient of $27.5 \text{ MPa}/\text{km}$ (e.g., Nibourel et al., 2021). Most of the exhumation of the RG occurs during the activity of D₂, reverse ductile shear zones, accommodating exhumation from peak conditions (590°C and 0.9 GPa) persisting until 20 Ma (based on the youngest U–Pb of garnet), to D₃ conditions (395°C and 0.4 GPa) at ~ 14 Ma (Rb–Sr on white mica and biotite). On average, exhumation occurred at a rate of $\sim 3.0 \text{ mm}/\text{yr}$, associated with a cooling rate of $\sim 30^{\circ}\text{C}/\text{Myr}$. Such exhumation rates are comparable to those retrieved from regional thermochronometry ($\sim 1\text{--}3 \text{ mm}/\text{yr}$; Glotzbach et al., 2010; Herwegh et al., 2020; Nibourel et al., 2021). Similarly, the high cooling rates recorded during D₂–D₃ exhumation are compatible with the estimates of $30\text{--}40^{\circ}\text{C}/\text{Myr}$ provided by Janots et al. (2009) for the eastern Gotthard nappe.

After the exhumation through the brittle-ductile transition, and the switch to regional transpression, the exhumation became much slower, as constrained from the comparison between D₃ Rb–Sr white mica/biotite ages and the youngest K–Ar illite age (3 Ma; Kralik et al., 1992; Pleuger et al., 2012) for gouge-bearing faults similar to D₄ structures. The obtained exhumation rate of $\sim 0.6 \text{ mm}/\text{yr}$ is associated with a cooling rate of $\sim 15^{\circ}\text{C}/\text{Myr}$ (Figure 9a), similarly to what previously reported from thermochronological constraints (Glotzbach et al., 2010; Herwegh et al., 2020).

5.2. Rheological Evolution of the Rotondo Granite During Alpine Collision

Here, we discuss the rheological evolution of the RG during the development of D₂ to D₄ deformation structures. The rheology of and mechanical significance of D₁ deformation structures needs further investigation and better constraints on the origin and geometry of the brittle deformation zones. A detailed analysis of faulting mechanisms and stress conditions during D₁ deformation will be addressed in a separate paper. Despite the severe misorientation of D₁ structures with the Alpine shortening direction, they have influenced the rheology and localized strain throughout the whole D₂–D₄ evolution across different P – T and rheological conditions of the crystalline basement (Figures 10d–10f). In the following sections, we constrain the rheology, as well as the stress and fluid regimes at which D₁ to D₄ structures likely formed.

During D₂–D₃ retrograde shearing at amphibolite-to-upper greenschist facies conditions ($T = 400\text{--}520^{\circ}\text{C}$, $P = 0.4\text{--}0.8 \text{ GPa}$), the granite was characterized by a network of high strain shear zones, localized on magmatic (aplitic, mafic dykes) and tectonic (D₁) precursors, delimiting low strain domains of relatively undeformed granite. D₂ shear zones are oriented at high angle ($70\text{--}80^{\circ}$) to the principal stress σ_1 (Figure 10d). In addition, the pervasive occurrence of tensional veins in the undeformed granite indicates high (quasi-supralithostatic) P_f and limited $\Delta\sigma$ ($< 4 \text{ Ts}$, in the range $36\text{--}60 \text{ MPa}$ for granite, Cox, 2010; Etheridge, 1983; Sibson, 2003; Sibson et al., 1988). Considering all these factors, shearing on D₂ planes occurred at low resolved shear stress close to $\sim 10 \text{ MPa}$ (Figure 10d).

Shear zone strength was controlled by reaction-weakening processes related to plagioclase destabilization, which led to the activation of fluid-mediated grain-size sensitive deformation mechanisms, as observed in similar granitoid ductile shear zones (Ceccato, Goncalves, & Pennacchioni, 2020, 2022; Oliot et al., 2014). This localized weakening might have been related to the higher fluid content of D₁ structures exploited by D₂ compared to the host rock. Ductile shear zones acted as fluid pathways during D₂ deformation as can be inferred from the occurrence of sheared Qz + Cal veins (Figure 4a), including Cal porphyroclasts (Figure 11 of Rast et al., 2022).

Cal porphyroclasts in Qz-mylonites have been constrained to develop during ductile shearing at low $\Delta\sigma$ (<10 MPa) at amphibolite facies, fluid-rich conditions (Mancktelow & Pennacchioni, 2010). Considering the misorientation of D_2 planes of 80° with respect to σ_1 , and the concomitant formation of (likely extensional) veins, the resolved shear stress during D_2 ductile shearing was very low, in the order of few MPa (Figure 10d). Therefore, during D_2 – D_3 ductile shearing: (a) there is a difference in the maximum $\Delta\sigma$ of ~25–50 MPa between weak shear zones (<10 MPa) and in the low-strain granite (<60 MPa); (b) the strength of both high-strain shear zone and low strain granite domains is limited by tensional veining related to fluid overpressure, which in turn implies (c) low permeability in the low strain granite during ductile deformation. D_2 ductile shear zones might have acted as higher-permeability fluid conduits, but overall the permeability was not high enough to allow the dissipation of P_f build-up to supralithostatic conditions.

During further cooling and exhumation ($T < 350^\circ\text{C}$, $P = 0.2$ – 0.4 GPa), pre-existent misoriented structures failed to be reactivated (Figure 10e). The development of new conjugate faults (D_{3B}) suggests increasing $\Delta\sigma$ (>5.6 Ts) and decreasing P_f (probably close to hydrostatic conditions) during strike-slip deformation across the brittle-ductile transition. The decreased P_f was also related to the increased porosity and permeability of the granite during this deformation stage. High permeability of such deformation structures is documented by the pervasive occurrence of mineralized open veins at fault steps and jogs (Figure 5a), as well as by the occurrence of high-porosity hydrothermal alteration and the development of episyenites (Pennacchioni et al., 2016; Rossi & Rolland, 2014).

At shallow crustal levels ($T < 200^\circ\text{C}$, $P < 0.2$ GPa), D_4 zeolite- and gouge-bearing faults reactivated the rock fabrics and pre-existent structural heterogeneities (Figure 10f). Fluids leading to the crystallization of zeolites percolated through the highly permeable network of pre-existing fractures and structural heterogeneities. Similar zeolite-bearing fractures and faults are reported from the granitoid plutons of the Central and Eastern Alps (Adamello: Pennacchioni et al., 2006; Rieserferner: Ceccato & Pennacchioni, 2018), as well as from all the crystalline massifs of the Central Alps (Weissenberger & Bucher, 2010). For instance, in the Adamello pluton zeolite veins and gouges are observed to intrude and reactivate the pre-existent fracture and fault network (Pennacchioni et al., 2006). The complex kinematics of zeolite-bearing, reactivated faults and gouges were interpreted to be the result of earthquake swarm activity at shallow crustal levels, related to mineralizing fluids injection (Dempsey et al., 2014). In the RG, D_4 structures localize on misoriented D_1 – D_2 – D_3 structures. Mylonites and brittle-ductile faults are characterized by phyllosilicate-bearing fabrics reducing frictional and cohesion properties of the shearing planes at brittle conditions (Bistacchi et al., 2012; Pozzi et al., 2022; Volpe et al., 2022). Thus, in the RG, likely local P_f variations have promoted the reactivation of misoriented planes with low friction and low cohesion (Figure 10f). Furthermore, the low friction of the fault gouges might have promoted the following localization of brittle faulting on highly misoriented, and otherwise frictionally locked, fault planes during the latest stages of Alpine brittle deformation (Bistacchi et al., 2012; Collettini et al., 2019; Volpe et al., 2023).

6. Conclusions

The P – T – t evolution of the RG is recorded by a brittle-ductile-brittle structural evolution. D_1 breccias and cataclasites develop in the RG before the attainment of the Alpine peak metamorphic conditions, the latter occurring between 34 and 20 Ma and recorded by U–Pb in garnet. Peak metamorphic conditions are closer to the amphibolite facies ($T > 550^\circ\text{C}$, $P > 0.7$ GPa) than those previously proposed for the ECMs and the Gotthard nappe ($T < 450$ – 500°C , $P < 0.5$ – 0.6 GPa; Todd & Engi, 1997). The peak metamorphic conditions promoting garnet growth likely persisted for more than 10 Myr, consistently to what is commonly reported from the Lepontine dome further south (Rubatto et al., 2009; Tagliaferri et al., 2023). Exhumation was then controlled by reverse ductile shearing on D_2 ductile shear zones, localized on pre-existent structural and compositional heterogeneities. The very limited shear strength of D_2 ductile shear zones allowed it to accommodate fast exhumation of the Gotthard nappe at ~3 km/Myr between 20 and 14 Ma. Further exhumation was accommodated at slower rates by D_3 greenschist facies ($T < 400^\circ\text{C}$, $P < 0.4$ GPa) ductile and brittle-ductile shear zones, developed as a local response to the regional strike-slip activity of the Simplon-Rhone fault system.

The RG represents a rather strong granitic intrusion in otherwise weak polymetamorphic units of the Gotthard nappe. Thus, its rheological and strength evolution likely represents a “strong” upper bound to the overall strength of the embedding Gotthard nappe. Based on the common structural and tectonometamorphic history of the RG and the other ECMs in the Central and Western Alps, we can extrapolate the results obtained from the Rotondo to infer fundamental implications for the rheology of the European continental upper crust during Alpine collision:

- The European continental upper crust, now exposed in the ECMs, was extremely weak, that is, capable of sustaining only very low differential and shear stresses, during Alpine continental collision and deformation at amphibolite-to-greenschist facies.
- The occurrence of inherited tectonic and primary (e.g., magmatic) fabrics and structures, although highly misoriented, clearly controlled strain geometry and localization throughout their entire rheological and metamorphic evolution.
- The main weakening event occurred at retrograde conditions.

The weakness of the European continental crust during Alpine collision allowed it to focus and localize collisional strain in the external domains of the orogen, promoting the localized and fast exhumation of the crystalline massifs ahead of the advancing dry and strong Adriatic lower crust. At the scale of the orogen, collisional shortening was therefore accommodated through the localized, and fast exhumation of the ECMs, by means of the activity of weak ductile shear zones, localized on pre-existing tectonic and primary fabrics.

Data Availability Statement

Chemical and geochronological data supporting the conclusions of the present study are reported in the main text and in the Supplementary Information files. The data set is also available at ETH Zurich Research Collection via <https://doi.org/10.3929/ethz-b-000644819> with Creative Commons Attribution 4.0 International license (Ceccato et al., 2023—[Dataset]).

Acknowledgments

The authors thank two anonymous Reviewers and Loïc Labrousse for their constructive comments on the first version of the manuscript. This paper is FEAR Publication #19. This research is supported by ETH Zurich research funds to WMB and the European Research Council Synergy project “Fault Activation and Earthquake Rupture” (FEAR) (Grant 856559) granted to Domenico Giardini, Stefan Wiemer, Massimo Cocco, and Florian Amman. Markus Rast is warmly thanked for the introduction to the Bedretto tunnel geology and for providing the sample B19-1417 and thin sections adopted for Rb–Sr analysis of biotite. Neil Mancktelow and Bernardo Cesare are thanked for fruitful discussions about regional geology and garnet petrography. The authors acknowledge Luiz G. Morales and ScopeM for their support and assistance during SEM data acquisition and processing. Julien Allaz is thanked for EPMA data acquisition and technical assistance. Remy Lüchinger and Lydia Zender are gratefully acknowledged for their support during thin section preparation and XRF analyses, respectively. The crew of the Capanna Piansecco and SAC-CAS Bellinzona & Valli are warmly thanked for their logistic support, and their friendly and unconditional hospitality during fieldwork in the Rotondo area. The authors acknowledge the logistic support from the Bedretto Lab team. Open access funding provided by Eidgenössische Technische Hochschule Zurich.

References

- Audet, P., & Bürgmann, R. (2011). Dominant role of tectonic inheritance in supercontinent cycles. *Nature Geoscience*, 4(3), 184–187. <https://doi.org/10.1038/ngeo1080>
- Aysal, N., Guillong, M., Bayanova, T., Fukuyama, M., Leonard, N., Yılmaz, İ., et al. (2023). A new natural secondary reference material for garnet U–Pb dating by TIMS and LA–ICP–MS. *Geostandards and Geoanalytical Research*, 47(2), 297–310. <https://doi.org/10.1111/ggr.12493>
- Ballèvre, M., Manzotti, P., & Dal Piaz, G. V. (2018). Pre-Alpine (Variscan) inheritance: A key for the location of the future Valais basin (Western Alps). *Tectonics*, 37(3), 786–817. <https://doi.org/10.1002/2017tc004633>
- Behr, W. M., & Platt, J. P. (2013). Rheological evolution of a Mediterranean subduction complex. *Journal of Structural Geology*, 54, 136–155. <https://doi.org/10.1016/j.jsg.2013.07.012>
- Behr, W. M., & Platt, J. P. (2014). Brittle faults are weak, yet the ductile middle crust is strong: Implications for lithospheric mechanics. *Geophysical Research Letters*, 41(22), 8067–8075. <https://doi.org/10.1002/2014GL061349>
- Bellahsen, N., Mouthereau, F., Boutoux, A., Bellanger, M., Lacombe, O., Jolivet, L., & Rolland, Y. (2014). Collision kinematics in the western external Alps. *Tectonics*, 33(6), 1055–1088. <https://doi.org/10.1002/2013TC003453>
- Bellanger, M., Bellahsen, N., Jolivet, L., Baudin, T., Augier, R., & Boutoux, A. (2014). Basement shear zones development and shortening kinematics in the Ecrins Massif, Western Alps. *Tectonics*, 33(2), 84–111. <https://doi.org/10.1002/2013TC003294>
- Beltrando, M., Frasca, G., Compagnoni, R., & Vitale-Brovarone, A. (2012). The Valais controversy revisited: Multi-stage folding of a Mesozoic hyper-extended margin in the Petit St. Bernard pass area (Western Alps). *Tectonophysics*, 579, 17–36. <https://doi.org/10.1016/J.TECTO.2012.02.010>
- Beltrando, M., Manatschal, G., Mohn, G., Dal Piaz, G. V., Vitale Brovarone, A., & Masini, E. (2014). Recognizing remnants of magma-poor rifted margins in high-pressure orogenic belts: The Alpine case study. *Earth-Science Reviews*, 131, 88–115. <https://doi.org/10.1016/J.EARSCIREV.2014.01.001>
- Bergemann, C., Gnoss, E., Berger, A., Janots, E., & Whitehouse, M. J. (2020). Dating tectonic activity in the Lepontine Dome and Rhone-Simplon Fault regions through hydrothermal monazite–(Ce). *Solid Earth*, 11(1), 199–222. <https://doi.org/10.5194/se-11-199-2020>
- Berger, A., Engi, M., Erne-Schmid, S., Glotzbach, C., Spiegel, C., de Goede, R., & Herwegh, M. (2020). The relation between peak metamorphic temperatures and subsequent cooling during continent–continent collision (western Central Alps, Switzerland). *Swiss Journal of Geosciences*, 113(1), 1–18. <https://doi.org/10.1186/S00015-020-00356-4>
- Berger, A., Mercolli, I. P., Herwegh, M., & Gnoss, E. (2017). *Explanatory notes to the geological map of the Aar massif, Tavetsch and Gotthard nappes, geological special map No. 129*. Federal Office of Topography swisstopo.
- Bertini, G., Marcucci, M., Nevini, R., Passerini, P., & Sguazzoni, G. (1985). Patterns of faulting in the Mont Blanc granite. *Tectonophysics*, 111(1–2), 65–106. [https://doi.org/10.1016/0040-1951\(85\)90066-6](https://doi.org/10.1016/0040-1951(85)90066-6)
- Bistacchi, A., Massironi, M., Menegon, L., Bolognesi, F., & Donghi, V. (2012). On the nucleation of non-Andersonian faults along phyllosilicate-rich mylonite belts. *Geological Society, London, Special Publications*, 367(1), 185–199. <https://doi.org/10.1144/SP367.13>
- Boston, K. R., Rubatto, D., Hermann, J., Engi, M., & Amelin, Y. (2017). Geochronology of accessory allanite and monazite in the Barrovian metamorphic sequence of the Central Alps, Switzerland. *Lithos*, 286–287, 502–518. <https://doi.org/10.1016/J.LITHOS.2017.06.025>
- Bühler, M., Zurbiggen, R., Berger, A., Herwegh, M., & Rubatto, D. (2022). Late Carboniferous Schlingen in the Gotthard nappe (Central Alps) and their relation to the Variscan evolution. *International Journal of Earth Sciences*, 112(2), 1–26. <https://doi.org/10.1007/S00531-022-02247-5>
- Bürgmann, R., & Dresen, G. (2008). Rheology of the lower crust and upper mantle: Evidence from rock mechanics, geodesy, and field observations. *Annual Review of Earth and Planetary Sciences*, 36(1), 531–567. <https://doi.org/10.1146/annurev.earth.36.031207.124326>
- Bussien, D., Bussy, F., Masson, H., Magna, T., & Rodionov, N. (2008). Variscan lamprophyres in the Lower Penninic domain (Central Alps): Age and tectonic significance. *Bulletin de la Société Géologique de France*, 179(4), 369–381. <https://doi.org/10.2113/gssgfbull.179.4.369>
- Campani, M., Mancktelow, N., Seward, D., Rolland, Y., Müller, W., & Guerra, I. (2010). Geochronological evidence for continuous exhumation through the ductile–brittle transition along a crustal-scale low-angle normal fault: Simplon Fault Zone, central Alps. *Tectonics*, 29(3). <https://doi.org/10.1029/2009TC002582>

- Candioti, L. G., Duretz, T., Moulas, E., & Schmalholz, S. M. (2021). Buoyancy versus shear forces in building orogenic wedges. *Solid Earth*, 12(8), 1749–1775. <https://doi.org/10.5194/se-12-1749-2021>
- Ceccato, A., Behr, W. M., Zappone, A. S., Tavazzani, L., & Giuliani, A. (2023). Datasets for "Structural evolution, exhumation rates, and rheology of the European crust during Alpine collision: Constraints from the Rotondo granite – Gotthard nappe [Dataset]. *ETH Zurich Research Collection*. <https://doi.org/10.3929/ethz-b-000644819>
- Ceccato, A., Goncalves, P., & Menegon, L. (2022). On the petrology and microstructures of small-scale ductile shear zones in granitoid rocks: An overview. *Journal of Structural Geology*, 161(June), 104667. <https://doi.org/10.1016/j.jsg.2022.104667>
- Ceccato, A., Goncalves, P., & Pennacchioni, G. (2020). Temperature, fluid content and rheology of localized ductile shear zones in subsolidus cooling plutons. *Journal of Metamorphic Geology*, 38(8), 881–903. <https://doi.org/10.1111/jmg.12553>
- Ceccato, A., Menegon, L., Warren, C. J., & Halton, A. M. (2020). Structural and metamorphic inheritance controls strain partitioning during orogenic shortening (Kalak Nappe Complex, Norwegian Caledonides). *Journal of Structural Geology*, 136, 104057. <https://doi.org/10.1016/j.jsg.2020.104057>
- Ceccato, A., & Pennacchioni, G. (2018). Structural evolution of the Rieserferner pluton in the framework of the Oligo-Miocene tectonics of the Eastern Alps. *Journal of Structural Geology*, 116, 64–80. <https://doi.org/10.1016/j.jsg.2018.08.004>
- Célini, N., Mouthereau, F., Lahfid, A., Gout, C., & Callot, J. P. (2023). Rift thermal inheritance in the SW Alps (France): Insights from RSCM thermometry and 1D thermal numerical modelling. *Solid Earth*, 14(1), 1–16. <https://doi.org/10.5194/SE-14-1-2023>
- Kenki-Tok, B., Darling, J. R., Rolland, Y., Dhuime, B., & Storey, C. D. (2014). Direct dating of mid-crustal shear zones with synkinematic allanite: New in situ U-Th-Pb geochronological approaches applied to the Mont Blanc massif. *Terra Nova*, 26(1), 29–37. <https://doi.org/10.1111/TER.12066>
- Cesare, B., Nestola, F., Johnson, T., Mugnaioli, E., Della Ventura, G., Peruzzo, L., et al. (2019). Garnet, the archetypal cubic mineral, grows tetragonal. *Scientific Reports*, 9(1), 14672. <https://doi.org/10.1038/s41598-019-51214-9>
- Challandes, N., Marquer, D., & Villa, I. M. (2008). P-T-t modelling, fluid circulation, and 39Ar-40Ar and Rb-Sr mica ages in the Aar Massif shear zones (Swiss Alps). *Swiss Journal of Geosciences*, 101(2), 269–288. <https://doi.org/10.1007/s00015-008-1260-6>
- Colletini, C., Tesi, T., Scuderi, M. M., Carpenter, B. M., & Viti, C. (2019). Beyond Byerlee friction, weak faults and implications for slip behavior. *Earth and Planetary Science Letters*, 519, 245–263. <https://doi.org/10.1016/j.epsl.2019.05.011>
- Connolly, J. A. D. (2005). Computation of phase equilibria by linear programming: A tool for geodynamic modeling and its application to subduction zone decarbonation. *Earth and Planetary Science Letters*, 236(1–2), 524–541. <https://doi.org/10.1016/j.epsl.2005.04.033>
- Cook, K. L., & Royden, L. H. (2008). The role of crustal strength variations in shaping orogenic plateaus, with application to Tibet. *Journal of Geophysical Research*, 113(B8). <https://doi.org/10.1029/2007JB005547>
- Cox, S. F. (2010). The application of failure mode diagrams for exploring the roles of fluid pressure and stress states in controlling styles of fracture-controlled permeability enhancement in faults and shear zones. *Geofluids*, 10(1–2), 217–233. <https://doi.org/10.1111/j.1468-8123.2010.00281.x>
- Dall'Asta, N., Hoareau, G., Manatschal, G., Centrella, S., Denèle, Y., Ribes, C., & Kalifi, A. (2022). Structural and petrological characteristics of a Jurassic detachment fault from the Mont-Blanc massif (Col du Bonhomme area, France). *Journal of Structural Geology*, 159(March), 104593. <https://doi.org/10.1016/j.jsg.2022.104593>
- Dal Piaz, G. V., Bistacchi, A., & Massironi, M. (2003). Geological outline of the Alps. *Episodes*, 26(3), 175–180. <https://doi.org/10.18814/epiugs/2003/v26i3/004>
- Dempsey, E. D., Holdsworth, R. E., Imber, J., Bistacchi, A., & Di Toro, G. (2014). A geological explanation for intraplate earthquake clustering complexity: The zeolite-bearing fault/fracture networks in the Adamello Massif (Southern Italian Alps). *Journal of Structural Geology*, 66, 58–74. <https://doi.org/10.1016/j.jsg.2014.04.009>
- Deng, X.-D., Li, J.-W., Luo, T., & Wang, H.-Q. (2017). Dating magmatic and hydrothermal processes using andradite-rich garnet U–Pb geochronometry. *Contributions to Mineralogy and Petrology*, 172(9), 71. <https://doi.org/10.1007/s00410-017-1389-2>
- Egli, D., Mancktelow, N., & Spikings, R. (2017). Constraints from 40Ar/39Ar geochronology on the timing of Alpine shear zones in the Mont Blanc-Aiguilles Rouges region of the European Alps. *Tectonics*, 36(4), 730–748. <https://doi.org/10.1002/2016TC004450>
- Egli, D., Müller, W., & Mancktelow, N. (2016). Laser-cut Rb–Sr microsampling dating of deformational events in the Mont Blanc-Aiguilles Rouges region (European Alps). *Terra Nova*, 28(1), 35–42. <https://doi.org/10.1111/ter.12184>
- Etheridge, M. A. (1983). Differential stress magnitudes during regional deformation and metamorphism: Upper bound imposed by tensile fracturing. *Geology*, 11(4), 231–234. [https://doi.org/10.1130/0091-7613\(1984\)12](https://doi.org/10.1130/0091-7613(1984)12)
- Faccenda, M., Gerya, T. V., & Chakraborty, S. (2008). Styles of post-subduction collisional orogeny: Influence of convergence velocity, crustal rheology and radiogenic heat production. *Lithos*, 103(1–2), 257–287. <https://doi.org/10.1016/j.lithos.2007.09.009>
- Fitzpayne, A., Giuliani, A., Hergt, J., Woodhead, J. D., & Maas, R. (2020). Isotopic analyses of clinopyroxenes demonstrate the effects of kimberlite melt metasomatism upon the lithospheric mantle. *Lithos*, 370, 105595. <https://doi.org/10.1016/j.lithos.2020.105595>
- Galli, A., Mancktelow, N., Reusser, E., & Caddick, M. (2007). Structural geology and petrography of the Naret region (northern Valle Maggia, N. Ticino, Switzerland). *Swiss Journal of Geosciences*, 100(1), 53–70. <https://doi.org/10.1007/s00015-007-1211-7>
- Gapais, D., Bale, P., Choukroune, P., Cobbold, P. R., Mahjoub, Y., & Marquer, D. (1987). Bulk kinematics from shear zone patterns: Some field examples. *Journal of Structural Geology*, 9(5–6), 635–646. [https://doi.org/10.1016/0191-8141\(87\)90148-9](https://doi.org/10.1016/0191-8141(87)90148-9)
- Girault, J. B., Bellahsen, N., Boutoux, A., Rosenberg, C. L., Nanni, U., Verlaquet, A., & Beyssac, O. (2020). The 3-D thermal structure of the Helvetic Nappes of the European Alps: Implications for collisional processes. *Tectonics*, 39(3), e2018TC005334. <https://doi.org/10.1029/2018TC005334>
- Giuliani, A., Oesch, S., Guillon, M., & Howarth, G. H. (2024). Mica RbSr dating by laser ablation ICP-MS/MS using an isochronous calibration material and application to West African kimberlites. *Chemical Geology*, 121982. <https://doi.org/10.1016/j.chemgeo.2024.121982>
- Glottzbach, C., Reinecker, J., Danišik, M., Rahn, M., Frisch, W., & Spiegel, C. (2010). Thermal history of the central Gotthard and Aar massifs, European Alps: Evidence for steady state, long-term exhumation. *Journal of Geophysical Research*, 115(F3). <https://doi.org/10.1029/2009JF001304>
- Goncalves, P., Oliot, E., Marquer, D., & Connolly, J. A. D. (2012). Role of chemical processes on shear zone formation: An example from the Grimsel metagranodiorite (Aar massif, Central Alps). *Journal of Metamorphic Geology*, 30(7), 703–722. <https://doi.org/10.1111/j.1525-1314.2012.00991.x>
- Goncalves, P., Poilvet, J. C., Oliot, E., Trap, P., & Marquer, D. (2016). How does shear zone nucleate? An example from the Suretta nappe (Swiss Eastern Alps). *Journal of Structural Geology*, 86, 166–180. <https://doi.org/10.1016/j.jsg.2016.02.015>
- Green, E., Holland, T., & Powell, R. (2007). An order-disorder model for omphacitic pyroxenes in the system jadeite-diopside-hedenbergite-acmite, with applications to eclogitic rocks. *American Mineralogist*, 92(7), 1181–1189. <https://doi.org/10.2138/AM.2007.2401>

- Groome, W. G., Koons, P. O., & Johnson, S. E. (2008). Metamorphism, transient mid-crustal rheology, strain localization and the exhumation of high-grade metamorphic rocks. *Tectonics*, 27(1). <https://doi.org/10.1029/2006TC001992>
- Guermani, A., & Pennacchioni, G. (1998). Brittle precursors of plastic deformation in a granite: An example from the Mont Blanc massif (Helvetic, western Alps). *Journal of Structural Geology*, 20(2–3), 135–148. [https://doi.org/10.1016/S0191-8141\(97\)00080-1](https://doi.org/10.1016/S0191-8141(97)00080-1)
- Guillong, M., Hametner, K., Reusser, E., Wilson, S. A., & Günther, D. (2005). Preliminary characterisation of new glass reference materials (GSA-1G, GSC-1G, GSD-1G and GSE-1G) by laser ablation-inductively coupled plasma-mass spectrometry using 193 nm, 213 nm and 266 nm wavelengths. *Geostandards and Geoanalytical Research*, 29(3), 315–331. <https://doi.org/10.1111/j.1751-908X.2005.tb00903.x>
- Guillong, M., von Quadt, A., Sakata, S., Peytcheva, I., & Bachmann, O. (2014). LA-ICP-MS Pb–U dating of young zircons from the Kos–Nisyros volcanic centre, SE Aegean arc. *Journal of Analytical Atomic Spectrometry*, 29(6), 963–970. <https://doi.org/10.1039/C4JA00009A>
- Hafner, S. (1958). Petrographie des südwestlichen Gotthardmassives (zwischen St. Gotthardpass und Nufenenpass). *Schweizerische Mineralogische und Petrographische Mitteilungen*, 38. <https://doi.org/10.3929/ethz-a-000097546>
- Handy, M. R., M. Schmid, S., Bousquet, R., Kissling, E., & Bernoulli, D. (2010). Reconciling plate-tectonic reconstructions of Alpine Tethys with the geological-geophysical record of spreading and subduction in the Alps. *Earth-Science Reviews*, 102(3–4), 121–158. <https://doi.org/10.1016/j.earscirev.2010.06.002>
- Hentschel, F., Janots, E., Trepmann, C. A., Magnin, V., & Lanari, P. (2020). Corona formation around monazite and xenotime during greenschist-facies metamorphism and deformation. *European Journal of Mineralogy*, 32(5), 521–544. <https://doi.org/10.5194/ejm-32-521-2020>
- Herwegh, M., Berger, A., Baumberger, R., Wehrens, P., & Kissling, E. (2017). Large-scale crustal-block-extrusion during late Alpine collision. *Scientific Reports*, 7(1), 1–10. <https://doi.org/10.1038/s41598-017-00440-0>
- Herwegh, M., Berger, A., Glotzbach, C., Wangenheim, C., Mock, S., Wehrens, P., et al. (2020). Late stages of continent-continent collision: Timing, kinematic evolution, and exhumation of the Northern rim (Aar Massif) of the Alps. *Earth-Science Reviews*, 200, 102959. <https://doi.org/10.1016/J.EARSCIREV.2019.102959>
- Holland, T. J. B., & Powell, R. (1998). An internally consistent thermodynamic data set for phases of petrological interest. *Journal of Metamorphic Geology*, 16(3), 309–343. <https://doi.org/10.1111/J.1525-1314.1998.00140.X>
- Holland, T. J. B., & Powell, R. (2011). An improved and extended internally consistent thermodynamic dataset for phases of petrological interest, involving a new equation of state for solids. *Journal of Metamorphic Geology*, 29(3), 333–383. <https://doi.org/10.1111/J.1525-1314.2010.00923.X>
- Horstwood, M. S. A., Košler, J., Gehrels, G., Jackson, S. E., McLean, N. M., Paton, C., et al. (2016). Community-derived standards for LA – ICP – MS U–(Th)–Pb geochronology – Uncertainty propagation, age interpretation and data reporting. *Geostandards and Geoanalytical Research*, 40(3), 311–332. <https://doi.org/10.1111/j.1751-908X.2016.00379.x>
- Jamveit, B., Petley-Ragan, A., Incel, S., Dunkel, K. G., Aupart, C., Austrheim, H., et al. (2019). The effects of earthquakes and fluids on the metamorphism of the lower continental crust. *Journal of Geophysical Research: Solid Earth*, 124(8), 7725–7755. <https://doi.org/10.1029/2018JB016461>
- Janots, E., Engi, M., Berger, A., Allaz, J., Schwarz, J. O., & Spandler, C. (2008). Prograde metamorphic sequence of REE minerals in pelitic rocks of the Central Alps: Implications for allanite-monazite-xenotime phase relations from 250 to 610°C. *Journal of Metamorphic Geology*, 26(5), 509–526. <https://doi.org/10.1111/j.1525-1314.2008.00774.x>
- Janots, E., Engi, M., Rubatto, D., Berger, A., Gregory, C., & Rahn, M. (2009). Metamorphic rates in collisional orogeny from in situ allanite and monazite dating. *Geology*, 37(1), 11–14. <https://doi.org/10.1130/G25192A.1>
- Janots, E., & Rubatto, D. (2014). U–Th–Pb dating of collision in the external Alpine domains (Urseren zone, Switzerland) using low temperature allanite and monazite. *Lithos*, 184–187, 155–166. <https://doi.org/10.1016/j.lithos.2013.10.036>
- Jenkin, G. R., Ellam, R. M., Rogers, G., & Stuart, F. M. (2001). An investigation of closure temperature of the biotite Rb–Sr system: The importance of cation exchange. *Geochimica et Cosmochimica Acta*, 65(7), 1141–1160. [https://doi.org/10.1016/S0016-7037\(00\)00560-3](https://doi.org/10.1016/S0016-7037(00)00560-3)
- Kralik, M., Clauer, N., Holnsteiner, R., Huemer, H., & Kappel, F. (1992). Recurrent fault activity in the Grimsel Test Site (GTS, Switzerland): Revealed by Rb–Sr, K–Ar and tritium isotope techniques. *Journal of the Geological Society*, 149(2), 293–301. <https://doi.org/10.1144/GSJS.149.2.0293>
- Lacombe, O., & Mouthereau, F. (2002). Basement-involved shortening and deep detachment tectonics in forelands of orogens: Insights from recent collision belts (Taiwan, Western Alps, Pyrenees). *Tectonics*, 21(4), 1–22. <https://doi.org/10.1029/2001TC901018>
- Lemoine, M., Bas, T., Arnaud-Vanneau, A., Arnaud, H., Dumont, T., Gidon, M., et al. (1986). The continental margin of the Mesozoic Tethys in the Western Alps. *Marine and Petroleum Geology*, 3(3), 179–199. [https://doi.org/10.1016/0264-8172\(86\)90044-9](https://doi.org/10.1016/0264-8172(86)90044-9)
- Leydier, T., Goncalves, P., Lanari, P., & Oliot, E. (2019). On the petrology of brittle precursors of shear zones – An expression of concomitant brittle deformation and fluid–rock interactions in the ‘ductile’ continental crust? *Journal of Metamorphic Geology*, 37(8), 1129–1149. <https://doi.org/10.1111/jmg.12504>
- Lützenkirchen, V., & Loew, S. (2011). Late Alpine brittle faulting in the Rotondo granite (Switzerland): Deformation mechanisms and fault evolution. *Swiss Journal of Geosciences*, 104(1), 31–54. <https://doi.org/10.1007/s00015-010-0050-0>
- Ma, X., Hertrich, M., Amann, F., Bröker, K., Doonechaly, N. G., Gischig, V., et al. (2022). Multi-disciplinary characterizations of the BedrettoLab-A new underground geoscience research facility. *Solid Earth*, 13(2), 301–322. <https://doi.org/10.5194/se-13-301-2022>
- Mancktelow, N., & Pennacchioni, G. (2010). Why calcite can be stronger than quartz. *Journal of Geophysical Research*, 115(10), 1–16. <https://doi.org/10.1029/2009JB006526>
- Mancktelow, N., & Pennacchioni, G. (2020). Intermittent fracturing in the middle continental crust as evidence for transient switching of principal stress axes associated with the subduction zone earthquake cycle. *Geology*, 48(11), 1072–1076. <https://doi.org/10.1130/G47625.1>
- McDonough, W. F., & Sun, S. S. (1995). The composition of the Earth. *Chemical Geology*, 120(3–4), 223–253. [https://doi.org/10.1016/0009-2541\(94\)00140-4](https://doi.org/10.1016/0009-2541(94)00140-4)
- Menegon, L., Nasipuri, P., Stünitz, H., Behrens, H., & Ravna, E. (2011). Dry and strong quartz during deformation of the lower crust in the presence of melt. *Journal of Geophysical Research*, 116(B10), B10410. <https://doi.org/10.1029/2011JB008371>
- Menegon, L., & Pennacchioni, G. (2010). Local shear zone pattern and bulk deformation in the Gran Paradiso metagranite (NW Italian Alps). *International Journal of Earth Sciences*, 99(8), 1805–1825. <https://doi.org/10.1007/S00531-009-0485-6>
- Mohn, G., Manatschal, G., Beltrando, M., & Hauptert, I. (2014). The role of rift-inherited hyper-extension in Alpine-type orogens. *Terra Nova*, 26(5), 347–353. <https://doi.org/10.1111/ter.12104>
- Mouthereau, F., Watts, A. B., & Burov, E. (2013). Structure of orogenic belts controlled by lithosphere age. *Nature Geoscience*, 6(9), 785–789. <https://doi.org/10.1038/ngeo1902>
- Musso Piantelli, F., Mair, D., Berger, A., Schlunegger, F., Wiederkehr, M., Kurmann, E., et al. (2022). 4D reconstruction of the Doldenhorn nappe-basement system in the Aar massif: Insights into late-stage continent-continent collision in the Swiss Alps. *Tectonophysics*, 843, 229586. <https://doi.org/10.1016/J.TECTO.2022.229586>

- Newton, R. C., Charlu, T. V., & Kleppa, O. J. (1980). Thermochemistry of the high structural state plagioclases. *Geochimica et Cosmochimica Acta*, 44(7), 933–941. [https://doi.org/10.1016/0016-7037\(80\)90283-5](https://doi.org/10.1016/0016-7037(80)90283-5)
- Nibourel, L., Berger, A., Egli, D., Heuberger, S., & Herwegh, M. (2021). Structural and thermal evolution of the eastern Aar Massif: Insights from structural field work and Raman thermometry. *Swiss Journal of Geosciences*, 114(1), 9. <https://doi.org/10.1186/s00015-020-00381-3>
- Oliot, E., Goncalves, P., & Marquer, D. (2010). Role of plagioclase and reaction softening in a metagranite shear zone at mid-crustal conditions (Gotthard Massif, Swiss Central Alps). *Journal of Metamorphic Geology*, 28(8), 849–871. <https://doi.org/10.1111/j.1525-1314.2010.00897.x>
- Oliot, E., Goncalves, P., Schulmann, K., Marquer, D., & Lexa, O. (2014). Mid-crustal shear zone formation in granitic rocks: Constraints from quantitative textural and crystallographic preferred orientations analyses. *Tectonophysics*, 612–613, 63–80. <https://doi.org/10.1016/j.tecto.2013.11.032>
- Paton, C., Hellstrom, J., Paul, B., Woodhead, J., & Hergt, J. (2011). Iolite: Freeware for the visualisation and processing of mass spectrometric data. *Journal of Analytical Atomic Spectrometry*, 26(12), 2508. <https://doi.org/10.1039/c1ja10172b>
- Pennacchioni, G., Ceccato, A., Fioretti, A. M., Mazzoli, C., Zorzi, F., & Ferretti, P. (2016). Episyenites in meta-granitoids of the Tauern Window (Eastern Alps): Unpredictable? *Journal of Geodynamics*, 101, 73–87. <https://doi.org/10.1016/j.jog.2016.04.001>
- Pennacchioni, G., Di Toro, G., Brack, P., Menegon, L., & Villa, I. M. (2006). Brittle–ductile–brittle deformation during cooling of tonalite (Adamello, Southern Italian Alps). *Tectonophysics*, 427(1–4), 171–197. <https://doi.org/10.1016/j.tecto.2006.05.019>
- Pfiffner, O. A. (2016). Basement-involved thin-skinned and thick-skinned tectonics in the Alps. *Geological Magazine*, 153(5–6), 1085–1109. <https://doi.org/10.1017/S0016756815001090>
- Piccolo, A., Faccenda, M., Carosi, R., Montomoli, C., & Visonà, D. (2018). Crustal strength control on structures and metamorphism in collisional orogens. *Tectonophysics*, 746, 470–492. <https://doi.org/10.1016/j.tecto.2017.09.018>
- Pleuger, J., Mancktelow, N., Zwingmann, H., & Manser, M. (2012). K–Ar dating of synkinematic clay gouges from Nealpine faults of the Central, Western and Eastern Alps. *Tectonophysics*, 550–553, 1–16. <https://doi.org/10.1016/j.tecto.2012.05.001>
- Pozzi, G., Scuderi, M. M., Tinti, E., Nazzari, M., & Colletini, C. (2022). The role of fault rock fabric in the dynamics of laboratory faults. *Journal of Geophysical Research: Solid Earth*, 127(6), e2021JB023779. <https://doi.org/10.1029/2021JB023779>
- Rast, M., Galli, A., Ruh, J. B., Guillong, M., & Madonna, C. (2022). Geology along the Bedretto tunnel: Kinematic and geochronological constraints on the evolution of the Gotthard Massif (Central Alps). *Swiss Journal of Geosciences*, 115(1), 8. <https://doi.org/10.1186/s00015-022-00409-w>
- Ribeiro, B. V., Kirkland, C. L., Kelsey, D. E., Reddy, S. M., Hartnady, M. I., Faleiros, F. M., et al. (2023). Time-strain evolution of shear zones from petrographically constrained Rb–Sr muscovite analysis. *Earth and Planetary Science Letters*, 602, 117969. <https://doi.org/10.1016/j.epsl.2022.117969>
- Ricchi, E., Bergemann, C., Gnoss, E., Berger, A., Rubatto, D., & Whitehouse, M. J. (2019). Constraining deformation phases in the Aar Massif and the Gotthard Nappe (Switzerland) using Th–Pb crystallization ages of fissure monazite–(Ce). *Lithos*, 342–343, 223–238. <https://doi.org/10.1016/j.lithos.2019.04.014>
- Rolland, Y., Cox, S. F., & Corsini, M. (2009). Constraining deformation stages in brittle-ductile shear zones from combined field mapping and 40Ar/39Ar dating: The structural evolution of the Grimsel Pass area (Aar Massif, Swiss Alps). *Journal of Structural Geology*, 31(11), 1377–1394. <https://doi.org/10.1016/j.jsg.2009.08.003>
- Rolland, Y., & Rossi, M. (2016). Two-stage fluid flow and element transfers in shear zones during collision burial-exhumation cycle: Insights from the Mont Blanc Crystalline Massif (Western Alps). *Journal of Geodynamics*, 101, 88–108. <https://doi.org/10.1016/j.jog.2016.03.016>
- Rolland, Y., Rossi, M., Cox, S. F., Corsini, M., Mancktelow, N., Pennacchioni, G., et al. (2008). 40Ar/39Ar dating of synkinematic white mica: Insights from fluid-rock reaction in low-grade shear zones (Mont Blanc Massif) and constraints on timing of deformation in the NW external Alps (Vol. 299, pp. 293–315). Geological Society Special Publication. <https://doi.org/10.1144/SP299.18>
- Rosenberg, C. L., & Kissling, E. (2013). Three-dimensional insight into Central-Alpine collision: Lower-plate or upper-plate indentation? *Geology*, 41(12), 1219–1222. <https://doi.org/10.1130/G34584.1>
- Rossi, M., & Rolland, Y. (2014). Stable isotope and Ar/Ar evidence of prolonged multiscale fluid flow during exhumation of orogenic crust: Example from the Mont Blanc and Aar Massifs (NW Alps). *Tectonics*, 33(9), 1681–1709. <https://doi.org/10.1002/2013TC003438>
- Rubatto, D., Hermann, J., Berger, A., & Engi, M. (2009). Protracted fluid-induced melting during Barrovian metamorphism in the Central Alps. *Contributions to Mineralogy and Petrology*, 158(6), 703–722. <https://doi.org/10.1007/s00410-009-0406-5>
- Sanchez, G., Rolland, Y., Schneider, J., Corsini, M., Oliot, E., Goncalves, P., et al. (2011). Dating low-temperature deformation by 40Ar/39Ar on white mica, insights from the Argentera-Mercantour Massif (SW Alps). *Lithos*, 125(1–2), 521–536. <https://doi.org/10.1016/j.lithos.2011.03.009>
- Sarkar, S., Giuliani, A., Dalton, H., Phillips, D., Ghosh, S., Misev, S., & Maas, R. (2023). Derivation of lamproites and kimberlites from a common evolving source in the convective mantle: The case for southern African ‘transitional kimberlites’. *Journal of Petrology*, 64(7), egad043. <https://doi.org/10.1093/petrology/egad043>
- Schmid, S. M., Fügenschuh, B., Kissling, E., & Schuster, R. (2004). Tectonic map and overall architecture of the Alpine orogen. *Eclogae Geologicae Helveticae*, 97(1), 93–117. <https://doi.org/10.1007/S00015-004-1113-X>
- Schwartz, S., Rolland, Y., Nouibat, A., Boschetti, L., Bienvegnant, D., Dumont, T., et al. (2024). Role of mantle indentation in collisional deformation evidenced by deep geophysical imaging of Western Alps. *Communications Earth & Environment*, 5(1), 17. <https://doi.org/10.1038/s43247-023-01180-y>
- Seman, S., Stockli, D. F., & McLean, N. M. (2017). U–Pb geochronology of grossular-andradite garnet. *Chemical Geology*, 460, 106–116. <https://doi.org/10.1016/j.chemgeo.2017.04.020>
- Sibson, R. H. (2003). Brittle-failure controls on maximum sustainable overpressure in different tectonic regimes. *American Association of Petroleum Geologists Bulletin*, 87(6), 901–908. <https://doi.org/10.1306/01290300181>
- Sibson, R. H., Robert, F., & Poulsen, K. H. (1988). High-angle reverse faults, fluid-pressure cycling, and mesothermal gold-quartz deposits. *Geology*, 16(June), 551–555. [https://doi.org/10.1130/0091-7613\(1988\)016%3C0551:HARFFP%3E2.3.CO;2](https://doi.org/10.1130/0091-7613(1988)016%3C0551:HARFFP%3E2.3.CO;2)
- Simonetti, M., Carosi, R., Montomoli, C., Cottle, J. M., & Law, R. D. (2020). Transpressive deformation in the southern European Variscan Belt: New insights from the Aiguilles Rouges Massif (Western Alps). *Tectonics*, 39(6), e2020TC006153. <https://doi.org/10.1029/2020TC006153>
- Spear, F. S. (2010). Monazite–allanite phase relations in metapelites. *Chemical Geology*, 279(1–2), 55–62. <https://doi.org/10.1016/j.chemgeo.2010.10.004>
- Steck, A. (1976). Albit-Oligoklas-Mineralgesellschaften der Peristeritlücke aus alpinmetamorphen Granitgneisen des Gotthardmassivs. *Schweizerische Mineralogische Und Petrographische Mitteilungen*, 56, 269–292. <https://doi.org/10.5169/seals-43687>
- Steck, A., & Burri, G. (1971). Chemismus und Paragenesen von Granaten aus Granitgneisen der Grünschiefer-und Amphiholitfazies der Zentralalpen. *Schweizerische Mineralogische Und Petrographische Mitteilungen*, 51(2–3), 534–538.

- Tagliaferri, A., Schenker, F. L., Ulianov, A., Maino, M., & Schmalholz, S. M. (2023). Implications of new geological mapping and U-Pb zircon dating for the Barrovian tectono-metamorphic evolution of the Lepontine dome (Central European Alps). *Geochemistry, Geophysics, Geosystems*, 24(3), e2022GC010772. <https://doi.org/10.1029/2022GC010772>
- Todd, C. S., & Engi, M. (1997). Metamorphic field gradients in the Central Alps. *Journal of Metamorphic Geology*, 15(4), 513–530. <https://doi.org/10.1111/j.1525-1314.1997.00038.x>
- Vanardois, J., Trap, P., Roger, F., Melleton, J., Marquer, D., Paquette, J. L., et al. (2022). Deformation, crustal melting and magmatism in the crustal-scale East-Variscan Shear Zone (Aiguilles-Rouges and Mont-Blanc massifs, Western Alps). *Journal of Structural Geology*, 163, 104724. <https://doi.org/10.1016/j.jsg.2022.104724>
- Vermesch, P. (2018). IsoplotR: A free and open toolbox for geochronology. *Geoscience Frontiers*, 9(5), 1479–1493. <https://doi.org/10.1016/j.gsf.2018.04.001>
- Vogt, K., Willingshofer, E., Matenco, L., Sokoutis, D., Gerya, T., & Cloetingh, S. (2018). The role of lateral strength contrasts in orogenesis: A 2D numerical study. *Tectonophysics*, 746, 549–561. <https://doi.org/10.1016/j.tecto.2017.08.010>
- Volpe, G., Pozzi, G., & Collettini, C. (2022). YBPR or SCC'? Suggestion for the nomenclature of experimental brittle fault fabric in phyllosilicate-granular mixtures. *Journal of Structural Geology*, 165, 104743. <https://doi.org/10.1016/j.jsg.2022.104743>
- Volpe, G., Pozzi, G., Collettini, C., Spagnuolo, E., Achtziger-Zupančič, P., Zappone, A., et al. (2023). Laboratory simulation of fault reactivation by fluid injection and implications for induced seismicity at the BedrettoLab, Swiss Alps. *Tectonophysics*, 862, 229987. <https://doi.org/10.1016/j.tecto.2023.229987>
- Waldbaum, D. R., & Thompson, J. B., Jr. (1968). Mixing properties of sanidine crystalline solutions; [Part] 2, Calculations based on volume data. *American Mineralogist*.
- Wehrens, P., Berger, A., Peters, M., Spillmann, T., & Herwegh, M. (2016). Deformation at the frictional-viscous transition: Evidence for cycles of fluid-assisted embrittlement and ductile deformation in the granitoid crust. *Tectonophysics*, 693, 66–84. <https://doi.org/10.1016/j.tecto.2016.10.022>
- Weisenberger, T., & Bucher, K. (2010). Zeolites in fissures of granites and gneisses of the Central Alps. *Journal of Metamorphic Geology*, 28(8), 825–847. <https://doi.org/10.1111/j.1525-1314.2010.00895.x>
- White, R. W., Powell, R., Holland, T. J. B., Johnson, T. E., & Green, E. C. R. (2014). New mineral activity–composition relations for thermodynamic calculations in metapelitic systems. *Journal of Metamorphic Geology*, 32(3), 261–286. <https://doi.org/10.1111/JMG.12071>
- Whitney, D. L., & Evans, B. W. (2010). Abbreviations for names of rock-forming minerals. *American Mineralogist*, 95(1), 185–187. <https://doi.org/10.2138/am.2010.3371>
- Wiederkehr, M., Bousquet, R., Ziemann, M. A., Berger, A., & Schmid, S. M. (2011). 3-D assessment of peak-metamorphic conditions by Raman spectroscopy of carbonaceous material: An example from the margin of the Lepontine dome (Swiss Central Alps). *International Journal of Earth Sciences*, 100(5), 1029–1063. <https://doi.org/10.1007/S00531-010-0622-2>
- Willingshofer, E., Sokoutis, D., & Burg, J. P. (2005). Lithospheric-scale analogue modelling of collision zones with a pre-existing weak zone. *Geological Society, London, Special Publications*, 243(1), 277–294. <https://doi.org/10.1144/GSL.SP.2005.243.01.18>
- Wolf, S. G., Huismans, R. S., Braun, J., & Yuan, X. (2022). Topography of mountain belts controlled by rheology and surface processes. *Nature*, 606(7914), 516–521. <https://doi.org/10.1038/s41586-022-04700-6>
- Zertani, S., Menegon, L., Pennacchioni, G., Buisman, I., Corfu, F., & Jamtveit, B. (2023). Protracted localization of metamorphism and deformation in a heterogeneous lower-crustal shear zone. *Journal of Structural Geology*, 104960, 104960. <https://doi.org/10.1016/j.jsg.2023.104960>

References From the Supporting Information

- Redaa, A., Farkaš, J., Gilbert, S., Collins, A. S., Lohr, S., Vasegh, D., et al. (2023). Testing nano-powder and fused-glass mineral reference materials for in situ Rb-Sr dating of glauconite, phlogopite, biotite and feldspar via LA-ICP-MS/MS. *Geostandards and Geoanalytical Research*, 47(1), 23–48. <https://doi.org/10.1111/ggr.12467>